

# Journal Pre-proof

Molecular characterization of *Giardia* spp. and *Cryptosporidium* spp. from dogs and coyotes in an urban landscape suggests infrequent occurrence of zoonotic genotypes



Anya F. Smith<ce:contributor-role>Conceptualization) (Data curation) (Formal analysis) (Funding acquisition) (Investigation) (Methodology) (Project administration) (Validation) (Visualization) (Writing - original draft), Norman Neumann (Conceptualization) (Funding acquisition) (Methodology) (Resources) (Supervision) (Validation) (Writing - review and editing), Graham Banting (Formal analysis) (Investigation) (Validation) (Writing - review and editing), Claudia Klein (Funding acquisition) (Investigation) (Methodology) (Resources) (Validation) (Writing - review and editing), Stefano Liccioli (Resources) (Investigation) (Writing - review and editing), Alessandro Massolo (Conceptualization) (Funding acquisition) (Methodology) (Resources) (Supervision) (Writing - review and editing)

PII: S0304-4017(20)30095-9

DOI: <https://doi.org/10.1016/j.vetpar.2020.109115>

Reference: VETPAR 109115

To appear in: *Veterinary Parasitology*

Received Date: 27 October 2019

Revised Date: 15 April 2020

Accepted Date: 16 April 2020

Please cite this article as: Smith AF, Neumann N, Banting G, Klein C, Liccioli S, Massolo A, Molecular characterization of *Giardia* spp. and *Cryptosporidium* spp. from dogs and coyotes in an urban landscape suggests infrequent occurrence of zoonotic genotypes, *Veterinary Parasitology* (2020), doi: <https://doi.org/10.1016/j.vetpar.2020.109115>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Published by Elsevier.

**Molecular characterization of *Giardia* spp. and *Cryptosporidium* spp. from dogs and coyotes in an urban landscape suggests infrequent occurrence of zoonotic genotypes**

\* Anya F. Smith<sup>a,b</sup>, Norman Neumann<sup>c</sup>, Graham Banting<sup>c</sup>, Claudia Klein<sup>d</sup>, Stefano Liccioli<sup>a,e</sup>,  
Alessandro Massolo<sup>a,f,g</sup>.

<sup>a</sup> Department of Ecosystem and Public Health, Faculty of Veterinary Medicine, University of Calgary, Calgary, Alberta, Canada

<sup>b</sup> Clinical Service Innovation, Child Health BC, Provincial Health Services Authority, Vancouver, BC, Canada

<sup>c</sup> Environmental Health, School of Public Health, University of Alberta, Edmonton, Alberta, Canada

<sup>d</sup> Department of Veterinary Clinical and Diagnostic Sciences, Faculty of Veterinary Medicine, University of Calgary, Calgary, Alberta, Canada

<sup>e</sup> Grasslands National Park, Parks Canada Agency, Val Marie, Saskatchewan, Canada

<sup>f</sup> Ethology Unit, Department of Biology, University of Pisa, Pisa, Province of Pisa, Italy

<sup>g</sup> UMR CNRS 6249 Chrono-Environnement, Université Bourgogne Franche-Comté, Besancon, France.

\*Corresponding author (for both the manuscript and reprints).

Email address: [anya.smith@cw.bc.ca](mailto:anya.smith@cw.bc.ca)

Postal address: Provincial Health Services Authority, Child Health BC, 1770 West 7<sup>th</sup> Avenue  
Vancouver, BC, Canada, V6J 4Y6

## HIGHLIGHTS

- Dogs and coyotes were infected primarily with canid-specific genotypes
- There exists the potential for transmission among canid host species in urban areas

- Dogs and coyotes present a relatively low concern for zoonotic transmission
- Immunocompromised individuals and children should take precautions
- Follow up with advanced techniques would further inform health significance

## ABSTRACT

*Giardia* spp. and *Cryptosporidium* spp. are common gastrointestinal parasites with the potential for zoonotic transmission. This study aimed to (1) determine the genotypes occurring in dogs and coyotes occupying a similar urban area; (2) determine if these hosts were infected with potentially zoonotic genotypes; (3) provide baseline molecular data. In August and September 2012, 860 dog owners living in neighborhoods bordering six urban parks in Calgary, Alberta, Canada, provided faecal samples from their dogs. From March 2012 through July 2013, 193 coyote faeces were also collected from five of six of the same parks. Direct immunofluorescence microscopy (DFA) indicated that *Giardia* spp. and *Cryptosporidium* spp. infected a total of 64 (7.4%) and 21 (2.4%) dogs, as well as 15 (7.8%) and three (1.6%) coyotes, respectively. Semi-nested, polymerase chain reactions targeting the 16S small-subunit ribosomal ribonucleic acid (SSU rRNA) and 18S SSU rRNA genes of *Giardia* spp. and *Cryptosporidium* spp., respectively, were conducted on samples that screened positive by DFA, and products were sequenced and genotyped. Dogs were infected with *Giardia intestinalis* canid-associated assemblages C (n=14), D (n=13), and *Cryptosporidium canis* (n=3). Similarly, *G. intestinalis* assemblages C (n=1), D (n=1) and *C. canis* (n=1), were detected in coyotes, as well as *G. intestinalis* assemblage A (n=1) and *Cryptosporidium* vole genotype (n=1). Dogs and coyotes were predominantly infected with host-specific genotypes and few potentially zoonotic genotypes, suggesting that they may not

A.F. Smith *et al.*

*Giardia* and *Cryptosporidium* infecting canids

represent a significant risk for zoonotic transmission of these parasites in urban areas where these hosts are sympatric.

**Keywords:** *Giardia* spp., *Cryptosporidium* spp., genotype, dogs, coyotes, humans.

Journal Pre-proof

## 1. INTRODUCTION

Dogs (*Canis lupus familiaris*) are common companion animals in North America, with an estimated six million owned dogs in Canada and 83 million in the United States (Humane Society of the United States, 1954; Perrin, 2009). Among wild carnivores, coyotes (*Canis latrans*) have experienced the largest range expansion across North America (Laliberte and Ripple, 2004), and are now increasingly common in urban landscapes of US and Canada (Gehrt *et al.*, 2009). Spatial overlap of canids and humans in urban landscapes introduces the possibility of disease transmission among these hosts.

*Giardia* spp. and *Cryptosporidium* spp. are two of the most common gastrointestinal (GI) parasites to infect humans and other host species (Xiao and Fayer, 2008) and have frequently been reported in dogs (Bryan *et al.*, 2011; Bugg *et al.*, 1999; Fontanarroza *et al.*, 2006; Gaunt and Carr, 2011; Joffe *et al.*, 2011; Little *et al.*, 2009; Smith *et al.*, 2015; Smith *et al.*, 2014) and coyotes (Liccioli *et al.*, 2012; Thompson *et al.*, 2009) across geographic localities. Infective *Giardia* spp. and *Cryptosporidium* spp. (oo)cysts can persist in the environment (soil and water) for long periods, and can be transmitted directly through faecal-oral routes, or indirectly through the ingestion of contaminated water or food (Feng and Xiao, 2011; Ryan *et al.*, 2014; Xiao and Fayer, 2008).

*Giardia intestinalis* is considered a species complex composed of eight assemblages, A – H (Feng and Xiao, 2011); assemblages C and D most often infect canids, appear largely host-specific, and are rarely detected in humans (Sprong *et al.*, 2009). Conversely, assemblages A and B infect the widest range of host species, are the primary genotypes found to infect humans (Ballweber *et al.*, 2010; Volotao *et al.*, 2007), and also occur in dogs and coyotes (Ballweber *et*

al., 2010; Thompson et al., 2009; Trout et al., 2006; Volotao et al., 2007). These protozoa are considered potentially zoonotic but predominantly host-specific. However, debate continues over the degree of risk for zoonotic transmission due to varying results from sub-typing analyses and epidemiological studies (Ballweber et al., 2010; Bowman and Lucio-Forster, 2010; Marangi et al., 2009; Ryan et al., 2014; Volotao et al., 2007; Xiao et al., 2007).

The most commonly reported etiological agents of human cryptosporidiosis are *Cryptosporidium hominis* and *Cryptosporidium parvum* (Ryan and Hijjawi, 2015). In canids, there is no evidence of *C. hominis* occurrence, and infrequent reports of *C. parvum* (Lucio-Forster et al., 2010; Smith et al., 2009; Sotiriadou et al., 2013). Dogs and coyotes are typically infected with *Cryptosporidium canis* (Lucio-Forster et al., 2010; Ryan et al., 2014; Thompson et al., 2009; Trout et al., 2006; Xiao et al., 2002), and although this parasite species has rarely been documented in humans (Ryan et al., 2014), immunocompromised persons and children may be at heightened risk for infection (Bowman and Lucio-Forster, 2010; Xiao et al., 2007).

Despite the uncertainty around the risk for zoonotic transmission of some *G. intestinalis* assemblages and *Cryptosporidium* species, the co-existence of dogs and coyotes within urban areas and the ubiquity of these parasites as infectious agents of multiple host species, introduces the question of transmission potential among human and canid hosts. Few known studies from Canada have focused on the molecular characterization of *Giardia* spp. and *Cryptosporidium* spp. infecting dogs (McDowall et al., 2011; Uehlinger et al., 2013), and few from coyotes, overall (Thompson et al., 2009; Trout et al., 2006; Xiao et al., 2002). While previous research allowed us to quantify the prevalence of *Giardia* spp. and *Cryptosporidium* spp. infection in domestic dogs and identify risk factors associated to dog-walking behaviour (Smith et al., 2015;

Smith et al., 2014), parasite assemblage and species identification was not obtained. By genotyping parasite species at the SSU rRNA (small-subunit ribosomal ribonucleic acid), this study aimed to (1) conduct a preliminary assessment of (a) genotypes circulating in urban dogs and coyotes and; (b) assess the occurrence of zoonotic genotypes and; (2) provide baseline molecular data for ongoing research and surveillance.

## **2. MATERIALS AND METHODS**

### **2.1 Study design and study areas**

We used an observational, cross-sectional study design. The study recruited participants living in residential communities surrounding the following six urban parks in Calgary, Alberta, Canada (51°50N, 114°55'W): Bowmont; Fish Creek Provincial; Nosehill; River, Southland; Weaselhead. The faeces from dogs residing in these surrounding communities were collected, and coyote faecal samples were collected from the environment within five of six of these parks, with the exception of River Park. The collection of coyote faeces was largely unsuccessful in River Park, and therefore was excluded as a study site for this species. By targeting these residential communities for the collection of dog faeces, we aimed to maximize the potential for sampling faeces from dogs and coyotes that accessed and/or resided in similar geographic areas, including parks.

### **2.2 Faecal sampling protocol**

#### **2.2.1 Dogs**

A total of 860 dog faecal samples were collected from dog-owning residences in August and September 2012, following online questionnaire addressing dog and owner demographics and dog-walking and husbandry practices, as described in detail in Smith et al. (2015).



### 2.2.2 Coyotes

Coyote faeces were initially collected from the ground in parks from March 2012 to July 2013 as described by Liccioli *et al.* (2012). A total of 193 coyote faeces collected within this timeframe were selected in order to coincide as much as possible with the collection of samples from dogs, while maximizing sample size. All faeces collected over this time period and estimated to be between 2-5 days old (Liccioli *et al.*, 2012) were selected for the current study.

The faecal samples were kept at -80° C for 72 hours to inactivate *Echinococcus* spp. eggs (Veit *et al.*, 1995) and then transferred to -20° C until laboratory analysis.

## **2.3 Laboratory analysis**

### 2.3.1 Microscopy

Two grams of faeces were prepared for direct immunofluorescence (DFA) microscopy using a modified version of the manufacturer's instructions (Waterborne Inc., New Orleans, LA, USA) for detection of *Giardia* spp. and *Cryptosporidium* spp. Faeces were homogenized in phosphate-buffered saline (PBS), strained through double-layered cheesecloth, and centrifuged for five minutes at 528 x g to isolate (oo)cysts. The majority of supernatant was discarded, apart from 1.5 ml used to homogenize sediment. For each sample, 20 µl from 1.5 ml of homogenized sediment was transferred to a microscope slide, dried, and mounted with 20 µl of Aqua-Glo™ (20X) containing *Giardia* spp. and *Cryptosporidium* spp. fluorescein-labeled monoclonal antibodies. Meanwhile, 200 µl of homogenized sediment from each sample was transferred separately to a microcentrifuge tube in preparation for deoxyribonucleic acid (DNA) extraction. Positive samples were identified as those containing (oo)cysts that were within correct size, shape, and colour parameters (United States Environmental Protection Agency, 2005). Presence or absence

of *Giardia* spp. or *Cryptosporidium* spp. (oo)cysts, as well as the number of (oo)cysts per microscope slide were recorded.

Parasite prevalence was calculated based on results of DFA microscopy and confidence intervals (CI) were calculated using the 95% CI Sterne estimator (Santner and Duffy, 1989). The number of cysts per gram (cpg) or oocysts per gram (opg) were estimated (and rounded up to the nearest whole number) by extrapolating the number of (oo)cysts counted per microscope slide to the volume of homogenized sediment sourced from two grams of faeces and median and upper and lower limits reported. A chi-square exact test (Good, 2005) was used to look for significant differences of *Giardia* spp. and *Cryptosporidium* spp. prevalence in dogs versus coyotes. Differences of median intensities of *Giardia* spp. and *Cryptosporidium* spp. between dogs and coyotes were compared using the Mann–Whitney Test for two independent samples (Good, 2005). Additional prevalence values were calculated to account for the potential of re-sampling coyotes ('Data quality control measures' in Supplementary Material). All statistical analyses were conducted using Statistical Package for the Social Sciences version 20.0 (SPSS Inc., Armonk, NY, USA).

### 2.3.2 Molecular analysis

Samples positive for (oo)cysts were prepared for DNA extraction using the 200 µl of homogenized sediment set aside during preparation for microscopy. Faecal sediment was first exposed to five rounds of freeze/thaw (dry ice with 95% ethanol and thawing at 70°C) to fracture (oo)cysts. The remaining DNA extraction procedure followed the QIAamp DNA Stool protocol (Qiagen, Toronto, ON, Canada) yielding 200 µl elution volumes. In addition to positive faecal specimens, the following negative controls were extracted to account for possible contamination

and to assess synchrony of negatives using DFA and polymerase chain reaction (PCR): DNA lysis buffer and a sub-sample of faecal specimens that had screened DFA negative for *Giardia* spp. and *Cryptosporidium* spp. Positive controls were extracted to determine extraction efficiency using aliquots of 50 *Cryptosporidium muris* oocysts (Waterborne Inc.).

An internal amplification control (IAC) was initially used to determine and control for the level of PCR inhibition in dog and coyote faecal specimens screening positive for *Giardia* spp. and/or *Cryptosporidium* spp. using DFA ('Data quality control measures' in Supplementary Material). A 292 bp fragment of the 16S SSU rRNA (Hopkins et al., 1997) and 840bp fragment of the 18S SSU rRNA gene (Xiao et al., 2001) were subsequently targeted for detection of *Giardia* spp. and *Cryptosporidium* spp., respectively. Coyote and dog faecal specimens DFA positive for *Giardia* spp. and/or *Cryptosporidium* spp., as well as negative and positive controls were analysed using semi-nested PCR. Faecal DNA extracts were analysed in replicates of five and negative and positive controls in replicates of three. Water was used as an additional negative control on all PCR plates.

The *Giardia* spp. assay was a semi-nested reaction and used a 25 µl reaction volume for the primary reaction and contained the following: five µl of template, 1.1 µl of water, 1X Maxima Hot Start PCR Mastermix (Thermo Fisher, Waltham, MA, USA), 1M Betaine (Sigma-Aldrich, St. Louis, MO, USA), 400 ng/µL bovine serum albumin (BSA, Sigma-Aldrich), and primer concentrations of 0.8 µM each (high-performance liquid chromatography–HPLC purified, Thermo Fisher). The secondary reaction used five µl of template from the primary reaction and similar reagents and volumes. A previously published forward primer sequence was used for both the primary and secondary reactions (Hopkins et al., 1997) and the following reverse primer

for the primary reaction: 5'- TTG GAT GTG CGG GCC GTC TCG CA- 3'. The reverse primer for the secondary reaction used was: 5' - GTC GAA CCC TGA TTC TCC GCC AG – 3'. PCR cycling conditions were as follows: 95°C for 4 minutes, and 35 cycles of 95°C for 20 seconds, 59°C for 30 seconds, 72°C for 40 seconds, and a final elongation step of 72°C for 7 minutes (Applied Biosystems, 2720 thermocycler). Products were separated on 2.5% agarose gels comprised of 1X TAE buffer and visualized using SYBR® Safe (Thermo Fisher) under ultraviolet (UV) light.

Previously published primer sequences and a PCR assay validated for several *Cryptosporidium* genera described by Ruecker et al. (2011) were used to detect *Cryptosporidium* spp.

*Giardia* spp. PCR amplicons were cut from gels and purified using a QIAquick Gel Extraction Kit (Qiagen). Amplicons were bi-directionally sequenced with Sanger sequencing at Macrogen (Seoul, South Korea), using the same nested primers used for PCR. Multiple alignments using ClustalW were conducted with reference sequences for *G. intestinalis* assemblages A [GenBank: M54878], B [GenBank: AF199447], and C-G [GenBank: AF199449, AF199443, AF199448, AF199444, AF199450] published by Sprong *et al.* (2009). Alternate sources were used for the following reference sequences: *Giardia ardeae* [GenBank: Z17210], *Giardia muris* [GenBank: X65063] (van Keulen et al., 1993), *Giardia microti* [GenBank: AF006677] (van Keulen et al., 1998), and *Giardia psittaci* [GenBank: AF473853] (van Keulen et al., 2002). *Giardia* spp. was identified using polymorphic regions at the positions 22/23/24/38/44/45/63/75/95/170/175 (Table S1 in Supplementary Material), where the first base of the forward primer was considered bp one.

The sequences were cropped to the edge of the primer region and then aligned using ClustalW. For sequences without interpretable variants at indicative polymorphic positions, GenBank using BLAST (GenBank, 1982) was searched for the closest sequences. Query sequences confirmed by BLAST to be most highly identical to *Giardia* spp. were aligned to reference sequences using ClustalW. Percent identity scores >98% were considered *Giardia* spp. All *Giardia* spp. bioinformatics work was conducted using Geneious 7 (Biomatters Ltd., Auckland, New Zealand).

*Cryptosporidium* spp. PCR products were purified using QIAquick PCR Purification Kit (Qiagen) and sequenced bi-directionally at the University of Calgary DNA Sequencing and Genomics Analysis Laboratory with the same nested primers used for PCR (Ruecker *et al.*, 2011). Consensus sequences were generated from forward and reverse DNA trace sequences. Multiple alignments were conducted using BioNumerics software (Applied Maths, Sint-Martens-Latem, Belgium) with reference sequences for *Cryptosporidium* species and genotypes sourced from Ruecker *et al.* (2013; 2012), as well as sequences more recently added to the sequence library using methods described in Ruecker *et al.* (2012). For alignments with less than 100% identity, bases were scanned for alignment anomalies, and identified.

### 3. RESULTS

#### 3.1 Laboratory analysis

##### 3.1.1 Microscopy

There was no significant difference in *Giardia* spp. prevalence in dogs and coyotes ( $P_{\text{exact}} = 0.880$ , Table 1) or *Cryptosporidium* spp. ( $P_{\text{exact}} = 0.599$ , Table 1). Median *Giardia* spp. infection

intensity was higher in dogs than coyotes ( $P < 0.002$ , Table 1), and median *Cryptosporidium* spp. infection intensity was higher in coyotes than dogs ( $P_{\text{exact}} < 0.004$ , Table 1).

### 3.1.2 Molecular analysis

DNA was extracted from 64 dog and 15 coyote faecal specimens that screened DFA positive for *Giardia* spp., and 21 dog and three coyote specimens that screened DFA positive for *Cryptosporidium* spp. (total of 94 extracts that accounted for mixed *Giardia* spp. - *Cryptosporidium* spp. infections). Controls and additional samples extracted included 18 negative samples (DNA lysis buffer), 18 faecal specimens that screened DFA negative for *Giardia* spp. and *Cryptosporidium* spp., and six positive controls.

Of the dog and coyote faecal specimens screening DFA positive for *Giardia* spp. and that were analysed using PCR, 34/63 (54.0%) and 7/14 (50.0%) were also PCR positive, respectively (Figure 1a). Note that two samples analysed by DFA (from one dog and one coyote) were not analysed by PCR due to the insignificant quantities of biological sample remaining after DFA (and, consequently, inability to extract DNA). Of the dog and coyote faecal specimens screening DFA positive for *Cryptosporidium* spp. that were analysed using PCR, 3/21 (14.3%) and 1/3 (33.3%) were also PCR positive, respectively (Figure 1b). The large majority of negative controls screened PCR negative for both parasites. A single dog and coyote faecal specimen that were negative using DFA, were positive for *Giardia* spp. and *Cryptosporidium* spp. using PCR, respectively. The majority of *Cryptosporidium* spp. positive extraction controls were affirmative for all three replicates.

Of the dog and coyote faecal samples PCR positive and sequenced for *Giardia* spp., 27/33 (82%) and 3/7 (43%) returned sequences interpretable using discriminatory polymorphic nucleotides for dog and coyotes, respectively (Table 2). Of the 27 dog faeces successfully sequenced for *Giardia* spp., the majority was sourced from park-attending dogs (n=25/27). Of the interpretable *G. intestinalis* sequences from dogs, 20 were a 100% identity match to the reference sequences for assemblages C and D for at least one of two duplicate samples from the same individual. The remaining 7 interpretable sequences had percent identity values of over 99% apart from one dog sample with a 98.9% identity match value to *G. intestinalis* assemblage D (Table 2). The majority of base pair anomalies in all samples with identity scores less than 100% appeared to be due to polymerase errors, showing obscured peaks on at least one trace sequence, as well as potential single nucleotide polymorphisms (SNP) in four samples. For the six samples that were not identifiable using polymorphic regions of the 16S SSU rRNA gene, GenBank searches using BLAST returned five as *G. intestinalis*. Multiple alignments to reference sequences using ClustalW produced two higher identity scores of 90.1% and 85.5% to *G. intestinalis* assemblage C, the former was the DFA negative dog faecal specimen that screened PCR positive for *G. intestinalis*. The remaining three *G. intestinalis* samples produced identity scores between 38% and 65%. The one remaining dog sample searched in GenBank using BLAST returned results consistent with a microorganism other than *Giardia* spp., suggesting primer cross-reactivity. Three *G. intestinalis* sequences from coyotes were 100% identical to reference sequences. Three uninterpretable sequences returned GenBank matches to microorganisms other than *Giardia* spp., and one did not return results.

Of the dog and coyote faecal samples PCR positive and sequenced for *Cryptosporidium* spp., all of them returned interpretable sequences (Table 2). Two of three dog faeces successfully sequenced for *Cryptosporidium* spp. were sourced from park-attending dogs. All three dogs

PCR positive for *Cryptosporidium* spp. were infected with *C. canis* (Table 2). Two of these were 100% identical to GenBank: AB210854 and the third had one potential SNP. One coyote was infected with *C. canis* [99.7% match to GenBank: AB210854, one indel accounting for the 0.3% difference] and the coyote sample DFA negative but PCR positive for *Cryptosporidium* was carrying the vole genotype [100% identity match to GenBank: JQ178298].

#### 4. DISCUSSION

Our characterization of *G.intestinalis* assemblages and *Cryptosporidium* species using an SSU rRNA target locus suggested that dogs and coyotes were primarily infected with host-specific genotypes. While there is the potential for transmission of these parasites among sympatric dogs and coyotes, the potential for zoonotic transmission appeared relatively low.

The predominance of host-specific assemblages of *G. intestinalis* and *Cryptosporidium* species infecting dogs and coyotes found in the current study is not surprising, and has been reported for dogs globally, spanning research locales in Italy (Scaramozzino et al., 2009; Zanzani et al., 2014), Poland (Solarczyk and Majewska, 2010), the United States (Wang et al., 2012), Japan (Abe et al., 2002), Finland (Rimhanen-Finne et al., 2007), Australia (Palmer et al., 2008), and Canada (McDowall et al., 2011; Uehlinger et al., 2013). Despite the predominance of host-specific genotypes often reported for dogs, there are opposing accounts of assemblages A/B as the dominant genotypes present (Leonhard et al., 2007; Marangi et al., 2009; Sotiriadou et al., 2013; Thompson et al., 2008; Volotao et al., 2007). Not all of the studies targeted the same genetic loci or investigated sub-assemblages of A/B, making direct comparison difficult, but it does suggest a high frequency of potentially zoonotic genotypes in *G. intestinalis*-infected dogs in some locales. The absence of *C. parvum* detection in dogs in our study is supported by



literature (Lucio-Forster et al., 2010; Smith et al., 2009; Sotiriadou et al., 2013). Prevalence of *C. parvum* likely also varies among dogs geographically and may be higher in regions where dogs have access to human infected faeces or livestock (e.g. rural, farming communities), although no known studies have been conducted to compare *C. parvum* occurrence among these two host populations.

Assemblages of *G. intestinalis* and *Cryptosporidium* species infecting coyotes have been investigated less often than for dogs, but published studies indicate the occurrence of both host-associated and zoonotic genotypes (Thompson et al., 2009; Trout et al., 2006; Xiao et al., 2002). In our study, we detected only two non-host-specific genotypes of *G. intestinalis* and *Cryptosporidium* infecting coyotes: one *G. intestinalis* assemblage A, and the other, the *Cryptosporidium* vole genotype. *Giardia intestinalis* assemblage A was previously reported in coyotes by Thompson et al. (2009), but this is the first known report of the *Cryptosporidium* vole genotype, likely a result of coyote predation of co-habiting rodent species -a principal component of coyotes' diet in our study area (Liccioli et al., 2013; Morey et al., 2007).

It is possible that we underestimated the occurrence of some *G. intestinalis* assemblages (e.g. assemblages C/D). Notable SNPs in one or both 16S reverse primer regions for several *G. intestinalis* assemblages and *Giardia* species, excluding *G. intestinalis* assemblage A, suggests variable sensitivity of the assay (Figure S1 in Supplementary Material). Additional SNPs may also occur within the 16S forward primer region, however, for many *G. intestinalis* assemblages and *Giardia* species, lack of sequence data upstream of the forward primer region (5') renders this inconclusive. Unfortunately, GenBank 16S DNA sequence data related to *G. intestinalis*

assemblages, other than those represented by assemblages A (Healey et al., 1990; Prabhu et al., 2007)/B (van Keulen et al., 1995), originates within the forward primers themselves – a problem since even mis-matched forward primer sequences are artificially amplified through the sequencing reactions (and subsequently entered into GenBank as confirmed sequence in the primer regions). The two other *Giardia* species for which 16S DNA sequences exist upstream of the forward primer region include *G.muris* (van Keulen et al., 1992; van Keulen et al., 1993) and *G. ardeae* (van Keulen et al., 1993), and both contain a SNP in this region, and SNPs within both reverse primer regions. Reduced sensitivity of the assay from SNPs within primer regions is demonstrated by our own data that shows a nearly 50-fold decrease in PCR method sensitivity against *G. muris* (Table S2 in Supplementary Material). The discrepant detection of *G. intestinalis* seen by PCR versus DFA in this study, may be partially attributed to the application of a PCR assay with potentially heightened sensitivity to assemblages A/B relative to assemblages C/D. The antibodies used in the DFA assay are able to cross-react to various *G. intestinalis* assemblages and *Cryptosporidium* species (Waterborne Inc., New Orleans, LA, USA), and therefore may have detected some positive cases that the PCR assay did not.

This *Giardia* PCR assay highly sensitive for *G. intestinalis* assemblages A/B (i.e. LOD<sub>95</sub>=10.0 template copies per reaction, CI=25.2 – 4.0, Table S2 in Supplementary Material), suggests a lower relative abundance of zoonotic than host-specific genotypes within *G. intestinalis* infections of dogs and coyotes in our study. Although *C. canis* is potentially zoonotic, the risk to human health is considered minimal, as few cases of *C. canis* have been documented in humans, overall (Lucio-Forster et al., 2010; Xiao, 2010). In extremely rare cases, children and the immunocompromised individuals represent the susceptible group (Bowman and Lucio-Forster, 2010; Marangi et al., 2009; Volotao et al., 2007; Xiao et al., 2007). It is possible that zoonotic *G.*

*intestinalis* assemblages and *Cryptosporidium* species were present within mixed infections and went undetected because they were not predominant; for *Cryptosporidium* spp., loss of detection has been demonstrated to occur most often when the concentration ratio among genotypes is high (Ruecker et al., 2011).

There are known limitations of using a single-locus approach for *G. intestinalis* assemblage differentiation; multilocus sequence typing (MLST) is recommended to confirm genotypes ascertained using SSU rRNA (Feng and Xiao, 2011). While some of the more advanced molecular methods (including MLST) also help to distinguish sub-genotypes, resolve mixed infections, and provide higher discriminatory power to assist with interpreting zoonotic transmission potential for both *G. intestinalis* and *Cryptosporidium* spp. (Feng and Xiao, 2011; Ruecker et al., 2011; Thompson and Ash, 2016), our study targeted the SSU rRNA to preliminarily assess the parasitic genotypes circulating in dogs and coyotes. Allelic sequence heterozygosity within *G. intestinalis* assemblages at identifying positions could also influence assemblage-based typing results, although this phenomenon is uncommon for this species (Sprong et al., 2009). Our study contributes to the international body of research pertaining to genotypes of *Giardia* spp. and *Cryptosporidium* spp. from dogs and coyotes, and provides preliminary, baseline molecular data for ongoing research and surveillance. Follow up studies using a combined epidemiological and MLST approach would provide a relevant confirmatory measure and help to further clarify public and animal health significance of *Giardia* spp. and *Cryptosporidium* spp. circulating in wild and domestic canids.

## 5. CONCLUSIONS

Although risk for zoonotic transmission cannot be excluded, our infrequent detection of zoonotic genotypes using this *Giardia* spp. PCR assay sensitive to assemblages A/B, and validated for multiple genera of *Cryptosporidium* spp. (Ruecker et al., 2011), supports the conclusion that dogs and coyotes in our study area present a relatively low concern for zoonotic transmission of these parasites.

## 6. ACKNOWLEDGEMENTS

Thank you to the City of Calgary, Animal & Bylaw Services for media and procedural support in recruiting participants; the University of Calgary Media Relations for organizing press releases; and the Alberta Provincial Laboratory for Public Health for laboratory support. Thank you, as well, to Jacquelyn Wollman, Monica Viapiana, Akshay Gurdita, and Lalita Bharadwaj for their help in the field and laboratory, and to Dr. Melanie Rock for sharing funding resources (re: grant numbers AHFMR-200600378 and MSH-83745 acknowledged below).

This study was funded by Alberta Innovates Health Solutions (A.S., grant numbers 201001195, 201001196), (M.R., non-author, grant number AHFMR-200600378); the University of Calgary (A.M.), Animal & Bylaw Services at the City of Calgary (A.M.); the O'Brien Institute for Public Health (A.M., grant number 10002391); and Canadian Institutes of Health Research (M.R., non-author, grant number MSH-83745). The funders had no role in study design, collection, analysis, or interpretation of the data, in the writing of the report, or decision to submit the article for publication.

## 8. DECLARATION OF INTEREST

None.

**Table 1.** Prevalence (%) and median intensities (#(oo)cysts/g) of *G. intestinalis* and *Cryptosporidium* genotypes occurring in dog and coyote host species.

Host Species	No. of cysts/g <sup>a</sup>	<i>Giardia</i> spp.			<i>Cryptosporidium</i> spp.			Overall		
		Infection intensity range <sup>b</sup>	No. (%) <sup>c</sup> of cases	CI <sup>d</sup>	No. of oocysts/g	Infection intensity range	No. (%) of cases	CI	No. (%) of cases	CI
Dog ( <i>n</i> =860)	1.0 x 10 <sup>4</sup>	3.8 x 10 <sup>1</sup> - 1.6 x 10 <sup>5</sup>	64(7.4)	5.9-9.4	3.8 x 10 <sup>1</sup>	3.8 x 10 <sup>1</sup> - 1.6 x 10 <sup>3</sup>	21(2.4)	1.5-3.7	79(9.2)	7.4- 1.1 x 10 <sup>1</sup>
Coyote ( <i>n</i> =193)	4.9 x 10 <sup>2</sup>	3.8 x 10 <sup>1</sup> - 6.0 x 10 <sup>3</sup>	15(7.8)	4.6- 1.2 x 10 <sup>1</sup>	4.2 x 10 <sup>3</sup>	4.5 x 10 <sup>2</sup> - 5.6 x 10 <sup>3</sup>	3(1.6)	4.0 x 10 <sup>-1</sup> - 4.6	15(7.8)	4.6- 1.2 x 10 <sup>1</sup>

<sup>a</sup>Median intensity values<sup>b</sup>Upper and lower limits of (oo)cysts per gram<sup>c</sup>Prevalence based on DFA microscopy from faecal samples<sup>d</sup>95% confidence intervals

**Table 2.** *Giardia* spp. and *Cryptosporidium* spp. genotypes infecting dog and coyotes.

Host Species	<i>G. intestinalis</i>		<i>Cryptosporidium</i> spp.	
	Assemblage	No. (%)	Species/Genotype	No. (%)
Dog	Assemblage C	14(51.9)	<i>C. canis</i> (n=3)	3(100.0)
	Assemblage D (n=27)	13(48.1)		
Coyote	Assemblage A	1(33.3)	<i>C. canis</i>	1(50.0)
	Assemblage C	1(33.3)	Vole genotype (n=2)	1(50.0)
	Assemblage D (n=3)	1(33.3)		

**Author Contributions****Anya F Smith:**

Conceptualization; Data curation; Formal analysis; Funding acquisition; Investigation; Methodology; Project administration; Validation; Visualization; Roles/Writing - original draft

**Norman Neumann:**

Conceptualization; Funding acquisition; Methodology; Resources; Supervision; Validation; Writing - review & editing

**Graham Banting:**

Formal analysis; Investigation; Validation; Writing - review & editing

**Claudia Klein:**

Funding acquisition; Investigation; Methodology; Resources; Validation; Writing - review & editing

**Stefano Liccioli:**

Resources; Investigation; Writing - review & editing

**Alessandro Massolo:**

Conceptualization; Funding acquisition; Methodology; Resources; Supervision; Writing - review & editing

## Declaration of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

## 9. REFERENCES

- Abe, N., Sawano, Y., Yamada, K., Kimata, I., Iseki, M., 2002. *Cryptosporidium* infection in dogs in Osaka, Japan. *Vet. Parasitol.* 108, 185-193.
- Ballweber, L.R., Xiao, L., Bowman, D.D., Kahn, G., Cama, V.A., 2010. Giardiasis in dogs and cats: update on epidemiology and public health significance. *Trends Parasitol.* 26, 180-189.
- Bowman, D.D., Lucio-Forster, A., 2010. Cryptosporidiosis and giardiasis in dogs and cats: veterinary and public health importance. *Exp. Parasitol.* 124, 121-127.
- Bryan, H.M., Darimont, C.T., Paquet, P.C., Ellis, J.A., Goji, N., Gouix, M., Smits, J.E., 2011. Exposure to infectious agents in dogs in remote coastal British Columbia: possible sentinels of diseases in wildlife and humans. *Can. J. Vet. Res.* 75, 11-17.
- Bugg, R.J., Robertson, I.D., Elliot, A.D., Thompson, R.C., 1999. Gastrointestinal parasites of urban dogs in Perth, Western Australia. *Vet. J.* 157, 295-301.
- Feng, Y., Xiao, L., 2011. Zoonotic potential and molecular epidemiology of *Giardia* species and giardiasis. *Clin. Microbiol. Rev.* 24, 110-140.

- Fontanarroza, M.F., Vezzani, D., Basabe, J., Eiras, D.F., 2006. An epidemiological study of gastrointestinal parasites of dogs from Southern Greater Buenos Aires (Argentina): age, gender, breed, mixed infections, and seasonal and spatial patterns. *Vet. Parasitol.* 136, 283-295.
- Gaunt, M.C., Carr, A.P., 2011. A survey of intestinal parasites in dogs from Saskatoon, Saskatchewan. *Can. Vet. J.* 52, 497-500.
- Gehrt, S.D., Anchor, C., White, L.A., 2009. Home range and landscape use of coyotes in a metropolitan landscape: conflict or coexistence? *J. Mammal.* 90, 1045-1057.
- GenBank: [https://blast.ncbi.nlm.nih.gov/Blast.cgi?PAGE\\_TYPE=BlastSearch](https://blast.ncbi.nlm.nih.gov/Blast.cgi?PAGE_TYPE=BlastSearch). Accessed 1 August 2014.
- Good, P., 2005. Permutation, parametric, and bootstrap tests of hypotheses, 3rd Edition. Springer Science and Business Media Inc., New York, 316 p.
- Healey, A., Mitchell, R., Upcroft, J.A., Boreham, P.F., Upcroft, P., 1990. Complete nucleotide sequence of the ribosomal RNA tandem repeat unit from *Giardia intestinalis*. *Nucleic Acids Res.* 18, 4006.
- Hopkins, R.M., Meloni, B.P., Groth, D.M., Wetherall, J.D., Reynoldson, J.A., Thompson, R.C., 1997. Ribosomal RNA sequencing reveals differences between the genotypes of *Giardia* isolates recovered from humans and dogs living in the same locality. *J. Parasitol.* 83, 44-51.
- Humane Society of the United States:  
[http://www.humanesociety.org/issues/pet\\_overpopulation/facts/pet\\_ownership\\_statistics.html](http://www.humanesociety.org/issues/pet_overpopulation/facts/pet_ownership_statistics.html). Accessed 1 November 2014.
- Joffe, D., Van Niekerk, D., Gagne, F., Gilleard, J., Kutz, S., Lobingier, R., 2011. The prevalence of intestinal parasites in dogs and cats in Calgary, Alberta. *Can. Vet. J.* 52, 1323-1328.



- Labierre, A.F., Ripple, W.J., 2004. Range contractions of North American carnivores and ungulates. *Bioscience* 54, 123-138.
- Leonhard, S., Pfister, K., Beelitz, P., Wielinga, C., Thompson, R.C., 2007. The molecular characterisation of *Giardia* from dogs in southern Germany. *Vet. Parasitol.* 150, 33-38.
- Liccioli, S., Catalano, S., Kutz, S.J., Lejeune, M., Verocai, G.G., Duignan, P.J., Fuentealba, C., Hart, M., Ruckstuhl, K.E., Massolo, A., 2012. Gastrointestinal parasites of coyotes (*Canis latrans*) in the metropolitan area of Calgary, Alberta, Canada. *Can. J. Zool.* 90, 1023-1030.
- Liccioli, S., Duignan, P.J., Lejeune, M., Deunk, J., Majid, S., Massolo, A., 2013. A new intermediate host for *Echinococcus multilocularis*: the southern red-backed vole (*Myodes gapperi*) in urban landscape in Calgary, Canada. *Parasitol. Int.* 62, 355-357.
- Little, S.E., Johnson, E.M., Lewis, D., Jaklitsch, R.P., Payton, M.E., Blagburn, B.L., Bowman, D.D., Moroff, S., Tams, T., Rich, L., Aucoin, D., 2009. Prevalence of intestinal parasites in pet dogs in the United States. *Vet. Parasitol.* 166, 144-152.
- Lucio-Forster, A., Griffiths, J.K., Cama, V.A., Xiao, L., Bowman, D.D., 2010. Minimal zoonotic risk of cryptosporidiosis from pet dogs and cats. *Trends Parasitol.* 26, 174-179.
- Marangi, M., Berrilli, F., Otranto, D., Giangaspero, A., 2009. Genotyping of *Giardia duodenalis* among children and dogs in a closed socially deprived community from Italy. *Zoonoses Public Health* 57, 54-58.
- McDowall, R.M., Peregrine, A.S., Leonard, E.K., Lacombe, C., Lake, M., Rebelo, A.R., Cai, H.Y., 2011. Evaluation of the zoonotic potential of *Giardia duodenalis* in fecal samples from dogs and cats in Ontario. *Can. Vet. J.* 52, 1329-1333.
- Morey, P.S., Gese, E.M., Gehrt, S.D., 2007. Spatial and temporal variation in the diet of coyotes in the Chicago metropolitan area. *American Midland Naturalist* 158, 147-161.

Palmer, C.S., Traub, R.J., Robertson, I.D., Devlin, G., Rees, R., Thompson, R.C., 2008.

Determining the zoonotic significance of *Giardia* and *Cryptosporidium* in Australian dogs and cats. *Vet. Parasitol.* 154, 142-147.

Perrin, T., 2009. The business of urban animals survey: the facts and statistics on companion animals in Canada. *Can. Vet. J.* 50, 48-52.

Prabhu, A., Morrison, H.G., Martinez, C.R., Adam, R.D., 2007. Characterisation of the subtelomeric regions of *Giardia lamblia* genome isolate WBC6. *Int. J. Parasitol.* 37, 503-513.

Rimhanen-Finne, R., Enemark, H.L., Kolehmainen, J., Toropainen, P., Hanninen, M.L., 2007. Evaluation of immunofluorescence microscopy and enzyme-linked immunosorbent assay in detection of *Cryptosporidium* and *Giardia* infections in asymptomatic dogs. *Vet. Parasitol.* 145, 345-348.

Ruecker, N.J., Hoffman, R.M., Chalmers, R.M., Neumann, N.F., 2011. Detection and resolution of *Cryptosporidium* species and species mixtures by genus-specific nested PCR-restriction fragment length polymorphism analysis, direct sequencing, and cloning. *Appl. Environ. Microbiol.* 77, 3998-4007.

Ruecker, N.J., Matsune, J.C., Lapen, D.R., Topp, E., Edge, T.A., Neumann, N.F., 2013. The detection of *Cryptosporidium* and the resolution of mixtures of species and genotypes from water. *Infect. Genet. Evol.* 15, 3-9.

Ruecker, N.J., Matsune, J.C., Wilkes, G., Lapen, D.R., Topp, E., Edge, T.A., Sensen, C.W., Xiao, L., Neumann, N.F., 2012. Molecular and phylogenetic approaches for assessing sources of *Cryptosporidium* contamination in water. *Water. Res.* 46, 5135-5150.

Ryan, U., Fayer, R., Xiao, L., 2014. *Cryptosporidium* species in humans and animals: current understanding and research needs. *Parasitol* 141, 1667-1685.

- Ryan, U., Hijjawi, N., 2015. New developments in *Cryptosporidium* research. *Int. J. Parasitol.* 45, 367-373.
- Santner, T.J., Duffy, D.E., 1989. The statistical analysis of discrete data. Springer-Verlag, New York, 367 p.
- Scaramozzino, P., Di Cave, D., Berrilli, F., D'Orazi, C., Spaziani, A., Mazzanti, S., Scholl, F., De Liberato, C., 2009. A study of the prevalence and genotypes of *Giardia duodenalis* infecting kennelled dogs. *Vet. J.* 182, 231-234.
- Smith, A.F., Rock, M., Neumann, N., Massolo, A., 2015. Urban park-related risks for *Giardia* spp. infection in dogs. *Epidemiol. Infect.* 143, 3277-3291.
- Smith, A.F., Semeniuk, C.A., Kutz, S.J., Massolo, A., 2014. Dog-walking behaviours affect gastrointestinal parasitism in park-attending dogs. *Parasit. Vectors* 7, 429.
- Smith, R.P., Chalmers, R.M., Elwin, K., Clifton-Hadley, F.A., Mueller-Doblies, D., Watkins, J., Paiba, G.A., Giles, M., 2009. Investigation of the role of companion animals in the zoonotic transmission of cryptosporidiosis. *Zoonoses Public Health* 56, 24-33.
- Solarczyk, P., Majewska, A.C., 2010. A survey of the prevalence and genotypes of *Giardia duodenalis* infecting household and sheltered dogs. *Parasitol. Res.* 106, 1015-1019.
- Sotiriadou, I., Pantchev, N., Gassmann, D., Karanis, P., 2013. Molecular identification of *Giardia* and *Cryptosporidium* from dogs and cats. *Parasite* 20, 1-7.
- Sprong, H., Caccio, S.M., van der Giessen, J.W., 2009. Identification of zoonotic genotypes of *Giardia duodenalis*. *PLoS Negl. Trop. Dis.* 3, e558.
- Thompson, R.C., Colwell, D.D., Shury, T., Appelbee, A.J., Read, C., Njiru, Z., Olson, M.E., 2009. The molecular epidemiology of *Cryptosporidium* and *Giardia* infections in coyotes from Alberta, Canada, and observations on some cohabiting parasites. *Vet. Parasitol.* 159, 167-170.

- Thompson, R.C., Palmer, C.S., O'Handley, R., 2008. The public health and clinical significance of *Giardia* and *Cryptosporidium* in domestic animals. *Vet. J.* 177, 18-25.
- Thompson, R.C.A., Ash, A., 2016. Molecular epidemiology of *Giardia* and *Cryptosporidium* infections. *Infect. Genet. Evol.* 40, 315-323.
- Trout, J.M., Santin, M., Fayer, R., 2006. *Giardia* and *Cryptosporidium* species and genotypes in coyotes (*Canis latrans*). *J. Zoo Wildl. Med.* 37, 141-144.
- Uehlinger, F.D., Greenwood, S.J., McClure, J.T., Conboy, G., O'Handley, R., Barkema, H.W., 2013. Zoonotic potential of *Giardia duodenalis* and *Cryptosporidium* spp. and prevalence of intestinal parasites in young dogs from different populations on Prince Edward Island, Canada. *Vet. Parasitol.* 196, 509-514.
- United States Environmental Protection Agency: Method 1622/1623: *Giardia* & *Cryptosporidium* <https://www.epa.gov/sites/production/files/2015-07/documents/epa-1623.pdf>. Accessed 1 November 2014.
- van Keulen, H., Feely, D.E., Macechko, P.T., Jarroll, E.L., Erlandsen, S.L., 1998. The sequence of *Giardia* small subunit rRNA shows that voles and muskrats are parasitized by a unique species *Giardia microti*. *J. Parasitol.* 84, 294-300.
- van Keulen, H., Gutell, R.R., Campbell, S.R., Erlandsen, S.L., Jarroll, E.L., 1992. The nucleotide sequence of the entire ribosomal DNA operon and the structure of the large subunit rRNA of *Giardia muris*. *J. Mol. Evol.* 35, 318-328.
- van Keulen, H., Gutell, R.R., Gates, M.A., Campbell, S.R., Erlandsen, S.L., Jarroll, E.L., Kulda, J., Meyer, E.A., 1993. Unique phylogenetic position of Diplomonadida based on the complete small subunit ribosomal RNA sequence of *Giardia ardeae*, *G. muris*, *G. duodenalis*, and *Hexamita* sp. *FASEB J.* 7, 223-231.

- van Keulen, H., Homan, W.L., Erlandsen, S.L., Jarroll, E.L., 1995. A three nucleotide signature sequence in small subunit rRNA divides human *Giardia* in two different genotypes. *J. Eukaryot. Microbiol.* 42, 392-394.
- van Keulen, H., Macechko, P.T., Wade, S., Schaaf, S., Wallis, P.M., Erlandsen, S.L., 2002. Presence of human *Giardia* in domestic, farm, and wild animals, and environmental samples suggests a zoonotic potential for giardiasis. *Vet. Parasitol.* 108, 97-107.
- Veit, P., Bilger, B., Schad, V., Schafer, J., Frank, W., Lucius, R., 1995. Influence of environmental factors on the infectivity of *Echinococcus multilocularis* eggs. *Parasitol.* 110 (Pt 1), 79-86.
- Volotao, A.C., Costa-Macedo, L.M., Haddad, F.S., Brandao, A., Peralta, J.M., Fernandes, O., 2007. Genotyping of *Giardia duodenalis* from human and animal samples from Brazil using beta-giardin gene: a phylogenetic analysis. *Acta Trop.* 102, 10-19.
- Wang, A., Ruch-Gallie, R., Scorza, V., Lin, P., Lappin, M.R., 2012. Prevalence of *Giardia* and *Cryptosporidium* species in dog park attending dogs compared to non-dog park attending dogs in one region of Colorado. *Vet. Parasitol.* 184, 335-340.
- Xiao, L., 2010. Molecular epidemiology of cryptosporidiosis: an update. *Exp. Parasitol.* 124, 80-89.
- Xiao, L., Cama, V.A., Cabrera, L., Ortega, Y., Pearson, J., Gilman, R.H., 2007. Possible transmission of *Cryptosporidium canis* among children and a dog in a household. *J. Clin. Microbiol.* 45, 2014-2016.
- Xiao, L., Fayer, R., 2008. Molecular characterisation of species and genotypes of *Cryptosporidium* and *Giardia* and assessment of zoonotic transmission. *Int. J. Parasitol.* 38, 1239-1255.

- Xiao, L., Singh, A., Limor, J., Graczyk, T.K., Gradus, S., Lal, A., 2001. Molecular characterization of *Cryptosporidium* oocysts in samples of raw surface water and wastewater. *Appl. Environ. Microbiol.* 67, 1097-1101.
- Xiao, L., Sulaiman, I.M., Ryan, U.M., Zhou, L., Atwill, E.R., Tischler, M.L., Zhang, X., Fayer, R., Lal, A.A., 2002. Host adaptation and host-parasite co-evolution in *Cryptosporidium*: implications for taxonomy and public health. *Int. J. Parasitol.* 32, 1773-1785.
- Zanzani, S.A., Gazzonis, A.L., Scarpa, P., Berrilli, F., Manfredi, M.T., 2014. Intestinal parasites of owned dogs and cats from metropolitan and micropolitan areas: prevalence, zoonotic risks, and pet owner awareness in northern Italy. *Biomed Res. Int.* 2014, 1-10.

**Figure Legends****Figure 1. Two images depicting results from repetitive nested PCR.**

(a) Four replicates of amplified *G. intestinalis* 16S SSU rRNA gene product (292bp) from a dog (sample number B38) (b) Two replicates of amplified *C. canis* 18S SSU rRNA gene product (800bp) from a dog (sample number W122). MW indicates a 100bp molecular weight ladder.

