

Weak solution to the Navier-Stokes equations constructed by semi-discretization are suitable

Luigi C. Berselli and Stefano Spirito

Dedicated to Hugo Beirão da Veiga on the occasion of his seventieth birthday

ABSTRACT. We consider the three dimensional Navier-Stokes equations and we prove that weak solutions constructed by approximating the time-derivative by finite differences are suitable. The so-called method of semi-discretization is of fundamental importance in the numerical analysis and it is one of the building blocks for the full discretization of the equations.

1. Introduction

In this paper we consider the three dimensional Navier-Stokes equations with unit viscosity and zero external force (assumptions which are unessential since corresponding result can be obtained by scaling or by adding a smooth enough external force), namely

$$(1.1) \quad \begin{aligned} \partial_t u - \Delta u + (u \cdot \nabla)u + \nabla p &= 0 & (t, x) \in]0, T[\times]0, 2\pi[^3, \\ \nabla \cdot u &= 0 & (t, x) \in]0, T[\times]0, 2\pi[^3, \\ u(0, x) &= u_0(x) & x \in]0, 2\pi[^3, \end{aligned}$$

and, for simplicity, we restrict to the space-periodic setting. More general, but more technical results in a bounded domain with Dirichlet boundary condition will appear in a forthcoming work, still in progress [7]. In this short paper we address just the space-periodic case to explain the main ideas needed to handle the time-discretization, a topic which seems not explored yet, in the context of construction of solutions satisfying the local energy inequality.

Starting from the results of Scheffer [22] and Caffarelli, Kohn, and Nirenberg [9] concerning the partial regularity for the Navier-Stokes equations, it turned out that *Suitable Weak Solution* (SWS in the sequel) are of paramount importance. We recall that the weak solutions constructed by Leray and Hopf (for the Cauchy problem and for the bounded domain with Dirichlet boundary conditions, respectively) satisfy the equations in a weak sense:

$$\int_0^{+\infty} (u, \partial_t \phi) - (\nabla u, \nabla \phi) - ((u \cdot \nabla)u, \phi) dt = -(u_0, \phi(0)),$$

for all smooth, periodic, and divergence-free functions $\phi : [0, +\infty[\times]0, 2\pi[^3 \rightarrow \mathbb{R}^3$, such that $\phi(t, x) = 0$, for all $t \geq T$; The initial datum is attained in the sense that

$$\lim_{t \rightarrow 0^+} \|u(t) - u_0\| = 0;$$

The velocity u satisfies also the energy inequality

$$\frac{1}{2}\|u(t)\|^2 + \int_0^t \|\nabla u(s)\|^2 ds \leq \frac{1}{2}\|u_0\|^2 \quad \text{for all } t \in [0, T],$$

where $\|\cdot\|$ denotes the $L^2([0, 2\pi[^3)$ -norm, and (\cdot, \cdot) the associated scalar product.

On the other hand, the celebrated results of partial regularity from [9], concern solutions with a more stringent condition: Solutions need to be also suitable weak solutions, that is certain regularity on the pressure p is requested and the local-energy inequality

$$\partial_t \left(\frac{1}{2}|u|^2 \right) + \nabla \cdot \left(\left(\frac{1}{2}|u|^2 + p \right) u \right) - \Delta \left(\frac{1}{2}|u|^2 \right) + |\nabla u|^2 \leq 0,$$

has to be satisfied in the sense of distributions.

In [9] an approximation scheme with retarded mollifiers, which resemble the ‘‘Tonelli approximation’’ for ordinary differential equations, has been introduced in order to prove existence of SWS. This opened immediately the question whether Leray-Hopf weak solutions are suitable or not, hence if the partial regularity is valid for them. Two different techniques producing SWS have been later introduced by Beirão da Veiga in 1985. In a couple of companion papers [1, 2] (but see also [3]) he addressed in two elegant ways the problem, showing that the approximation by the bi-Laplacian (Δ^2) and also another (very general) one for the convective term lead to SWS. At present it is known that most of the infinite-dimensional approximations produce suitable weak solutions, even if proofs can become very technical (or in some case are still unknown) in the case of a bounded domain with Dirichlet boundary conditions. In this respect we want to recall the results on Leray- α approximation [16], those for the compressible approximation in [11], and the forthcoming analysis of the Voigt approximation and the compressible approximation [5, 6], with particular emphasis on the treatment of solid boundaries. We also recall the existence of SWS via smoothing with Yosida approximation by Farwig, Kozono, and Sohr [12] in very general (also unbounded and with non-compact boundary) domains.

Probably even more importantly, in the introduction of [2], Beirão da Veiga highlighted the fact that:

There is no evidence that solutions obtained by the Faedo-Galerkin method verify the local energy inequality.

This is a fundamental question, since approximations by finite dimensional spaces are the most relevant for the numerical analysis of the problem. The first partial solution to this problem came with the two companion papers by Guermond [13, 14], where he proved that if projectors over the finite element spaces used to discretize (with respect to x) velocity and pressure satisfy certain *commutation* properties, then the weak solutions constructed in the limit of vanishing mesh-size are suitable. In particular, these results cover the MINI element and the Taylor-Hood one. What is still missing is the case of the Fourier-Galerkin method in the three-dimensional torus. Further partial results in this case have been later obtained by Biryuk, Craig, and Ibrahim [8], but the problem –in its full-generality– is still unsolved. The links

and connections between the local energy inequality, the numerical approximations, and also the theory of turbulence and its description, raised in the last decade. Beside the role for partial regularity results, the notion of SWS represents now also a paradigm in the behavior of reasonable Large Eddy Simulation approximations to the Navier-Stokes equations, see Guermond *et al.* [15, 17]. In fact, making a parallel with the notion of entropy solutions, it is suggested that LES models should select “physically relevant” solutions of the Navier-Stokes equations, namely those which satisfy the local energy inequality.

In this paper, we continue in the spirit of connecting results from analysis, with those requested to design efficient and stable schemes for the numerical approximation of the 3D Navier-Stokes equations. In particular, we focus on the problem whether approximations of the Navier-Stokes by numerical methods produce SWS in the limit of the mesh-size (or discretization parameter) going to zero. More precisely, we focus here only on the time-discretization and we consider approximation of the time-derivative obtained by finite differences. A time-step size $\kappa > 0$ is given and $M = \lceil T/\kappa \rceil \in \mathbb{N}$ is the total number of time-steps, while $I^M := \{t_m\}_{m=0}^M$, with $t_m := m\kappa$, is the corresponding net. We then make the following backward finite-difference approximation for the time-derivative in the interval $]t_{m-1}, t_m[$

$$\partial_t u \sim d_t u^m := \frac{u^m - u^{m-1}}{\kappa}.$$

This produces for each $1 \leq m \leq M$ an elliptic problem, which is properly determined by the type of numerical algorithm chosen. This technique of discretization in time, while the problem remains continuous in space, is called Rothe method (or method of lines, or method of semi-discretization), see for instance Rektorys [21] and Kačur [18]. A more specific analysis for the Navier-Stokes equations is done by Temam [23, Ch. III, § 4] and Rautmann [20]. The time discretization can be also coupled with the space one, in order to produce a full numerical approximation.

In particular, we first analyze the following scheme, briefly discussing related numerical methods.

Algorithm. (Euler implicit) Given a time-step-size $\kappa > 0$ and the corresponding net $I^M = \{t_m\}_{m=0}^M$, for $m \geq 1$ and for u^{m-1} given from the previous step with $u^0 = u_0$, compute the iterate u^m as follows: Solve

$$(NS^k) \quad \begin{cases} d_t u^m - \Delta u^m + (u^m \cdot \nabla) u^m + \nabla p^m = 0, \\ \nabla \cdot u^m = 0, \end{cases}$$

endowed with periodic boundary conditions

REMARK 1.1. The scheme (NS^k) is an Euler implicit scheme and, at each step, a fully implicit equation has to be solved. Observe that for each given t_m we have to solve a problem very close to the stationary Navier-Stokes one, for which we have standard results of existence of weak solutions u^m , but the most important part is obtaining estimates on $\{u^m\}_{m=1, \dots, M}$ independent of κ .

For each m , we also associate to u^m a corresponding pressure p^m and then, to the sequence $\{u^m, p^m\}_{m=1, \dots, M}$ we can associate the functions (v_M, u_M, q_M) defined

in $[0, T]$, as follows for $m = 1, \dots, M$:

$$(1.2) \quad \begin{cases} v_M(t) = u^{m-1} + \frac{t - t_{m-1}}{\kappa} (u^m - u^{m-1}) & \text{for } t \in [t_{m-1}, t_m[, \\ v_M(t) = u^M & \text{for } t = t_M, \\ u_M(t) = u^0 & \text{for } t = t_0, \\ u_M(t) = u^m & \text{for } t \in]t_{m-1}, t_m], \\ q_M(t) = p^m & \text{for } t \in]t_{m-1}, t_m], \end{cases}$$

in such a way that $v_M(t_m) = u_M(t_m)$, for all $m = 0, \dots, M$.

The main result of this paper is the following:

THEOREM 1.2. *Let be given $u_0 \in H^1$. Then, there exists $(v, q) \in L^\infty(0, T; L^2) \cap L^2(0, T; H^1) \times L^{5/3}(0, T; L^{5/3})$ and a sequence $\kappa \rightarrow 0$ ($M \rightarrow +\infty$) such that the functions v_M and u_M both converge to u , and q_M converges to q . The functions v is a Leray-Hopf weak solutions to the Navier-Stokes equations (1.1) and moreover, for all $\phi \in C^\infty(\mathbb{T} \times [0, T])$ such that $\phi \geq 0$ and $\phi(0, x) = 0$, the couple (v, q) satisfies the local energy inequality*

$$(1.3) \quad \begin{aligned} & \int_{\Omega} |v(x, t)|^2 \phi(x, t) dx + 2 \int_0^t \int_{\Omega} |\nabla v(x, \tau)|^2 \phi(x, \tau) dx d\tau \\ & \leq \int_0^t \int_{\Omega} \left(|v(x, \tau)|^2 (\partial_t \phi(x, \tau) - \Delta \phi(x, \tau)) + (|v(x, \tau)|^2 \right. \\ & \quad \left. + 2q(x, \tau)) v(x, \tau) \cdot \nabla \phi(x, \tau) \right) dx d\tau. \end{aligned}$$

The existence part is rather standard, see Temam [23, Ch. III, § 4], while the original contribution of this paper is the verification of the local energy inequality (1.3).

REMARK 1.3. The hypothesis on the initial datum can be relaxed to the more natural condition of square integrability. We are assuming more regularity to keep the proof as simple as possible and without technicalities. We also observe that this point (see also [21, Ch. 13]) is generally not treated or overlooked in the literature and requires to handle a further technical part, which will be appropriately detailed in [7].

Beside the proof being elementary, this result will represent the core of more general results which will correlate the full space-time discretization of the Navier-Stokes equations, with the notion of local energy inequality [7]. In particular, in a forthcoming paper we shall consider space-time discretizations as those in Quarteroni and Valli [19, § 13.4], but following a different path: Time-discretization will be performed before numerical discretization of the space variables.

2. Notation

We briefly fix the notation, which is typical of space-periodic problems. In the sequel we shall use the customary Lebesgue spaces $L^p(\Omega)$ and Sobolev spaces

$W^{k,p}(\Omega)$ and $H^s(\Omega) := W^{s,2}(\Omega)$, with $\Omega :=]0, 2\pi[^3$; for simplicity we shall do not distinguish between scalar and vector valued functions. Since we shall work with periodic boundary conditions the spaces are made of periodic functions. In the Hilbertian case $p = 2$ we can easily characterize the divergence-free spaces by using Fourier Series on the 3D torus. We denote by (e_1, e_2, e_3) the orthonormal basis of \mathbb{R}^3 , and by $x := (x_1, x_2, x_3)$ the generic point in \mathbb{R}^3 . Let \mathbb{T} be the torus defined by $\mathbb{T} := \mathbb{R}^3/2\pi\mathbb{Z}^3$. We use $\|\cdot\|$ to denote the $L^2(\mathbb{T})$ norm and we impose the zero mean condition $\int_{\Omega} \phi dx = 0$ on velocity and pressure. We also define, for an exponent $s \geq 0$,

$$H^s := \left\{ w : \mathbb{T} \rightarrow \mathbb{R}^3, w \in H^s(\mathbb{T})^3, \nabla \cdot w = 0, \int_{\mathbb{T}} w dx = 0 \right\},$$

where $H^s(\mathbb{T})^3 := [H^s(\mathbb{T})]^3$ and if $0 \leq s < 1$ the condition $\nabla \cdot w = 0$ must be understood in a weak sense. For $w \in H^s$, we can expand the velocity field with Fourier series $w(x) = \sum_{k \neq 0} \widehat{w}_k e^{ik \cdot x}$, where k is the wave-vector and the Fourier coefficients are given by $\widehat{w}_k = \frac{1}{(2\pi)^3} \int_{\mathbb{T}} w(x) e^{-ik \cdot x} dx$. If $|k| = \sqrt{|k_1|^2 + |k_2|^2 + |k_3|^2}$, then the H^s norm is defined by

$$\|w\|_{H^s}^2 := \sum_{k \neq 0} |k|^{2s} |\widehat{w}_k|^2,$$

where, as above, $\|w\|_{H^0} := \|w\|$. We finally characterize $H^s \subset H^s(\mathbb{T})$ as follows:

$$H^s := \left\{ w = \sum_{k \neq 0} \widehat{w}_k e^{ik \cdot x} : \sum_{k \neq 0} |k|^{2s} |\widehat{w}_k|^2 < \infty, k \cdot \widehat{w}_k = 0, \widehat{w}_{-k} = \overline{\widehat{w}_k} \right\}.$$

Together with the classical Bochner spaces $L^p(0, T; X)$, endowed with the norm

$$\|f\|_{L^p(0, T; X)} := \begin{cases} \left(\int_0^T \|f(s)\|_X \right)^{1/p} & \text{if } 1 \leq p < \infty, \\ \sup_{0 \leq s \leq T} \|f(s)\|_X & \text{if } p = +\infty, \end{cases}$$

to deal with discrete problems we shall make use of the natural weighted spaces $l^p(I^M; X)$. The discrete counterpart of $L^p(0, T; X)$ consists of X -valued sequences $\{a_m\}_{m=0}^M$, endowed with the norm

$$\|a_m\|_{l^p(I^M; X)} := \begin{cases} \left(\kappa \sum_{m=0}^M \|a_m\|_X^p \right)^{1/p} & \text{if } 1 \leq p < \infty, \\ \max_{0 \leq m \leq M} \|a_m\|_X & \text{if } p = +\infty. \end{cases}$$

We first recall a lemma about existence of solutions, see for instance Temam [23, Ch. III, Lem. 4.3].

LEMMA 2.1. *There exists at least one sequence $\{u^m\}_{m=0}^M$ defined by the algorithm (NS^k) with $u^0 = u_0$.*

PROOF. Observe that we have $M - 1$ coupled systems and in particular, given $u^{m-1} \in H^1$ and $\kappa > 0$, the function $u^m \in H^1$ can be obtained as solution of the

modified steady Navier-Stokes system

$$\begin{cases} \frac{u^m}{\kappa} - \Delta u^m + (u^m \cdot \nabla) u^m + \nabla p^m = \frac{u^{m-1}}{\kappa}, \\ \nabla \cdot u^m = 0, \end{cases}$$

By testing with u^m itself (reasoning which can be made rigorous by a Faedo-Galerkin approximation) one obtains

$$\frac{\|u^m\|^2}{2\kappa} + \|\nabla u^m\|^2 \leq \frac{\|u^{m-1}\|^2}{2\kappa},$$

and it is enough to use the Brouwer fixed point theorem to infer existence of at least one solution $u^m \in H^1$. The existence of the associated p^m is then obtained by De Rham theorem. Observe that we do not have neither uniqueness, neither estimates independent of $\kappa > 0$. \square

REMARK 2.2. In many problems of numerical analysis it is unpleasant to have non-uniqueness for the approximate discrete problem. In the perspective of trying to have better properties of regularity on u^m , which could traduce in better convergence, it is also worth considering the following scheme:

Algorithm. (Euler semi-implicit) Given a time-step-size $\kappa > 0$ and the corresponding net $I^M = \{t_m\}_{m=0}^M$, for $m \geq 1$ and u^{m-1} given from the previous step with $u^0 = u_0$, compute the iterate u^m as follows: Solve

$$(2.1) \quad \begin{cases} d_t u^m - \Delta u^m + (u^{m-1} \cdot \nabla) u^m + \nabla p^m = 0, \\ \nabla \cdot u^m = 0, \end{cases}$$

endowed with periodic boundary conditions

The role of the semi-implicit approximation is emphasized for instance in [4], where it represents a critical tool to obtain optimal convergence rates for shear-dependent fluids with nonlinear viscosities (and consequently with the lack of the standard regularity results known for strong solutions to the Navier-Stokes equations). See also Diening, Ebmeyer, and Růžička [10] for general parabolic system.

For system (2.1) we have to solve for each $m = 1, \dots, M$ the linear system

$$\begin{cases} \frac{u^m}{\kappa} - \Delta u^m + (u^{m-1} \cdot \nabla) u^m + \nabla p^m = \frac{u^{m-1}}{\kappa}, \\ \nabla \cdot u^m = 0, \end{cases}$$

for which the *a-priori* estimate is the same as for the implicit scheme (NS^k), since $\int_{\mathbb{T}} (u^{m-1} \cdot \nabla) u^m \cdot u^m dx = 0$, and uniqueness follows from linearity of the problem.

3. Proof of the main result

In order to show that solutions constructed by the algorithm (NS^k) converge to a SWS, we need to pass to the limit as $\kappa \rightarrow 0^+$ (and consequently $M \rightarrow +\infty$), hence estimates independent of κ are requested. To this end, we multiply the equations (NS^k) by u^m itself and we use a slightly different argument to prove the following lemma. Observe that the same argument will also work for the algorithm (2.1). We mainly consider (NS^k), since it is the first one found in many texts when proposing alternate proofs of existence of weak solutions by semi discretization.

LEMMA 3.1. *Let be given $u_0 \in H^1$. Then, there exists a constant $C > 0$, (independent of κ) such that*

$$(3.1) \quad \begin{aligned} \|u^m\|_{\ell^\infty(I^M; H^0) \cap \ell^2(I^M; H^1)} &\leq C, \\ \|p^m\|_{\ell^{5/3}(I^M; L^{5/3}(\mathbb{T}))} &\leq C. \end{aligned}$$

PROOF. We test the equations (NS^k) by u^m . By integration by parts and with the elementary algebra equality

$$(a - b, a) = \frac{a^2 - b^2}{2} + \frac{(a - b)^2}{2} \quad a, b \in \mathbb{R},$$

we easily get

$$\frac{1}{2} d_t \|u^m\|^2 + \frac{\kappa}{2} \|d_t u^m\|^2 + \|\nabla u^m\|^2 = 0.$$

Next, by multiplying by $\kappa > 0$ and, summing up over $m = 1, \dots, M$, we obtain

$$(3.2) \quad \frac{1}{2} \|u^M\|^2 + \frac{\kappa^2}{2} \sum_{m=1}^M \|d_t u^m\|^2 + \kappa \sum_{m=1}^M \|\nabla u^m\|^2 \leq \frac{1}{2} \|u_0\|^2.$$

From the last inequality (considering also sums up to any number smaller or equal than M) we deduce

$$u^m \in \ell^\infty(I^M; H^0) \cap \ell^2(I^M; H^1).$$

The definition of the weighted spaces $\ell^p(I^M; X)$ allows us to use the standard Hölder inequality. By using also the usual interpolation results one directly obtains

$$u^m \in \ell^{10/3}(I^M; L^{10/3}(\mathbb{T})).$$

As we previously observed in Lemma 2.1, to each u^m we can also associate a discrete pressure p^m , by using De Rham theorem. Here we want to have a more precise information about the regularity of p^m and this is the crucial point where we use the space periodicity. By taking the divergence of (NS^k) and since $\nabla \cdot u^m = 0$, we get the following Poisson equation (endowed with periodic boundary conditions)

$$(3.3) \quad -\Delta p^m = \nabla \cdot (u^m \cdot \nabla) u^m = \sum_{i,j=1}^3 \frac{\partial}{\partial x_i} \frac{\partial}{\partial x_i} u_i^m u_j^m, \quad m = 1, \dots, M,$$

where u_l^m denotes the l -component of the vector u^m . By using that $u_i^m u_j^m \in \ell^{5/3}(I^M; L^{5/3}(\mathbb{T}))$, the unique solution p^m with zero mean value of (3.3) satisfies

$$p^m \in \ell^{5/3}(I^M; L^{5/3}(\mathbb{T})),$$

since it is obtained just inverting the Laplace operator for each $m = 1, \dots, M$. \square

From the estimate (3.2) we have the following inequality holds. We state it as a Lemma, because it will be crucial in the proof of the Theorem 1.2.

LEMMA 3.2. *The following estimate holds true*

$$\kappa^2 \sum_{m=1}^M \|d_t u^m\|^2 = \sum_{m=1}^M \|u^m - u^{m-1}\|^2 \leq \|u_0\|^2.$$

We observe that this estimate is obtained because the “*natural multiplier*” (the one which cancels out the convective term) is u^m , and the estimate comes from algebraic manipulation of the integral $\int_{\mathbb{T}} d_t u^m u^m dx$. In presence of different schemes this argument may fail and explains why a different treatment is required by other numerical methods, see Section 4. On the other hand, exactly the same estimates can be proved for the semi-implicit scheme (2.1), since again the correct multiplier is u^m .

We show now some properties of the step function u_M and of the piecewise linear function v_M .

LEMMA 3.3. *Let $u_0 \in H^1$ be given. Then, there exists a constant $C > 0$ (independent of κ) such that*

$$\begin{aligned} \|v_M\|_{L^\infty(0,T;H^0) \cap L^2(0,T;H^1)} &\leq C, \\ \|u_M\|_{L^\infty(0,T;H^0) \cap L^2(0,T;H^1)} &\leq C, \\ \|q_M\|_{L^{5/3}(0,T;L^{5/3}(\mathbb{T}))} &\leq C, \\ \|\partial_t v_M\|_{L^{4/3}(0,T;H^{-1})} &\leq C, \text{ where for any } s > 0, H^{-s} := (H^s)'. \end{aligned}$$

Moreover, we also have the following identity

$$(3.4) \quad \|u_M - w_M\|_{L^2(0,T;H^0)}^2 = \frac{\kappa}{3} \sum_{m=1}^M \|u^m - u^{m-1}\|^2.$$

PROOF. The results of Lemma 3.1 imply that the functions v_M defined in (1.2) are, for each positive M , Lipschitz functions $[0, T] \mapsto H^1$. For each M , the function v_M satisfies, in the sense of distributions over $]0, T[$, the following equality

$$(3.5) \quad \frac{d}{dt}(v_M, \psi) + (\nabla u_M, \nabla \psi) + ((u_M \cdot \nabla) u_M, \psi) dt = 0 \quad \forall \psi \in H^1.$$

Note that, by their definition (v_M, u_M, q_M) have the regularity stated in the lemma, which derive directly from the analogous one valid for the sequence $\{u^m, p^m\}$.

REMARK 3.4. Observe that the proof of existence of weak solutions in [23] is based on estimates for the discrete time-derivative in $L^2(H^{-s})$, for $s \geq 3/2$. The idea of obtaining such an estimates in negative spaces (but also considering negative Sobolev spaces in the time variable) represents also the core of the results in [13, 14]. Here, due to the particular setting, we can follow a more standard path.

The estimate $\partial_t v_M \in L^{4/3}(0, T; (H^{-1}))$ is obtained, as usual, by a comparison argument. It remains to prove estimate (3.4). We have

$$v_M(t) - u_M(t) = \frac{(t - t_{m-1})}{k} (u^m - u^{m-1}) + u^{m-1} - u^m \quad \forall t \in]t_{m-1}, t_m].$$

Then

$$\begin{aligned}
\int_0^T \|v_M - u_M\|^2 dt &= \sum_{m=1}^M \int_{t_{m-1}}^{t_m} \|v_M(t) - u_M(t)\|^2 dt \\
&= \sum_{m=1}^M \|u^m - u^{m-1}\|^2 \int_{t_{m-1}}^{t_m} \left(\frac{t - t_{m-1}}{k} - 1 \right)^2 dt \\
&= \frac{\kappa}{3} \sum_{m=1}^M \|u^m - u^{m-1}\|^2.
\end{aligned}$$

□

We are now ready to prove the main result of this paper.

PROOF OF THEOREM 1.2. From a standard application of the Aubin-Lions compactness argument, we get that from the sequence $\{v_M\}_M$ bounded uniformly in $L^2(0, T; H^1)$ and such that $\partial_t v_M \in L^{4/3}(0, T; (H^{-1}))$, again with bound independent of M , we can extract a (relabelled) sub-sequence $v_M \rightarrow v$ in $L^2(0, T; H^0)$. Moreover, by using (3.4) and Lemma 3.2 we have also that

$$u_M - v_M \rightarrow 0 \text{ in } L^2(0, T; H^0),$$

hence also the sequence u_M converges strongly to v in $L^2(0, T; H^0)$. By using standard interpolation inequality and the previous strong convergence, it is common to show also that

$$\begin{aligned}
v_M &\rightarrow v \quad \text{strongly in } L^3(0, T; L^3(\mathbb{T})), \\
u_M &\rightarrow v \quad \text{strongly in } L^3(0, T; L^3(\mathbb{T})).
\end{aligned}$$

Moreover, since q_M is uniformly bounded in $L^{5/3}((0, T) \times \mathbb{T})$, up to extraction of a further sub-sequence, we have that

$$q_M \rightarrow q \text{ weakly in } L^{5/3}((0, T) \times \mathbb{T}).$$

Finally, as in [23] we have that v is a weak solutions of the Navier-Stokes equations with associated pressure q . We show now that (v, q) satisfies the local energy inequality. To this end we test the equations (3.5) by $u_M \phi$, where $\phi \geq 0$ is periodic in the space variable, smooth, and vanishes for $t = 0, T$. The first term regarding the time-derivative is the most relevant for our purposes. We have

$$\begin{aligned}
\int_0^T (\partial_t v_M, u_M \phi) dt &= \int_0^T (\partial_t v_M, (v_M - v_M + u_M) \phi) dt \\
&= \int_0^T (\partial_t v_M, v_M) \phi dt + \int_0^T (\partial_t v_M, (u_M - v_M) \phi) dt \\
&= I_1 + I_2
\end{aligned}$$

We start with the first term I_1 . By splitting the integral over $[0, T]$ with the sum of integrals over $[t_{m-1}, t_m]$ and, by performing integration by parts, we immediately

obtain

$$\begin{aligned} \int_0^T (\partial_t v_M, v_M \phi) dt &= \sum_{m=1}^M \int_{t_{m-1}}^{t_m} (\partial_t v_M, v_M \phi) dt = \sum_{m=1}^M \int_{t_{m-1}}^{t_m} \left(\frac{1}{2} \partial_t |v_M|^2, \phi \right) dt \\ &= \frac{1}{2} \sum_{m=1}^M (|u^m|^2, \phi(t_m, x)) - (|u^{m-1}|^2, \phi(t_{m-1}, x)) - \sum_{m=1}^M \int_{t_{m-1}}^{t_m} \left(\frac{1}{2} |v_M|^2, \partial_t \phi \right) dt, \end{aligned}$$

where we used that $\partial_t v_M(t) = \frac{u^m - u^{m-1}}{\kappa}$, for $t \in [t_{m-1}, t_m[$. Next, we observe that the sum telescopes and consequently we have

$$\begin{aligned} \int_0^T (\partial_t v_M, v_M \phi) dt &= \frac{1}{2} (|u^M|^2, \phi(T, x)) - \frac{1}{2} (|u_0|^2, \phi(0, x)) - \sum_{m=1}^M \int_{t_{m-1}}^{t_m} \left(\frac{1}{2} |v_M|^2, \partial_t \phi \right) dt \\ &= - \int_0^T \left(\frac{1}{2} |v_M|^2, \partial_t \phi \right) dt. \end{aligned}$$

By the strong convergence of $v_M \rightarrow v$ in $L^2(0, T; L^2(\mathbb{T}))$ we can conclude that

$$\lim_{M \rightarrow +\infty} \int_0^T \int_{\mathbb{T}} \partial_t v_M v_M \phi dx dt = - \frac{1}{2} \int_0^T \int_{\mathbb{T}} |v|^2 \partial_t \phi dx dt.$$

Then, we consider the second term. Since u_M is constant on the interval $[t_{m-1}, t_m[$ we can write

$$\begin{aligned} \int_0^T (\partial_t v_M, (u_M - v_M) \phi) dt &= - \sum_{m=1}^M \int_{t_{m-1}}^{t_m} (\partial_t (v_M - u_M), (v_M - u_M) \phi) dt \\ &= - \sum_{m=1}^M \int_{t_{m-1}}^{t_m} \left(\partial_t \left(\frac{|v_M - u_M|^2}{2} \right), \phi \right) dt \\ &= \sum_{m=1}^M \int_{t_{m-1}}^{t_m} \left(\frac{|v_M - u_M|^2}{2}, \partial_t \phi \right) dt, \end{aligned}$$

where in the last line we have used the fact we do not have boundary terms because $v_M(t_m) = u_M(t_m)$ for all $m = 0, \dots, M$. Then, since $u_M - v_M$ goes to 0 strongly in $L^2(0, T; H^0)$, we get that $I_2 \rightarrow 0$ as $M \rightarrow +\infty$ (or $\kappa \rightarrow 0$).

By the usual reasoning we have (in this term the integration by parts is in space, so there is no need for a special treatment) that

$$\begin{aligned} - \int_0^T \int_{\mathbb{T}} \Delta u_M u_M \phi dx dt &= \int_0^T \int_{\mathbb{T}} |\nabla u_M|^2 \phi dx dt + \frac{1}{2} \int_0^T \int_{\mathbb{T}} \nabla |u_M|^2 \nabla \phi dx dt \\ &= \int_0^T \int_{\mathbb{T}} |\nabla u_M|^2 \phi dx dt - \frac{1}{2} \int_0^T \int_{\mathbb{T}} |u_M|^2 \Delta \phi dx dt, \end{aligned}$$

and integration by parts are possible due to the space-periodicity. By using the lower semi-continuity of the norm and $\phi \geq 0$, we obtain:

$$\lim_{M \rightarrow +\infty} \int_0^T \int_{\mathbb{T}} |\nabla u_M|^2 \phi dx \geq \int_0^T \int_{\mathbb{T}} |\nabla v|^2 \phi dx,$$

while again by the strong convergence in $L^2(0, T; L^2(\mathbb{T}))$

$$\lim_{M \rightarrow +\infty} \frac{1}{2} \int_0^T \int_{\mathbb{T}} |u_M|^2 \Delta \phi \, dx dt = \frac{1}{2} \int_0^T \int_{\mathbb{T}} |v|^2 \Delta \phi \, dx dt.$$

The convective term is treated again by integrating by parts. In fact, we obtain

$$\begin{aligned} \int_0^T \int_{\mathbb{T}} (u_M \cdot \nabla) u_M u_M \phi \, dx &= \frac{1}{2} \int_0^T \int_{\mathbb{T}} \nabla |u_M|^2 u_M \phi \, dx \\ &= -\frac{1}{2} \int_0^T \int_{\mathbb{T}} |u_M|^2 u_M \nabla \phi \, dx, \end{aligned}$$

and, by the strong convergence $u_M \rightarrow v$ in $L^3(0, T; L^3(\mathbb{T}))$, we get

$$\lim_{M \rightarrow +\infty} \int_0^T \int_{\mathbb{T}} (u_M \cdot \nabla) u_M u_M \phi \, dx = \frac{1}{2} \int_0^T \int_{\mathbb{T}} |v|^2 v \nabla \phi \, dx.$$

Finally, the term with the pressure is integrated by parts

$$\int_0^T \int_{\mathbb{T}} \nabla q_M u_M \phi \, dx = - \int_0^T \int_{\mathbb{T}} q_M u_M \nabla \phi \, dx,$$

and, thanks to the weak convergence $q_M \rightarrow q$ in $L^{5/3}(0, T; L^{5/3}(\mathbb{T}))$ and again the strong convergence of u_M in $L^3(0, T; L^3(\mathbb{T}))$, in we get

$$\lim_{M \rightarrow +\infty} \int_0^T \int_{\mathbb{T}} \nabla q_M u_M \phi \, dx = \int_0^T \int_{\mathbb{T}} q v \nabla \phi \, dx.$$

We finally proved that

$$\begin{aligned} &2 \int_0^T \int_{\mathbb{T}} |\nabla v(x, \tau)|^2 \phi(x, \tau) \, dx d\tau \\ &\leq \int_0^T \int_{\mathbb{T}} \left(|v(x, \tau)|^2 (\partial_t \phi(x, \tau) - \Delta \phi(x, \tau)) + (|v(x, \tau)|^2 \right. \\ &\quad \left. + 2q(x, \tau)) v(x, \tau) \cdot \nabla \phi(x, \tau) \right) \, dx d\tau, \end{aligned}$$

for all smooth and non-negative ϕ , which are space-periodic, and with compact support with respect to time.

By following the argument detailed in [9, p. 13], by with a further test-function only of the time variable and approximating a Dirac's delta at a given time $t \in]0, T]$, one can easily deduce from the latter the validity of (1.3) for a.e. $t \in [0, T]$. \square

4. Remarks on other algorithms

The algorithm (NS^k) analyzed in the previous section is a particular case of a more general algorithm, the so called θ -scheme, defined below (cf. [19, § 13.4]):

Algorithm. (θ -scheme) Let $\theta \in [0, 1]$. Given a time-step-size $\kappa > 0$ and the corresponding net $I^M = \{t_m\}_{m=0}^M$, for $m \geq 1$ and for u^{m-1} given from the previous step with $u^0 = u_0$, compute the iterate u^m as follows:

$$(NS_{\theta}^k) \quad \begin{cases} d_t u^m - \Delta u_{\theta}^m + (u_{\theta}^m \cdot \nabla) u_{\theta}^m + \nabla p_{\theta}^m = 0, \\ \nabla \cdot u_{\theta}^m = 0, \end{cases}$$

References

1. H. Beirão da Veiga, *On the suitable weak solutions to the Navier-Stokes equations in the whole space*, J. Math. Pures Appl. (9) **64** (1985), no. 1, 77–86. MR 802384 (87a:35149)
2. ———, *On the construction of suitable weak solutions to the Navier-Stokes equations via a general approximation theorem*, J. Math. Pures Appl. (9) **64** (1985), no. 3, 321–334. MR 823407 (87h:35268)
3. ———, *Local energy inequality and singular set of weak solutions of the boundary non-homogeneous navier-stokes problem*, Current topics in partial differential equations (Y. Ohya, K. Kasahara, and N. Shimakura, eds.), Kinokuniya Company Ltd., Tokyo, 1986, Papers dedicated to Professor Sigeru Mizohata on the occasion of his sixtieth birthday.
4. L. C. Berselli, L. Diening, and M. Růžička, *Optimal error estimates for a semi-implicit Euler scheme for incompressible fluids with shear dependent viscosities*, SIAM J. Numer. Anal. **47** (2009), no. 3, 2177–2202. MR 2519599
5. L. C. Berselli and S. Spirito, *Suitable weak solutions of the Navier-Stokes equations obtained with the voigt approximation in a bounded domain*, 2014, In preparation.
6. ———, *Suitable weak solutions of the Navier-Stokes equations as limit of artificial compressibility in a bounded domain*, 2014, In preparation.
7. ———, *Certain finite-difference/finite element space-time discretization of the Navier-Stokes equations are suitable*, 2014, In preparation.
8. A. Biryuk, W. Craig, and S. Ibrahim, *Construction of suitable weak solutions of the Navier-Stokes equations*, Stochastic analysis and partial differential equations, Contemp. Math., vol. 429, Amer. Math. Soc., Providence, RI, 2007, pp. 1–18. MR 2391525 (2009c:35337)
9. L. Caffarelli, R. Kohn, and L. Nirenberg, *Partial regularity of suitable weak solutions of the Navier-Stokes equations*, Comm. Pure Appl. Math. **35** (1982), no. 6, 771–831. MR 673830 (84m:35097)
10. L. Diening, C. Ebmeyer, and M. Růžička, *Optimal convergence for the implicit space-time discretization of parabolic systems with p -structure*, SIAM J. Numer. Anal. **45** (2007), no. 2, 457–472 (electronic). MR 2300281
11. D. Donatelli and S. Spirito, *Weak solutions of Navier-Stokes equations constructed by artificial compressibility method are suitable*, J. Hyperbolic Differ. Equ. **8** (2011), no. 1, 101–113. MR 2796051 (2012k:35387)
12. R. Farwig, H. Kozono, and H. Sohr, *An L^q -approach to Stokes and Navier-Stokes equations in general domains*, Acta Math. **195** (2005), 21–53. MR 2233684 (2007d:35207)
13. J.-L. Guermond, *Finite-element-based Faedo-Galerkin weak solutions to the Navier-Stokes equations in the three-dimensional torus are suitable*, J. Math. Pures Appl. (9) **85** (2006), no. 3, 451–464. MR 2210084 (2006m:35277)
14. ———, *Faedo-Galerkin weak solutions of the Navier-Stokes equations with Dirichlet boundary conditions are suitable*, J. Math. Pures Appl. (9) **88** (2007), no. 1, 87–106. MR 2334774 (2008f:35283)
15. ———, *On the use of the notion of suitable weak solutions in CFD*, Internat. J. Numer. Methods Fluids **57** (2008), no. 9, 1153–1170. MR 2435087 (2009j:76064)
16. J.-L. Guermond, J. T. Oden, and S. Prudhomme, *Mathematical perspectives on large eddy simulation models for turbulent flows*, J. Math. Fluid Mech. **6** (2004), no. 2, 194–248. MR 2053583 (2005b:76078)
17. J.-L. Guermond and S. Prudhomme, *On the construction of suitable solutions to the Navier-Stokes equations and questions regarding the definition of large eddy simulation*, Phys. D **207** (2005), no. 1-2, 64–78. MR 2166975 (2006e:76081)
18. J. Kačur, *Method of Rothe in evolution equations*, Teubner-Texte zur Mathematik, vol. 80, BSB B. G. Teubner Verlagsgesellschaft, Leipzig, 1985, With German, French and Russian summaries. MR 834176 (87j:35004)
19. A. Quarteroni and A. Valli, *Numerical approximation of partial differential equations*, Springer Series in Computational Mathematics, vol. 23, Springer-Verlag, Berlin, 1994. MR 1299729 (95i:65005)
20. R. Rautmann, *H^2 -convergence of Rothe's scheme to the Navier-Stokes equations*, Nonlinear Anal. **24** (1995), no. 7, 1081–1102. MR 1321746 (96a:76028)
21. K. Rektorys, *The method of discretization in time and partial differential equations*, Mathematics and Its Applications (East European Series), vol. 4, D. Reidel Publishing Co.,

- Dordrecht-Boston, Mass., 1982, Translated from the Czech by the author. MR 689712 (84k:65104)
22. V. Scheffer, *Hausdorff measure and the Navier-Stokes equations*, *Comm. Math. Phys.* **55** (1977), no. 2, 97–112. MR 0510154 (58 #23176)
 23. R. Temam, *Navier-Stokes equations. Theory and numerical analysis*, North-Holland Publishing Co., Amsterdam-New York-Oxford, 1977, *Studies in Mathematics and its Applications*, Vol. 2. MR 0609732 (58 #29439)

LUIGI C. BERSELLI, DIPARTIMENTO DI MATEMATICA, VIA BUONARROTI 1/C, I-56127 PISA, ITALY

E-mail address: berselli@dma.unipi.it

STEFANO SPIRITO, MATHEMATISCHES INSTITUT, UNIVERSITÄT BASEL, RHEINSPRUNG 21, CH-4051 BASEL

E-mail address: stefano.spirito@unibas.ch