



ATTI
DELLA
SOCIETÀ TOSCANA
DI
SCIENZE NATURALI

MEMORIE • SERIE A • VOLUME CXXXI • ANNO 2024



Edizioni ETS

ATTI DELLA SOCIETÀ TOSCANA DI SCIENZE NATURALI

MEMORIE

Via Santa Maria, 53 - 56126 Pisa

Rivista di proprietà della Società Toscana di Scienze Naturali. Fondata nel 1875, la rivista pubblica Memorie e Note originali, recensioni, corrispondenze e notiziari nel campo delle Scienze Naturali. È inviata ai soci e depositata in biblioteche corrispondenti. Tutti i contenuti dei volumi a stampa (articoli originali, comunicazioni brevi, notizie e recensioni) è liberamente disponibile in rete, in conformità all'Open Access, sulle pagine <http://www.stsn.it/en/memorie-serie-a.html> (Serie A) e <http://www.stsn.it/en/memorie-serie-b.html> (Serie B). *Journal owned by Società Toscana di Scienze Naturali. Founded in 1875, the journal publishes original papers, short communications, news and book reviews on Natural Sciences. The Journal is sent to Società Toscana di Scienze Naturali members and deposited in selected libraries. All content of the printed version (original papers, short communications, news and book reviews) is freely available online in accordance with the Open Access at <http://www.stsn.it/en/memorie-serie-a.html> (Serie A) and <http://www.stsn.it/en/memorie-serie-b.html> (Serie B).*

Gli Atti sono pubblicati in due volumi (Serie A - Abiologica, ISSN 0365-7655; Serie B - Biologica, ISSN 0365-7450) all'anno nel mese di dicembre. Possono essere pubblicati ulteriori volumi, definiti Supplementi, su temi specifici.

Atti are published yearly in two Issues (Serie A - Abiological, ISSN 0365-7655; Serie B - Biological, ISSN 0365-7450) in December. Some monographic volumes may be published as Supplementi.

Le lingue usate dalla rivista sono l'inglese o l'italiano // *The languages of the journal are English or Italian.*

Comitato scientifico (*Editorial Board*)

Serie A: G. Bianucci (Pisa, Italia), R. Blanco Chao, (Santiago de Compostela, Spagna), L. Disperati (Siena, Italia), W. Finsinger, (Montpellier, Francia), C. Montomoli (Torino, Italia), A. Oros Sršen (Zagabria, Croazia), E. Pandeli (Firenze, Italia), S. Richiano (Puerto Madrin, Argentina), E. Starnini (Pisa, Italia).

Serie B: A. Aguilera (Valencia, Spain), N.E. Baldaccini (Pisa, Italy), E. Biondi (Ancona, Italy), B. Foggi (Firenze, Italy), E. Palagi (Pisa, Italy), G. Paradis (Ajaccio, France), L. Peruzzi (Pisa, Italy), M. Zuffi (Pisa, Italy).

Direttore scientifico (*Editor in Chief*): Paolo Roberto Federici

Comitato di redazione (*Editorial Office*)

Serie A: A. Chelli (*Segretario agli Atti / Editor*)

Serie B: G. Bedini (*Segretario agli Atti / Editor*), G. Astuti, A. Carta, M. D'Antraccoli, L. Peruzzi, F. Roma-Marzio

La corrispondenza deve essere inviata ai Segretari agli Atti (per la Serie A: A. Chelli, Dipartimento di Chimica, Scienze della Vita e della Sostenibilità Ambientale, Parco Area delle Scienze, 157/A - 43124 Parma, email: alessandro.chelli@unipr.it; per la Serie B: G. Bedini, Dipartimento di Biologia, via Derna, 1 - 56126 Pisa, e-mail: gianni.bedini@unipi.it).

The correspondence must be sent to Editors (for Serie A: A. Chelli, Dipartimento di Chimica, Scienze della Vita e della Sostenibilità Ambientale, Parco Area delle Scienze, 157/A - 43124 Parma, email: alessandro.chelli@unipr.it; for Serie B: G. Bedini, Dipartimento di Biologia, via Derna, 1 - 56126 Pisa, e-mail: gianni.bedini@unipi.it).

Per ulteriori informazioni / *For further information:* <http://www.stsn.it/>

Per informazioni editoriali / *For editorial information:* Edizioni ETS - <https://www.edizioniets.com/>

SOCIETÀ TOSCANA DI SCIENZE NATURALI

Fondata nel (*founded in*) 1874

Via Santa Maria, 53 - 56126 Pisa

Consiglio Direttivo (*Executive Committee*)
(2023-2024)

Presidente

Vice Presidenti

Segretario generale

Segretari agli Atti (Editors)

Bibliotecario

Economo-Cassiere

P.R. Federici

W. Landini, M. Pappalardo

F. Rapetti

A. Chelli (Serie A),

G. Bedini (Serie B)

M. Zuffi

R. Narducci

Autorizzazione n. 17/56 del 26 luglio 1956, Trib. di Pisa

Direttore responsabile (Editor in Chief): Paolo Roberto Federici

© Copyright 2023-2024 by Società Toscana di Scienze Naturali.

Per tutti gli articoli pubblicati (articoli originali, comunicazioni brevi, notizie e recensioni) gli autori trasferiscono i diritti d'autore e di pubblicazione alla rivista.

For all published articles (original papers, short communications, news and book reviews) the authors transfer copyright and publishing rights to the Journal.

Tutti gli articoli pubblicati sugli Atti (articoli originali, comunicazioni brevi, notizie e recensioni) sono rilasciati con licenza CC BY 4.0. Essi possono essere condivisi e adattati, a condizione che venga dato credito adeguato, e possono essere utilizzati anche per scopi commerciali.

All published articles in Atti (original papers, short communications, news and book reviews) are released under CCL BY 4.0. They can be shared and adapted, provided that adequate credit is given, for any purpose, even commercially.

GIANCARLO MOLLI ⁽¹⁾

GEOLOGY OF THE ALPI APUANE AND THEIR MARBLES: AN UPDATE OVERVIEW

Abstract - G. MOLLI, *Geology of the Alpi Apuane and their marbles: An update overview*.

The Alpi Apuane are renowned globally for their marble deposits, notably the famed white Carrara marble, quarried there for over two millennia. The occurrence of marble within a region predominantly composed of non-metamorphic rock types has captivated geological interest since the early 19th century. Consequently, this area was the site of Italy's earliest modern geological maps and remains a crucial zone for structural and tectonic research. Following a general tectonic introduction, this study provides a current geological synthesis of the Alpi Apuane, emphasizing structures, tectono-metamorphic development, and the chronology of deformation events. Furthermore, the microstructural variability of the marbles will be illustrated and correlated with the different kinematic and dynamic stages of the wedge at the crustal scale.

Key words - Alpi Apuane, marbles, structural geology, tectonics, Apennines, Italy

Riassunto - G. MOLLI, *Geologia delle Alpi Apuane e dei loro marmi: una revisione aggiornata*.

Le Alpi Apuane sono rinomate a livello mondiale per i loro giacimenti di marmo, in particolare per il famoso marmo bianco di Carrara, estratto in quest'area da oltre due millenni. La presenza del marmo in una regione caratterizzata prevalentemente da rocce non metamorfiche ha suscitato interesse geologico fin dall'inizio del XIX secolo. Di conseguenza, le Alpi Apuane sono state oggetto delle prime carte geologiche moderne in Italia e da sempre rappresentano una zona cruciale per la ricerca geologico-strutturale e la comprensione dell'evoluzione tettonica della catena appenninica. Dopo un'introduzione tettonica regionale, questo studio fornisce una sintesi delle ricerche geologiche più recenti delle Alpi Apuane, sottolineando le strutture, lo sviluppo tettono-metamorfico e la cronologia degli eventi deformativi. Viene inoltre illustrata la variabilità microstrutturale dei marmi e correlata con le diverse fasi cinematiche e dinamiche di sviluppo della catena Appenninica.

Parole chiave - Alpi Apuane, marmi, geologia strutturale, tettonica, Appennino, Italia

INTRODUCTION AND REGIONAL TECTONIC FRAME

A unique morpho-structural domain exists in the northernmost portion of the Italian peninsula, extending from the Gulf of La Spezia to Monte Pisano (Federici & Raggi, 1975). This area, part of the inner northwestern Apennines (Fig. 1), encompasses the Alpi Apuane

a mountain range rising to approximately 2000 m near the coast (Baroni *et al.*, 2015). The Alpi Apuane has long intrigued naturalists due to its distinctive topography and the unusual presence of marble within an area of predominantly non-metamorphic rock-units.

The discovery of fossils (bivalvs and ammonites) at the beginning of '800 by Girolamo Guidoni (1794-1870), in the reliefs surroundings of La Spezia and in the Alpi Apuane, and their correct attribution to the Mesozoic (Secondary at that time) make this region special in the historical development of research and geological maps for the entire Apennines (Molli, 2023). As matter of facts the region represents at the end of the 19th century the first largest area of the entire Italian peninsula mapped at a scale of 1:25000 by Zaccagna and Lotti (1879-1897) in the frame of the first edition of the Italian geological map project.

These detailed maps provided the basis for cross-sections that supported the earliest nappe interpretations for the Apennines. In these models, developed by Lenciewicz (1917) and subsequently repropoed by Ippolito (1950) and Elter (1960), the Alpi Apuane play a key role being recognized as a tectonic window showing the lowermost structural units of the Apennines nappe stack.

Cross-section in Fig. 1 shows at a crustal scale the Alpi Apuane within the Northern Apennines tectonic frame. In the section it is possible to recognize a system of uppermost units (Ligurian and sub-Ligurian) representing the remnants of the pre-collisional accretionary wedge below which the distal Adria continental margin was underplated. In the internal zones of the range two main groups of Adria-derived units may be distinguished (Elter, 1975; Carmignani & Kligfield, 1990):

1. The Tuscan Nappe which consists of Mesozoic carbonate rocks and Tertiary deep-water and turbiditic sequences that are almost everywhere detached from their original basement (only locally recognized see Molli *et al.*, 2020) along the décollement levels of the former Carnian and Norian evaporites and dolostones of the "Calcare Cavernoso" (Conti *et al.*, 2020; Cornamusini *et al.*, 2024);

⁽¹⁾ Dipartimento Scienze della Terra, Università di Pisa, Via S. Maria, 53, Pisa 56126, Italy
Corresponding author: Giancarlo Molli (giancarlo.molli@unipi.it)

2. The Tuscan metamorphic units showing a Mesozoic–Tertiary metasedimentary sequences, like that of the Tuscan Nappe, which rest locally on a Permian cover or directly overlie a low-grade Variscan basement (Conti *et al.*, 2020). The peak metamorphic conditions and some differences in the Mesozoic to Tertiary metasedimentary sequences allowed to differentiate various metamorphic tectonic units (Molli 2023 and ref.).

THE STRATIGRAPHIC SEQUENCES AND THE MARBLE-TYPES

The lithostratigraphic successions observable in Alpi Apuane record the complete sedimentary evolution from breakup of Pangaea (Permian and Triassic) with rifting of the old Variscan crust to drifting (from late Jurassic onward) associated with basin deepening and widening of the Ligurian Tethys ocean. The inversion of the Europe-Adria plates kinematics brings during the Tertiary to the Apennines orogenesis (Elter, 1975; Bernoulli, 2001; Carmignani & Kligfield, 1990; Conti *et al.*, 2020 and references therein).

The exposed pre-Mesozoic basement observable in the Alpi Apuane (Massa and Apuane units) (Fig. 2) includes low-grade Variscan units (Conti *et al.*, 1991). These mainly contain albite-bearing chloritic phyllites and quartzites with lenses of mafic metavolcanics and calcareous schists (Lower Phyllites), felsic metavolcanics and metavolcanoclastic rocks (Porphyroids and porphyritic schists), metasandstones, quartzites

and phyllites (Upper Phyllites), graphitic schists and Orthoceras bearing dolostone. These units are considered as part of a Cambrian to Silurian succession, classically correlated with the Palaeozoic units of central-southern Sardinia (Conti *et al.*, 1991; Pandeli *et al.*, 1994). During Late Carboniferous/early Permian a regional scale trans-extensional regime (Rau, 1990; Molli *et al.*, 2020) affected the Adria crust. This history is recorded by the different Permian deposits recognized in the Massa unit (Patacca *et al.*, 2011; Conti *et al.*, 2020) and in the eastern Alpi Apuane (Fornovolasco area) and by the intrusions of early Permian subvolcanics felsic bodies which may be observed in Fornovolasco as well as in the Sant'Anna di Stazzema areas (Vezzoni *et al.*, 2018; Pieruccioni *et al.*, 2023). Evidence of a later crustal attenuation is provided by the Anisian-Ladinian extensional basin sequences in the Massa unit where marine platform sediments (Diplopora-bearing marbles) are associated with alkaline basaltic flows and breccias. This sequence well studied in the nearby area of Punta Bianca testifies a Mid-Triassic aborted rifting stage affected the future Adria-plate (Elter & Federici, 1964; Federici, 1966; Martini *et al.*, 1986; Storti, 1995).

The Upper Ladinian(?)–Carnian sedimentation is then witnessed by continental to marine deposits (Conti *et al.*, 2020) locally observable at the base of shallow water carbonates (Carnian (?)–Norian Dolomites of the “Grezzoni Fm.”) or dolomite-evaporites sequences (Calcare Cavernoso) formed during the renewal of the rifting process. Shallow water carbonate deposi-




Figure 1. Tectonic map of the Northern Apennines, with crustal scale (b) cross section (modified after Molli, 2008).




Figure 2. (a) Simplified geological-structural map and (b) composite cross sections across the Alpi Apuane. The dashed lines show the axial traces of the major D1 and D2 folds. Equal area lower hemisphere stereograms show the poles to the main foliation (Sp), stretching lineation (Lp), and the poles of late crenulation cleavage (St).

tion occurred all over the Tuscan continental margin (Calcare Massiccio platform) during the Rhaetian to early Liassic, locally interrupted by uplift, emersion and development of slope breccias (Rhaetian-Liassic boundary). Early to Middle Jurassic block faulting and progressive subsidence of the continental margin were associated with the dismembering of the carbonate platforms and the oceanization of the Ligurian Tethys (Malm) far west (Marroni *et al.*, 2017). Drowning of carbonate platform is then testified by Middle Jurassic-Cretaceous to Cenozoic pelagic carbonates and shales grading upward to Oligocene lower Miocene sandstones and shales (Pseudo-Macigno Fm).

The variability in the stratigraphy in the different areas of Alpi Apuane (Carmignani *et al.*, 1987; Coli & Fazzuoli, 1992; Conti *et al.*, 2020) relates to the syn-sedimentary extensional history started within Anisian-Ladinian as recorded in the main different stratigraphic sequences between Massa and Apuane units. The syn-sedimentary extensional history continued in the Carnian (Vinca Fm.'s facies variability, Fig. 3), and then in the Norian-Rhaetian (lagoonal metalimestones and black marbles "Nero di Colonnata" of the Western Apuane as opposite to a carbonate platform Megalodon-bearing marbles, Seravezza Breccia and Chloritoid Schists of the Central-Eastern Apuane).




Figure 3. (a) General stratigraphic sequence of the Apuane unit with some characteristic rock-types quarried as ornamental stones. In particular: b) Carnian impure marble part of the Vinca Fm. quarried as “Rosso Rubino”; (c) Breccia di Seravezza, Late Rhaetian scree-breccia also known with the historical name of “Breccia Medicea”; (d) “Carrara marble”: variety “Statuario” (upper part) and “Bardiglio” (grey marble) in the lowermost part; (e) marble breccia variety of “Arabescato”; (f) impure marble quarried with the name of “Zembrino”. This rock types occupy the stratigraphic position of the “Rosso Ammonitico” between Massiccio and Calcare Selcifero in the un-metamorphic Tuscan succession; (g) impure marble “Cipollino”; (h) Breccia Rossa, characteristic rock-units of the eastern Alpi Apuane, typical of “condensed” stratigraphic sequences; (i) Pseudomacigno Fm. quarried in the southernmost Alpi Apuane with the name of “Pietra del Cardoso”.

The Jurassic sequences well witness the presence of different paleoenvironments related to lagoons, tidal channels and emerged areas (White marble, Veined marble, Statuario) as well island-bounding scarps, documented by the Arabescato (Fig. 3e) and other marble breccias including paleokarsts remnants (Fig. 4a). Some other kinds of the marble breccias (Fig. 4b,c,d)

may be related to paleofractures forming Neptunian dykes and in some cases damage zones of paleofaults locally mineralized by basement-derived fluids (marble variety of the Paonazzo).

The demise and drowning of Jurassic carbonate platform toward deep-water condition of sedimentation is recorded in different but coeval lithostratigraphic




Figure 4. (a) Paleokarst and related marble breccias (Morlungho quarry); (b) Fault-related neptunian-dyke, Pianza quarry; (c) Veined marble interpreted as paleo-fault damage zone and related paleo-fault breccias; (d) marble variety of “Paonazzo” a breccia-variety characterized by Fe- and Mn-rich mineralized matrix.

terms explaining the variabilities in Late Jurassic-Tertiary sequences in the different parts of the Alpi Apuane giving way to the variable kinds of “Cipollino” and a red-greenish-coloured variety of marble breccia (Breccia Rossa in Carmignani *et al.*, 2007) (Fig. 3f,g). The end of history in the metasedimentary sequence exposed in the Alpi Apuane is recorded by the siliciclastic turbidites of the Pseudo-Macigno Fm. (Conti *et al.*, 2020) a fine to coarse grain metasandstone-metasilstones and black slates quarried as “Pietra del Cardoso” with the local merceological name (Fig. 3i).

D1 DEFORMATION AND THE DISTRIBUTION OF THE JURASSIC MARBLES

Map and cross-sections of Fig. 2 illustrate the large-scale structures in the Alpi Apuane metamorphic core which is characterized by kilometer- to hectometer-scale folds (main phase folds or D1) affecting the Paleozoic basement rocks and their Mesozoic to Cenozoic metasedimentary cover sequence (Carmignani *et al.*, 1985). In the northwestern part of the Alpi Apuane, the first-order structure is represented by the east-facing M. Brughiana antiform-anticline in the Massa unit. In its northernmost exposures, between Massa and Carrara, this structure has a core of Paleozoic basement and limbs exposing Ladinian-Carnian continental to marine sequences (Patacca *et al.*, 2011; Di Vincenzo *et al.*, 2022). Below the Massa unit, the Apuane unit shows two regional-scale structures, the Carrara syncline resting above a major east-facing fold, the Vinca-Forno anticline, with a core of Paleozoic basement. These two

regional-scale structures are separated by minor (hectometer-scale) antiform-synform pairs, the Pianza and Vallini folds, with stretched limbs and the core of Jurassic marbles and cherty metalimestone, respectively. Below the Vinca-Forno anticline a second kilometeric scale D1 isoclinal fold structure can be observed, the M. Tambura anticline, as the previous ones cored by Paleozoic basement rocks. The two major antiform-anticlines are separated by the Mesozoic metasediments of the Orto di Donna-M. Altissimo-M. Corchia synclines (Fig. 2). The M. Tambura anticline and the Orto di Donna-M. Altissimo synclines close to their southernmost exposures (Fig. 2) are characterized by a nearly 90° change in orientation of fold axes (Carmignani & Giglia, 1977; Carmignani *et al.*, 1978; Meccheri *et al.*, 2008; Molli & Meccheri, 2012), which defines the so-called “Virgazione di Arni” (Giglia, 1967) that will be discussed further. Below the M. Tambura anticline in the north-eastward part of the Alpi Apuane an high strain domain (Eastern Alpi Apuane) is characterized by tight to isoclinal folds (anticlines and synclines pairs) associated with pluridecameter-thick mylonitic zones affecting the whole sequence, from Jurassic-marble to Tertiary Pseudomacigno Fm. A main planar anisotropy (S1 foliation) of S-L type can be recognized across all the Alpi Apuane as the axial plane foliation of the isoclinal decimeter to kilometer scale folds (Fig. 5). This foliation has a composite origin (Carmignani & Kligfield, 1990; Molli & Meccheri, 2000) and bears a WSW-ENE trending mineral and extension lineation (Fig. 2) which appears to be parallel to the long axes of the stretched pebble clasts in marble breccias and in quartzitic metaconglomerates. D1 fold




Figure 5. (a) Panoramic view, looking north, from Foce di Giovo, northern Alpi Apuane. Two plurihctometer-scale folds in the hinge zone of the km-scale D1 Vinca-Forno Anticline are visible. D1 folds are refolded by D2 folds with a horizontal fold axial plane. Dark green Paleozoic schists making cusps within the metadolstones of the Grezzoni Fm.; (b) D2 west-vergent cascade-folds affecting the Grezzoni metadolstone in the normal limb of Vinca-Forno Anticline (Cima D'Uomo, Colonnata). (c) D1 meter-scale isoclinal fold and related axial planar foliation, overprint by D2 cleavage (Cherty metalimestone of Mt.Uccelliera); (d) meter-scale open D2 folds associated with a sub-horizontal crenulation cleavage (Scisti Sericitici Fm., Arni valley, eastern Alpi Apuane); (e) calc-mylonites derived from cherty metalimestones within a Late D1 shear zone. Colonnata valley, Carrara western Alpi Apuane; (f) decimeter-scale D2 shear zone in impure marble, Forno valley inland of Massa (for detailed microstructural description and fabric-types evolution see Oesterling *et al.*, 2007).

axes in the western area mainly trend NW-SE and are sub-horizontal (but see the area around Ponti di Vara, inland of Carrara) with a D1 lineation plunging down-dip within the main foliation at 90° from fold axis. In the north-eastern region, below the Mt. Tambura anticline, fold axes are parallel to sub-parallel to the down-dip stretching lineation and highly non-cylindric sheath folds appear (Carmignani & Giglia, 1984; Carmignani & Kligfield, 1990). This relationship between fold axis and stretching lineation has been proposed as an example of passive rotation of early formed folds into the extension direction in the higher strained domains during progressive simple shear (Kligfield *et al.*, 1981). The deformation geometries, strain patterns and kinematic data allowed to interpret the D1 history as the result of: (1) underthrusting and early nappe stacking within the Apenninic collisional wedge; (2) “antiformal stack phase” in which further shortening and a crustal scale duplex are realized (Carmignani & Kligfield, 1990; Molli *et al.*, 2000).

STRUCTURAL HERITAGE AND D1 DEFORMATION

The development of the large scale D1 structures is strongly controlled by the original paleotectonic setting and its lateral heterogeneities documenting the important role of structural inheritance (Molli & Meccheri, 2012). In the geological literature of the Alpi Apuane the concept of paleotectonic heritage can be traced back to Zaccagna (1897) and occurs throughout the literature until the '60s (e.g. Giglia, 1967; Nardi, 1969). More recently it was appreciated and illustrated by Carmignani *et al.* (1978), Kligfield (1979), Kligfield *et al.* (1981), Carmignani *et al.* (1987), Carmignani & Kligfield (1990) who clearly suggested a direct relationship between orogenic structures and Norian–Cretaceous paleotectonic ones. For these authors the first order folds of the Alpi Apuane (i.e. the “Vinca-Forno” and “Tambura” anticlines) were related to the geometry of the Mesozoic sedimentary basins and intervening highs of the external Tuscan domain (Elter, 1975). According to Carmignani *et al.* (1987) and

Carmignani & Kligfield (1990), the lateral and vertical thickness variabilities in the Mesozoic stratigraphy of the rifted Tuscan continental margin controlled the localization of the ramps in “prograde” ramp-flat tectonics, whose structures were successively modified at deeper structural levels in the metamorphic realm. This proposal has been slightly modified in the interpretation by Molli & Meccheri (2012), who applied a “modified overthrust model” proposed for the Helvetic nappes by Casey & Dietrich (1997) and used by Butler *et al.* (2006a; 2006b), Bellahsen *et al.* (2012) as key to understand inversion tectonic styles at mid-crustal depth. The interpretation considers the role of paleofault-related vertical thickness variabilities in the Mesozoic stratigraphy of the rifted Adria continental margin and the kinematic evolution of the investigated structures (Fig. 6). A localization of the low-amplitude initial folds by pre-existing fault domains may have served to create initial buckling instabilities associated with heterogeneous pure-shear basement deformation. During

progressive fold amplification, a modification of the initial buckle folds was produced by shearing with formation of the major regional folds later refolded during the anti-formal stacking (and D2 deformation see below) ending with the presently observable Alpi Apuane architecture (see Fig. 2, 6, 7). Molli & Meccheri (2012) underlined the point that, if the original buckling was controlled by the pattern of preexisting faults (Fig. 7d), the presence of fault segmentation and transfer zones might bring to a deviation of the velocity discontinuities with the frontal fold axis would become nearly parallel to the bulk shortening direction (Calassou *et al.*, 1993; Konstantinovskaya *et al.*, 2007) giving way to a transversal trend of folds. This was considered as the origin of the early stage of regional transversal trend of folding which would have been further modified by shearing leading to the antiapenninic “rotated” trends sub-parallel to the regional stretching lineation at kilometer-scale in the “Virgazione di Arni” (Carmignani & Giglia, 1983; Molli & Vaselli, 2006).




Figure 6. (a) General cross-section of the Alpi Apuane showing the regional-scale structures and their relationships with the original paleogeography modified after Carmignani *et al.* (1987) and Carmignani & Kligfield (1990); (b) White marble thrust above stratified impure pale greenish marble, scale bar ca. 2.5 m. Figured are details of thrust plane and meter-scale east-vergent folds developed in the footwall of the thrust; (c): c1 schematic line drawing showing the present-day geometry and lithology of the outcrop. Based on the local finite strain data and a simple shear strain path the original geometry can be reconstructed (c2); (d) schematic paleotectonic map view showing the possible segmentation of the former basin and transfer zones as well as their influence during the early “inversion”-related shortening; (e) regional tectonic frame with close up of vertical profiles and map views of early normal fault-controlled buckling during underthrusting of the external Tuscan Domain (Alpi Apuane zone) and later modification by shearing to produce the recumbent Alpi Apuane D1-fold structures.




Figure 7. (a) P-T estimates and paths for the Apuane and Massa units. Data used for boxes 1-4 are derived from quantitative classical thermo-barometry as well as pseudosections (Franceschelli *et al.*, 1986; Jolivet *et al.*, 1998; Molli *et al.*, 2000; Papeschi *et al.*, 2023). Data used to define box 4 also include temperatures based on Raman spectroscopy on carbonaceous material (RSCM) (Molli *et al.*, 2018). Red and blue lines outline the relative timing of deformation for D1 and D2 with changing metamorphic conditions. For more details see Di Vincenzo *et al.*, 2022; (b) Time-depth paths of the Massa and Apuane units and of the overlying Tuscan nappe (Macigno) modified after Fellin *et al.*, 2007 including data in Di Vincenzo *et al.*, 2022; (c) Paleothermal architecture of the Alpi Apuane metamorphic core complex in map view; (d) paleothermal architecture at window-scale in cross-sections view; (e) overall structural architecture of the Alpi Apuane metamorphic core. Also indicated and named the main boundary faults around the metamorphic core.

D2 DEFORMATION AND THE REWORKING OF THE EARLY STRUCTURES

All the D1 structures and tectonic contacts are overprinted by later structures of different generations and styles collectively referable to the regional D2 event. The early structures, syn-metamorphic, are represented by variously sized high strain zones and well-developed folds mainly associated with a low dipping to sub-horizontal axial planar foliation (S2), of crenulation type (Figs. 2, 5b,d). After these early generations of folds and shear zones, Late D2 structures formed. These are mainly represented by upright kinks and different generations of brittle faults (the earlier with a low angle attitude as opposite to the younger at a high angle) that accommodate the most recent tectonic history.

As documented since the end of '70s (Carmignani *et al.*, 1978; Carmignani & Giglia, 1979; Carmignani & Kligfield, 1990) a complex D2 mega-antiform with Apenninic trending axis (nearly N 130°-170°), and corresponding to the entire width of the Alpi

Apuane window, refold the early-formed D1 structures (Fig. 7). All around the antiform, second order asymmetric folds facing away from the dome crests are described. At the scale of the whole Alpi Apuane, large scale reverse drag-folds having "S" and "Z" sense of asymmetries can be observed on the southwestern and northeastern flanks, respectively. These structures form series of folds at different scale (from centimeters to kilometers) with variable morphologies related to rock competence and structural position within the folded multilayer but also from the characters of the previously developed D1 structures.

For this early D2 deformation, different and alternative interpretations were proposed in the literature.

A first model, frame the D2 history during a continuous shortening regime in which superimposed folding occurred (Carmignani *et al.*, 1978; Kligfield, 1979; Boccaletti *et al.*, 1983; Carosi *et al.*, 2002). In this model-groups may be also inserted the '90s proposal of Jolivet *et al.* (1998) which envisaged the south-west verging folds as related to local shear zones conjugate

to the major top-to-the-northeast thrusts, accommodating rotation of intra “Apuane” rigid domains within the major top-to-the-northeast contractional shearing. Differently, Carmignani & Kligfield (1990), proposed a kinematic model in which the D2 structures were related with conjugate top-to-the northeast and top-to-the southwest shear zones, in which cascade-type folds were produced within an overall extensional crustal kinematics.

As alternative to all the previously mentioned models, the one first proposed by Carmignani & Giglia (1979) and more recently re-proposed by Molli *et al.* (2022) based on strain fringes analyses, abides the only requirement necessary for the development of D2 deformation and strain features: a roughly vertical shortening of medium to steeply oriented layers. The vertical shortening-initiated buckle instabilities in more competent units and layer-parallel shearing on the limbs of previously developed Late D1 antiforms with an asymmetry related to the orientation of pre-existing composite layering and early isoclinal fold structures. This general architectural pattern is presently recognized as typical of region undergoing crustal scale extension/transtension (Froitzheim, 1992; Carosi *et al.*, 1996; Arden & Malavieille, 1999; Fossen *et al.*, 2013) and therefore may be considered as a key signature for geodynamic interpretations.

The Late D2 deformation history is then associated with cataclastic low-angle normal faults with the major structures localized by the reactivation of the window fault (Calcare Cavernoso Auct.) and their footwall splays, few hundreds of meters below the main fault (Conti *et al.*, 2020; Pieruccioni *et al.*, 2018).

This Late D2 low angle normal faults are overprinted by a further generation of structures related with deformation at shallower crustal depth. The Alpi Apuane are indeed an homogeneous regional scale brittle ‘low strain’ domain surrounded by large displacement faults (boundary faults) to the east, west, north and south. These high angle boundary faults (see Fig. 7e) divide the Alpi Apuane from the surroundings tectonic depressions of Lunigiana, Garfagnana, Versilia and Camaiore (Molli *et al.*, 2021 and references). Deformations associated with the boundary faults are witnessed by damage zones of variable width toward the external parts of the tectonic window and within the metamorphic core by an immature intra Alpi Apuane fault network. This fault network, characterized by a very low degree of interconnection, shows low displacement (from meter to pluridecameter) along every single structure. This is the reason why the brittle structures were neglected or missed in most of the available geological maps, although they are very important and significant for the quarrying activity, hydrogeo-

logy and karst network development (Piccini *et al.*, 2003; Cortopassi *et al.*, 2007; Isola *et al.*, 2021; Molli *et al.*, 2015).

THE METAMORPHIC HISTORY AND PALEOTHERMAL ARCHITECTURE

Regional scale differences in peak of metamorphism were envisaged in the Alpi Apuane since the mid of eighties (Di Pisa *et al.*, 1985; Franceschelli *et al.*, 1986). In metapelites of the Massa Unit metamorphic peak conditions (Fig. 7) are associated with kyanite + chloritoid + phengitic muscovite assemblages and related to ~1.0 GPa and 420-500°C (Franceschelli *et al.*, 1986; Jolivet *et al.*, 1998; Molli *et al.*, 2002; Di Vincenzo *et al.*, 2022; Papeschi *et al.*, 2023). In the Apuane Unit the peak assemblages of pyrophyllite + chloritoid + chlorite + phengitic muscovite in metapelites indicate peak metamorphism in the pressure range of 0.4-0.6 GPa and temperatures ranging between 350 and 450°C in different structural positions (Franceschelli *et al.*, 1986; Jolivet *et al.*, 1998; Molli *et al.*, 2002).

RSCM thermometry allowed to refine the T distribution at the scale of the entire region in the different pre-Mesozoic rock-units and in Mesozoic to Tertiary ones. The pre-Mesozoic basement rocks show a mean peak RSCM T of 445 °C whereas a mean of 386 °C is found in the metasedimentary Mesozoic to Tertiary cover (Molli *et al.*, 2018). The distribution of peak T at the scale of the whole Alpi Apuane (Fig. 7c) indicates a systematic arrangement with respect to the overall architecture of the regional structures. The north-west part of the Alpi Apuane is characterized by a SW to NE decrease of peak temperature from 495 °C in the west to 357 °C in the central and 336 °C in the easternmost position. A difference of 125 °C is observed with a structural distance of ~8 km, outlying an inverted field metamorphism with a T gradient of ~20°/km. The thermal architecture combined with the structural ones indicate an overall “contractional” model of antiformal stacking suggesting an orogenic wedge dominated by material advection realized by underplating of more external units (Molli *et al.*, 2018). In the south-east part of the Alpi Apuane (Fig. 7c) an “extensional” core-complex type thermal architecture and an apparent normal-type gradient is observed with transition from temperatures higher than 500 °C to less than 300 °C (projected base of the Tuscan Nappe) in less than 1,5 km (e.g. cross-section G). This thermal architecture, however, could be only apparent since the exposed geometrically deepest and cooler levels represented in the eastern Alpi Apuane are not at the surface due to the 3D dome shape structural architecture previously illustrated.

AGE OF DEFORMATION AND EXHUMATION HISTORY

Based on the geochronological data (Kligfield *et al.*, 1986; Di Vincenzo *et al.*, 2022), the available thermochronology (Abbate *et al.*, 1994; Fellin *et al.*, 2007) and geological constrains (Carmignani & Kligfield, 1990; Molli *et al.*, 2002; Molli *et al.*, 2021) the following deformation age and exhumation history may be reconstructed (Di Vincenzo *et al.*, 2022):

1. onset of metamorphism and deformation in the exposed units of the Alpi Apuane region may be constrained as having a minimum age of 20-21 Ma (Aquitainian/Burdigalian boundary), as documented by ^{40}Ar - ^{39}Ar data for the composite D1 foliation and related structures in the western parts of the Alpi Apuane (i.e., Massa unit). This is in line with the recently acquired data in the close area of Punta Bianca (Montomoli *et al.*, 2024) as well as at regional scale in the Adria-derived Tuscan metamorphic units in Southern Tuscany (Bianco *et al.*, 2015; Giuntoli & Viola, 2024).
2. D1 deformation, consistently with its composite character, occurred during a protracted (at least ~6.5 Ma) metamorphic evolution, lasting from the Aquitainian/Burdigalian to the Langhian/Serravallian boundary. A possible West-East diachronism for the development of D1 foliation is suggested by the minimum age of ~14.4 Ma obtained for the development of the D1 foliation in the easternmost sample of the Apuane Unit (Arni valley). Such a diachronism in the development of D1 structures appear to be significant from a wedge kinematic viewpoint being consistent with the West-East sense of transport, the stacking of the various units, and their internal deformation at the scale of the whole metamorphic core.
3. D2 deformation shows a common temporal evolution at the scale of the entire Alpi Apuane region and is recorded for a duration of at least ~2.5 Ma, starting not after the upper Serravallian and ending in the middle Tortonian, at ~10.5 Ma.

Interpreting the early D2 deformation structures, with their architectures and kinematics features as marking the switch from crustal contraction to crustal extension (Carmignani & Kligfield, 1990; Molli *et al.*, 2018), point (3) constraints the onset of the genuine lithospheric scale extensional history in this portion of the inner northern Apennines as developed since the upper Serravallian (~12.5-12.0 Ma) and recorded until c. 10 Ma (youngest Ar/Ar ages associated to the D2 fabrics in the Massa unit Di Vincenzo *et al.*, 2022). The available thermochronological data in Abbate *et al.* (1994) and Fellin *et al.* (2007) allowed to complete the timing of deformation and exhumation from the middle-upper crust (temperature c. 300 °C) up to the surface (Fig. 7a,b). The data regards the thermal evolution of the Tuscan Nappe in its uppermost stratigraphic

term of Macigno Fm. (the only lithotype useful for fission tracks dating) and for the metamorphic units the Palaeozoic volcanic-volcanoclastic of Porphyroids and Porphyric Schists as well as the siliciclastic rock-units of the Pseudomacigno Fm. The data show the vertical path of the tectonic units, analysed through the temperature ranges of 250, 200, 120, 70 °C (respectively the closure temperatures for fission tracks in the Zr, He-Zr, Ap, He-Ap; Fellin *et al.*, 2007).

Using the available cooling ages data sets, the time-depth path in Fig. 7b may be reconstructed. This suggests that after the first stage of exhumation (pre-7 Ma), c. 3.6 ± 0.5 km of tectonic unroofing of the Alpi Apuane core occurred in a narrow time interval between approximately 7 and 5 Ma (Upper Tortonian-Messinian). After this, the Alpi Apuane core and the Tuscan nappe were jointly exhumed as a single body. This occurred by the activity of boundary faults and erosion. The presence of metamorphic pebbles from the Alpi Apuane core in the early Pliocene Garfagnana, and Lunigiana intramontane basins (Molli *et al.*, 2021 and references) indicate that at c. 2.5 Ma the Alpi Apuane core rocks were at the surface in agreement with the U/Pb radiometric age of speleothems (Isola *et al.*, 2021).

MARBLE MICROSTRUCTURES AS RECORD OF THE TECTONIC HISTORY OF THE INNER NA

After the pioneering studies reported in Zaccagna (1932) and Bonatti (1938) the microstructures of Alpi Apuane marbles was the object of investigations by Crisci *et al.*, (1975), Di Pisa *et al.*, (1985) and Coli (1989a). Considering the main microstructural features and relationships with mesoscopic field structures (foliations, folds and shear zones), the marbles microstructures were subdivided by Molli *et al.*, (2000a) into three main group-types (Fig. 8) interpreted respectively product of:

- a) static recrystallization (type A microfabric);
- b) dynamic recrystallization (type B microfabric, further subdivided into two types B1 and B2);
- c) reworking during the late stage of deformation (type C microfabric).

These distinctions represent the end-member of a wide range of transitional types which in some cases can be observed superimposing each other (see detailed description in Molli & Heilbronner, 1999; Molli *et al.*, 2000; Oesterling *et al.*, 2007).

- The statically recrystallized microfabric or “annealed-type” (type-A microfabric) is characterized by equant polygonal grains (granoblastic or “foam” microstructure, Fig. 8a,b,c), with straight to slightly curved grain boundaries that meet in triple points at angles of nearly 120°. C-axis orientations show a random distribution or a weak crystallographic preferred orientation. These microfabrics are




Figure 8. Marbles microstructures a,b,c: Statically recrystallized or annealed microstructure (Type A). Sample (a) from the Pania unit, southern Alpi Apuane, structural domain characterized by RSCM temperature of 320 °C; sample (b) from the eastern Alpi Apuane, domain characterized by RSCM of 360 °C; sample (c) from the western Alpi Apuane, domain characterized RSCM of 480 °C. Dynamic recrystallized microstructures (Type B): (d) sample coming from a millimeter-scale rim tapering the main slip surface of an High Angle Normal Fault (Late D2 history); (e) Type B2-microstructure related to D2 shear zones, localized deformation during the switch from crustal-scale contraction to crustal-scale extension (early D2); (f) Type B1-microstructure related to a Late D1 shear zone, localized deformation during the antiformal stacking.

observable in marble levels belonging to km-scale D1 isoclinal folds, where also minor parasitic folds developed. The presence of such microstructures within D1 folds indicates that the grain growth which produced type A microfabric occurred after the main D1 folding phase and obliterated all earlier syntectonic microstructures associated with folding. However, the presence of a texture in some samples of impure marbles may be related to the pre-annealing deformation history (Leiss & Molli, 2003). Marbles with annealed microstructure can be observed in the western, central and eastern parts of the Alpi Apuane, with a medium grain size decreasing from west to east (300-150 μm to 100-80 μm) and from geometrically deeper to higher structural levels in well agreement with the paleothermal architecture (Fig. 8a,b,c).

- The dynamically recrystallized microfabrics (type-B microfabrics) were differentiated in two main groups, the first one (type B1) exhibiting strong shape preferred orientation, coarse grains and lobate grain boundaries whereas the second show microstructures with shape preferred orientation, smaller grain size and predominantly straight grain boundaries (type B2). Fig. 8e,f shows repre-

sentative examples of the two types of microfabrics. These two types of microstructures are both interpreted as related to high strain and high temperature (300-400 °C) crystal plastic deformation mechanisms (dislocation creep regime). Whereas grain boundary migration recrystallization can be considered as predominant in type B1 microfabric, an important contribution of both rotation recrystallization and grain boundary migration has been inferred to prevail in type B2 microfabric (Molli *et al.*, 2000; Oesterling *et al.*, 2007).

- Twinned microfabric (type-C microfabric). The third type of microfabric is related to low-strain and low-temperature crystal plastic deformation mechanisms. Characterized by thin straight e-twins, it occurs in all the marble outcrops of the Alpi Apuane region, overprinting both type A and type B microfabrics. It is mostly developed in coarse grained marble. This microstructural-type has been also found associated to the marbles involved in the damage zones of late high angle normal to oblique slip faults where are associated to extremely localized (millimeter-scale) recrystallized microstructures (Fig. 8d) tapering their principal slip surfaces (Molli *et al.*, 2011; Molli, 2016).

FINAL REMARKS

The Alpi Apuane represent one of the key areas for understanding the geologic evolution of the Northern Apennines as well documented by the literature of the last of two centuries. During the 1970's the area was the ground of the first applications of modern structural geology with the recognition of superimposed foliations and structures (Carmignani & Giglia 1975; Pertusati *et al.*, 1977; Kligfield, 1979; Boccaletti & Gosso, 1980; Boccaletti *et al.*, 1983), stretching fabrics and sheath folds (Carmignani *et al.*, 1978) and the studies of finite strain (Kligfield *et al.*, 1981). Moreover in the Alpi Apuane the first pioneering geochronological study by Giglia & Radicati di Brozolo (1970) was realized and after that one of the first world-wide attempts of radiometric dating of superimposed low-grade foliations (Kligfield *et al.*, 1986). During '90s the papers of Coli (1989), and above all, Carmignani & Kligfield (1990) underlined and linked the late stage of syn-metamorphic history of the Alpi Apuane to the extensional geodynamics characteristic of the Apennines-Tyrrhenian system. This make the Alpi Apuane a key and unique area of the Apennines where it is possible to track and document how the Adria continental crust was deformed at depth during the phases of continental subduction, but also how it record of the different stages of the syn- and post-collisional extension from the depth of the middle crust (c. 10-15 km) up to the surface. As matter of facts this complete cycle of the Apennines orogenic evolution may be also traced and followed by the microstructural variability of the Alpi Apuane marbles which document, as successive footprints, the different kinematics and dynamic stages of the Apennines wedge at the crustal scale.

ACKNOWLEDGEMENTS

I thank P.R. Federici for inviting me to contribute to this 150° Anniversary of the Società Toscana di Scienze Naturali. Paolo Conti and Fabrizio Storti are acknowledged for their revisions and suggestions as well as Alessandro Chelli for the editorial handling. A special thanks to Luigi Carmignani, Roy Kligfield, Paolo Conti and the late Marco Meccheri for the discussions and shared knowledge of the Alpi Apuane's geology.

CONFLICT OF INTEREST STATEMENT

The author declares that he has no conflict of interest neither known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

REFERENCES

- ABBATE E., BALESTRIERI M.L., BIGAZZI G., NORELLI P., QUERCIOLE C., 1994. *Fission-track datings and recent rapid denudation in northern Apennines, Italy*. Memorie Società Geologica Italiana 48: 579-585.
- ARDEEN D.G.A.M., MALAVIEILLE J., 1999. *Origin of a large-scale fold nappe in the Montagne Noire Variscan belt, France*. Journal of Structural Geology 21: 1321-1333.
- BARONI C., PIERUCCINI P., BINI M., COLTORTI M., FANTOZZI P.L., GUIDOBALDI G., NANNINI D., RIBOLINI A., SALVATORE M.C., 2015. *Geomorphological and Neotectonic Map of the Apuan Alps (Tuscany, Italy)*. Geografia Fisica e Dinamica del Quaternario 38 (2): 201-227.
- BELLAHSEN N., JOLIVET L., LACOMBE O., BELLANGER M., BOU-TOUX A., GARCIA S., MOUTHEREAU F., LE POURHIET L., GUMIAUX C., 2012. *Mechanisms of margin inversion in the external Western Alps: Implications for crustal rheology*. Tectonophysics 560-561: 62-83.
- BERNOULLI D., 2001. *Mesozoic-Tertiary carbonate platforms, slopes and basins of the external Apennines and Sicily*. In: G.B. Vai, I.P. Martini (eds), *Anatomy of an Orogen*: 307-326. Kluwer, Dordrecht-Boston-London.
- BIANCO C., BROGI A., CAGGIANELLI A., GIORGETTI G., LIOTTA D., MECCHERI M., 2015. *HP-LT metamorphism in Elba Island: Implications for the geodynamic evolution of the inner northern Apennines (Italy)*. Journal of Geodynamics 91: 13-25. doi: 10.1016/j.jog.2015.08.001
- BOCCALETTI M., GOSSO G., 1980. *Analisi della deformazione plicativa e rapporti con lo sviluppo della blastesi metamorfica nell'area di Campo Cecina-M. Pisanino delle Alpi Apuane Settentrionali*. Memorie della Società Geologica Italiana 21: 101-110.
- BOCCALETTI M., CAPITANI S., COLI M., FORNACE G., GOSSO G., GRANDINI G., MILANO P.F., MORATTI G., NAFISSI P., SANI F., 1983. *Caratteristiche deformative delle Alpi Apuane settentrionali*. Memorie Società Geologica Italiana 26: 527-534.
- BOCCALETTI M., GOSSO G., MORATTI G., 1981. *Indizi di paleocarsismo nel Marmo delle Alpi Apuane Settentrionale*. Rendiconti della Società Geologica Italiana 4: 315-316.
- BONATTI S., 1938. *Studio petrografico delle Alpi Apuane*. Memorie Descrittive della Carta Geologica d'Italia 26: 116.
- BUTLER R.W.H., TAVARNELLI E., GRASSO M., 2006. *Structural inheritance in mountain belts: An Alpine-Apennine perspective*. Journal of Structural Geology 28: 1893-1908.
- BUTLER R.W.H., MAZZOLI S., 2006. *Styles of continental contraction: A review and introduction*. Styles of Continental Contraction. Geological Society America Spec. Pap. 414: 1-10.
- CALASSOU S., LARROQUE C., MALAVIEILLE J., 1993. *Transfer zones of deformation in thrust wedges: an experimental study*. Tectonophysics 221: 325-344.
- CARMIGNANI L., CONTI P., FANTOZZI P., MANCINI S., MASSA G., MOLLI G., VASELLI L., 2007. *I marmi delle Alpi Apuane*. Geoitalia 21: 19-30.
- CARMIGNANI L., GATTIGLIO M., KALIN O., MECCHERI M., 1987. *Guida all'escursione sul Complesso Metamorfico delle Alpi Apuane*. Escursione conclusiva della "Summer School" di Geologia e Petrologia dei Basamenti Cristallini", settembre 1987. CNR - Università di Siena, Tipografia Editrice Pisana, Pisa.
- CARMIGNANI L., GIGLIA G., 1975. *Le fasi tettoniche terziarie dell'Autoctono Apuano: studio delle strutture minori della zona centro-meridionale*. Bollettino della Società Geologica Italiana 94: 1957-1981.
- CARMIGNANI L., GIGLIA G., 1979. *Large scale reverse "drag folds" in the late Alpine building of the Apuane Alps (N. Apennines)*. Atti della Società Toscana di Scienze Naturali, Memorie, Serie A 86: 109-126.

- CARMIGNANI L., GIGLIA G., 1983. *Il problema della doppia vergenza sulle Alpi Apuane e la struttura del monte Corchia*. Memorie della Società Geologica Italiana 26: 515-525.
- CARMIGNANI L., GIGLIA G., 1984. 'Autoctono Apuano' e Falda Toscana: sintesi dei dati e interpretazioni più recenti. In: Cento anni di Geologia Italiana. Volume Giubilare Società Geologica Italiana, Roma, 199-214.
- CARMIGNANI L., GIGLIA G.G. 1977. *Analisi mesostrutturale della zona occidentale delle Apuane metamorfiche*. Bollettino della Società Geologica Italiana 96 (3): 429-450.
- CARMIGNANI L., GIGLIA G., KLIGFIELD R., MECCHERI M., 1985. *Carta Geologico-Strutturale del Complesso Metamorfico delle Alpi Apuane, Foglio Nord*. Scala 1:25.000, Dipartimento Scienze della Terra Università di Pisa, Litografia Artistica Cartografica, Firenze.
- CARMIGNANI L., GIGLIA G., KLIGFIELD R., 1978. *Structural evolution of the Apuane Alps: An example of continental margin deformation in the Northern Apennine*. Journal of Geology 86: 487-504. <https://doi.org/10.1086/649714>
- CARMIGNANI L., KLIGFIELD R., 1990. *Crustal extension in the northern Apennines: The transition from compression to extension in the Alpi Apuane core complex*. Tectonics 9: 1275-1303. <https://doi.org/10.1029/TC009i006p01275>
- CAROSI R., CERBAI N., MONTOMOLI C., 1996. *The F2 folds in the Verrucano as record of extensional tectonics in the Northern Apennines (Italy)*. Comptes Rendu Académies des Sciences Paris (Serie IIA) 322: 773-780.
- CAROSI R., MONTOMOLI C., PERTUSATI P.C., 2002. *Late orogenic structures and orogen-parallel compression in the Northern Apennines*. Bollettino Società Geologica Italiana Volume Speciale 1: 167-180.
- CASEY M., DIETRICH D., 1997. *Overthrust shear in mountain building*. In: S. Segupta (ed.), Evolution of geological structures in micro- to macroscales: 119-142. Chapman and Hall, London.
- COLI M., 1989. *Litho-structural assemblage and deformation history of "Carrara marble"*. Bollettino della Società Geologica Italiana 108: 581-590.
- COLI M., 1989. *Time and mode of uplift of the Apuane Alps metamorphic complex*. Atti Ticinensi Scienze Terra 32: 47-56.
- COLI M., FAZZUOLI M., 1992. *Evoluzione sedimentaria dell'area Apuana nel Triassico-Liassico*. Studi Geologici Camerti Volume Speciale CROP1-1a: 193-201.
- CONTI P., CARMIGNANI L., MASSA G., MECCHERI M., PATACCA E., SCANDONE P., PIERUCCIONI D., 2020. *Note Illustrative CARG Foglio 249*. ISPRA, Massa.
- CONTI P., GATTIGLIO M., MECCHERI M. 1991. *The overprint of the Alpine tectonometamorphic evolution on the Hercynian orogen: An example from the Apuane Alps (Northern Apennines, Italy)*. Tectonophysics 191: 335-346.
- CORNAMUSINI G., MILANESCHI L., CONTI P., LIBERATO G.P., 2024. *Significance of the Calcare Cavernoso: Stratigraphic-structural setting and role as tectonic chaotic unit in the evolution of the Northern Apennines (Italy)*. Italian Journal Geoscience 143 (2): 187-209. <https://doi.org/10.3301/IJG.2024.17>
- CORTOPASSI A., MOLLI G., OTTRIA G., 2007. *Studio della deformazione fragile nel bacino marmifero di Fantiscritti (Alpi Apuane, Carrara) finalizzato alla ricostruzione del campo di paleostress*. Geologia tecnica ed ambientale 1-2: 27-45.
- CRISCI G.M., LEONI L., SBRANA A., 1975. *La formazione dei marmi delle Alpi Apuane (Toscana): studio petrografico, mineralogico e chimico*. Atti della Società Toscana di Scienze Naturali, Memorie, Serie A 82: 199-236.
- DI PISA A., FRANCESCHELLI M., LEONI L., MECCHERI M., 1985. *Regional variation of the metamorphic temperatures across the Tuscanid I Unit and its implications on the alpine metamorphism (Apuan Alps, N-Tuscany)*. Neues Jahrbuch für Mineralogie, Abhandlungen 151 (2): 197-211.
- DI VINCENZO G., GODARD G., MOLLI G., 2022. *Dating low-grade deformation: role of lithology and strain partitioning on Ar isotope records in the Alpi Apuane of northern Apennines (Italy)*. Tectonics 41: 1-25. <https://doi.org/10.1029/2022TC007248>
- ELTER P., 1960. *I lineamenti tettonici dell'Appennino a nord-ovest delle Apuane*. Bollettino della Società Geologica Italiana 79: 273-312.
- ELTER P., 1975. *Introduction à la géologie de l'Apennin septentrional*. Bulletin de la Société Géologique de France 7: 956-962. <https://doi.org/10.2113/gssgfbull.S7-XVII.6.956>
- ELTER P., FEDERICI P.R., 1964. *Sulla presenza di terreni attribuibili al Trias medio nel promontorio orientale del Golfo della Spezia*. Bollettino della Società Geologica Italiana 83: 395-4022.
- FEDERICI P.R., 1966. *Gasteropodi ladinici nei Calcari di Punta Bianca (Golfo della Spezia)*. Rivista Italiana di Paleontologia e Stratigrafia 72 (4): 994-1022.
- FEDERICI P.R., RAGGI G., 1975. *Una nuova interpretazione della tettonica dei monti della Spezia*. Bollettino della Società Geologica Italiana 94: 945-960.
- FELLIN M.G., REINERS P.W., BRANDON M.T., WUTHRICH E., BALESTRIERI M.L., 2007. *Thermochronologic evidence for exhumational history of the Alpi Apuane metamorphic core complex, northern Apennines, Italy*. Tectonics 26: TC6015.
- FOSSEN H., TEYSSIER C., WHITNEY D.L., 2013. *Transtensional folds*. Journal of Structural Geology 56: 89-102.
- FRANCESCHELLI M., LEONI L., MEMMI M., PUXEDDU M., 1986. *Regional distribution of Al-silicates and metamorphic zonation in the low-grade Verrucano metasediments from the Northern Apennines, Italy*. Journal of Metamorphic Geology 4: 309-321.
- FROITZHEIM N., 1992. *Formation of recumbent folds during syn-orogenic crustal extension (Austroalpine nappes, Switzerland)*. Geology 20: 923-926.
- GIGLIA G., RADICATI DI BROZOLO F., 1970. *K/Ar age of metamorphism in the Apuane Alps (Northern Tuscany)*. Bollettino della Società Geologica Italiana 89: 485-497.
- GIGLIA G., 1967. *Geologia dell'Alta Versilia settentrionale (Tav. M. Altissimo)*. Memorie Società Geologica Italiana 6: 67-95.
- GIUNTOLI F., VIOLA G., VILLA I.M., 2024. *Dating syn-orogenic exhumation of subducted continental crust: The case of the Northern Apennines*. Lithos: 488-489. <https://doi.org/10.1016/j.lithos.2024.107801>
- IPPOLITO F. 1950. *Le Alpi Apuane*. Memorie e Note dell'Istituto di Geologia Applicata dell'Università di Napoli 3: 49-95.
- ISOLA I., MAZZARINI F., MOLLI G., PICCINI L., ZANELLA E., ZANCETTA G., DRYSDALE R., HELLSTROM J., WOODHEAD J., RONCONI A., MILAZZO F., PIERUCCIONI P., REGATTIERI E., 2021. *New chronological constraints from hypogean deposits for Late Pliocene to recent morphotectonic history of the Alpi Apuane (NW Tuscany, Italy)*. Geosciences 11: 65. <https://doi.org/10.3390/geosciences11020065>

- JOLIVET L., FACCENNA C., GOFFÉ B., MATTEI M., ROSSETTI F., BRUNET C., STORTI F., FUNICIELLO R., CADET J.P., D'AGOSTINO N., PARRA T., 1998. *Midcrustal shear zones in postorogenic extension: Example from the northern Tyrrhenian Sea*. Journal of Geophysical Research 103:12,123-12,160.
- KLIGFIELD R., 1979. *The Northern Apennine as a collisional orogen*. American Journal Science 279: 679-681.
- KLIGFIELD R., CARMIGNANI L., OWENS W.H., 1981. *Strain analysis of a Northern Apennine shear zone using deformed marble breccias*. Journal of Structural Geology 3: 421-436. [https://doi.org/10.1016/0191-8141\(81\)90042-0](https://doi.org/10.1016/0191-8141(81)90042-0)
- KLIGFIELD R., HUNZIKER J., DALLMEYER R.D., SCHAMEL S., 1986. *Dating of deformational phases using K-Ar and $^{40}\text{Ar}/^{39}\text{Ar}$ techniques: results from the Northern Apennines*. Journal of Structural Geology 8: 781-798.
- KONSTANTINOVSKAYA E.A., HARRIS L.B., POULIN J., IVANOV G.M., 2007. *Transfer zones and fault reactivation in inverted rift basins: Insights from physical modeling*. Tectonophysics 441: 1-26.
- LEISS B., MOLLI G., 2003. *"High-temperature" texture in naturally deformed Carrara marble from the Alpi Apuane, Italy*. Journal of Structural Geology 25: 649-658.
- LENCEWICZ S., 1917. *Profilo geologiczne przez Apenin Toskanski (Sezioni geologiche dell'Appennino Toscano)*. Rendiconti Società di Scienze di Varsavia 10: 637-662.
- MARRONI M., MENEGHINI F., PANDOLFI L., 2017. *A revised subduction inception model to explain the Late Cretaceous, double vergent orogen in the precollisional Western Tethys: Evidence from the Northern Apennines*. Tectonics 36: 2227-2249. <https://doi.org/10.1002/2017TC004627>
- MARTINI I.P., RAU A., TONGIORGI M., 1986. *Syntectonic sedimentation in a Middle triassic Rift, Northern Apennines, Italy*. Sedimentary Geology 47: 191-219. [https://doi.org/10.1016/0037-0738\(86\)90084-9](https://doi.org/10.1016/0037-0738(86)90084-9)
- MECCHERI M., MOLLI G., VASELLI L., CONTI P., 2008. *La struttura polideformata delle Alpi Apuane centrali tra Arni e M. Corchia: conoscenze storiche, dati attuali e problemi aperti*. Rendiconti Online Società Geologica Italiana, 1 Note Brevi, www.socgeol.it: 112-117.
- MOLLI G., 2008. *Northern Apennine-Corsica orogenic system: An updated review*. In: S. Siegesmund, B. Fügenschuh, N. Froitzheim (eds), Tectonic Aspects of the Alpine-Dinaride-Carpathian System. Geological Society of London Special Publication 298: 413-442. <https://doi.org/10.1144/SP298.19>
- MOLLI G., 2016. *Switching deformation mode and mechanisms during natural faulting of Carrara marble*. Rendiconti Online Società Geologica Italiana, Suppl. n. 1 al Vol. 40. 88° Congresso SGI, Napoli 2016
- MOLLI G., 2023. *The Alpi Apuane and their surroundings: A tale of the origins of modern Italian geological maps and of a misused 'early recognition' of nappes in the Apennines*. In: R.W.H. Butler, T. Torvela, L. Williams (eds) Geological Mapping of Our World and Others. Geological Society, London, Special Publications 541. <https://doi.org/10.1144/SP541-2022-273>
- MOLLI G., BROGI A., CAGGIANELLI A., CAPEZZUOLI E., LIOTTA D., SPINA A., ZIBRA I. 2020. *Late Palaeozoic tectonics in Central Mediterranean: a reappraisal*. Swiss Journal of Geosciences 113: 23. <https://doi.org/10.1186/s00015-020-00375-1>
- MOLLI G., CONTI P., GIORGETTI G. MECCHERI M., OESTERLING N., 2000. *Microfabric studies on the deformation and thermal history of the Alpi Apuane marbles (Carrara marbles), Italy*. Journal of Structural Geology 22 (11-12): 1809-1825. [https://doi.org/10.1016/S0191-8141\(00\)00086-9](https://doi.org/10.1016/S0191-8141(00)00086-9)
- MOLLI G., DOVERI M., MANZELLA A., BONINI L., BOTTI F., MENICINI M., MONTANARI D., TRUMPY E., UNGARI A., VASELLI L., 2015. *Surface-subsurface structural architecture and groundwater flow of the Equi Terme hydrothermal area, Northern Tuscany Italy*. Italian Journal Geosciences 134 3: 442-457. doi: 10.3301/IJG.2014.25.
- MOLLI G., GIORGETTI G., MECCHERI M., 2002. *Tectono-metamorphic evolution of the Alpi Apuane Metamorphic Complex: New data and constraints for geodynamic models*. Bollettino della Società Geologica Italiana 1: 789-800.
- MOLLI G., HEILBRONNER P.R., 1999. *Microstructures associated with static and dynamic recrystallization of Carrara marble (Alpi Apuane, NW Tuscany Italy)*. Geologie en Mijnbouw 78: 119-126.
- MOLLI G., KOEHN D., ALLACORTA L., DANESI L., ZAMPELLI S., ZAZZERI M., 2022. *Superimposed structures, incremental strain and deformation path from field data to modelling: A case study from the Alpi Apuane metamorphic core complex (NW Tuscany, Italy)*. Journal of Structural Geology 161: 200-216. <https://doi.org/10.1016/j.jsg.2022.104676>
- MOLLI G., MANIGHETTI I., BENNETT R., MALAVIEILLE J., SERPELONI E., STORTI F., GIAMPIETRO T., BIGOT A., PINELLI G., GIACOMELLI S., LUCCA A., ANGELI L., PORTA L., 2021. *Active fault systems in the inner northwest Apennines, Italy: a reappraisal one century after the 1920 Mw ~6.5 Fivizzano earthquake*. Geosciences 11: 139. <https://doi.org/10.3390/geosciences11030139>.
- MOLLI G., MECCHERI M., 2000. *Deformazione polifasica e composta nelle Alpi Apuane (NW Toscana)*. Bollettino della Società Geologica Italiana 119: 379-394.
- MOLLI G., MECCHERI M., 2012. *Structural inheritance and style of reactivation at mid-crustal levels: a case study from the Alpi Apuane (Tuscany, Italy)*. Tectonophysics 579: 74-87. <https://doi.org/10.1016/j.tecto.2012.06.044>.
- MOLLI G., VASELLI L., 2006. *Structures, interference patterns and strain regime during mid-crustal deformation in the Alpi Apuane (Northern Apennines, Italy)*. Geological Society of America Special Paper, 414: 79-94.
- MOLLI G., VITALE BROVARONE A., BEYSSAC O., CINQUINI I., 2018. *RSCM thermometry in the Alpi Apuane (NW Tuscany, Italy): New constraints for the metamorphic and tectonic history of the inner northern Apennines*. Journal of Structural Geology 113: 2000-2016.
- MONTOMOLI C., IACCARINO S., DI VINCENZO G., LANARI P., PETROCCHIA A., CAROSI R., 2024. *Deformation and pressure-temperature-time history of the External Tuscan Units in the Northern Apennines (Italy): The case of the Punta Bianca Unit*. Journal of Geodynamics 161: 102045. <https://doi.org/10.1016/j.jog.2024.102045>
- NARDI R., 1969. *Schema stratigrafico e strutturale delle Apuane Nord-orientali*. Bollettino della Società Geologica Italiana 82: 1-18.
- OESTERLING N., HEILBRONNER R., STUNITZ H., BARNHOORN A., MOLLI G., 2007. *Strain dependent variation of microstructure and texture in naturally deformed Carrara marble*. Journal of Structural Geology 29: 681-696. <https://doi.org/10.1016/j.jsg.2006.10.007>
- PANDELI E., GIANELLI G., PUXEDDU M., ELTER F.M., 1994. *The Palaeozoic basement of the Northern Apennines: Stratigraphy, tectono-metamorphic evolution and alpine hydrothermal process*. Memorie Società Geologica Italiana 48: 627-654.

- PAPESCHI S., ROSSETTI F., WALTERS J.B., 2023. *Growth of kyanite and Fe-Mg chloritoid in Fe₂O₃-rich high-pressure–low-temperature metapelites and metapsammites: A case study from the Massa Unit (Alpi Apuane, Italy)*. *Journal of Metamorphic Geology* 41 (8): 1049-1079. <https://doi.org/10.1111/jmg.12736>
- PATACCA E., SCANDONE P., MECCHERI M., MASSA G., 2011. *Stratigraphic and structural revision of the Massa “Schuppenzone” (Alpi Apuane, Northern Apennines)*. *Rendiconti online Società Geologica Italiana* 15: 102-105.
- PERTUSATI P., PLESI G., CERRINA FERONI A., 1977. *Alcuni esempi di tettonica polifasata nella Falda toscana*. *Bollettino Società Geologica Italiana* 96: 587-603.
- PICCINI L., DRYSDALE R., HEIJNIS H., 2003. *Karst morphology and cave sediments as indicators of the uplift history in the Alpi Apuane (Tuscany, Italy)*. *Quaternary International* 101: 219-227.
- PIERUCCIONI D., GALANTI Y., BIAGIONI C., MOLLI G., 2018. *Geologic map of the Fornovolasco area, Alpi Apuane (Tuscany, Italy)*. *Journal of Maps* 14 (2): 357-367. <https://doi.org/10.1080/17445647.2018.1476985>
- PIERUCCIONI D., SPINA A., BROGI A., CAPEZZUOLI E., ZUCCHI M., VEZZONI S., LIOTTA D., SORCI A., MOLLI G., 2023. *The Fornovolasco area (Alpi Apuane, Northern Apennines): A review and update on its Palaeozoic succession, middle Permian magmatism, and tectonic setting*. *Italian Journal Geoscience* 142: 359-382. <https://doi.org/103301/IJ.2023.25>
- RAU A., 1990. *Evolution of the Tuscan Domain from Late Carboniferous to the Mid Triassic: A new hypothesis*. *Bollettino della Società Geologica Italiana* 109: 231-238.
- STORTI F., 1995. *Tectonics of the Punta Bianca promontory: Insights for the evolution of the Northern Apennines-Northern Tyrrhenian Sea basin*. *Tectonics* 14: 832-847. <https://doi.org/10.1029/95TC01203>
- VEZZONI S., BIAGIONI C., D’ORAZIO M., PIERUCCIONI D., GALANTI Y., PETRELLI M., MOLLI G., 2018. *Evidence of Permian magmatism in the Alpi Apuane metamorphic complex (Northern Apennines, Italy): New hints for the geological evolution of the basement of the Adria plate*. *Lithos* 318-319: 104-123.
- ZACCAGNA D., 1897. *Carta e sezioni geologiche delle Alpi Apuane*. *Bollettino del Regio Comitato Geologico Italiano* 28.
- ZACCAGNA D., 1932. *Descrizione Geologica delle Alpi Apuane*. *Memorie Descrittive della Carta Geologica d’Italia*, 25. Servizio Geologico d’Italia, Roma.

(ms. pres. 1 dicembre 2024; ult. bozze 30 dicembre 2024)

