

Abstract We shall prove a Poincaré-Bendixson theorem describing the asymptotic behavior of geodesics for a meromorphic connection on a compact Riemann surface. We shall also briefly discuss the case of non-compact Riemann surfaces, and study in detail the geodesics for a holomorphic connection on a complex torus.

Keywords Meromorphic connections, geodesics, Poincaré-Bendixson Theorem

Mathematics Subject Classification (2000) 32H50 · 34M03 · 34M40 · 37F99

A Poincaré-Bendixson theorem for meromorphic connections on Riemann surfaces

Marco Abate · Fabrizio Bianchi

the date of receipt and acceptance should be inserted later

1 Introduction

The main goal of this paper is to prove a Poincaré-Bendixson theorem describing the asymptotic behavior of geodesics for a meromorphic connection on a compact Riemann surface.

Roughly speaking (see Section 2 for more details), a *meromorphic connection* on a Riemann surface S is a \mathbb{C} -linear operator $\nabla: \mathcal{T}(S) \rightarrow \mathcal{M}^1(S) \otimes \mathcal{T}(S)$, where $\mathcal{T}(S)$ is the space of holomorphic vector fields on S and $\mathcal{M}^1(S)$ is the space of meromorphic 1-forms on S , satisfying a Leibniz rule $\nabla(\phi s) = \phi \nabla s + d\phi \otimes s$ for every $s \in \mathcal{T}(S)$ and every meromorphic function $\phi \in \mathcal{M}(S)$. Meromorphic connections are a classical subject, related for instance to linear differential systems (see, e.g., [IY08]); but in this paper we shall study them following a less classical point of view, introduced in [AT11].

A meromorphic connection ∇ on a Riemann surface S (again, see Section 2 for details) can be represented in local coordinates by a meromorphic 1-form; it turns out that the poles (and the associated residues) of this form do not depend on the coordinates but only on the connection. If $p \in S$ is not a pole, as for ordinary connections in differential geometry, it is possible to use ∇ for differentiating along a direction $v \in T_p S$ any vector field s defined along a curve in S tangent to v , obtaining a tangent vector $\nabla_v s \in T_p S$. In particular, if $\sigma: (a, b) \rightarrow S$ is a smooth curve with image

Marco Abate
Dipartimento di Matematica, Università di Pisa
Largo Pontecorvo 5, 56127 Pisa, Italy
+39 050 2213 230
E-mail: abate@dm.unipi.it

Fabrizio Bianchi
Institut de Mathématiques de Toulouse, Université Paul Sabatier and Dipartimento di Matematica, Università di Pisa
118 route de Narbonne, F-31062 Toulouse, France
E-mail: fbianchi@math.univ-toulouse.fr

contained in the complement of the poles, it makes sense to consider the vector field $\nabla_{\sigma'}\sigma'$ along σ ; and we shall say that σ is a *geodesic* for ∇ if $\nabla_{\sigma'}\sigma' \equiv 0$.

As far as we know, geodesics for meromorphic connections in this sense were first introduced in [AT11]. As shown there, locally they behave as Riemannian geodesics of a flat metric; but their global behavior can be very different from the global behavior of Riemannian geodesics. Thus they are an interesting object of study *per se*; but they also have dynamical applications, in particular in the theory of local discrete holomorphic dynamical systems of several variables — explaining why we are particularly interested in their asymptotic behavior.

One of the main open problems in local dynamics of several complex variables is the understanding of the dynamics, in a full neighbourhood of the origin, of holomorphic germs tangent to the identity, that is of germs of holomorphic endomorphisms of \mathbb{C}^n fixing the origin and with differential there equal to the identity. In dimension one, the Leau-Fatou flower theorem (see, e.g., [A10] or [M06]) provides exactly such an understanding; and building on this theorem Camacho ([C78]; see also [S82]) in 1978 has been able to prove that every holomorphic germ tangent to the identity in dimension one is locally topologically conjugated to the time-1 map of a homogeneous vector field. In other words, time-1 maps of homogeneous vector fields provide a complete list of models for the local topological dynamics of one-dimensional holomorphic germs tangent to the identity.

In recent years, many authors have begun to study the local dynamics of germs tangent to the identity in several complex variables; see, e.g., Écalle ([É81a], [É81b], [É85]), Hakim ([H97], [H98]), Abate, Bracci, Tovena ([A01], [AT03], [ABT04], [AT11]), Rong [R10], Molino [M09], Vivas [V11], Arizzi, Raissy [AR14], and others. A few generalizations to several variables of the Leau-Fatou flower theorem have been proved, but none of them was strong enough to be able to describe the dynamics in a full neighbourhood of the origin; furthermore, examples of unexpected phenomena not appearing in the one-dimensional case have been found. Thus it is only natural to try and study the dynamics of meaningful classes of examples, with the aim of extracting ideas applicable to a general setting; and Camacho's theorem suggests that a particularly interesting class of examples is provided by time-1 maps of homogeneous vector fields. (Actually, the evidence collected so far strongly suggests that a several variable version of Camacho's theorem might hold: generic germs tangent to the identity should be locally topologically conjugated to time-1 maps of homogeneous vector fields. But we shall not pursue this topic here. See also [A15, Sections 5, 6], [B03], [Br08], [BCL08] and [LS14] for applications of time-1 maps in related contexts.)

This is the approach initiated in [AT11], where we discovered that there is a strong relationship between the dynamics of the time-1 map of homogeneous vector fields and the dynamics of geodesics for meromorphic connections on Riemann surfaces. To describe this relationship, we need to introduce a few notations and definitions.

Let Q be a homogeneous vector field on \mathbb{C}^n of degree $v+1 \geq 2$. First of all, notice that the orbits of its time-1 map are contained in the real integral curves of Q ; so we are interested in studying the dynamics of the *real* integral curves of the *complex* homogeneous vector field Q . (Actually, it turns out that complex integral curves of a

homogeneous vector field are related to — classically studied— sections which are horizontal with respect to a meromorphic connection; see [AT11] for details.)

A *characteristic direction* for Q is a direction $v \in \mathbb{P}^{n-1}(\mathbb{C})$ such that the complex line (the *characteristic leaf*) issuing from the origin in the direction v is Q -invariant. An integral curve issuing from a point of a characteristic leaf stays in that leaf forever; so the dynamics in a characteristic leaf is one-dimensional, and thus completely known. In particular, if the vector field Q is a multiple of the radial field (we shall say that Q is *dicritical*) every direction is characteristic, the dynamics is one-dimensional and completely understood. So, we are mainly interested in understanding the dynamics of integral curves outside the characteristic leaves of non-dicritical vector fields. Then in [AT11] we proved the following result:

Theorem 1.1 (Abate-Tovena [AT11]) *Let Q be a non-dicritical homogeneous vector field of degree $v + 1 \geq 2$ in \mathbb{C}^n and let M_Q be the complement in \mathbb{C}^n of the characteristic leaves of Q . Let $[\cdot]: \mathbb{C}^n \setminus \{O\} \rightarrow \mathbb{P}^{n-1}(\mathbb{C})$ denote the canonical projection. Then there exists a singular holomorphic foliation \mathcal{F} of $\mathbb{P}^{n-1}(\mathbb{C})$ in Riemann surfaces, and a partial meromorphic connection ∇ inducing a meromorphic connection on each leaf of \mathcal{F} , whose poles coincide with the characteristic directions of Q , such that the following hold:*

- (i) *if $\gamma: I \rightarrow M_Q$ is an integral curve of Q then the image of $[\gamma]$ is contained in a leaf S of \mathcal{F} and it is a geodesic for ∇ in S ;*
- (ii) *conversely, if $\sigma: I \rightarrow \mathbb{P}^{n-1}(\mathbb{C})$ is a geodesic for ∇ in a leaf S of \mathcal{F} then there are exactly v integral curves $\gamma_1, \dots, \gamma_v: I \rightarrow M_Q$ such that $[\gamma_j] = \sigma$ for $j = 1, \dots, v$.*

Thanks to this result, we see that the study of integral curves for a homogeneous vector field in \mathbb{C}^n is reduced to the study of geodesics for meromorphic connections on a Riemann surface S (obtained as a leaf of the foliation \mathcal{F}).

In [AT11] the latter study has been carried out in the case $S = \mathbb{P}^1(\mathbb{C})$, which is the only case arising when $n = 2$ (and indeed it has led to a fairly extensive understanding of the dynamics of homogeneous vector fields in \mathbb{C}^2 , including the description of the dynamics in a full neighbourhood of the origin for a substantial class of examples); the main goal of this paper is to extend this study from $\mathbb{P}^1(\mathbb{C})$ to a generic compact Riemann surface, with the hope of applying in the future our results to the study of the dynamics of homogeneous vector fields in \mathbb{C}^n with $n \geq 3$.

More precisely, we shall prove a Poincaré-Bendixson theorem describing the ω -limit set of the geodesics for a meromorphic connection on a generic compact Riemann surface. We recall that, in general, a “Poincaré-Bendixson theorem” is a result describing recurrence properties of a class of dynamical systems (see [C12] for a survey on the subject). We shall prove (see Theorems 4.3 and 5.1) that the ω -limit of a not self-intersecting geodesic whose interior is empty may only be a pole for the connection, a closed geodesic, or a boundary graph of saddle connections, i.e., a topological graph on the surface whose vertices are poles and whose arcs are disjoint geodesics connecting two poles and such that it is the boundary of an open subset of the whole surface (see Definition 4.4).

An important ingredient in our proof will be a Poincaré-Bendixson theorem due to Hounie ([H81]), describing the minimal sets for smooth singular line fields on compact orientable surfaces (which in turn follows from a similar statement for smooth

vector fields, see [S63]). Indeed, in Theorem 4.1 we shall show that it is possible to consider a not self-intersecting geodesic for a meromorphic connection as part of an integral curve for a suitable line field on the surface, which, in a neighbourhood of the support of the geodesic, is singular exactly on the poles of the connection, allowing us to apply Hounie's theorem.

The paper is organized as follows. In Section 2 we collect all the preliminary results that we shall need. In Section 3 we generalize results obtained in [AT11] concerning geodesics for meromorphic connections on the Riemann sphere to geodesics for meromorphic connections on a generic compact Riemann surface. In Sections 4 and 5 we prove our main theorems, using the result of Hounie mentioned above to give a topological description of the possible minimal sets, and then concluding the proof by applying the theory developed in the previous sections. In Section 6 we shall briefly discuss what happens on non-compact Riemann surfaces. Finally, in Section 7 we present a detailed study of the geodesics for holomorphic connections on complex tori (the only compact Riemann surfaces admitting meromorphic connections without poles).

2 Preliminary notions

Throughout this paper we shall denote by S a Riemann surface and by $p: TS \rightarrow S$ the tangent bundle on S . Furthermore, \mathcal{O}_S will be the structure sheaf of S , i.e., the sheaf of germs of holomorphic functions on S , and \mathcal{M}_S the sheaf of germs of meromorphic functions. $\mathcal{T}\mathcal{S}$ will denote the sheaf of germs of holomorphic sections of TS and \mathcal{M}_{TS} the sheaf of germs of its meromorphic sections. Finally, Ω_S^1 will denote the sheaf of germs of holomorphic 1-forms, and \mathcal{M}_S^1 the sheaf of germs of meromorphic 1-forms.

We start recalling the definitions and the first properties of holomorphic and meromorphic connections on $p: TS \rightarrow S$. We refer to [AT11] and [IY08] for details.

Definition 2.1 A *holomorphic connection* on the tangent bundle $p: TS \rightarrow S$ over a Riemann surface is a \mathbb{C} -linear map $\nabla: \mathcal{T}\mathcal{S} \rightarrow \Omega_S^1 \otimes \mathcal{T}\mathcal{S}$ satisfying the Leibniz rule

$$\nabla(fs) = df \otimes s + f\nabla s$$

for all $s \in \mathcal{T}\mathcal{S}$ and $f \in \mathcal{O}_S$. We shall often write $\nabla_{u,s}$ for $\nabla s(u)$.

A *geodesic* for a holomorphic connection ∇ is a real curve $\sigma: I \rightarrow S$, with $I \subseteq \mathbb{R}$ an interval, such that $\nabla_{\sigma'} \sigma' \equiv 0$.

Let us see what this definition means in local coordinates. Given a holomorphic atlas $\{(U_\alpha, z_\alpha)\}$ on S , we denote by $\partial_\alpha := \partial/\partial z_\alpha$ the induced local generator of TS over U_α . We shall always suppose that the U_α 's are simply connected. On each U_α we can find a holomorphic 1-form η_α such that

$$\nabla \partial_\alpha = \eta_\alpha \otimes \partial_\alpha,$$

and we shall say that the form η_α represents the connection ∇ on U_α . In fact we see that, for a general section s of TS , we can locally compute ∇s by representing $s|_{U_\alpha}$ as

$s_\alpha \partial_\alpha$ for some holomorphic function s_α on U_α and writing

$$\nabla(s_\alpha \partial_\alpha) = ds_\alpha \otimes \partial_\alpha + s_\alpha \nabla(\partial_\alpha) = ds_\alpha \otimes \partial_\alpha + s_\alpha \eta_\alpha \otimes \partial_\alpha = (ds_\alpha + \eta_\alpha) \otimes \partial_\alpha .$$

Definition 2.2 Let g be an Hermitian metric on $p: TS \rightarrow S$. We say that a connection ∇ on TS is *adapted to*, or *compatible with*, the metric g if

$$d\langle R, T \rangle = \langle \nabla R, T \rangle + \langle R, \bar{\nabla} T \rangle$$

for every pair of smooth vector fields R, T , that is if

$$X\langle R, T \rangle = \langle \nabla_X R, T \rangle + \langle R, \nabla_{\bar{X}} T \rangle$$

for every triple of smooth vector fields X, R and T on S .

It is known that it is possible to associate to any Hermitian metric a (not necessarily holomorphic) connection on the tangent bundle adapted to it, the *Chern connection*. Locally, it is also possible to solve the converse problem, i.e., to construct local metrics adapted to a given holomorphic connection on TS . To do this, it is convenient to consider the local real function $n_\alpha: U_\alpha \rightarrow \mathbb{R}^+$ given by

$$n_\alpha(p) = g_p(\partial_\alpha, \partial_\alpha) .$$

It is straightforward to see that n_α is a smooth function and, conversely, we see that the function n_α uniquely determines the metric on U_α .

With these notations in place it is not difficult to see that the compatibility of a metric g with a holomorphic connection ∇ on the domain U_α of a local chart is equivalent to

$$\partial n_\alpha = n_\alpha \eta_\alpha . \tag{2.1}$$

Equation (2.1) can actually be solved:

Proposition 2.1 ([AT11], **Proposition 1.1**) *Let $\nabla: \mathcal{T}\mathcal{S} \rightarrow \Omega_S^1 \otimes \mathcal{T}\mathcal{S}$ be a holomorphic connection on a Riemann surface S . Let (U_α, z_α) be a local chart for S and let $\eta_\alpha \in \Omega_S^1(U_\alpha)$ be the 1-form representing ∇ on U_α . Given a holomorphic primitive K_α of η_α on U_α , then*

$$n_\alpha = \exp(2\operatorname{Re} K_\alpha) = \exp(K_\alpha + \bar{K}_\alpha)$$

is a positive solution of (2.1).

Conversely, if n_α is a positive solution of (2.1) then for any $z_0 \in U_\alpha$ and any simply connected neighbourhood $U \subseteq U_\alpha$ of z_0 there exists a holomorphic primitive $K_\alpha \in \mathcal{O}(U)$ of η_α over U such that $n_\alpha = \exp(2\operatorname{Re} K_\alpha)$ in U . Furthermore, K_α is unique up to a purely imaginary additive constant.

Finally, two (local) solutions of (2.1) differ (locally) by a positive multiplicative constant.

Remark 2.1 It is important to notice that Proposition 2.1 gives only *local* metrics adapted to ∇ ; a *global* metric adapted to ∇ might not exist (see [AT11, Proposition 1.2]).

So we can associate to a holomorphic connection ∇ a conformal family of compatible local metrics. It turns out (see [AT11]) that these local metrics are locally isometric to the Euclidean metric on \mathbb{C} . In fact, given any holomorphic primitive K_α of η_α , let the function $J_\alpha: U_\alpha \rightarrow \mathbb{C}$ be a holomorphic primitive of $\exp(K_\alpha)$. We immediately remark that J_α actually exists, because U_α is simply connected, and that it is locally invertible, because $J'_\alpha = \exp(K_\alpha)$. In the following Proposition we summarize the main properties of J_α .

Proposition 2.2 ([AT11]) *Let $\nabla: \mathcal{F}\mathcal{S} \rightarrow \Omega_S^1 \otimes \mathcal{F}\mathcal{S}$ be a holomorphic connection on a Riemann surface S . Let (U_α, z_α) be a local chart for S with U_α simply connected, η_α the 1-form representing ∇ on U_α , and K_α a holomorphic primitive of η_α on U_α . Then every primitive $J_\alpha: U_\alpha \rightarrow \mathbb{C}$ of $\exp(K_\alpha)$ is a local isometry between U_α , endowed with the metric represented by $n_\alpha = \exp(2\operatorname{Re} K_\alpha)$, and \mathbb{C} , endowed with the Euclidean metric. In particular, every local metric adapted to ∇ is flat, that is its Gaussian curvature vanishes identically.*

Moreover, a smooth curve $\sigma: I \rightarrow U_\alpha$ is a geodesic for ∇ if and only if there are two constants c_0 and $w_0 \in \mathbb{C}$ such that $J_\alpha(\sigma(t)) = c_0 t + w_0$. In particular, the geodesic with $\sigma(0) = z_0$ and $\sigma'(0) = v_0 \in \mathbb{C}^$ is given by*

$$\sigma(t) = J_\alpha^{-1}(c_0 t + J_\alpha(z_0)), \quad (2.2)$$

where $c_0 = \exp(K_\alpha(z_0))v_0$ and J_α^{-1} is the analytic continuation of the local inverse of J_α near $J_\alpha(z_0)$ such that $J_\alpha^{-1}(J_\alpha(z_0)) = z_0$.

Finally, a curve $\sigma: [0, \varepsilon) \rightarrow U_\alpha$ is a geodesic for ∇ if and only if

$$\sigma'(t) = \exp(-K_\alpha(\sigma(t))) \exp(K_\alpha(\sigma(0))) \sigma'(0), \quad (2.3)$$

if and only if

$$J_\alpha(\sigma(t)) = \exp(K_\alpha(\sigma(0))) \sigma'(0)t + J_\alpha(\sigma(0)). \quad (2.4)$$

Remark 2.2 Another consequence of the existence of a conformal family of local metrics adapted to ∇ is that we can introduce on each tangent space $T_p S$ a well-defined notion of *angle* between tangent vectors, clearly independent of the particular local metric used to compute it.

We shall now give the official definition of *meromorphic connection*.

Definition 2.3 A *meromorphic connection* on the tangent bundle $p: TS \rightarrow S$ of a Riemann surface S is a \mathbb{C} -linear map $\nabla: \mathcal{M}_{TS} \rightarrow \mathcal{M}_S^1 \otimes_{\mathcal{M}(S)} \mathcal{M}_{TS}$ satisfying the Leibniz rule

$$\nabla(\tilde{f}\tilde{s}) = d\tilde{f} \otimes \tilde{s} + \tilde{f}\nabla\tilde{s}$$

for all $\tilde{s} \in \mathcal{M}_{TS}$ and $\tilde{f} \in \mathcal{M}_S$.

It is easy to see that on a local chart (U_α, z_α) a meromorphic connection is represented by a meromorphic 1-form, that we continue to call η_α , such that

$$\nabla(\partial_\alpha) = \eta_\alpha \otimes \partial_\alpha$$

and so

$$\nabla(\tilde{s}) = \nabla(\tilde{s}_\alpha \partial_\alpha) = (d\tilde{s}_\alpha + \eta_\alpha) \otimes \partial_\alpha$$

for every local meromorphic section $\tilde{s}|_{U_\alpha} = \tilde{s}_\alpha \partial_\alpha$ on U_α , exactly as it happens for holomorphic connections.

In particular, all the forms η_α 's are holomorphic if and only if ∇ is a holomorphic connection. More precisely, if we say that $p \in S$ is a *pole* for a meromorphic connection ∇ if p is a pole of η_α for some (and hence any) local chart U_α at p , then ∇ is a holomorphic connection on the complement S^0 of the poles of ∇ in S .

This allows to define *geodesics* for a meromorphic connections as we did for holomorphic connections.

Definition 2.4 A *geodesic* for a meromorphic connection ∇ on $p: TS \rightarrow S$ is a real curve $\sigma: I \rightarrow S^0$, with $I \subseteq \mathbb{R}$ an interval, such that $\nabla_{\sigma'} \sigma' \equiv 0$.

The following definition will be of primary importance in the sequel.

Definition 2.5 The *residue* $\text{Res}_p \nabla$ of a meromorphic connection ∇ at a point $p \in S$ is the residue of any 1-form η_α representing ∇ on a local chart (U_α, z_α) at p . Clearly, $\text{Res}_p \nabla \neq 0$ only if p is a pole of ∇ .

It is easy to see that the residue of a meromorphic connection at a point $p \in S$ is well defined, i.e., it does not depend on the particular chart U_α . Moreover, by the Riemann-Roch formula, the sum of the residues of a meromorphic connection ∇ is independent from the particular connection, as stated in the following classical Theorem (see [IY08] for a proof).

Theorem 2.1 *Let S be a compact Riemann surface and ∇ a meromorphic connection on $p: TS \rightarrow S$. Then*

$$\sum_{p \in S} \text{Res}_p \nabla = -\chi_S, \quad (2.5)$$

where χ_S is the Euler characteristic of S .

Our main result is a classification of the possible ω -limit sets for the geodesics of meromorphic connections on the tangent bundle of a compact Riemann surface. We recall that the ω -limit set $\omega(\sigma)$ of a curve $\sigma: [0, \varepsilon) \rightarrow S^0$ is given by the points p of S for which there exists a sequence $\{t_n\}$, with $t_n \rightarrow \varepsilon$, such that $\sigma(t_n) \rightarrow p$. It is easy to see that

$$\omega(\sigma) = \bigcap_{\varepsilon' < \varepsilon} \overline{\{\sigma(t) : t > \varepsilon'\}}.$$

The main part of the proof will consist in extending the tangent field of the geodesic to a smooth line field (a rank-1 real foliation) on S , singular only on the poles of the connection. Then we shall be able to use the following theorem by Hounie ([H81]) to study the minimal sets of this line field, where a *minimal set* for a line field Λ is a closed, non-empty, Λ -invariant subset of S without proper subsets having the same properties:

Theorem 2.2 (Hounie [H81]) *Let S be a compact connected smooth real manifold of dimension two (e.g., a Riemann surface) and let Λ be a smooth line field with singularities on S . Then a Λ -minimal set Ω must be one of the following:*

1. *a singularity of Λ ;*
2. *a closed integral curve of Λ , homeomorphic to S^1 ;*
3. *all of S , and in this case Λ is equivalent to an irrational line field on the torus (i.e., there exists a homeomorphism $\phi: S \rightarrow T$ between S and a torus T transforming the given foliation into one induced by an irrational line field).*

3 Meromorphic connections on the tangent bundle

In this section, we study in more detail geodesics for meromorphic connections on the tangent bundle of a compact Riemann surface. In particular, we extend results contained in Section 4 of [AT11] from $\mathbb{P}^1(\mathbb{C})$ to the case of a generic compact Riemann surface S .

To do so, we start introducing the following definitions/notations.

Definition 3.1 Let S be a compact Riemann surface. Let ∇ be a meromorphic connection on TS and let $S^0 \subseteq S$ be the complement of the poles.

- A *geodesic (n -)cycle* is the union of n geodesic segments $\sigma_j: [0, 1] \rightarrow S^0$, disjoint except for the conditions $\sigma_j(0) = \sigma_{j-1}(1)$ for $j = 2, \dots, n$ and $\sigma_1(0) = \sigma_n(1)$. The points $\sigma_j(0)$ will be called *vertices* of the geodesic cycle.
- A *(m -)multicurve* is a union of m disjoint geodesic cycles. A multicurve will be said to be *disconnecting* if it disconnects S , *non-disconnecting* otherwise.
- A *part P* is the closure of a connected open subset of S whose boundary is a multicurve γ . A component σ of γ is *surrounded* if the interior of P contains both sides of a tubular neighbourhood of σ in S ; it is *free* otherwise. The *filling \hat{P}* of a part P is the compact surface obtained by glueing a disk along each of the free components of γ , and not removing any of the surrounded components of γ .

We remark that a part may be all of S , when the associated multicurve is non-disconnecting (which is equivalent to saying that the multicurve has no free components). Moreover, we see that every disconnecting multicurve contains the boundary of a part $P \subsetneq S$.

As recalled in the previous section, we can associate to a meromorphic connection ∇ conformal families of local metrics on the regular part $S^0 \subseteq S$ of the Riemann surface, and we have a well-defined notion of angle between tangent vectors. In particular, it makes sense to speak of the *external angle ε* at a vertex $\sigma_j(0)$ of a geodesic cycle as the angle between the tangent vectors $\sigma'_{j-1}(1)$ and $\sigma'_j(0)$; by definition, $\varepsilon \in (-\pi, \pi)$.

Again using the conformal families of local metric it is possible to define the *geodesic curvature k_g* of a curve contained in S^0 . In particular, if $p \in S$ is a pole of ∇ and τ is a small (clockwise) circle around p , not containing other poles, we have

$$\int_{\tau} k_g = -2\pi(1 + \operatorname{Re} \operatorname{Res}_p(\nabla)); \quad (3.1)$$

see [AT11, Theorem 4.1].

With all these ingredients at our disposal, it becomes natural to try and apply a Gauss-Bonnet Theorem to study the relation between the residues of the connection ∇ in a part P of S , the external angles at the vertices of the multicurve bounding P and the topology of S (cp. [AT11, Theorem 4.1]).

Theorem 3.1 *Let ∇ be a meromorphic connection on a compact Riemann surface S , with poles $\{p_1, \dots, p_r\}$ and set $S^0 := S \setminus \{p_1, \dots, p_r\}$. Let P be a part of S whose boundary multicurve $\gamma \subset S^0$ has $m_f \geq 1$ free components, positively oriented with respect to P . Let z_1, \dots, z_s denote the vertices of the free components of γ , and $\varepsilon_j \in (-\pi, \pi)$ the external angle at z_j . Suppose that P contains the poles $\{p_1, \dots, p_g\}$ and denote by $g_{\hat{P}}$ the genus of the filling \hat{P} of P . Then*

$$\sum_{j=1}^s \varepsilon_j = 2\pi \left(2 - m_f - 2g_{\hat{P}} + \sum_{j=1}^g \operatorname{Re} \operatorname{Res}_{p_j}(\nabla) \right). \quad (3.2)$$

Proof For $j = 1, \dots, g$ let τ_j be a small clockwise circle bounding a disk in P centered at p_j , and let k_g^j be the geodesic curvature of τ_j .

Applying the Gauss-Bonnet Theorem as in [AT11, Theorem 4.1] to the complement P^0 in P of the small disks bounded by τ_1, \dots, τ_g we find that

$$\sum_{j=1}^g \int_{\tau_j} k_g^j + \sum_{j=1}^s \varepsilon_j = 2\pi \chi_{P^0} = 2\pi(2 - m_f - g - 2g_{\hat{P}}). \quad (3.3)$$

But from (3.1) we get

$$\sum_{j=1}^g \int_{\tau_j} k_g^j = -2\pi g - 2\pi \sum_{j=1}^g \operatorname{Re} \operatorname{Res}_{p_j}(\nabla). \quad (3.4)$$

Comparing (3.3) and (3.4) we get the assertion. \square

Remark 3.1 When the multicurve γ does not disconnect S then Theorem 3.1 reduces to Theorem 2.1. Indeed, in this case γ has no free components, and thus (3.2) becomes

$$0 = 2\pi \left(2 - 2g_S + \sum_{j=1}^g \operatorname{Re} \operatorname{Res}_{p_j}(\nabla) \right),$$

which is equivalent to (2.5).

In the next two Corollaries we highlight what happens when the disconnecting multicurve is made up by a single geodesic or by a single geodesic cycle composed by two geodesics (cp. [AT11, Corollaries 4.2 and 4.3]).

Corollary 3.1 *Let ∇ be a meromorphic connection on a compact Riemann surface S , with poles $\{p_1, \dots, p_r\}$ and set $S^0 := S \setminus \{p_1, \dots, p_r\}$. Let σ be a disconnecting closed geodesic. Let P be one of the two parts in which S is disconnected by σ . Then*

$$\sum_{p_j \in P} \operatorname{Re} \operatorname{Res}_{p_j}(\nabla) = -1 + 2g_{\hat{P}}.$$

Corollary 3.2 *Let ∇ be a meromorphic connection on a compact Riemann surface S , with poles $\{p_1, \dots, p_r\}$ and set $S^0 := S \setminus \{p_1, \dots, p_r\}$. Let γ be a disconnecting geodesic 2-cycle. Let P be one of the two parts in which S is disconnected by γ , and $\varepsilon_0, \varepsilon_1 \in (-\pi, \pi)$ the two external angles of γ . Then*

$$\varepsilon_0 + \varepsilon_1 = 2\pi \left(1 - 2g_{\hat{P}} + \sum_{p_j \in P} \operatorname{Re} \operatorname{Res}_{p_j}(\nabla) \right),$$

and hence

$$\sum_{p_j \in P} \operatorname{Re} \operatorname{Res}_{p_j}(\nabla) \in (-2 + 2g_{\hat{P}}, 2g_{\hat{P}}).$$

Remark 3.2 In particular, a part of S bounded by a disconnecting 1-cycle or by a disconnecting 2-cycle must necessarily contain a pole, because $0 \notin (-2 + 2g_{\hat{P}}, 2g_{\hat{P}})$.

In the following we shall need to consider *closed* and *periodic* geodesics for a meromorphic connection.

Definition 3.2 A geodesic $\sigma: [0, l] \rightarrow S$ is *closed* if $\sigma(l) = \sigma(0)$ and $\sigma'(l)$ is a positive multiple of $\sigma'(0)$; it is *periodic* if $\sigma(l) = \sigma(0)$ and $\sigma'(l) = \sigma'(0)$.

By Corollary 3.1 we immediately see that a disconnecting geodesic 1-cycle σ is a closed geodesic if and only if for every part P of S bounded by σ we have

$$\sum_{p_j \in P} \operatorname{Re} \operatorname{Res}_{p_j}(\nabla) = -1 + 2g_{\hat{P}}. \quad (3.5)$$

Contrarily to the Riemannian case, closed geodesics are not necessarily periodic; for examples (and more) in the Riemann sphere see [AT11], and for examples on a complex torus see Section 7.

4 ω -limits sets of geodesics

In this section we prove our main Theorem, the classification of the possible ω -limit sets of geodesics for a meromorphic connection on a compact Riemann surface. The main idea will be to see a not self-intersecting geodesic σ as part of an integral curve of a suitable line field Λ on S , and then to apply Theorem 2.2 to get some information about the minimal sets for Λ contained in the ω -limit set of σ . Then we shall use these information to discuss the shape of the ω -limit set itself.

Let us start by studying the local structure of the ω -limit set of a not self-intersecting geodesic σ .

Proposition 4.1 *Let S be a Riemann surface, ∇ a meromorphic connection on S , and S^0 the complement of the poles for ∇ . Let $\sigma: I \rightarrow S^0$ be a maximal not self-intersecting geodesic for ∇ and denote by W its ω -limit set. Let $z \in W \cap S^0$. Then:*

- (i) *there exists a unique direction v at z such that for every sequence $\{z_n\} \subset \sigma(I)$ with $z_n \rightarrow z$ the directions $\sigma'(z_n)$ converge to v ; moreover, the image of the maximal geodesic σ_z issuing from z in the direction v is contained in W ;*

- (ii) if W contains a curve τ passing through z transversal to σ_z then $z \in \mathring{W}$;
 (iii) if $z \in \partial W$ then σ_z is the unique geodesic segment through z contained in W .

Proof (i) The existence of ν follows from the flatness of the metric and the fact that σ does not self-intersect. Using a local isometry J defined in a neighbourhood of z transforming geodesic segments in Euclidean segments we immediately see that the image of the geodesic segment issuing from z in the direction ν must be contained in W , and the maximality follows from the maximality of σ .

(ii) Up to using a local isometry J we can assume that all geodesic segments in a neighbourhood of z are Euclidean segments.

Consider a point $z' \neq z$ belonging to τ . Since $z' \in W \subset S^0$, (i) gives us a geodesic segment $\sigma_{z'}$ through z' contained in W . Notice that $\sigma_{z'} \neq \tau$ as soon as z' is close enough to z . In fact, since we have segments of σ accumulating σ_z , the segments of σ accumulating $\sigma_{z'}$ cannot converge to τ without forcing σ to self-intersect, impossible. The mapping $z' \mapsto \sigma_{z'}$ is continuous, again because σ cannot self-intersect; thus the geodesic segments $\sigma_{z'}$ fill an open neighbourhood of z contained in W , and $z \in \mathring{W}$ as claimed.

(iii) It immediately follows from (ii). □

Definition 4.1 Let S be a Riemann surface, ∇ a meromorphic connection on S , and S^0 the complement of the poles of ∇ . Let $\sigma: I \rightarrow S^0$ be a maximal not self-intersecting geodesic for ∇ , with ω -limit $W \subseteq S$. Given $z \in W \cap S^0$, the maximal geodesic σ_z given by Proposition 4.1.(i) issuing from z and contained in W is the *distinguished geodesic* issuing from z .

The following two lemmas contain basic properties of distinguished geodesics.

Lemma 4.1 Let S be a Riemann surface, ∇ a meromorphic connection on S , and S^0 the complement of the poles of ∇ . Let $\sigma: I \rightarrow S^0$ be a maximal not self-intersecting geodesic for ∇ , with ω -limit $W \subseteq S$. Let σ_z be the distinguished geodesic issuing from $z \in W \cap S^0$. Then either $\sigma_z = \sigma$ or σ does not intersect σ_z .

Proof If σ intersects σ_z , either $\sigma = \sigma_z$ or the intersection is transversal; but σ accumulates σ_z , and thus in the latter case σ must self-intersect, contradiction. □

Lemma 4.2 Let S be a Riemann surface, ∇ a meromorphic connection on S , and S^0 the complement of the poles of ∇ . Let $\sigma: I \rightarrow S^0$ be a maximal not self-intersecting geodesic for ∇ , with ω -limit $W \subseteq S$. Assume there exists a closed distinguished geodesic $\alpha \subseteq W$. Then $W = \alpha$.

Proof If $\alpha = \sigma$ then the assertion is trivial; so by the previous lemma we can assume that σ does not intersect α .

For each $w \in \alpha$ we can find a neighbourhood $A_w \subset S^0$ of w , contained in a tubular neighbourhood of α , and an isometry $J_w: A_w \rightarrow B_w$ such that:

- (a) B_w is a quadrilateral containing the origin, with two opposite sides parallel to the imaginary axis;
- (b) $J_w(w) = O$ and $J_w(\alpha \cap A_w)$ is the intersection of B_w with the real axis;

(c) the J_w -images of the connected components of the intersection of σ with A_w contained in the lower half-plane intersect both vertical sides of B_w and accumulate the real axis (this can be achieved using Proposition 4.1.(i)).

By compactness, we can cover α with finitely many A_{w_0}, \dots, A_{w_r} of such neighbourhoods; furthermore, we can also assume that each A_{w_j} intersects only $A_{w_{j-1}}$ and $A_{w_{j+1}}$, with the usual convention $w_{-1} = w_r$ and $w_{r+1} = w_0$; let $A = A_0 \cup \dots \cup A_r$. Furthermore, since A is contained in a tubular neighbourhood of α and S is orientable, $A \setminus \alpha$ has exactly two connected components; and we can assume that the segments of σ mapped by the J_{w_j} in the lower half-plane (and thus accumulating α) are all contained in the same connected component of $A \setminus \alpha$.

Let τ be the segment of geodesic issuing from w_0 such that $J_{w_0}(\tau)$ is the intersection of B_{w_0} with the negative imaginary axis. Denote by $\{z_j\}$ the infinitely many intersections of σ with τ , accumulating w_0 . Again by compactness, we can assume that if we follow σ starting from a z_1 close enough to w_0 we stay in A until we get to the next intersection z_2 . Furthermore, by property (c) the segment σ_1 of σ from z_1 to z_2 must intersect all A_{w_k} 's either in clockwise or in counter-clockwise order. Let $D \subset A$ denote the domain bounded by α , the geodesic segment σ_1 and the geodesic segment of τ from z_1 to z_2 .

Assume first that z_2 is closer to w_0 than z_1 . In this case, if we follow σ starting from z_2 property (c) forces σ to remain inside D because it cannot intersect itself nor α . Furthermore, property (c) again implies that the next intersection z_3 should be closer to w_0 than z_2 . We can then repeat the argument, and we find that successive intersections of σ with τ monotonically converge to w_0 . This implies that σ accumulates α and nothing else, that is $W = \alpha$ as claimed.

If instead z_2 is farther away from w_0 than z_1 , if we follow σ starting from z_2 we may possibly leave A ; but since σ accumulates w_0 we must sooner or later get back to D , and the only way is intersecting τ in a point z_3 between z_1 and z_2 . But now following σ starting from z_3 we are forced to stay into D until we intersect τ in a point z_4 necessarily closer to w_0 than z_1 — and a fortiori closer than z_3 . We can then repeat the previous argument using z_3 and z_4 instead of z_1 and z_2 to get again the assertion. \square

The main tool for the study of the global behavior of not self-intersecting geodesics is the following Theorem, providing a smooth line field Λ such that σ is (part of) an integral curve of Λ .

Theorem 4.1 *Let S be a Riemann surface and ∇ a meromorphic connection on S , with poles $p_1, \dots, p_r \in S$. Let $S^0 = S \setminus \{p_1, \dots, p_r\}$. Let $\sigma: I \rightarrow S^0$ be a geodesic for ∇ without self-intersections, maximal in both forward and backward time. Then there exists a smooth line field Λ with singularities on S which has σ as integral curve and, in a neighbourhood of σ , is singular exactly on the poles of ∇ . Furthermore, the ω -limit W of σ is Λ -invariant, and $\Lambda|_W$ is uniquely determined.*

Proof If $A \subset S^0$ is open, simply connected and small enough, we can find a metric g on A compatible with ∇ and an isometry J between A endowed with g and an open set in \mathbb{C} endowed with the euclidean metric; g is unique up to a positive multiple (see Propositions 2.1 and 2.2).

We consider an open cover $\mathcal{A} = \{A_i\}$ of S^0 with the following properties:

- (a) each A_i is endowed with a metric g_i compatible with ∇ , and with an isometry $J_i: A_i \rightarrow B_i \subseteq \mathbb{C}$, with B_i convex;
- (b) if σ intersects A_i then the angle between any pair of lines in $\mathbb{C} \supset B_i$ containing a component (which necessarily is a line segment) of the J_i -image of σ is less than $\frac{\pi}{8}$;
- (c) \mathcal{A} is locally finite.

In order to do this, we construct for every point $p \in S^0$ a neighbourhood satisfying the properties (a)-(b) and then we recover (c) by extracting a locally finite cover. The construction of the neighbourhood can be achieved in the following way. We start with an A endowed with the local metric and isometric to a convex $B \subset \mathbb{C}$, as in (a). Then we can shrink it (holding the convexity property on B) in order to recover (b) because the smooth dependence of the solution of the geodesic equation from the initial conditions, the fact that the isometry is smooth, and the property of σ of being non self-intersecting force the images by J of the intersections of σ and A to be quasi-parallel near the image of a given point in $A \cap \sigma$.

We now start building a line field on every open set A_i of \mathcal{A} . Then we shall show how to use them to find a global line field Λ on S^0 having σ as integral curve, and finally we shall extend Λ to all of S .

Let us then choose an open set $A \in \mathcal{A}$, together with its image B . We shall now construct a smooth flow of curves in B that will correspond to a smooth flow of curves, and so to a line field, in A .

If σ does not cross A , we put on B any smooth vector field which is never zero and consider its associated (regular) foliation.

If σ crosses A , we consider the segments $\sigma_n \subset B$ which are the images of the connected components of the intersection of σ with A (recall that J sends geodesic segments in A to Euclidean segments in B). By our assumption on A , the angle between σ_i and σ_j is bounded by $\frac{\pi}{8}$ for every pair (i, j) .

By the convexity of B and the maximality of σ , the σ_n 's subdivide B in connected components. We describe now how to construct the flow in all these components.

1. If a connected component C is open and its boundary (in B) consists of a unique σ_0 , then we define our flow on C by means of lines parallel to σ_0 and take the associated line field.
2. If a connected component C is open and it is bounded by two segments σ_0 and σ_1 we define the vector field in the following way, exploiting the convexity property of the cover. Denote by $\tilde{\sigma}_i$ the line containing σ_i . If $\tilde{\sigma}_0$ and $\tilde{\sigma}_1$ are parallel we define the line field by means of parallel lines to them, too. If $\tilde{\sigma}_0$ and $\tilde{\sigma}_1$ are not parallel, their intersection point p_0 is outside B , because B is convex and σ_0 and σ_1 are disjoint. We take two points $y_0 \in \tilde{\sigma}_0$ and $y_1 \in \tilde{\sigma}_1$, both close enough to p_0 , such that the segment joining them is completely outside B and contained in the bounded part of plane bounded by $\tilde{\sigma}_0$, $\tilde{\sigma}_1$ and the boundary of B . This can be done again thanks to the convexity of B and the fact that σ does not self-intersect. We parametrize this segment $[y_0, y_1]$ by an affine map $\tau: [0, 1] \rightarrow \mathbb{C}$ with $\tau(0) = y_0$ and $\tau(1) = y_1$. For every $t \in [0, 1]$, we consider the line \tilde{l}_t passing through $\tau(t)$ forming an angle $\phi(t)\theta_1$ with $\tilde{\sigma}_0$, where $\phi: [0, 1] \rightarrow [0, 1]$ is a suitable smooth not decreasing function which is 0 in a neighbourhood of 0 and 1 in a neighbourhood

of 1, and θ_1 is the angle between $\tilde{\sigma}_0$ and $\tilde{\sigma}_1$. We see that the intersections between two distinct such lines must be contained in the half-plane bounded by the line extending $[y_0, y_1]$ and containing p_0 , and so it is outside B . It follows that the intersections $l_i = \tilde{l}_i \cap B$ form a smooth flow on C , and this flow is clearly smooth also at the boundary of C (i.e. near σ_0 and σ_1).

3. Since the σ_n 's are disjoint, maximal and with angles bounded by $\pi/8$, the only missing case is a component C whose boundary is a segment τ accumulated by a subsequence of σ_n outside C ; we then add this segment to the line field and proceed as in the previous cases.

In this way, thanks to the smooth dependence of geodesics on initial conditions, we have defined a smooth and never vanishing line field on B and hence, via J , on A . Notice that the image in B of a component of the intersection of the ω -limit set W with A must be a segment as in case 3; thus $W \cap A$ is invariant with respect to this local line field, which is uniquely defined there.

The next step consists in glueing the local line fields we have built on the A_i to a global field on S^0 . This means that we must specify, for every point $p \in S^0$, a direction $\lambda(p)$ in $T_p S^0$ such that the correspondence $p \mapsto \lambda(p)$ is smooth. To do so, we consider a partition of unity $\{\rho_i\}$ subordinated to the cover \mathcal{A} . If p belongs to a unique A_i , we use as $\lambda(p)$ the one given by the local construction above. Otherwise, if p belongs to a finite number of A_i 's (recall that the cover is locally finite) we do the following. Suppose that $p \in A_1 \cap \dots \cap A_n$. We have n lines in $T_p S^0$, given by the local constructions on the B_i 's. We use the partition of unity to do a convex combination of (the angles with respect to any fixed line of) these lines, thus obtaining a line in $T_p S^0$. We remark that, by Remark 2.2, the angles are the same on all A_1, \dots, A_n , and the bounds on the angles ensure that the convex combination yields a well-defined line.

We have thus obtained a smooth line field on S^0 having σ as integral curve. Since it is non-singular along σ , by smoothness it is non-singular in a neighbourhood of σ in S^0 .

Finally, we extend this line field to all of S , adding the poles as singular points. The resulting field satisfies the requests of the Theorem, and we are done. \square

Remark 4.1 The reason we have to use a line field instead of a vector field is step 3 in the previous proof: a priori, the geodesic σ might accumulate the segment C from both sides along opposite directions, preventing the identification of a non-singular vector field along C .

Definition 4.2 Let S be a Riemann surface, ∇ a meromorphic connection on S , and $S^0 \subset S$ the complement of the poles. Let $\sigma: I \rightarrow S^0$ be a maximal geodesic for ∇ without self-intersections, maximal in both forward and backward time. A smooth line field Λ with singularities on S as in the statement of Theorem 4.1 is called *associated* to σ .

The uniqueness of an associated line field on the ω -limit set suggests the following definition. We recall that a *minimal set* for a line field Λ is a minimal element (with respect to inclusion) in the family of closed Λ -invariant subsets.

Definition 4.3 Let S be a Riemann surface, ∇ a meromorphic connection on S , and S^0 the complement of the poles of ∇ . Let $\sigma: I \rightarrow S^0$ be a maximal not self-intersecting geodesic for ∇ , with ω -limit set $W \subseteq S$. Then a *minimal set* for σ is a minimal set contained in W of any line field associated to σ .

The following result characterizes the possible minimal sets for a maximal not self-intersecting geodesic in a compact Riemann surface.

Theorem 4.2 *Let S be a compact Riemann surface and ∇ a meromorphic connection on S . Let S^0 be the complement of the poles of ∇ . Let $\sigma: [0, \varepsilon_0) \rightarrow S^0$ be a maximal not self-intersecting geodesic for ∇ . Then the possible minimal sets for σ are the following:*

1. a pole of ∇ ;
2. a closed curve, homeomorphic to S^1 , which is a closed geodesic for ∇ , and in this case the ω -limit set of σ coincides with this closed geodesic;
3. all of S , and in this case S is a torus and ∇ is holomorphic everywhere.

Proof Call $p_0 = \sigma(0)$ the starting point of σ , and W the ω -limit set of σ .

We apply the construction in Theorem 4.1, considering p_0 as a virtual pole. In this way, σ becomes maximal in both forward and backward time and the construction can be carried out as before. We thus build a line field Λ with singularities on S , which in a neighbourhood of σ is singular exactly on the poles of ∇ and on p_0 , and having σ as integral curve.

Applying Theorem 2.2 to Λ we see that the minimal sets for σ can be: all of S (and in this case S is a torus and ∇ has no poles), a closed curve homeomorphic to S^1 (which is a closed geodesic thanks to Proposition 4.1), a pole of ∇ or p_0 . This last possibility is excluded by considering a new starting point p'_0 for σ close enough to p_0 and noticing that now p_0 is not a pole for the new line field.

Finally, the closed geodesic in case 2 being Λ -invariant is necessarily distinguished; we can then end the proof by quoting Lemma 4.2. \square

To describe the possible ω -limit sets of a ∇ -geodesic, we need a definition.

Definition 4.4 A *saddle connection* for a meromorphic connection on a Riemann surface S is a maximal geodesic $\sigma: (-\varepsilon_-, \varepsilon_+) \rightarrow S^0$ such that $\sigma(t)$ tends to a pole for both $t \rightarrow -\varepsilon_-$ and $t \rightarrow \varepsilon_+$.

A *graph of saddle connections* is a connected planar graph in S whose vertices are poles and whose arcs are disjoint saddle connections. A *spike* is a saddle connection of a graph which does not belong to any cycle of the graph.

A *boundary graph of saddle connections* (or *boundary graph*) is a graph of saddle connections which is also the boundary of a connected open subset of S . A boundary graph is *disconnecting* if its complement in S is not connected.

We can now state and prove our main result.

Theorem 4.3 *Let S be a compact Riemann surface and ∇ a meromorphic connection on S , with poles $p_1, \dots, p_r \in S$. Let $S^0 = S \setminus \{p_1, \dots, p_r\}$. Let $\sigma: [0, \varepsilon_0) \rightarrow S^0$ be a maximal geodesic for ∇ . Then either*

1. $\sigma(t)$ tends to a pole of ∇ as $t \rightarrow \varepsilon_0$; or
2. σ is closed; or
3. the ω -limit set of σ is the support of a closed geodesic; or
4. the ω -limit set of σ in S is a boundary graph of saddle connections; or
5. the ω -limit set of σ is all of S ; or
6. the ω -limit set of σ has non-empty interior and non-empty boundary, and each component of its boundary is a graph of saddle connections with no spikes and at least one pole (i.e., it cannot be a closed geodesic); or
7. σ intersects itself infinitely many times.

Furthermore, in cases 2 or 3 the support of σ is contained in only one of the components of the complement of the ω -limit set, which is a part P of S having the ω -limit set as boundary.

Proof Suppose σ is not closed, nor with infinitely many self-intersections. Then up to changing the starting point of σ we can assume that σ does not self-intersect. Let W be the ω -limit set of σ .

By Zorn's lemma, W must contain a minimal set, that by Proposition 4.2 must be either a pole, a closed geodesic or all of S (and in this case $W = S$, which gives 5).

If W reduces to a pole or to a closed geodesic, we are in cases 1 or 3. So we can assume that W properly contains a minimal set and is properly contained in S ; we have to prove that we are in case 4 or 6.

The interior of W may be empty or not empty; we consider the two cases separately. Let us start with $\mathring{W} = \emptyset$. First, remark that σ cannot intersect W , because otherwise (recall Proposition 4.1) σ would intersect itself. As a consequence, σ is contained in only one connected component P of $S \setminus W$, and $W = \partial P$. Then, since W does not reduce to a single pole and is connected (because S is compact), we must have $W \cap S^0 \neq \emptyset$. Assume by contradiction that W contains a closed geodesic α . Since $\mathring{W} = \emptyset$, Proposition 4.1.(iii) implies that α is distinguished; but then Lemma 4.2 implies $W = \alpha$, impossible. So all the minimal sets contained in W are poles.

By Proposition 4.1, every point in $W \cap S^0$ must be contained in a unique maximal distinguished geodesic contained in W ; moreover the ω -limit sets of these geodesic segments must be contained in W , because W is closed; we are going to prove that the ω -limit sets of these geodesics must be poles, completing the proof that W is a boundary graph of saddle connections.

Let then ρ be a distinguished geodesic and assume by contradiction that a point $w \in S^0$ belongs to the ω -limit set $\omega(\rho)$ of ρ . By the local form of the geodesics, and since $\omega(\rho)$, being contained in W , has empty interior, by Proposition 4.1 we find a unique geodesic segment α through w contained in $\omega(\rho)$. Take another geodesic segment transversal to α in w . Since ρ must accumulate α , and ρ is accumulated by σ , it follows that σ must intersect the transversal arbitrarily near w with both the directions of intersection. This means that we have infinitely many disconnecting geodesic 2-cycles, with disjoint interiors and arbitrarily close to $\omega(\rho)$. But by Remark 3.2 all these cycles must contain a pole, which is impossible.

Now let us consider the case $\mathring{W} \neq \emptyset$. We know that W is a closed union of leaves (and singular points) for Λ , and the same is true for its boundary ∂W , with $\partial W \neq \emptyset$ because we are assuming $W \neq S$. The minimal sets contained in ∂W must be poles or

closed geodesics, and ∂W cannot consist of poles only because otherwise we would have $W = S$. So, $\partial W \cap S^0 \neq \emptyset$ and, by Proposition 4.1, every $z \in \partial W \cap S^0$ is contained in a unique geodesic segment contained in ∂W . In particular, if by contradiction ∂W contained a closed geodesic we would have $W = \alpha$ (Lemma 4.2), impossible; so arguing as above we see that each connected component of ∂W is a graph of saddle connections, that clearly cannot contain spikes (because σ cannot cross ∂W), and we are done. \square

Remark 4.2 We have examples for all the cases of Theorem 4.3 (see Section 7 and [AT11]) except for cases 4 and 6. Notice that in these cases if p_0 is a pole belonging to the boundary of the ω -limit set of the geodesic σ then there must exist a neighbourhood U of p_0 containing infinitely many distinct segments of σ (because σ is accumulating p_0 but it is not converging to it). The study of the local behavior of geodesics nearby a pole performed in [AT11] implies that p_0 must then be an irregular singularity or a Fuchsian singularity having $\operatorname{Re} \operatorname{Res}_{p_0}(\nabla) \geq -1$ (see [AT11] for the definitions, recalling that the connection we denote by ∇ here is denoted by ∇^0 in [AT11]). Conversely, if all poles of ∇ are either apparent singularities or Fuchsian singularities with real part of the residues less than -1 then cases 4 and 6 cannot happen.

Remark 4.3 There are examples of smooth line fields (and even vector fields) on Riemann surfaces having integral curves accumulating graphs of saddle connections, or having an ω -limit set with not empty interior (but not covering everything); see, for example, [ST88].

Remark 4.4 When S is the Riemann sphere, case 6 cannot happen. Indeed, assume by contradiction that a not self-intersecting geodesic $\sigma \subset S$ has an ω -limit set W with not-empty interior. Since σ cannot cross ∂W , we can assume that σ is contained in \dot{W} . Let $w_0 \in \partial W$ be a smooth point of the boundary, and $\tau \subset W$ a geodesic segment issuing from w_0 . The intersections between σ and τ must be dense in τ ; then using suitable segments of σ and τ we can construct infinitely many disjoint geodesic 2-cycles. But in the Riemann sphere every 2-cycle is disconnecting; thus by Remark 3.2 each of them must contain at least one pole. Using the finiteness of the number of poles one readily gets a contradiction.

5 ω -limit sets and residues

Corollary 3.1 immediately gives a necessary condition for the existence of geodesics having as ω -limit set a disconnecting closed geodesic:

Corollary 5.1 *Let S be a compact Riemann surface and ∇ a meromorphic connection on S , with poles $p_1, \dots, p_r \in S$, and set $S^0 = S \setminus \{p_1, \dots, p_r\}$. Let $\sigma: [0, \varepsilon_0) \rightarrow S^0$ be a maximal geodesic for ∇ , either closed or having as ω -limit set a closed geodesic. Assume that either σ or its ω -limit set disconnects S , and let P be the part of S containing σ . Then*

$$\sum_{p_i \in P} \operatorname{Re} \operatorname{Res}_{p_i}(\nabla) = -1 + 2g_{\hat{P}}, \quad (5.1)$$

where \hat{P} is the filling of P .

Furthermore, using Theorem 3.1 as in [AT11, Theorem 4.6] one can easily get conditions on the residues that must be satisfied for the existence of self-intersecting geodesics. Our next aim is to find a similar necessary condition for the existence of ω -limit sets which are boundary graphs of saddle connections.

To do this, we shall need a slightly more refined notion of disconnecting graph, to avoid trivialities. To have an idea of the kind of situations we would like to avoid, consider a boundary graph consisting in a non-disconnecting cycle and a small (necessarily disconnecting) loop attached to a vertex. This boundary graph is disconnecting only because of the trivial small loop, and if we shrink the loop to the vertex we recover a non-disconnecting boundary graph. To define a class of graphs where this cannot happen we first introduce a procedure that we shall call *desingularization of a graph* G (obtained as ω -limit set of a geodesic) *to a curve* γ .

Let σ be a ∇ -geodesic without self-intersections and suppose that its ω -limit set G is a boundary graph. If G is tree (that is, it does not contain cycles) then it does not disconnect S ; so let us assume that G is not a tree (and thus it contains at least one cycle; but this does not imply that G disconnects S). In particular, if U is a small enough open neighbourhood of G then $U \setminus G$ is not connected.

Remark 5.1 If a boundary graph contains a spike, it can be the boundary of only one connected open set.

The following Lemma describes the asymptotic behavior of σ near G .

Lemma 5.1 *Let S be a compact Riemann surface and ∇ a meromorphic connection on S , with poles $p_1, \dots, p_r \in S$. Let $S^0 = S \setminus \{p_1, \dots, p_r\}$. Let $\sigma: [0, \varepsilon_0) \rightarrow S^0$ be a maximal geodesic for ∇ . Assume that σ has no self-intersections and that its ω -limit set G is a boundary graph containing at least one cycle. Let U be a small connected open neighbourhood of G , with $U \setminus G$ disconnected and contained in S^0 . Then the support of σ is definitively contained in only one connected component U_σ of $U \setminus G$.*

Proof If, by contradiction, σ definitely intersects two different connected components of $U \setminus G$, then it must intersect ∂U infinitely many times; being ∂U compact, σ then would have a limit point in ∂U which is disjoint from G , impossible. \square

We shall denote by U_σ the connected component of $U \setminus G$ given by Lemma 5.1. In particular, notice that we must have $G \subset \partial U_\sigma$, and that U_σ must be contained in the part P of S containing the support of σ whose boundary is G .

Let us now describe the desingularization of G . For every vertex p_j of G , let B_j be a small open ball centered at p_j . Moreover, for each spike $s_i \subseteq G$ let C_i be a small connected open neighbourhood of s_i with smooth boundary. We can assume that all the B_j and C_i are contained in an open neighbourhood U satisfying the hypotheses of Lemma 5.1. Then the union of the following three sets is a Jordan curve γ in S^0 , that we call a *desingularization of G* :

- $G \setminus (\cup_j B_j \cup \cup_i C_i)$;
- $(\cup_j \partial B_j \cap U_\sigma) \setminus \cup_i C_i$;
- $(\cup_i \partial C_i \cap U_\sigma) \setminus \cup_j B_j$.

The rationale behind this definition is the following: we take the graph and the boundary of the neighbourhoods of the spikes outside the small balls at the vertices and we connect the pieces with small arcs (which are contained in the boundaries of the small balls). We see that, in particular, we can (uniformly) approximate the graph G with desingularizing curves, with respect to any global metric on the compact Riemann surface S .

We are now ready to give the following definitions.

Definition 5.1 A boundary graph G of saddle connections, which is the ω -limit set of a not self-intersecting ∇ -geodesic, is *essentially disconnecting* if every sufficiently close desingularization of G disconnects S .

It is clear that, for sufficiently close desingularizations, if one of them is disconnecting then all of them are. Because of this we may give the previous definition without specifying which particular desingularization we are using.

Furthermore, if G is essentially disconnecting, then every sufficiently close desingularization γ is the boundary of a well-defined part P_γ of S , the (closure of the) connected component of $S \setminus \gamma$ intersecting the part P whose boundary is G and containing the support of σ . We can then consider the filling \hat{P}_γ obtained by glueing a disk along γ . It is clear that, by continuity, the genus of \hat{P}_γ is the same for all close enough desingularizations of G .

Definition 5.2 With a slight abuse of language we shall call *genus of the filling of P* this common value, and we shall denote it by $g_{\hat{P}}$.

Before further investigating (essentially) disconnecting cycles, we give some examples and counterexamples of possible boundary graphs, together with their desingularizations.

Example 5.1 Consider a single cycle γ_1 of saddle connections. Then, by Lemma 5.1, γ_1 locally disconnects S , and a geodesic σ having γ_1 as ω -limit set accumulates γ_1 locally on one side (say, staying inside an open set U_1). The definition of the desingularization of γ_1 gives a curve ω_1 in the same homology class of γ_1 , contained in (the closure of) U_1 . In particular, γ_1 disconnects S if and only if ω_1 does. So, for this cycle the properties of being disconnecting and of being essentially disconnecting are equivalent (see Lemma 5.2 below for a more general result).

Example 5.2 Let now γ_2 be the union of a non-disconnecting cycle γ_2^a and a small, homotopically trivial, disconnecting cycle γ_2^b , intersecting only at a common pole. We see that γ_2 disconnects S in two parts (the two given by γ_2^b). By Lemma 5.1, a geodesic σ_2 having γ_2 as ω -limit set must be contained in only one of these components. Because of the fact that it must accumulate both γ_2^a and γ_2^b , it cannot be contained in the small region bounded only by γ_2^b , and so it must be in the other one. So, we see that a desingularization of γ_2 is a curve ω_2 contained in the closure of this last region, and in particular that γ_2 is not essentially disconnecting.

Example 5.3 Let now γ_3 be the union of two non-disconnecting cycles γ_3^a and γ_3^b in the same homology class with exactly one vertex in common. We see that γ_3 is




Fig. 5.1: Desingularization of a graph

disconnecting, and in particular that the fundamental group of one of the two components (that we call U_3^1) has rank zero, while the fundamental group of the other component has at least rank 1 (since on the sphere we cannot find non-disconnecting cycles). A small neighbourhood of γ_3 is disconnected by it in three connected components, one contained in the component U_3^1 and two in the other component. The only one adherent to both γ_3^a and γ_3^b is the one contained in U_3^1 , and so a geodesic σ_3 having γ_3 as ω -limit set must live in U_3^1 . So, we see that the construction of the desingularization of γ_3 gives a curve ω_3 contained in the closure of U_3^1 . In particular, ω_3 disconnects S , and so γ_3 is essentially disconnecting (which is coherent with the idea that small perturbations of γ_3 still disconnect S).

Example 5.4 Finally, consider the cycle γ_3 of the previous example, and attach to it at the intersection point, as in Example 5.2, a small homotopically trivial disconnecting cycle γ_4^b not contained in U_3^1 . Lemma 5.1 then implies that the graph γ_4 so obtained cannot be the ω -limit set of any geodesic σ_4 on S . In fact, by the argument of the previous example, σ_4 should be contained in U_3^1 , and thus it could not accumulate γ_4^b , contradiction.

The next lemma describes an important class of essentially disconnecting boundary graphs:

Lemma 5.2 *Let G be a boundary graph of saddle connections in S , obtained as ω -limit set of a not self-intersecting geodesic, and such that every cycle of G disconnects S . Then G is essentially disconnecting.*

Proof Let U_σ be as in Lemma 5.1. Since, by assumption, every cycle in G disconnects S , the complement $S \setminus \mathcal{C}$ of any cycle \mathcal{C} in G has exactly two connected components, only one disjoint from U_σ ; let us color this component in black.

Furthermore, the fact that G is a boundary graph implies that every edge of G is contained in at most one cycle; as a consequence, $S \setminus G$ is formed by the part P of S containing U_σ with G as a boundary, and by the black connected components determined by the cycles in G .

Now we desingularize G to a curve γ in the following way, clearly equivalent to the construction above: near a pole connecting two (or more) cycles, we paint in black a little ball; and we replace each spike by a little black strip following its path. In this way we get a connected black region (see Figure 5.1).

Let γ be the boundary of the black region we have constructed in this way. Clearly γ is (homotopic to) a desingularization of G and, because it separates a black region from a white one, by definition γ disconnects S . It follows that G is essentially disconnecting. \square

So, we know that if every cycle is disconnecting then the graph is essentially disconnecting, while (generalizing Example 5.2 above) we can construct examples with an arbitrarily high number of disconnecting cycles (and at least one non disconnecting cycle) which are not essentially disconnecting.

We can now state the analogous of Equation (5.1) for case 4 of Theorem 4.3.

Theorem 5.1 *Let S be a compact Riemann surface and ∇ a meromorphic connection on S , with poles $p_1, \dots, p_r \in S$. Let $S^0 = S \setminus \{p_1, \dots, p_r\}$. Let $\sigma: [0, \varepsilon_0) \rightarrow S^0$ be a maximal geodesic for ∇ having as ω -limit set an essentially disconnecting boundary graph of saddle connections G . Let P the part of S having G as boundary and containing the support of σ . Then*

$$\sum_{p_i \in P} \operatorname{Re} \operatorname{Res}_{p_i}(\nabla) = -1 + 2g_{\hat{p}}, \quad (5.2)$$

where $g_{\hat{p}}$ is the genus of the filling of P (see Definition 5.2).

Proof Let U be a connected open neighbourhood of the support of σ as in Lemma 5.1; without loss of generality we can assume that the support of σ is completely contained in U_σ .

Consider a generic point $z_0 \in G \cap S^0$. Using the local isometry J near z_0 we see that we can find a geodesic $\tau: [0, \varepsilon) \rightarrow U_\sigma$ issuing from z_0 intersecting σ transversally infinitely many times and making an angle α close to $\pi/2$ with the geodesic segment in G passing through z_0 .

Let $p_0 \in U_\sigma$ be a point of intersection between σ and τ , and let $p_1 \in U_\sigma$ be the next (following σ) point of intersection between σ and τ ; necessarily $p_1 \neq p_0$ because σ has no self-intersections. The union of the segments of σ and τ between p_0 and p_1 is a geodesic 2-cycle γ contained in U_σ . The first observation is that the two external angles of γ must have opposite signs. Indeed, if they had the same sign (roughly speaking, if σ leaves and comes back to τ on the same side of τ) then γ would be the boundary of a part of S contained in U_σ , and thus containing no poles, against Remark 3.2. The second observation is that p_1 must be closer (along τ) to G than p_0 . Indeed, γ disconnects S , because G is essentially disconnecting; if p_1 were farther away from G than p_0 then the support of σ after p_1 would be contained in the connected component of $S \setminus \gamma$ disjoint from G , impossible (notice that after p_1 the

geodesic σ cannot intersect γ anymore, because σ has no self-intersections and any further intersection with the segment of τ contained between p_0 and p_1 would be on the same side of τ and thus can be ruled out as before).

Repeating this argument starting from p_1 we get a sequence $\{p_n = \sigma(t_n) = \tau(s_n)\}$ of points of intersections between σ and τ , with t_n increasing, s_n decreasing, and $p_n \rightarrow z_0$. Furthermore, if we denote by γ_n the geodesic 2-cycle bounded by the segments of σ and τ between p_n and p_{n+1} , we also know that the external angles of γ_n have opposite signs.

Since G is essentially disconnecting, for n large enough γ_n disconnects S ; let P_n be the connected component of $S \setminus \gamma_n$ not containing G . Notice that the poles contained in P_n are exactly the same contained in P , because $\gamma_n \subset U_\sigma \setminus G$. Furthermore, since the γ_n are eventually homotopic to desingularizations of G , for n large enough the genus of the filling of P_n becomes equal to $g_{\hat{p}}$, the genus of the filling of P .

Up to a subsequence, we can assume that the directions of $\sigma'(t_n)$ and $\sigma'(t_{n+1})$ converge to a direction in z_0 , necessarily the direction of the geodesic passing through z_0 contained in G . Since the external angles of γ_n have opposite signs, it follows that their sum must converge to 0. But, by Corollary 3.2 and the arguments above, this sum must be eventually constant; so it must be 0, and (5.2) follows recalling again Corollary 3.2. \square

In particular, [AT11, Theorem 4.6] is now a consequence of Theorems 4.3 and 5.1 (and Remark 4.4).

The formula (5.2) gives a necessary condition for the existence of geodesics on S having as ω -limit set an essentially disconnecting boundary graph of saddle connections. In particular, if no sum of real parts of residues is an odd integer number of the form $-1 + 2g$ then no such geodesic can exist on S .

The next statement contains a similar necessary condition for the existence of isolated vertices in a boundary graph of saddle connections:

Proposition 5.1 *Let S be a compact Riemann surface and ∇ a meromorphic connection on S . Let p be a pole belonging to a boundary graph of saddle connections G which is an ω -limit set for a not self-intersecting geodesic σ . Suppose that p is the vertex of only one arc ρ of the graph. Then*

$$\operatorname{Re} \operatorname{Res}_p(\nabla) = -\frac{1}{2}.$$

Proof Let τ be a geodesic crossing transversally ρ at a point q near p . The geodesic σ must intersect both sides (with respect to ρ) of τ infinitely many times. Indeed, if it would intersect eventually only one side, arguing as in the previous proof we would see that the external angles at consecutive intersections would have opposite signs; and this would force the existence of points in G close to p but not belonging to ρ , impossible.

So we can construct a sequence $\{\sigma_n\}$ of segments of σ connecting a point p_n on one side of τ to a point q_n on the other side of τ , with both points of intersection converging to q . Let γ_n be the geodesic 2-cycle consisting in σ_n and the segment of τ between p_n and q_n ; eventually, γ_n must be contained in a simply connected neighbourhood of p , and thus it is the boundary of a part P_n of S containing p and no other poles. Notice that the genus of the filling of P_n is zero.

An argument similar to the one used above shows that the external angles of γ_n must have the same sign. Up to a subsequence, we can also assume that the direction of σ both at p_n and at q_n converge to the same direction (which is the direction of ρ at q); thus the sum of the external angles of γ_n must tend to π . Applying Corollary 3.2 to P_n we get (that this sum must be constant and) that

$$\pi = 2\pi(1 + \operatorname{Re} \operatorname{Res}_p(\nabla)) ,$$

which gives the assertion. \square

So, in particular, if all the poles have the real part of the residue different from $-1/2$, the graph cannot have vertices belonging to one arc only — and thus in particular it cannot have spikes.

Remark 5.2 With the same argument as in Proposition 5.1 we see that if p is a vertex belonging only to two arcs that are spikes, then the real part of its residue must be zero.

6 Non-compact Riemann surfaces

In this section we shall generalize some of the results of Section 4 to non-compact Riemann surfaces. We shall limit ourselves to consider Riemann surfaces S that can be realized as an open subset of a compact Riemann surface \hat{S} . In this setting, all the results depending only on the local structure of geodesics still hold; for instance, Proposition 4.1, Lemmas 4.1 and 4.2, and Theorem 4.1 go through without any changes.

It is also easy to adapt Theorem 4.2. Indeed, the definition of singular line field on a compact Riemann surface \hat{S} used in [H81] is that of regular line field defined on an open subset S^0 of \hat{S} , and the singular set is by definition $\hat{S} \setminus S^0$. In particular, if we consider the ω -limit set W of a geodesic $\sigma: I \rightarrow S^0$ in \hat{S} (and not only in S) then $W \cap \partial S$ is automatically invariant with respect to any line field associated to σ . Therefore arguing as in the proof of Theorem 4.2 we get the following result.

Theorem 6.1 *Let S be an open connected subset of a compact Riemann surface \hat{S} . Let ∇ be a meromorphic connection on S , and denote by $S^0 \subseteq S$ the complement in S of the poles of ∇ . Let $\sigma: [0, \varepsilon_0) \rightarrow S^0$ be a maximal not self-intersecting geodesic for ∇ . Then the possible minimal sets for σ are the following:*

1. a pole of ∇ ;
2. a closed simple curve contained in S^0 , which is a closed geodesic for ∇ , and in this case the ω -limit set of σ coincides with this closed geodesic;
3. a point in $\partial S \subset \hat{S}$;
4. all of \hat{S} , and in this case \hat{S} is a torus.

Finally, in order to generalize Theorem 4.3 we need to consider saddle connections escaping to infinity, i.e., leaving any compact subset of S .

Definition 6.1 A *diverging saddle connection* for a meromorphic connection on a Riemann surface S is a maximal geodesic $\sigma: (-\varepsilon_-, \varepsilon_+) \rightarrow S^0$ satisfying one of the following conditions:

- (a) $\sigma(t)$ leaves every compact subset of S for both $t \rightarrow -\varepsilon_-$ and $t \rightarrow \varepsilon_+$; or
- (b) $\sigma(t)$ leaves every compact subset of S for $t \rightarrow -\varepsilon_-$ and tends to a pole for $t \rightarrow \varepsilon_+$;
or
- (c) $\sigma(t)$ tends to a pole for $t \rightarrow -\varepsilon_-$ and leaves every compact for $t \rightarrow \varepsilon_+$.

A *diverging graph of saddle connections* is a connected planar graph in S whose vertices are poles and whose arcs are disjoint saddle connections, with at least one of them diverging. A *spike* is a saddle connection of a graph which does not belong to any cycle of the graph.

A *diverging boundary graph of saddle connections* (or *diverging boundary graph*) is a diverging graph of saddle connections which is also the boundary of a connected open subset of S . A diverging boundary graph is *disconnecting* if its complement in S is not connected.

Using these definitions, the following theorem generalizes Theorem 4.3 describing the possible ω -limit sets of geodesics for a meromorphic connection on a non-compact Riemann surface of this kind.

Theorem 6.2 *Let S be an open connected subset of a compact Riemann surface \hat{S} . Let ∇ be a meromorphic connection on S , and denote by $S^0 \subseteq S$ the complement in S of the poles of ∇ . Let $\sigma: [0, \varepsilon_0) \rightarrow S^0$ be a maximal geodesic for ∇ . Then one of the following possibilities occurs:*

1. $\sigma(t)$ definitely leaves every compact subset of S (i.e., it tends to the boundary of S in \hat{S}); or
2. $\sigma(t)$ tends to a pole in S ; or
3. σ is closed; or
4. the ω -limit set of σ is the support of a closed geodesic; or
5. the ω -limit set of σ is a (possibly diverging) boundary graph of saddle connections; or
6. the ω -limit set of σ is all of $\bar{S} = S \cup \partial S$; or
7. the ω -limit set of σ has non-empty interior and non-empty boundary in S , and each component of its boundary is a (possibly diverging) graph of saddle connections with no spikes and at least one pole; or
8. σ intersects itself infinitely many times.

Furthermore, in cases 3 or 4 the support of σ is contained in only one of the components of the complement of the ω -limit set, which is a part P of S having the ω -limit set as boundary.

Proof Suppose that σ is not closed (case 3) and does not self-intersect infinitely many times (case 8). Up to changing the starting point, we can suppose that $\sigma: [0, \varepsilon_0) \rightarrow S^0$ does not self-intersect at all.

If σ definitely leaves every compact set of S , we are in case 1 and we are done. Otherwise, the ω -limit set W of σ must intersect S . We can then argue as in the proof of Theorem 4.3 using Theorems 4.1 and 6.1, and we are done. \square

Remark 6.1 When the ω -limit set is a non diverging graph it is possible to find conditions on the residues of the poles involved analogous to those described in Section 5, assuming it is disconnecting and the boundary of a relatively compact part.

7 Geodesics on the torus

In this section we study in detail the geodesics for holomorphic connections on the torus. We remark that, by Theorem 2.1, we cannot have holomorphic connections on Riemann surfaces different from the torus. Moreover, Proposition 4.2 tells us that this is the only case in which all the surface S can be a minimal set (and a minimal set for an associated line field, and thus Theorem 2.2 implies that the connection must be holomorphic).

So, our goals are: to characterize holomorphic connections on a torus, and to study their geodesics.

We recall that a complex torus can be realized as a quotient of \mathbb{C} under the action of a rank-2 lattice, generated over \mathbb{R} by two elements $\lambda_1, \lambda_2 \in \mathbb{C}$. Without loss of generality we can assume that one of the two generators is 1 and the other is a $\lambda \in \mathbb{C}$ with $\text{Im} \lambda > 0$. We shall denote by T_λ the torus associated to λ .

The next elementary result characterizes holomorphic connections on a torus.

Lemma 7.1 *Every holomorphic connection on a torus is the projection of a holomorphic connection on the cover \mathbb{C} represented by a constant global form adz , for a suitable $a \in \mathbb{C}$.*

Proof Let ∇ be a holomorphic connection on a torus T_λ . Since the tangent bundle of a torus is trivial, we can find a global holomorphic $(1,0)$ -form η representing ∇ by setting $\nabla e = \eta \otimes e$, where e is a global nowhere vanishing section of the tangent bundle.

Let $\pi: \mathbb{C} \rightarrow T_\lambda$ be the universal covering map, and let $\tilde{\eta} = \pi^* \eta$; it is a holomorphic $(1,0)$ -form on \mathbb{C} . Thus we can write $\tilde{\eta} = f dz$, for a suitable entire function f ; we have to prove that f is constant.

Let $\tau_t: \mathbb{C} \rightarrow \mathbb{C}$ be given by $\tau_t(z) = z + t$. Since $\pi \circ \tau_1 = \pi \circ \tau_\lambda = \pi$, it follows that $\tau_1^* \tilde{\eta} = \tau_\lambda^* \tilde{\eta} = \tilde{\eta}$. But this is equivalent to $f(z+1) = f(z+\lambda) = f(z)$ for all $z \in \mathbb{C}$, and thus f must be a bounded entire function, that is a constant. \square

The next step consists in studying the geodesics for a holomorphic connection ∇ on a torus. To do this, we shall study geodesics for the associated connection $\tilde{\nabla}$ on \mathbb{C} represented by a constant form adz and then project them to the torus (all geodesics on the torus are obtained in this way: see [AT11, Proposition 3.1]).

Let $\tilde{\sigma}$ be a geodesic for $\tilde{\nabla}$ issuing from a point $z_0 \in \mathbb{C}$ with tangent vector v_0 . We first consider the trivial case $a = 0$. In this case, the local isometry J is given by $J(z) = cz$, with $c \in \mathbb{C}^*$. So, applying equation (2.4) in Proposition 2.2 we obtain

$$c\tilde{\sigma}(t) = cv_0t + cz_0,$$

which means that $\tilde{\sigma}(t) = v_0t + z_0$ and the geodesics are the euclidean ones. Thus in this trivial case the geodesics on T_λ are exactly projections of straight lines.

If $a \neq 0$, the local isometry J is given by $J(z) = \frac{1}{a} \exp(az)$. We apply again equation (2.4) in Proposition 2.2 to get

$$\frac{1}{a} \exp(a\tilde{\sigma}(t)) = \exp(az_0)v_0t + \frac{1}{a} \exp(az_0),$$

which we can solve to obtain

$$\tilde{\sigma}(t) = z_0 + \frac{1}{a} \log(1 + av_0 t), \quad (7.1)$$

where \log is the branch of the logarithm with $\log 1 = 0$ defined along the half-line $t \mapsto 1 + av_0 t$.

So in general the geodesics for $\tilde{\nabla}$ are not straight lines, unless $av_0 \in \mathbb{R}$. However, every (projection of a) straight line in a torus is the support of a geodesic for a suitable (non-trivial) holomorphic connection:

Proposition 7.1 *Let T_λ be a complex torus, and $\pi: \mathbb{C} \rightarrow T_\lambda$ the usual covering map. Let ℓ be a straight line in \mathbb{C} issuing from $z_0 \in \mathbb{C}$ and tangent to $v_0 \in \mathbb{C}^*$. Then:*

- $\pi(\ell)$ is the support of a geodesic σ for the holomorphic connection ∇ represented by the global form $\eta = a dz$ with $a \in \mathbb{C}^*$ if and only if $v_0 \in \mathbb{R}\bar{a}$;
- $\pi(\ell)$ is the support of a closed geodesic σ for the holomorphic connection ∇ represented by the global form $\eta = a dz$ if and only if $v_0 \in \mathbb{R}\bar{a}$ and $\bar{a} \in \mathbb{R}^*(\mathbb{Z} \oplus \mathbb{Z}\lambda)$.

Proof For any $a \in \mathbb{C}^*$, the unique geodesic σ in T_λ issuing from $\pi(z_0)$ and tangent to $d\pi_{z_0}(v_0)$ is $\sigma = \pi \circ \tilde{\sigma}$, where $\tilde{\sigma}$ is given by (7.1). Then the support of σ coincides with $\pi(\ell)$ if and only if $\tilde{\sigma}$ is a parametrization of ℓ , and this happens if and only if $av_0 \in \mathbb{R}$, which is equivalent to $v_0 \in \mathbb{R}\bar{a}$.

Assume now $v_0 \in \mathbb{R}\bar{a}$. Then σ is closed if and only if $\tilde{\sigma}(t_0) = z_0 + m + n\lambda$ for suitable $t_0 \in \mathbb{R}^*$ and $m, n \in \mathbb{Z}$, that is if and only if

$$a(m + n\lambda) = \log(1 + av_0 t_0).$$

Recalling that $av_0 \in \mathbb{R}$, this can happen if and only if $\bar{a} = r(m + n\lambda)$ for a suitable $r \in \mathbb{R}^*$, and we are done. \square

Some of the geodesics given by Proposition 7.1 are closed, but none of them can be periodic. We prove this in the next proposition, where we characterize closed geodesics.

Proposition 7.2 *Let ∇ be a holomorphic connection on the torus T_λ represented by a global form $\eta = a dz$. Let $\sigma: [0, \infty) \rightarrow T_\lambda$ be a (non-constant) closed geodesic for ∇ . Then:*

- if $a \neq 0$ then the support of σ is the projection of a straight line parallel to \bar{a} in the covering space \mathbb{C} , and σ cannot be periodic;
- if $a = 0$ then σ is periodic.

Proof If $a = 0$, the geodesics are the euclidean ones and so, once σ is closed, it must also be periodic.

So let us study the problem when $a \neq 0$. In order for σ to be closed and non-trivial, (7.1) implies that we must have

$$\frac{1}{a} \log(1 + av_0 \bar{t}) = n + m\lambda$$

for suitable $\bar{t} \in \mathbb{R}^*$ and $n, m \in \mathbb{Z}$, not both zero, where $v_0 = \tilde{\sigma}'(0) \neq 0$ and $\tilde{\sigma}$ is the lifting of σ given by (7.1). Furthermore, since σ is closed then $\tilde{\sigma}'(\bar{t})$ must be parallel to $\tilde{\sigma}'(0)$. The derivative $\tilde{\sigma}'$ is given by

$$\tilde{\sigma}'(t) = \frac{v_0}{1 + av_0 t}; \quad (7.2)$$

so $1 + av_0 \bar{t} = e^{a(n+m\lambda)}$ must be real. Since \bar{t} is real too, it follows that av_0 must be real, and hence $\tilde{\sigma}$ is a parametrization of a straight line parallel to \bar{a} , as claimed.

Finally, σ is periodic if and only if $v_0 = \tilde{\sigma}'(0) = \tilde{\sigma}'(\bar{t})$; but by (7.2) this can happen only if $a = 0$, contradiction. \square

Remark 7.1 If the support of a geodesic σ of a holomorphic connection on a torus is the projection of a straight line, then σ is either closed or everywhere dense. In particular, Proposition 7.1 provides examples of cases 2 and 5 in Theorem 4.3.

In the last theorem of this section we study the geodesics which are not the projection of a straight line, and give a complete description of the possible ω -limit sets of geodesics for holomorphic connections on a torus.

Theorem 7.1 *Let ∇ be a holomorphic connection on the complex torus T_λ , represented by the global 1-form $\eta = a dz$. Let $\sigma: [0, T) \rightarrow T_\lambda$ be a maximal (non constant) geodesic for ∇ .*

- *If $a \neq 0$ then the ω -limit set of σ is (the closure of) the projection on T_λ of a straight line $l: \mathbb{R} \rightarrow \mathbb{C}$ of the form $l(t) = \bar{a}t + b$ for a suitable $b \in \mathbb{C}$. In particular:*
 1. *if $\bar{a} \in \mathbb{R}^*(\mathbb{Z} \oplus \lambda\mathbb{Z}) \setminus \{0\}$ then either*
 - σ *is closed, non periodic, and its support is the projection of the line l ;*
 - or*
 - σ *is not closed, but its ω -limit set is a closed geodesic whose support is the projection of the line l ;*
 2. *if $\bar{a} \notin \mathbb{R}^*(\mathbb{Z} \oplus \lambda\mathbb{Z}) \setminus \{0\}$ then the ω -limit set of σ is all of T_λ .*
- *If $a = 0$ then σ is the projection of a line of the form $l(t) = v_0 t + b$, where $v_0 = \tilde{\sigma}'(0)$. In particular:*
 - *if $v_0 \in \mathbb{R}^+(\mathbb{Z} \oplus \lambda\mathbb{Z})$ then σ is closed and periodic;*
 - *if $v_0 \notin \mathbb{R}^+(\mathbb{Z} \oplus \lambda\mathbb{Z})$ then σ is not closed and its ω -limit set is all of T_λ .*

Proof If $a = 0$, we know that the geodesics are the projection of euclidean straight lines, and the statement follows. So let us assume $a \neq 0$.

Let $\tilde{\sigma}: [0, T) \rightarrow \mathbb{C}$ be the lifting of σ to the covering space \mathbb{C} . By (7.1) the lifting $\tilde{\sigma}$ is given by

$$\tilde{\sigma}(t) = z_0 + \frac{1}{a} \log(1 + av_0 t) = z_0 + \frac{\bar{a} \log(1 + av_0 t)}{|a|^2},$$

where $v_0 = \tilde{\sigma}'(0)$. When $t \rightarrow T$ it is easy to see that $\text{Im} \log(1 + av_0 t) = \text{Arg}(1 + av_0 t)$ converges to a finite limit, while $\text{Re} \log(1 + av_0 t) = \log|1 + av_0 t|$ tends to $+\infty$ if $av_0 \notin \mathbb{R}^-$, to $-\infty$ otherwise. It follows that $\tilde{\sigma}$ is asymptotic to a line of the form $l(t) = \bar{a}t + b$ for a suitable $b \in \mathbb{C}$, and thus the ω -limit of σ is the closure of the projection of this line in T_λ . The rest of the assertion follows from Propositions 7.1 and 7.2. \square

Acknowledgements Both authors were partially supported by the FIRB 2012 project *Differential geometry and geometric function theory*. The first author has also been partially supported by GNSAGA (INdAM).

References

- [A01] M. Abate. The residual index and the dynamics of holomorphic maps tangent to the identity. *Duke Math. J.*, 107(1):173–207, 2001.
- [A10] M. Abate. Local discrete holomorphic dynamical systems. In *Holomorphic dynamical systems*. G. Gentili, J. Guenot, G. Patrizio eds., Lecture Notes in Math., Springer, Berlin, vol. 1998, pp. 1–55, 2010.
- [A15] M. Abate. Fatou flowers and parabolic curves. Preprint arXiv:1501.02176, 2015. To appear in *Complex Analysis and Geometry*. Bracci, F. et al eds, Springer Proceedings in Mathematics & Statistics, Vol. 144.
- [ABT04] M. Abate, F. Bracci and F. Tovena. Index theorems for holomorphic self-maps. *Annals of Math.*, 159(2):819–864, 2004.
- [ABT08] M. Abate, F. Bracci and F. Tovena. Index theorems for holomorphic maps and foliations. *Indiana Univ. Math. J.*, 57: 2999–3048, 2008.
- [AT03] M. Abate and F. Tovena. Parabolic curves in \mathbb{C}^3 . *Abstr. Appl. Anal.*, 2003(5):275–294, 2003.
- [AT11] M. Abate and F. Tovena. Poincaré-Bendixson theorems for meromorphic connections and homogeneous vector fields. *J. Differential Equations*, 251:2612–2684, 2011.
- [AR14] M. Arizzi and J. Raissy. On Écalle-Hakims theorems in holomorphic dynamics. In *Frontiers in complex dynamics*. Eds. A. Bonifant, M. Lyubich, S. Sutherland. Princeton University Press, Princeton, 2014, pp. 387–449.
- [B03] F. Bracci. The dynamics of holomorphic maps near curves of fixed points. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)*, 2(3), 493–520, 2003.
- [Br08] F.E. Brochero-Martínez. Dinámica de difeomorfismos dicríticos en $(\mathbb{C}^n, 0)$. *Rev. Semin. Iberoam. Mat.*, 3, 33–40, 2008.
- [BCL08] F.E. Brochero-Martínez, F. Cano and L. López-Hernanz. Parabolic curves for diffeomorphisms in \mathbb{C}^2 . *Publ. Mat.*, 52, 189–194, 2008.
- [C78] C. Camacho. On the local structure of conformal mappings and holomorphic vector fields in \mathbb{C}^2 . *Astérisque*, 59(60):83–94, 1978.
- [C12] K. Ciesielski. The Poincaré-Bendixson Theorem: from Poincaré to the XXIst century. *Cent. Eur. J. Math.*, 10(6):2110–2128, 2012.
- [É81a] J. Écalle. *Les fonctions résurgentes. Tome I: Les algèbres e fonctions résurgentes*, volume 81-05. Publ. Math. Orsay, 1981.
- [É81b] J. Écalle. *Les fonctions résurgentes. Tome II: Les fonctions résurgentes appliquées à l’itération*, volume 81-06. Publ. Math. Orsay, 1981.
- [É85] J. Écalle. *Les fonctions résurgentes. Tome III: L’équation du pont et la classification analytique des objets locaux*, volume 85-05. Publ. Math. Orsay, 1985.
- [H97] M. Hakim. Transformations tangent to the identity. Stable pieces of manifolds. *Université de Paris-Sud, Preprint*, 145, 1997.
- [H98] M. Hakim. Analytic transformations of $(\mathbb{C}^p, 0)$ tangent to the identity. *Duke Mathematical Journal*, 92(2):403–428, 1998.
- [H81] J. Hounie. Minimal sets of families of vector fields on compact surfaces. *J. Differential Geom.*, 16(4):739–744, 1981.
- [IY08] Y. Ilyashenko and S. Yakovenko. *Lectures on analytic differential equations*, volume 86. American Mathematical Society, Providence, RI, 2008.
- [LS14] L. López-Hernanz, F.S. Sánchez. Parabolic curves of diffeomorphisms asymptotic to formal invariant curves. Preprint arXiv:1411.2945, 2014
- [M06] J. Milnor. *Dynamics in one complex variable, 3rd ed.* Princeton University Press, Princeton, 2006.
- [M09] L. Molino. The dynamics of maps tangent to the identity and with nonvanishing index. *Trans. Amer. Math. Soc.*, 361(3):1597–1623, 2009.
- [R10] F. Rong. Absolutely isolated singularities of holomorphic maps of \mathbb{C}^n tangent to the identity. *Pacific J. Math.*, 246(2):421–433, 2010.

-
- [S63] A. J. Schwartz. A generalization of a Poincaré-Bendixson theorem to closed two-dimensional manifolds. *Amer. J. Math.*, 85(3):453–458, 1963.
- [S82] A.A. Shcherbakov. Topological classification of germs of conformal mappings with identical linear part. *Moscow Univ. Math. Bull.*, 37:60–65, 1982.
- [ST88] R.A. Smith and E.S. Thomas. Transitive flows on two-dimensional manifolds. *J. London Math. Soc. (2)*, 37:569–576, 1988.
- [V11] L. Vivas. Degenerate characteristic directions for maps tangent to the identity. *Indiana Univ. Math. J.*, 61:2019–2040, 2012.