



## Data in Brief

## Transcript profiling in the milk of dairy ewes fed extruded linseed



Tommaso Giordani, Alberto Vangelisti, Giuseppe Conte, Andrea Serra, Lucia Natali, Annamaria Ranieri, Marcello Mele, Andrea Cavallini \*

Department of Agriculture, Food and Environment, University of Pisa, Via del Borghetto 80, I-56124 Pisa, Italy

## ARTICLE INFO

## Article history:

Received 24 October 2016

Received in revised form 10 November 2016

Accepted 18 November 2016

Available online 19 November 2016

## ABSTRACT

The identification of genes regulating milk secretion by mammary glands is a key-step for exploiting changes in milk composition induced by different diet regimens. Linseed supplementation is a reliable feeding strategy to enhance polyunsaturated fatty acid content in milk fat from sheep, increasing milk quality and nutraceutical value. To investigate the molecular bases of diet induced differences in milk composition, we collected milk from dairy ewes at 90 day of lactation and after 3 weeks of diet supplementation with extruded linseed. The milk of dairy ewes contains milk somatic cells mostly derived by mammary glands. After isolating milk somatic cells by centrifugation, RNAs were purified from these cells, and Illumina RNA sequencing was performed to analyze RNA synthesis. Our data provide a resource (available at Gene Expression Omnibus database under GSE89163) to be employed for comparative analyses of gene expression in milk somatic cells in different breeds and different diets, with the long-term aim of developing strategies to improve sheep milk quality.

© 2016 Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Specifications	
Organism/cell line/tissue	<i>Ovis aries</i> /Comisana breed/milk somatic cells
Sex	F
Sequencer or array type	Illumina HiSeq2000
Data format	Raw data: FASTQ files, analyzed data: txt files
Experimental factors	Milk somatic cells before vs. after 3 weeks linseed feeding
Experimental features	RNA-Seq dataset for gene expression profiling in sheep milk somatic cells
Consent	Full consent
Sample source location	Perugia, Italy

## 1. Direct link to deposited data

Deposited data can be found here: <http://www.ncbi.nlm.nih.gov/geo/info/linking.html>.

\* Corresponding author.

E-mail address: [andrea.cavallini@unipi.it](mailto:andrea.cavallini@unipi.it) (A. Cavallini).

## 2. Experimental design, materials and methods

## 2.1. Sample collection and RNA isolation

To analyze transcriptome in the mammary gland a biopsy would be necessary, but this technique is particularly invasive and it can cause udder damage and infection [1]. Using milk somatic cells (MSC) is preferable for RNA isolation [2]. For transcriptomic analyses on sheep milk somatic cells, five Comisana mid-lactation sheep were used. Starting in February 2015, each sheep was fed grass hay and a commercial concentrate for three weeks, then the same sheep were fed grass hay and a concentrate containing 20% of extruded linseed on dry matter basis for 3 weeks. In both periods, the diets were formulated according to the nutrient requirements of an ewe weighing 65 kg and producing 1 kg of milk at 6.5% fat. Animal health was constantly monitored and since some ewes turned sick, they were excluded from the experiment. Milk from each sheep was collected at the beginning and at the end of the experiment. Total RNA was isolated from MSC, according to the method described by Wickramasinghe et al. [3], with minor modifications [4]. Briefly, MSC were pelleted from 250 ml fresh milk at 1100 g for 10 min at 4 °C. The cell pellet was washed twice in phosphate-buffered saline solution, pH 7.2 with 0.5 mM EDTA and resuspended in 3 ml of Trizol reagent (Life Technologies). RNA extraction continued following the Trizol protocol, except that all centrifugations were performed at 4 °C and precipitation in isopropanol was carried on at –80 °C for 30 min. Then, a DNase I (Roche) treatment was performed according to the manufacturer's instructions, to completely remove genomic

**Table 1**

Number of raw and high-quality Illumina reads, and number of reads matching the *Ovis aries* transcriptome for each library (C, control; L, after 3 weeks linseed feeding).

Library	Total reads	HQ reads	Aligned HQ reads	Aligned HQ reads (%)
C1	28,098,370	25,682,138	22,644,050	80.59
C2	15,834,738	12,902,452	8,805,705	55.61
C3	18,805,590	14,994,960	10,602,801	56.38
L1	17,380,388	5,973,508	4,941,935	28.43
L2	20,894,652	8,478,368	5,978,026	28.61
L3	26,732,292	24,451,846	19,912,996	74.49
Total	127,746,030	92,483,272	72,885,513	57.06

DNA contamination. Finally, RNA was purified by phenol/chloroform extraction and precipitated following standard procedures.

The handling of the animals was carried out according to the guidelines of the EU Directive 2010/63/EU for animal experiments and to the Institutional Animal Care and Use Committee of the University of Perugia.

## 2.2. Generation of RNA-Seq data

RNA-Seq libraries were generated using the TruSeq RNA-Seq Sample Prep kit according to the manufacturer's protocol (Illumina Inc., San Diego, CA). Poly-A RNAs were isolated from total RNA and chemically fragmented. First and second strand cDNA synthesis were followed by end-repair and adenines were added to the 3' ends. Adapters were ligated to the cDNA and 200 ± 25 bp fragments were gel purified and enriched by PCR. The libraries were quantified using Bioanalyzer 2100 (Agilent Technologies, Santa Clara, CA) and run on the Illumina HiSeq2000 (Illumina Inc.) using version 3 reagents. Paired-read sequences, 125 nt in length, were collected. Whole RNA-Seq data were submitted to NCBI Sequence Read Archive and Gene Expression Omnibus (series accession number GSE89163).

## 2.3. Data processing

Raw single reads (in FASTQ format) were subjected to sequence quality control using FastQC (<http://www.bioinformatics.babraham.ac.uk/projects/fastqc/>). FastQC performed a series of analysis modules on raw data and created a report with statistics for the data analyzed. For each library, FastQC showed high per base sequence quality, exceeding 30 on Phred scale (<1/1000 chance of a base being wrong) and detected adapter contamination, matching the reads to known adapter sequences. Then, raw reads were trimmed using Trimmomatic [5], version 0.33, with the following parameters: CROP:115 HEADCROP:15 MINLEN:100. After trimming, ribosomal reads were mapped against

*Ovis aries* and *Bos taurus* ribosomal DNA sequences and rDNA matching reads were removed.

Finally, FastQC was used again to examine the characteristics of the libraries and to verify trimming efficiency, which resulted into removal of a total of 54,860,517 (42.94%) low quality or ribosomal sequences (Table 1).

## 2.4. Differential expression quantification


Reads were aligned to the *Ovis aries* transcriptome (Oar), version 4.0, available at the NCBI site (<https://www.ncbi.nlm.nih.gov/genome/?term=ovis%20aries>) [6], with a tolerance of up to 2 mismatches, using CLC-BIO Genomic Workbench 8.0.3 (CLC). The percentage of aligned reads per each sample is reported in Table 1.

The sample cluster of linseed feeding in determining gene expression differences was verified applying a Multi Dimensional Scaling (MDS) plot to the gene expression data of the whole set of genes (Fig. 1). MSC before and after linseed feeding formed two separate groups.


Gene expression level was calculated and expressed as Reads Per Kilobase per Million reads mapped (RPKM [7]). Expression data were evaluated considering RPKM values before and after linseed feeding using likelihood test on edgeR [8]. The log fold changes between treatments were considered as significant when the weight of a sample was at least one-fold higher or lower than another, with an FDR corrected *p*-value ≤ 0.05. The analysis was limited to genes showing RPKM > 1 in at least one sample, as used in other studies [9]. Of a total of 22,328 analyzed genes, 1421 resulted over-expressed and 1343 under-expressed in ewes after linseed feeding compared to the same sheep before linseed feeding. Genes affected by treatment were annotated by gene ontology analysis using BLAST2GO [10]. The most represented GOs are reported in Fig. 2.

## 3. Discussion

It is known that feeding regimen plays a pivot role in affecting milk nutrient composition. Also milk nutraceutical compounds (products derived from food sources that are purported to provide extra health benefits) can change with animal diet. For example, linseed supplementation in the diet of dairy ewes resulted in an increase of milk polyunsaturated fatty acids, including conjugated linoleic acids (CLA) and alpha-linolenic acid (ALA) [11]. Cheese from sheep milk enriched with CLA and ALA was proven to be effective in reducing plasma cholesterol and endocannabinoids in human [12]. These data prompted us to study mammary gland gene expression of dairy ewes, as affected by linseed-enriched diet.



**Fig. 1.** Multidimensional scaling plot of sheep libraries showing cluster of samples. (C, ewes before linseed feeding; L, ewes after linseed feeding). Each ewe is identified by a number.



**Fig. 2.** The most represented ( $n > 10$ ) gene ontologies of differentially over- (above) and under-expressed genes (below) in milk somatic cells after 3 weeks of linseed feeding. Columns are separated into molecular functions (blue), biological processes (green), and cellular components (yellow).

The expression of 22,328 genes included in the *O. aries* transcriptome database was studied in milk somatic cells to infer the effect of linseed feeding on gene expression in mammary glands. Linseed feeding determined deep variation in the overall expression of genes, as indicated by MDS plot. Most detected genes were expressed before and after linseed feeding, however a number of genes were significantly induced or repressed by linseed, similar to results obtained in the liver of cows fed with CLA [13].

Expressed genes were characterised by gene ontology. Considering the three principal gene ontologies, molecular functions, biological processes, and cellular components, the most represented ontologies of genes induced by linseed feeding were Ion Binding, Response to Stress, and Protein Complex, respectively. On the other hand, the most common ontologies of repressed genes were Ion Binding, Biosynthetic Process, and Cytoplasm, respectively.

The biological significance of this finding remains to be elucidated in the future. The dataset of expression data will help improving our understanding of molecular mechanisms regulating sheep milk quality as affected by different diets.

### Conflict of interest

Authors declare no conflict of interest.

### Acknowledgements

This research work was supported by University of Pisa, Italy, project “Progetto di Ricerca di Ateneo: Metodi innovativi per l’analisi del trascrittoma mammario di pecore alimentate con semi di lino”. Thanks are due to Prof. Mariano Pauselli (University of Perugia) for ensuring availability of sheep and to Dr. Giulia Pasqualetto (University of Pisa) for help in RNA isolation, respectively.

### References

- [1] C.M. Murrieta, B.W. Hess, E.J. Scholljegerdes, T.E. Engle, K.L. Hossner, G.E. Mossand, D.C. Rule, Evaluation of milk somatic cells as a source of mRNA for study of lipogenesis in the mammary gland of lactating beef cows supplemented with dietary high-linoleate safflower seeds. *J. Anim. Sci.* 84 (2006) 2399–2405.
- [2] M.C. Mura, C. Daga, S. Bodano, M. Paludo, S. Luridiana, M. Pazzola, M.L. Dettori, G.M. Vacca, V. Carcangiu, Development of a RNA extraction method from milk for gene expression study in the mammary gland of sheep. *Mol. Biol. Rep.* 40 (2013) 2169–2173.
- [3] S. Wickramasinghe, G. Rincon, A. Islas-Trejo, J.F. Medrano, Transcriptional profiling of bovine milk using RNA sequencing. *BMC Genomics* 13 (2012) 45.
- [4] A. Vangelisti, T. Giordani, L. Natali, A. Serra, G. Conte, M. Mele, A. Ranieri, A. Cavallini, RNA isolation from the milk of dairy ewes for transcriptome analysis. *Agrochimica* 60 (2016) 320–324.
- [5] A.M. Bolger, M. Lohse, B. Usadel, Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30 (2014) 2114–2120.
- [6] The International Sheep Genomic Consortium, The sheep genome reference sequence: a work in progress. *Anim. Genet.* 5 (2010) 449–453.
- [7] A. Mortazavi, B.A. Williams, K. McCue, L. Schaeffer, B. Wold, Mapping and quantifying mammalian transcriptomes by RNA-Seq. *Nat. Methods* 5 (2008) 621–628.
- [8] M.D. Robinson, D.J. McCarthy, G.K. Smith, edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* 26 (2010) 139–140.
- [9] R.M. Cossu, T. Giordani, A. Cavallini, L. Natali, High-throughput analysis of transcriptome variation during water deficit in a poplar hybrid: a general overview. *Tree Genet. Genomes* 10 (2014) 53–66.
- [10] A. Conesa, S. Götz, J.M. Garcia-Gomez, J. Terol, M. Talon, M. Robles, Blast2GO: a universal tool for annotation, visualization, and analysis in functional genomics research. *Bioinformatics* 10 (2005) 3674–3676.
- [11] M. Mele, G. Contarini, L. Cercaci, A. Serra, A. Buccioni, M. Povolò, G. Conte, A. Funaro, S. Banni, G. Lercker, P. Secchiari, Enrichment of Pecorino cheese with conjugated linoleic acid by feeding dairy ewes with extruded linseed: effect on fatty acid and triglycerides composition and on oxidative stability. *Int. Dairy J.* 21 (2011) 365–372.
- [12] S. Pintus, E. Murru, G. Carta, L. Cordeddu, B. Batetta, S. Accossu, D. Pistis, S. Uda, G. Elena, M. Ghiani, M. Mele, P. Secchiari, G. Almerighi, P. Pintus, S. Banni, Sheep cheese naturally enriched in  $\alpha$ -linolenic, conjugated linoleic and vaccenic acids improves the lipid profile and reduces anandamide in the plasma of hypercholesterolaemic subjects. *British J. Nutr.* 109 (2013) 1453–1462.
- [13] R. Ringseis, W. Windisch, K. Eder, Transcript profiling in the liver of early-lactating dairy cows fed conjugated linoleic acid. *Genomics Data* 10 (2016) 101–103.