



# The Open Medicinal Chemistry Journal

Content list available at: [www.benthamopen.com/TOMCJ/](http://www.benthamopen.com/TOMCJ/)

DOI: 10.2174/187410450171101????



## RESEARCH ARTICLE

# Synthesis and Functional Evaluation of Novel Aldose Reductase Inhibitors Bearing a Spirobenzopyran Scaffold

Maria Digiacoimo<sup>§</sup>, Stefania Sartini<sup>§</sup>, Giulia Nesi, Simona Sestito, Vito Coviello, Concettina La Motta<sup>\*</sup> and Simona Rapposelli<sup>\*</sup>

Department of Pharmacy, University of Pisa, Via Bonanno 6, 56126 Pisa, Italy

Received: October 31, 2016

Revised: December 22, 2016

Accepted: January 09, 2017

### Abstract:

#### Background:

Aldose reductase, the first enzyme of the polyol pathway, is the key determinant for the pathogenesis of long term diabetic complications. Accordingly, its inhibition represents the major therapeutic strategy to treat this kind of pathologies.

#### Objectives:

In this work we describe the synthesis and the functional evaluation of a number of spiro-oxazolidinone and spiro-morpholinone acetic acid derivatives, and their benzyloxy analogs, developed as aldose reductase inhibitors.

#### Results:

Most of them proved to inhibit the target enzyme, showing  $IC_{50}$  values in the micromolar/low micromolar range. SARs observed among the three different series allowed to highlight their key pharmacophoric elements, thus creating sound basis for the design of novel and more effective inhibitors.

#### Conclusion:

Although further substitution patterns are needed, the novel compounds here proposed represent a good starting point for the development of novel and effective ARIs.

**Keywords:** Aldose reductase enzyme, Diabetic complications, Spiromorpholinone derivatives, Aldose reductase inhibitors (ARI), Spiro-oxazolidinones, Spirobenzopyran derivatives, ARL2.

## INTRODUCTION


Diabetes mellitus (DM) is one of the most common chronic metabolic disorders, characterized by elevated levels of blood glucose (hyperglycemia). According to “WHO Global report on diabetes”, the number of people with diabetes has risen from 108 million in 1980 to 422 million in 2014 [1]. The DM etiopathology is very complex and is closely related to the onset of chronic complications such as neuropathy, nephropathy, cataracts, retinopathy, accelerated atherosclerosis, and increased cardiovascular risk [2]. Therefore, nowadays, this disease is reasonably considered as a public health issue.

Many studies suggested the inhibition of aldose reductase (ARL2) enzyme as the key therapeutic strategy to prevent

\* Address correspondence to these authors at the Department of Pharmacy, University of Pisa via Bonanno, 6 56126 Pisa, Italy; Tel: (+) 390502219582, (+) 390502219593; E-mails: [simona.rapposelli@unipi.it](mailto:simona.rapposelli@unipi.it), [concettina.lamotta@unipi.it](mailto:concettina.lamotta@unipi.it)

<sup>§</sup> These authors equally contributed.

and/or ameliorate long-term diabetic complications [2]. ARL2 is the first enzyme of the polyol pathway; it is located in cytosol and it was found in eye lens, retina kidney, adrenal gland and some reproductive organs. ARL2 catalyzes the NADPH-dependent conversion of glucose to sorbitol, the rate-limiting step of the polyol pathway Fig. (1) [3]; then the sorbitol produced is converted into fructose by sorbitol dehydrogenase (SDH). Since its hydrophilic nature, this sugar alcohol does not pass easily through cell membranes, thus accumulating within the cells while being slowly converted into fructose. The accumulation of sorbitol leads to osmotic stress in cells, event strongly related to the increasing of  $\text{NAD}^+/\text{NADH}$  ratios. As a consequence, an impairment of cellular defense against oxidative injury occurs, leading to the onset of both micro- and macrovascular complications such as retinopathy, peripheral vascular disease and coronary artery disease. These events are frequently found in DM and are well known as “complications of secondary diabetes”. Moreover, many studies demonstrated an ARL2-dependent increase of ROS levels in cells cultured under high glucose conditions, thus confirming the leading role of this enzyme in the pathogenesis of many diabetic complications [4].



**Fig. (1).** Polyol pathway of glucose metabolism.

Therefore, ARL2 is considered an interesting target to counteract DM complications. In the last years, many ARL2 inhibitors (ARI) have been developed [5, 6] and some of them reached clinical trials to prove their efficacy in preventing and/or modulating/ameliorating DM complications. The most important ARIs can be classified into two chemical groups: (a) *cyclic imides*, such as sorbinil, fidarestat, minalrestat and ranirestat (AS-3201) [7] and (b) *acetic acid derivatives*, such as tolrestat, epalrestat, zopolrestat and ponalrestat [8]. To date, only the carboxylic acid epalrestat is available on the market and used for the treatment of diabetic neuropathy in Japan, India and China [9]. Most of the ARIs that showed to be active in *in vitro* tests failed the subsequent clinical trials, mainly because of their pharmacokinetic drawbacks and, consequently, low *in vivo* efficacy, besides the onset of adverse side effects. Notably, the side effects are principally due to the lack of selectivity of ARI toward aldehyde reductase (ALR1, EC 1.1.1.2). This last enzyme, ALR1, plays a detoxification role in specifically metabolizing toxic aldehydes such as hydroxynonenal (HNE), 3-deoxyglucosone, and methylglyoxal [10]. Zopolrestat, which belongs to the carboxylic acid derivatives, is quite selective for ALR2 vs ALR1 and other enzymes. However, the carboxylic class of agents becomes highly protein bound *in vivo* thus limiting their efficacy *in vivo*. On the other hand, ARIs bearing a hydantoin core, including sorbinil, are relatively non-selective and inhibit ALR1 and ALR2 with comparable efficacies.

Despite being chemically different, ARIs respond to some structural requirements represented by an acidic group, which interacts with the anionic site of the catalytic site, and a lipophilic group, binding the hydrophobic pocket, necessary for the activity and selectivity towards the enzyme [11].

In a previous work we synthesized new spirobenzopyran derivatives (compounds type **A**, Fig. 2), as effective ARIs [12]. The previously synthesized ALR2 inhibitors possessed  $\text{IC}_{50}$  values in the submicromolar range and a good selectivity against the target enzyme.

Starting from these encouraging results, herein we performed a SAR study to investigate the effects on the ARL2 inhibitory activity induced by small structural modifications on the scaffold of type **A** compounds (Fig. 2).

In particular, we synthesized the analogues with *gem*-dimethyl group in 2-position of the chromane-scaffold (**5-10**) to evaluate the influence of the steric hindrance in this position on the interaction with the active site of the enzyme. Then, in order to assess the importance of the benzopyran scaffold and the steric effects of substituents in the 2-position

of the core, we synthesized compounds **11-13**, in which the C2-C3 bond of the benzopyran nucleus has been cleaved, and compounds **5-10**, in which the pendant phenyl ring on the benzopyran core has been replaced by *gem*-dimethyl groups.





Fig. (2). Medicinal chemistry optimization of spirobenzopyran derivatives.

The new molecules synthesized were evaluated *in vitro* for their affinity and potency against the target enzyme, ARL2.

## RESULTS AND DISCUSSION

### Chemistry

The acetic acid derivatives **3,4** have been synthesized as reported in Scheme (1). Briefly, the reaction of aminoalcohols **14,15** [12] and chloroacetyl chloride in a heterogeneous phase yielded the chloroacetamides **16,17** which were submitted to a base-catalyzed (*t*-BuOK) cyclization to afford the spiro-morpholone derivatives **18,19**. The subsequent reaction of **18,19** with ethyl bromoacetate and NaH in DMF gave the corresponding esters, which were then cleaved in high yields to the carboxylic acids **3** and **4**, respectively.




Scheme 1. Reagents and Conditions: (a) Chloroacetyl chloride, NaOH, CH<sub>2</sub>Cl<sub>2</sub>/H<sub>2</sub>O, rt, 1 h; (b) *t*-BuOK, toluene, rt, 3 h; (c) NaH, ethyl bromoacetate, DMF, N<sub>2</sub>, 0°C, rt, 1.5 h; (d) KOH, MeOH, reflux, 2 h.

A tandem reaction of spiro-oxazolidinones **22,23** [13] and spiro-morpholones **24,25** [14] with ethyl bromoacetate and the following cleavage of the intermediates (**26-29**), performed with KOH in MeOH, yielded the carboxylic acids **5-8** as the final products (Scheme 2A).


Spiromorpholine derivative **30** was obtained from **24** by reduction with LiAlH<sub>4</sub>. Notably, to avoid the possible halogen/metal exchange induced by LiAlH<sub>4</sub>, the 6-bromide-spiromorpholine **31** was obtained by a selective reduction of **25** [15] performed with a borane-methyl sulfide complex which can tolerate different substituents such as halogens or nitro-groups [16 - 19]. Subsequently, the reaction of products **30,31** with ethyl bromoacetate in the presence of K<sub>2</sub>CO<sub>3</sub> gave compounds **32,33**, which were then cleaved to the corresponding carboxylic acids **9,10** by the use of KOH/MeOH

mixture (Scheme 2B).



**Scheme 2.** Reagents and Conditions: (a) Ethyl bromoacetate, NaH, DMF, N<sub>2</sub>, 0°C, rt, 1.5 h; (b) KOH, MeOH, reflux, 2 h; (c) Ethyl bromoacetate, K<sub>2</sub>CO<sub>3</sub>, MeCN, reflux, 12 h.

Compounds **11-13** were synthesized starting from 2-hydroxyacetophenone and the appropriate benzyl chloride. Benzyloxyphenylethanones **34-36** were subjected to a nucleophilic addition with trimethylsilyl cyanide (TMSCN), in the presence of zinc iodide (ZnI<sub>2</sub>), to give the corresponding trimethylsilyl cyanohydrins **37-39**. The reduction of **37-39**, followed by a cyclization reaction with carbonyl diimidazole (CDI) gave the spirooxalidinone derivatives **43-45**, which were then submitted to the same reaction of alkylation and cleavage above reported for compounds **9,10** to yield the final products as carboxylic acid **11-13** (Scheme 3).



**Scheme 3.** Reagents and Conditions: (a) Appropriate methoxybenzyl chloride, KOH, DMSO, rt, 4 h; (b) TMSCN, ZnI<sub>2</sub>, CH<sub>2</sub>Cl<sub>2</sub>, rt, 4 h; (c) LiAlH<sub>4</sub>, THF, rt, 4 h; (d) CDI, THF, rt, 4 h; (e) Ethyl bromoacetate, *n*-BuLi, THF, N<sub>2</sub>, -78 °C, rt, 12 h; (f) KOH, MeOH, reflux, 2 h.

## Biological Evaluation

Recently, we developed a new class of small molecules bearing a benzopyran scaffold that displayed inhibitory activity against ARL2, with a high degree of selectivity when compared to the effects induced against ARL1 [12].

Among the previously studied derivatives, compounds **1** ( $IC_{50}$  2.25  $\mu$ M) and **2** ( $IC_{50}$  0.58  $\mu$ M), depicted in Fig. (3), shown the best results in terms of potency against the target enzyme.




Fig. (3). Spirooxazolidine Derivatives **1** and **2**.

Previous studies highlighted that the ARL2 inhibition activity was influenced by the presence of both a bromine atom on the benzopyran scaffold and the substituent in the 2-position of the chromane nucleus. With the aim to extend the structure-activity relationship (SAR) study of this class of compounds, herein we describe the synthesis of new molecules, **3-13**, in which a series of structural modifications have been performed as following reported: (a) replacement of the spirooxazolidine ring in the core structure of **1** and **2** with a spiro-morpholone (derivatives **3** and **4**) or with a spiro-morpholine nucleus (derivatives **9** and **10**); (b) replacement of the phenyl ring in the 2-position of the benzopyran scaffold with small lipophilic groups such as the methyl ones (derivatives **5-10**) and (c) change in the conformational freedom of spirooxazolidine derivatives **1** and **2** through the cleavage of the C2-C3 bond within the benzopyran core (derivatives **11-13**).

All the synthesized compounds were evaluated for their inhibitory activity against ARL2 and for their selectivity against aldehyde dehydrogenase (ALR1). Sorbinil, a well-known ARL2 inhibitor characterized by a spiro-chromane core, was used as the reference standard. Results obtained are reported in (Tables **1** and **2**), expressed as  $IC_{50}$  values.

Table 1. ARL2 inhibitory activities of compounds 1-10.


Compounds	N	Y	X	R	R'	$IC_{50}$ ( $\mu$ M) <sup>a</sup>
Sorbinil						0.65
<b>1</b> <sup>b</sup>	0	CO	H	4-OMe-Ph	H	2.25
<b>2</b> <sup>b</sup>	0	CO	Br	4-OMe-Ph	H	0.58
<b>3</b>	1	CO	H	4-OMe-Ph	Me	n.a.
<b>4</b>	1	CO	Br	4-OMe-Ph	Me	4.97
<b>5</b>	0	CO	H	Me	Me	42.3
<b>6</b>	0	CO	Br	Me	Me	17.3
<b>7</b>	1	CO	H	Me	Me	2.00
<b>8</b>	1	CO	Br	Me	Me	n.a.
<b>9</b>	1	CH <sub>2</sub>	H	Me	Me	n.a.
<b>10</b>	1	CH <sub>2</sub>	Br	Me	Me	4.17

<sup>a</sup> $IC_{50}$  values represent the concentration required to produce 50% enzyme inhibition. Standard error of the mean (SEM) is  $\leq 10$ . <sup>b</sup>See reference [17]. n.a.: not active. No inhibition was observed up to 100  $\mu$ M of test compounds.

The synthesized compounds exhibited  $IC_{50}$  values in the micromolar/low micromolar range, with the only exception of compounds **3**, **8** and **9** which proved to be completely inactive.

Noteworthy, results depicted in Table 1 show that the replacement of the spirooxazolidine nucleus of the leads, **1** and **2** [17], with the wider spiomorpholone core, like in compounds **3** and **4**, induced a decrease of inhibitory potency against ARL2. Actually, compound **3**, the higher homologue of derivative **1**, turned out to be devoid of any inhibitory activity while compound **4**, the higher homologue of **2**, showed an almost 8-fold reduction in inhibitory efficacy when compared to the smaller parent (**4**,  $IC_{50}$  4.97 vs **2**,  $IC_{50}$  0.58  $\mu$ M).

**Table 2. ARL2 inhibitory activities of compounds 10-12.**

			
Compounds	R	R <sup>1</sup>	$IC_{50}(\mu\text{M})^a$
Sorbinil			0.65
<b>11</b>	H	H	4.13
<b>12</b>	H	OMe	nt
<b>13</b>	OMe	OMe	3.19

<sup>a</sup> $IC_{50}$  values represent the concentration required to produce 50% enzyme inhibition. Standard error of the mean (SEM) is  $\leq 10$ . n.t.: not tested.

We then decided to further investigate the effect of the substituent in the 2-position of the benzopyran core. In particular, we replaced the *p*-methoxyphenyl ring with a *gem*-substitution, in which two methyl groups were added. Regarding the spirooxazolidine series, this kind of chemical manipulation induced a significant decrease in the inhibitory potency of the resulting compounds, **5** and **6**, even if the presence of a 6-bromine atom on the chromane core, like in **6**, helped to keep the activity in the low micromolar range ( $IC_{50}$  17.3  $\mu$ M). On the contrary, the same substitution pattern on the spiomorpholone series (derivatives **7** and **8**) gave rise to contrasting results. Actually, while the *gem*-dimethyl substituents conferred a significant inhibitory activity to compound **7** ( $IC_{50}$  2.0  $\mu$ M), the concomitant presence of a 6-bromine atom on the chromane core, like in **8**, induced a dramatic loss of activity.

With the aim of reducing the conformational restriction of the spiomorpholone derivatives **7** and **8**, we synthesized the spiomorpholine analogs, **9** and **10**. This kind of structural modification was able to restore the inhibitory activity of the 6-bromine substituted **10** ( $IC_{50}$  =4.17 $\mu$ M). This result let us speculate that further improvement in the conformational freedom of this kind of compounds could enhance their inhibitory potency. On this basis, we replaced the benzopyran scaffold with the more flexible benzyloxy nucleus, thus synthesizing compounds **11-13**.

As reported in Table 2, the novel compounds turned out to be almost 1.5-fold less potent than the parent **1**, demonstrating that the cleavage of the C2-C3 bond of the previous synthesized compound **1** induced only a slight decrease of ARL2 inhibitory potency. Reasonably, the inhibitory activity of the novel series could be modulated by introducing suitable substituents on the benzyloxy nucleus. Accordingly, further investigation with both electron-withdrawing and electron-donating groups is needed in order to investigate their role in the identification of novel and more effective ARL2 inhibitors.

All the synthesized compounds, **3-13**, were also tested for their ability to inhibit ALR1. Unlike the spiro-derivative sorbinil, that showed a significant inhibitory activity against ALR1 ( $IC_{50}$  0.029  $\mu$ M), no inhibition was observed up to 100  $\mu$ M of test compounds (data not shown), thus our compounds proved to be completely selective for the target ALR2.

## MATERIALS AND METHODS

### Chemistry

Melting points were determined on a Kofler hot-stage apparatus and are uncorrected. <sup>1</sup>H NMR spectra were obtained with a Varian Gemini 200 MHz spectrometer whereas <sup>13</sup>C NMR were obtained with a Bruker TopSpin 3.2 400 MHz spectrometer. Chemical shifts (δ) are reported in parts per million (ppm) downfield from tetramethylsilane and referenced from solvent references. The elemental compositions of the compounds agreed to within 0.4% of the calculated value. Chromatographic separation was performed on silica gel columns by flash (Kieselgel 40, 0.040–0.063 mm; Merck). Reactions were followed by thin layer chromatography (TLC) on Merck aluminum silica gel (60 F254) sheets that were visualized under a UV lamp. Evaporation was performed in vacuo (rotating evaporator). Sodium sulfate was always used as the drying agent. Commercially available chemicals were purchased from Sigma-Aldrich.

### General Procedure for the Synthesis of Compounds 3-13

To a stirred solution of appropriate ester **20-21**, **26-29**, **32-33**, **46-48** (0.77 mmol) in MeOH (3 mL) was added KOH 50% (0.005 mL) and the resulting mixture was refluxed for 2 h. The solvent was evaporated under vacuum and then HCl 1N was added (pH = 3). The precipitate was filtered and dried.

#### 2-(2-(4-Methoxyphenyl)-2-methyl-5'-oxospiro[chroman-4,2'-morpholin]-4'-yl) acetic acid3

Compound **3** was obtained from **20** (0.47 g, 1.10 mmol) following procedure previously described without further purification. **3** (0.09 g, 0.23mmol, yield 21%). Mp 175-178°C. <sup>1</sup>H-NMR (DMSO-*d*<sub>6</sub>): δ 1.63 (s, 3H, Me); 2.30 (d, 1H, *J* = 12.6 Hz, CH<sub>2</sub>); 2.42 (d, 1H, *J* = 14.3 Hz, CH<sub>2</sub>); 2.92 (d, 1H, *J* = 14.3 Hz, CH<sub>2</sub>); 3.25 (d, 1H, *J* = 17.2 Hz, CH<sub>2</sub>); 3.62 (d, 1H, *J* = 12.6 Hz, CH<sub>2</sub>); 3.70 (s, 3H, OMe); 4.16 (d, 1H, *J* = 17.2 Hz, CH<sub>2</sub>); 4.22 (d, 1H, *J* = 17.2 Hz, CH<sub>2</sub>); 4.35 (d, 1H, *J* = 17.2 Hz, CH<sub>2</sub>); 6.88 (d, 2H, *J* = 8.7 Hz, Ar); 6.94-7.04 (m, 2H, Ar); 7.26-7.42 (m, 4H, Ar) ppm. <sup>13</sup>C NMR (400 MHz; DMSO-*d*<sub>6</sub>): δ 169.83, 166.13, 158.21, 153.34, 136.73, 130.18, 127.68, 125.98, 122.51, 120.68, 116.98, 113.80, 78.04, 68.51, 63.21, 55.30, 55.09, 48.12, 38.42, 29.33 ppm. Anal. (C<sub>22</sub>H<sub>23</sub>NO<sub>6</sub>) C, H, N. % Calcd: 66.49 (C); 5.83 (H); 3.52 (N). % Found: 66.65 (C); 6.00 (H); 3.63 (N).

#### 2-(6-Bromo-2-(4-methoxyphenyl)-2-methyl-5'-oxospiro[chroman-4,2'-morpholin]-4'-yl) acetic acid4

Compound **4** was obtained from **21** (0.2 g, 0.4 mmol) following the procedure previously described. The crude product was purified by crystallization from (*i*Pr)<sub>2</sub>O to give **4** (0.03 g, 0.07 mmol, yield 18%) <sup>1</sup>H-NMR (DMSO-*d*<sub>6</sub>): δ 1.64 (s, 3H, Me); 2.30-2.39 (m, 2H, CH<sub>2</sub>); 2.92 (d, 1H, *J* = 14.6 Hz, CH<sub>2</sub>); 3.25-3.38 (m, 2H, CH<sub>2</sub>); 3.54-3.65 (m, 1H, CH<sub>2</sub>); 3.72 (s, 3H, OMe); 4.08-4.40 (m, 2H, CH<sub>2</sub>); 6.89 (d, 2H, *J* = 8.8 Hz, Ar); 7.03 (d, 1H, *J* = 8.6 Hz, Ar); 7.31 (d, 2H, *J* = 8.8 Hz, Ar); 7.48 (dd, 1H, *J* = 2.4, 8.6 Hz, Ar); 7.55 (d, 1H, *J* = 2.4 Hz, Ar) ppm. Anal. (C<sub>22</sub>H<sub>22</sub>BrNO<sub>6</sub>) C, H, N. Anal. % Calcd: 55.47 (C); 4.66 (H); 2.94 (N). % Found: 55.52 (C); 4.66 (H); 2.94 (N).

#### 2-(2,2-Dimethyl-2'-oxospiro[chroman-4,5'-oxazolidin]-3'-yl) acetic acid5

Compound **5** was obtained from **26** (0.23 g, 0.56 mmol) following procedure previously described without further purification. **5** (0.04 g, 0.16 mmol, yield 29%). Mp 160-165 °C. <sup>1</sup>H-NMR (DMSO-*d*<sub>6</sub>): δ 1.32 (s, 3H, Me); 1.36 (s, 3H, Me); 2.27 (d, 1H, *J* = 14.8 Hz, CH<sub>2</sub>); 2.37 (d, 1H, *J* = 14.8 Hz, CH<sub>2</sub>); 3.68 (d, 1H, *J* = 9.1 Hz, CH<sub>2</sub>); 3.87 (d, 1H, *J* = 9.1 Hz, CH<sub>2</sub>); 3.95 (d, 1H, *J* = 17.7 Hz, CH<sub>2</sub>); 4.08 (d, 1H, *J* = 17.7 Hz, CH<sub>2</sub>); 6.80 (d, 1H, *J* = 8.7 Hz, Ar); 6.98 (m, 1H, Ar); 7.26 (m, 1H, Ar); 7.58 (d, 1H, *J* = 8.7 Hz, Ar) ppm. Anal. (C<sub>15</sub>H<sub>17</sub>NO<sub>5</sub>) C, H, N. Anal. % Calcd: 61.85 (C); 5.88 (H); 4.81 (N). % Found: 61.60 (C); 5.85 (H); 4.79 (N).

#### 2-(6-Bromo-2,2-dimethyl-2'-oxospiro[chroman-4,5'-oxazolidin]-3'-yl) acetic acid6

Compound **6** was obtained from **27** (0.33 g, 0.67 mmol) following procedure previously described without further purification. **6** (0.12 g, 0.35 mmol, 53%yield). Mp 165-170°C. <sup>1</sup>H-NMR (DMSO-*d*<sub>6</sub>): δ 1.32 (s, 3H, Me); 1.36 (s, 3H, Me); 2.27 (d, 1H, *J* = 15.0 Hz, CH<sub>2</sub>); 2.37 (d, 1H, *J* = 15.0 Hz, CH<sub>2</sub>); 3.68 (d, 1H, *J* = 9.2 Hz, CH<sub>2</sub>); 3.88 (d, 1H, *J* = 9.2 Hz, CH<sub>2</sub>); 3.90 (d, 1H, *J* = 18.1 Hz, CH<sub>2</sub>); 4.00 (d, 1H, *J* = 18.1 Hz, CH<sub>2</sub>); 6.79 (d, 1H, *J* = 8.7 Hz, Ar); 7.43 (dd, 1H, *J* = 2.3, 8.7 Hz, Ar); 7.85 (d, 1H, *J* = 2.3 Hz, Ar) ppm. Anal. (C<sub>15</sub>H<sub>16</sub>BrNO<sub>5</sub>) C, H, N. % Calcd: 48.67 (C); 4.36 (H); 3.78 (N). % Found: 48.3 (C); 4.70 (H); 3.50 (N).

**2-(2,2-Dimethyl-5'-oxospiro[chroman-4,2'-morpholin]-4'-yl) acetic acid7**

Compound **7** was obtained from **28** (0.26 g, 0.77 mmol) following procedure previously described. The crude product was collected without further purification to yield **7** (0.14 g, 0.46 mmol, 60% yield). Mp 163-165°C. <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.41 (s, 6H, Me); 2.19 (d, 1H, *J* = 14.6 Hz, CH<sub>2</sub>); 2.42 (d, 1H, *J* = 14.6 Hz, CH<sub>2</sub>); 3.26 (d, 1H, *J* = 12.1 Hz, CH<sub>2</sub>); 4.02- 4.10 (m, 2H, CH<sub>2</sub>); 4.35- 4.46 (m, 3H, CH<sub>2</sub>); 6.84- 7.0 (m, 2H, Ar); 7.22-7.29 (m, 1H, Ar); 7.46- 7.47 (m, 1H, Ar) ppm. <sup>13</sup>C NMR (400 MHz; CDCl<sub>3</sub>): δ 171.88, 168.30, 153.95, 130.58, 127.53, 121.34, 120.95, 118.42, 74.46, 69.43, 63.42, 57.81, 48.10, 39.65, 26.19 ppm. Anal. (C<sub>16</sub>H<sub>19</sub>NO<sub>5</sub>) C, H, N. % Calcd: 62.94 (C); 6.27 (H); 4.59 (N). % Found: 62.40 (C); 5.83 (H); 4.09 (N).

**2-(6-Bromo-2,2-dimethyl-5'-oxospiro[chroman-4,2'-morpholin]-4'-yl) acetic acid8**

Compound **8** was obtained from **29** (0.39 g, 0.96 mmol) following the general procedure. The crude product was collected without further purification to yield **8** (0.12 g, 0.32 mmol, 34% yield). Mp 132-134°C. <sup>1</sup>H-NMR (DMSO-*d*<sub>6</sub>): δ 1.32 (s, 3H, Me); 1.35 (s, 3H, Me); 2.14 (d, 1H, *J* = 15.0 Hz, CH<sub>2</sub>); 2.42- 2.52 (m, 1H, CH<sub>2</sub>); 3.96- 4.24 (m, 6H, CH<sub>2</sub>); 6.79 (d, 1H, *J* = 8.7 Hz, Ar); 7.40 (dd, 1H, *J* = 2.4, 8.7 Hz, Ar); 7.60 (d, 1H, *J* = 2.4 Hz, Ar) ppm. Anal. (C<sub>16</sub>H<sub>18</sub>BrNO<sub>5</sub>) C, H, N. % Calcd: 50.02 (C); 4.72 (H); 3.65 (N). % Found: 49.00 (C); 4.18 (H); 2.58 (N).

**2-(2,2-Dimethylspiro[chroman-4,2'-morpholin]-4'-yl) acetic acid9**

Compound **9** was obtained from **32** (0.55 g, 1.90 mmol) following procedure previously described. The crude product was collected and purified by precipitation from CHCl<sub>3</sub>/*n*-Hexane to yield **9** (0.01 g, 0.04 mmol, yield 2%). <sup>1</sup>H-NMR (CD<sub>3</sub>OD-*d*<sub>4</sub>): δ 1.35 (s, 3H, Me); 1.45 (s, 3H, Me); 2.20 (d, 1H, *J* = 14.8 Hz, CH<sub>2</sub>); 2.65 (d, 1H, *J* = 14.8 Hz, CH<sub>2</sub>); 3.26-3.40 (m, 2H, CH<sub>2</sub>); 3.70-4.01 (m, 3H, CH<sub>2</sub>); 4.07-4.24 (m, 1H, CH<sub>2</sub>); 4.90 (s, 2H, CH<sub>2</sub>); 6.78- 6.83 (m, 1H, Ar); 6.93- 7.00 (m, 1H, Ar); 7.20- 7.28 (m, 1H, Ar); 7.55 (dd, 1H, *J* = 1.5, 7.9 Hz, Ar) ppm. Anal. (C<sub>16</sub>H<sub>21</sub>NO<sub>4</sub>) C, H, N. % Calcd: 65.96 (C); 7.27 (H); 4.81 (N). % Found: 65.69 (C); 7.24 (H); 4.79 (N).

**2-(6-Bromo-2,2-dimethylspiro[chroman-4,2'-morpholin]-4'-yl) acetic acid10**

Compound **10** was obtained from **33** (0.03 g, 0.11 mmol) and chloroacetic acid (0.01 g, 0.11 mmol) following the general procedure. The crude was used without further purification. **10** (0.005 g, 0.07 mmol, yield 67%). <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.33 (s, 3H, Me); 1.40 (s, 3H, Me); 2.14 (d, 1H, *J* = 14.5 Hz, CH<sub>2</sub>); 2.54 (d, 1H, *J* = 14.5 Hz, CH<sub>2</sub>); 2.70-3.00 (m, 4H, CH<sub>2</sub>); 3.19-3.42 (m, 2H, CH<sub>2</sub>); 3.78-3.84 (m, 1H, CH<sub>2</sub>); 3.99-4.10 (m, 1H, CH<sub>2</sub>); 6.71 (d, 1H, *J* = 8.8 Hz, Ar); 7.29 (dd, 1H, *J* = 2.4, 8.8 Hz, Ar); 7.64 (d, 1H, *J* = 2.4 Hz, Ar) ppm. Anal. (C<sub>16</sub>H<sub>20</sub>BrNO<sub>4</sub>) C, H, N. Anal. (C<sub>16</sub>H<sub>21</sub>NO<sub>4</sub>) C, H, N. % Calcd: 51.90 (C); 5.44 (H); 3.78 (N). % Found: 51.74 (C); 5.42 (H); 3.76 (N).

**2-(5-(2-((3-Methoxybenzyl)oxy)phenyl)-5-methyl-2-oxooxazolidin-3-yl)acetic acid11**

Compound **11** was obtained from **46** (0.05 g, 0.12 mmol) following procedure previously described. The crude product was collected and purified by precipitation from EtOAc/*n*-Hexane to yield **11** (0.02 g, 0.06 mmol, 49% yield). Mp 117-119 °C. <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.75 (s, 3H, Me); 3.67-3.79 (m, 2H, CH<sub>2</sub>); 3.83 (s, 3H, OMe); 4.02-4.11 (m, 2H, CH<sub>2</sub>); 5.00 (s, 2H, CH<sub>2</sub>); 6.91-7.02 (m, 4H, Ar); 7.24-7.32 (m, 3H, Ar); 7.58 (dd, 1H, *J* = 1.8, 8.1 Hz, Ar) ppm. Anal. (C<sub>20</sub>H<sub>21</sub>NO<sub>6</sub>) C, H, N. % Calcd: 64.68 (C); 5.70 (H); 3.77 (N). % Found: 64.98 (C); 5.84 (H); 3.57 (N).

**2-(5-(2-((3,5-Dimethoxybenzyl)oxy)phenyl)-5-methyl-2-oxooxazolidin-3-yl)acetic acid12**

Compound **12** was obtained from **47** (0.05 g, 0.12 mmol) following same procedure described above. The crude was used without further purification. **12** (0.04 g, 0.11 mmol, 90% yield). Mp 118-120 °C. <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.79 (s, 3H, Me); 3.65-3.90 (m, 5H, CH<sub>2</sub>, OMe); 4.03-4.17 (m, 2H, CH<sub>2</sub>); 4.94-5.08 (m, 2H, CH<sub>2</sub>); 6.83-7.04 (m, 5H, Ar); 7.26-7.33 (m, 1H, Ar); 7.59-7.63 (m, 1H, Ar) ppm. Anal. (C<sub>21</sub>H<sub>23</sub>NO<sub>7</sub>) C, H, N. % Calcd: 62.83 (C); 5.78 (H); 3.49 (N). % Found: 63.02 (C); 5.67 (H); 3.44 (N).

**2-(5-Methyl-2-oxo-5-(2-((3,4,5-trimethoxybenzyl)oxy)phenyl)oxazolidin-3-yl)acetic acid13**

Compound **13** was obtained from **48** (0.54g, 1.18 mmol) following procedure previously described. The crude product was dissolved in EtOAc and washed with NaHCO<sub>3</sub> and then extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layers were dried and concentrated under vacuum to give **13** (0.07 g, 0.15 mmol, 13% yield). <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.83 (s, 3H, Me);



3.57-4.18 (m, 7H, CH<sub>2</sub>, OMe); 5.02 (s, 2H, CH<sub>2</sub>); 6.61 (s, 2H, Ar); 6.95-7.05 (m, 2H, Ar); 7.26-7.34 (m, 1H, Ar); 7.59-7.63 (m, 1H, Ar) ppm. <sup>13</sup>C NMR (400 MHz; CDCl<sub>3</sub>): δ 171.92, 157.54, 154.16, 153.58, 137.83, 132.12, 131.37, 129.41, 125.79, 121.33, 112.05, 105.19, 80.66, 70.46, 61.03, 57.43, 56.27, 44.95, 27.15 ppm. Anal. (C<sub>22</sub>H<sub>25</sub>NO<sub>8</sub>) C, H, N. % Calcd: 61.25 (C); 5.84 (H); 3.25 (N). % Found: 61.06 (C); 5.82 (H); 3.24 (N).

#### General Procedure for Synthesis of Compounds 6-17

A solution of NaOH (0.19 g, 4.75 mmol) in H<sub>2</sub>O (8.5 mL) was added to a solution appropriate amine **14-15** (4.00 mmol) [12] in CH<sub>2</sub>Cl<sub>2</sub> (12 mL). The mixture was stirred, cooled to 0° C and treated dropwise with chloroacetyl chloride (0.73 g, 6.5 mmol). At the end of the addition, the mixture was stirred vigorously at r.t. for 1 h. The layers were separated and the CH<sub>2</sub>Cl<sub>2</sub> solution was washed with diluted HCl and H<sub>2</sub>O, dried and evaporated.

#### 2-Chloro-N-((4-hydroxy-2-(4-methoxyphenyl)-2-methylchroman-4-yl)methyl)acetamide 16

Compound **16** was obtained from **14** (1.6 g, 5.35 mmol) and chloroacetyl chloride (0.86 g, 7.49 mmol) following the general procedure. The crude product was directly used in the next step without further purification. **16** (1.30 g, 3.47 mmol, yield 65%). <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.67 (s, 3H, Me); 2.29 (d, 1H, *J* = 14.2 Hz, CH<sub>2</sub>); 2.63 (d, 1H, *J* = 14.2 Hz, CH<sub>2</sub>); 3.45 (dd, 1H, *J* = 5.8, 14.4 Hz, CH<sub>2</sub>); 3.74 (d, 1H, *J* = 14.4 Hz, CH<sub>2</sub>); 3.75 (s, 3H, OMe); 3.99 (d, 1H, *J* = 15.5 Hz, CH<sub>2</sub>); 4.09 (d, 1H, *J* = 15.5 Hz, CH<sub>2</sub>); 6.79-6.90 (m, 3H, Ar); 6.94-7.02 (m, 1H, Ar); 7.18-7.36 (m, 3H, Ar); 7.45 (d, 1H, *J* = 7.6 Hz, Ar) ppm.

#### N-((6-Bromo-4-hydroxy-2-(4-methoxyphenyl)-2-methylchroman-4-yl)methyl)-2-chloroacetamide 17

Compound **17** was obtained from **15** (0.8 g, 2.11 mmol) and chloroacetyl chloride (0.34 g, 2.95 mmol) following the general procedure. The crude product was directly used in the next step without further purification. **17** (0.8 g, 1.76 mmol, yield 47%). <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.69 (s, 3H, Me); 2.24-2.36 (m, 2H, CH<sub>2</sub>); 2.66 (d, 1H, *J* = 14.2 Hz, CH<sub>2</sub>); 3.38-3.48 (m, 1H, CH<sub>2</sub>); 3.78 (s, 3H, OMe); 4.03 (d, 1H, *J* = 15.7 Hz, CH<sub>2</sub>); 4.13 (d, 1H, *J* = 15.7 Hz, CH<sub>2</sub>); 6.80-6.94 (m, 3H, Ar); 7.24-7.38 (m, 3H, Ar); 7.61 (d, 1H, *J* = 2.6 Hz, Ar) ppm.

#### General Procedure for Synthesis of Compounds 18-19

Potassium *tert*-butoxide (1.45 g, 13 mmol) was added portion wise over 1 h to a stirred solution of appropriate alcohol **16-17** (2.5 mmol) in toluene (20 mL). The reaction mixture was stirred at r.t. for 2 h, then the solvent was evaporated. The residue was diluted with EtOAc and the organic layer washed with water, dried and evaporated.

#### 2-(4-Methoxyphenyl)-2-methylspiro[chroman-4,2'-morpholin]-5'-one 18

Compound **18** was obtained from **16** (1.30 g, 3.47 mmol) following the procedure previously described. The crude product was directly used in the next step without further purification. **18** (0.73 g, 2.16 mmol, yield 63%). <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.72 (s, 3H, Me); 2.24 (d, 1H, *J* = 12.8 Hz, CH<sub>2</sub>); 2.33 (d, 1H, *J* = 13.7 Hz, CH<sub>2</sub>); 3.01 (d, 1H, *J* = 13.7 Hz, CH<sub>2</sub>); 3.33 (d, 1H, *J* = 12.8 Hz, CH<sub>2</sub>); 3.76 (s, 3H, OMe); 4.38 (s, 2H, CH<sub>2</sub>); 6.80 (d, 2H, *J* = 7.6 Hz, Ar); 6.92-6.99 (m, 1H, Ar); 7.03-7.07 (m, 1H, Ar); 7.15-7.39 (m, 4H, Ar) ppm.

#### 6-Bromo-2-(4-methoxyphenyl)-2-methylspiro[chroman-4,2'-morpholin]-5'-one 19

Compound **19** was obtained from **17** (0.8 g, 1.76 mmol) following the procedure previously described. The crude product was directly used in the next step without further purification. **19** (0.63 g, 1.5 mmol, yield 85%). <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.74 (s, 3H, Me); 2.29 (d, 1H, *J* = 12.8 Hz, CH<sub>2</sub>); 2.35 (d, 1H, *J* = 13.7 Hz, CH<sub>2</sub>); 3.05 (d, 1H, *J* = 13.7 Hz, CH<sub>2</sub>); 3.40 (d, 1H, *J* = 12.8 Hz, CH<sub>2</sub>); 3.78 (s, 3H, OMe); 4.40 (s, 2H, CH<sub>2</sub>); 6.82 (d, 2H, *J* = 8.8 Hz, Ar); 7.03 (d, 1H, *J* = 8.6 Hz, Ar); 7.29-7.35 (m, 2H, Ar); 7.45-7.50 (m, 1H, Ar); 7.55 (d, 1H, *J* = 2.4 Hz, Ar) ppm.

#### General Procedure for Synthesis of Compounds 20-21, 26-29

To a stirred solution of NaH (0.07 g, 3.00 mmol, 60% dispersion in mineral oil) in dry DMF (10 mL) was added the appropriate derivatives **18-19, 22-25** (4.00 mmol) under N<sub>2</sub> atmosphere. After 30 min, the reaction mixture was cooled at 0 °C and ethyl bromoacetate (0.83 g, 5.00 mmol) was added. The reaction mixture was allowed to warm at 25 °C and stirred for 1h before being quenched with water and extracted with EtOAc. The combined organic layers were dried, filtered, and concentrated under vacuum.

**Ethyl 2-(2-(4-methoxyphenyl)-2-methyl-5'-oxospiro[chroman-4,2'-morpholin]-4'-yl) acetate20**

Compound **20** was obtained from **18** (0.73 g, 2.16 mmol) and ethyl bromoacetate (0.78 g, 2.16 mmol) following the general procedure. The crude product was purified by flash column chromatography eluting *n*-Hexane/EtOAc (1:1) to give **20** (0.47 g, 1.10 mmol, yield 51%): <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.26 (t, 3H, *J* = 7.1 Hz, Me); 1.73 (s, 3H, Me); 2.21 (d, 1H, *J* = 12.6 Hz, CH<sub>2</sub>); 2.39 (d, 1H, *J* = 14.2 Hz, CH<sub>2</sub>); 3.04 (d, 1H, *J* = 14.2 Hz, CH<sub>2</sub>); 3.18 (d, 1H, *J* = 17.3 Hz, CH<sub>2</sub>); 3.64 (d, 1H, *J* = 12.6 Hz, CH<sub>2</sub>); 3.78 (s, 3H, OMe); 4.16 (q, 2H, *J* = 7.1 Hz, CH<sub>2</sub>); 4.34 (d, 1H, *J* = 17.3 Hz, CH<sub>2</sub>); 4.41 (s, 2H, CH<sub>2</sub>); 6.82 (d, 2H, *J* = 8.7 Hz, Ar); 6.93-7.08 (m, 2H, Ar); 7.21-7.36 (m, 3H, Ar); 7.44 (dd, 1H, *J* = 8.7, 1.4 Hz, Ar) ppm.

**Ethyl 2-(6-bromo-2-(4-methoxyphenyl)-2-methyl-5'-oxospiro[chroman-4,2'-morpholin]-4'-yl) acetate21**

Compound **21** was obtained from **19** (0.3 g, 0.72 mmol) and ethyl bromoacetate (0.12 g, 0.72 mmol) following the general procedure. The crude product was purified by flash column chromatography eluting *n*-Hexane/EtOAc (1:1) to give **21** (0.22 g, 0.4 mmol, yield 55%). <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.30 (t, 3H, *J* = 7.0 Hz, Me); 1.75 (s, 3H, Me); 2.23 (d, 1H, *J* = 12.4 Hz, CH<sub>2</sub>); 2.40 (d, 1H, *J* = 13.2 Hz, CH<sub>2</sub>); 3.01 (d, 1H, *J* = 13.2 Hz, CH<sub>2</sub>); 3.15 (d, 1H, *J* = 17.0 Hz, CH<sub>2</sub>); 3.61 (d, 1H, *J* = 12.4 Hz, CH<sub>2</sub>); 3.78 (s, 3H, OMe); 4.20 (q, 2H, *J* = 7.0 Hz, CH<sub>2</sub>); 4.30 (d, 1H, *J* = 17.0 Hz, CH<sub>2</sub>); 4.35 (s, 2H, CH<sub>2</sub>); 6.72 (d, 2H, *J* = 8.6 Hz, Ar); 6.97 (d, 1H, *J* = 8.3 Hz, Ar); 7.00 (d, 2H, *J* = 8.6 Hz, Ar); 7.60 (dd, 1H, *J* = 2.5, 8.3 Hz, Ar); 7.65 (d, 1H, *J* = 2.5 Hz, Ar) ppm.

**Ethyl 2-(2,2-dimethyl-2'-oxospiro[chroman-4,5'-oxazolidin]-3'-yl) acetate26**

Compound **26** was obtained from **22** (0.44 g, 1.37 mmol) following procedure previously described. The crude product was directly used for the next reaction without further purification. **26** (0.23 g, 0.56 mmol, yield 41%): <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.33 (t, 3H, *J* = 7.1 Hz, Me); 1.42 (s, 6H, Me); 2.17 (d, 1H, *J* = 14.6 Hz, CH<sub>2</sub>); 2.46 (d, 1H, *J* = 14.6 Hz, CH<sub>2</sub>); 3.64 (d, 1H, *J* = 8.4 Hz, CH<sub>2</sub>); 3.98 (d, 1H, *J* = 8.4 Hz, CH<sub>2</sub>); 4.02 (d, 1H, *J* = 18.0 Hz, CH<sub>2</sub>); 4.20 (d, 1H, *J* = 18.0 Hz, CH<sub>2</sub>); 4.25 (q, 2H, *J* = 7.1 Hz, CH<sub>2</sub>); 6.82 (d, 1H, *J* = 8.2 Hz, Ar); 6.94-7.02 (m, 1H, Ar); 7.20-7.29 (m, 1H, Ar); 7.58 (dd, 1H, *J* = 8.0 Hz, Ar) ppm.

**Ethyl 2-(6-bromo-2,2-dimethyl-2'-oxospiro[chroman-4,5'-oxazolidin]-3'-yl) acetate27**

Compound **27** was obtained from **23** (0.55 g, 1.37 mmol) following procedure previously described. The crude product was directly used for the next reaction without further purification. **27** (0.33 g, 0.67 mmol, yield 49%): <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.32 (t, 3H, *J* = 7.1 Hz, Me); 1.41 (s, 6H, Me); 2.16 (d, 1H, *J* = 14.7 Hz, CH<sub>2</sub>); 2.45 (d, 1H, *J* = 14.7 Hz, CH<sub>2</sub>); 3.67 (d, 1H, *J* = 8.7 Hz, CH<sub>2</sub>); 3.93 (d, 1H, *J* = 8.7 Hz, CH<sub>2</sub>); 4.01-4.33 (m, 4H, CH<sub>2</sub>); 6.70 (d, 1H, *J* = 8.7 Hz, Ar); 7.33 (dd, 1H, *J* = 2.4, 8.7 Hz, Ar); 7.68 (d, 1H, *J* = 2.4 Hz, Ar) ppm.

**Ethyl 2-(2,2-dimethyl-5'-oxospiro[chroman-4,2'-morpholin]-4'-yl) acetate28**

Compound **28** was obtained from **24** (0.6 g, 2.42 mmol) and ethyl bromo acetate (0.48 g, 2.90 mmol) following the procedure previously described. The crude product was purified by flash column chromatography eluting *n*-Hexane/EtOAc (7:3) to give **27** (0.26 g, 0.77 mmol, yield 32%): <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.21- 1.41 (m, 9H, Me); 2.21 (d, 1H, *J* = 14.7 Hz, CH<sub>2</sub>); 2.42 (d, 1H, *J* = 14.7 Hz, CH<sub>2</sub>); 3.22 (d, 1H, *J* = 12.1 Hz, CH<sub>2</sub>); 3.93- 4.41 (m, 7H, CH<sub>2</sub>); 6.84 (dd, 1H, *J* = 1.4; 8.0 Hz, Ar); 6.90- 6.98 (m, 1H, Ar); 7.19- 7.28 (m, 1H, Ar); 7.49 (dd, 1H, *J* = 1.4; 8.0 Hz, Ar) ppm.

**Ethyl 2-(6-bromo-2,2-dimethyl-5'-oxospiro[chroman-4,2'-morpholin]-4'-yl) acetate29**

Compound **29** was obtained from **25** (0.33 g, 1.00 mmol) and ethyl bromo acetate (0.20 g, 1.20 mmol) following the procedure previously described. The crude product was purified by flash column chromatography eluting *n*-Hexane/EtOAc (7:3) to give **29** (0.18 g, 0.44 mmol, yield 44%) <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.20-1.41 (m, 9H, Me); 2.20-2.45 (m, 2H CH<sub>2</sub>); 3.20 (d, 1H, *J* = 12.2 Hz, CH<sub>2</sub>); 3.90- 4.44 (m, 7H, CH<sub>2</sub>); 6.73 (d, 1H, *J* = 8.7 Hz, Ar); 7.32 (dd, 1H, *J* = 2.5, 8.7 Hz, Ar); 7.61 (d, 1H, *J* = 2.5 Hz, Ar) ppm.

**General Procedure for Synthesis of Compounds 32-33**

To a solution of appropriate derivatives **30-31** (0.57 mmol) in MeCN (5 mL) was added K<sub>2</sub>CO<sub>3</sub> (0.09 g, 0.64mmol) and ethyl bromoacetate (0.09 g, 5.00 mmol). The resulting mixture was refluxed for 12 h, then, after cooling, was filtered and the solvent evaporated.

**Ethyl 2-(2,2-dimethylspiro[chroman-4,2'-morpholin]-4'-yl) acetate32**

Compound **32** was obtained from **30** (0.45 g, 1.92 mmol) and ethyl bromoacetate (0.32 g, 1.92 mmol) following the same procedure described in the general procedure. The crude product was used for the next step without further purification. **32** (0.55 g, 1.90 mmol, yield 99%): <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.22-1.33 (m, 3H, Me); 1.36 (s, 3H, Me); 1.41 (s, 3H, Me); 2.28 (d, 1H, *J* = 14.8 Hz, CH<sub>2</sub>); 2.56 (d, 1H, *J* = 14.8 Hz, CH<sub>2</sub>); 2.65-2.90 (m, 2H, CH<sub>2</sub>); 3.19 (d, 1H, *J* = 16.7 Hz, CH<sub>2</sub>); 3.37 (d, 1H, *J* = 16.7 Hz, CH<sub>2</sub>); 3.73-3.82 (m, 3H, CH<sub>2</sub>); 4.01-4.28 (m, 3H, CH<sub>2</sub>); 6.78-6.96 (m, 2H, Ar); 7.14-7.23 (m, 1H, Ar); 7.56 (dd, 1H, *J* = 1.5, 7.8 Hz, Ar) ppm.

**Ethyl 2-(6-bromo-2,2-dimethylspiro[chroman-4,2'-morpholin]-4'-yl) acetate33**

Compound **33** was obtained from **31** (0.6 g, 2.42 mmol) and ethyl bromoacetate (0.40 g, 2.42 mmol) following the procedure previously described. The crude product was purified by flash column chromatography eluting *n*-Hexane/EtOAc (7:3) to give **33** (0.56 g, 1.98 mmol, yield 97%): <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.20- 1.30 (m, 3H, Me); 1.33 (s, 3H, Me); 1.39 (s, 3H, Me); 2.25 (d, 1H, *J* = 14.5 Hz, CH<sub>2</sub>); 2.56 (d, 1H, *J* = 14.5 Hz, CH<sub>2</sub>); 2.67 (d, 1H, *J* = 10.0 Hz, CH<sub>2</sub>); 2.93 (d, 1H, *J* = 10.0 Hz, CH<sub>2</sub>); 3.19 (d, 1H, *J* = 16.4 Hz, CH<sub>2</sub>); 3.37 (d, 1H, *J* = 16.4 Hz, CH<sub>2</sub>); 3.69-3.80 (m, 3H, CH<sub>2</sub>); 3.99-4.11 (m, 3H, CH<sub>2</sub>); 6.79 (d, 1H, *J* = 8.7 Hz, Ar); 7.4 (dd, 1H, *J* = 2.5, 8.7 Hz, Ar); 7.6 (d, 1H, *J* = 2.5 Hz, Ar) ppm.

**General Procedure for Synthesis of Compounds 34-36**

To a solution of 2-hydroxyacetophenone (1.5 g, 11.03 mmol) in DMSO (7 mL) was added KOH (1.86 g, 33.09 mmol) and the resulting mixture was stirred at 50°C for 15 min. After cooling to r.t. the appropriate methoxybenzyl chloride (11.03 mmol) was added. The reaction mixture was stirred at r.t. for 4h and then washed with H<sub>2</sub>O and NaHCO<sub>3</sub>sat. The organic layer was dried, filtered and evaporated under vacuum.

**1-(2-((3-Methoxybenzyl)oxy)phenyl)ethanone34**

Compound **34** was synthesized from 2-hydroxyacetophenone (1.5 g, 11.03 mmol) and 4-methoxybenzyl chloride (1.73 g, 11.03 mmol) following the procedure previously described. The crude product was directly used in the next step without further purification. **34** (1.5 g, 5.83 mmol, yield 53%): <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 2.57 (s, 3H, Me); 3.83 (s, 3H, OMe); 5.09 (s, 2H, CH<sub>2</sub>); 6.93 (d, 2H, *J* = 8.6 Hz, Ar); 6.96-7.05 (m, 2H, Ar); 7.36 (d, 2H, *J* = 8.6 Hz, Ar); 7.40-7.49 (m, 1H, Ar); 7.75 (dd, 1H, *J* = 2.0, 7.7, Hz, Ar) ppm.

**1-(2-((3,5-Dimethoxybenzyl)oxy)phenyl)ethanone35**

Compound **35** was synthesised from 2-hydroxyacetophenone (1.5 g, 11.03 mmol) and 3,4-dimethoxybenzyl chloride (2.06 g, 11.03 mmol) following the same procedure described in general procedure. The crude product was directly used for the next reaction without further purification. **35** (1.83 g, 9.81 mmol, yield 89%): <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 2.59 (s, 3H, Me); 3.89 (s, 3H, OMe); 3.90 (s, 3H, OMe); 5.09 (s, 2H, CH<sub>2</sub>); 6.85-7.05 (m, 5H, Ar); 7.41-7.49 (m, 1H, Ar); 7.74 (dd, 1H, *J* = 1.8, 7.5 Hz, Ar) ppm.

**1-(2-((3,4,5-Trimethoxybenzyl)oxy)phenyl)ethanone36**

Compound **36** was synthesised from 2-hydroxyacetophenone (0.9 g, 6.61 mmol) and 3,4,5-trimethoxybenzyl chloride (1.5 g, 6.93 mmol) following the same procedure previously described. The crude product was directly used for the next reaction without further purification. **36** (1.9 g, 6.03 mmol, yield 87%): <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 2.63 (s, 3H, Me); 3.86 (s, 9H, OMe); 5.09 (s, 2H, CH<sub>2</sub>); 6.67 (s, 2H, Ar); 6.99-7.06 (m, 2H, Ar); 7.41-7.50 (m, 1H, Ar); 7.74 (dd, 1H, *J* = 1.7, 8.0 Hz, Ar) ppm.

**General Procedure for Synthesis of Compounds 37-39**

To a solution of appropriate derivatives **37-39** (1.23 g, 7 mmol) in CH<sub>2</sub>Cl<sub>2</sub> was added trimethylsilylcyanide (TMSCN) (1.4 mL, 10.5 mmol) and ZnI<sub>2</sub> (0.33 g, 1.05 mmol). The mixture was stirred at r.t. for 4 h, then CH<sub>2</sub>Cl<sub>2</sub> was added and the solution was washed with water. The organic layer was dried and evaporated.

**3-(2-((3-Methoxybenzyl)oxy)phenyl)-3-((trimethylsilyl)oxy)butanenitrile37**

Compound **37** was synthesised from **34** (1.5 g, 5.83 mmol) following the procedure described in general procedure. The crude product was directly used in the next step without further purification. **37** (1.74 g, 4.88 mmol, yield

84%). <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 0.31 (s, 9H, Me); 1.89 (s, 3H, Me); 3.82 (s, 3H, OMe); 5.05-5.16 (m, 2H, CH<sub>2</sub>); 6.77-7.04 (m, 4H, Ar); 7.19-7.35 (m, 1H, Ar); 7.46 (d, 2H, *J* = 8.8 Hz, Ar); 7.56 (dd, 1H, *J* = 2.0, 8.8 Hz, Ar) ppm.

#### 3-(2-((3,5-Dimethoxybenzyl)oxy)phenyl)-3-((trimethylsilyl)oxy)butanenitrile 38

Compound **38** was synthesised from **35** (1.09 g, 5.83 mmol) following the procedure previously described. The crude product was directly used in the next step without further purification. **38** (1.91 g, 4.95 mmol, yield 85%): <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 0.29 (s, 9H, Me); 1.89 (s, 3H, Me); 3.89 (s, 3H, OMe); 3.90 (s, 3H, OMe); 5.03-5.18 (m, 2H, CH<sub>2</sub>); 6.67-7.03 (m, 5H, Ar); 7.18-7.41 (m, 1H, Ar); 7.56 (dd, 1H, *J* = 1.8, 8.1 Hz) ppm.

#### 3-(2-((3,4,5-Trimethoxybenzyl)oxy)phenyl)-3-((trimethylsilyl)oxy)butanenitrile 39

Compound **39** was synthesised from **36** (1.90 g, 6.03 mmol) following the same procedure described in general procedure. The crude product was directly used in the next step without further purification. **39** (2.22 g, 5.37 mmol, yield 89%): <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 0.26 (s, 9H, Me); 2.01 (s, 3H, Me); 3.88 (s, 9H, OMe); 5.18 (s, 2H, CH<sub>2</sub>); 6.80 (s, 2H, Ar); 7.01-7.08 (m, 2H, Ar); 7.35-7.42 (m, 2H, Ar) ppm.

#### General Procedure for Synthesis of Compounds 40-42

A solution of appropriate trimethylsilyl cyanohydrins **37-39** (5.00 mmol) in THF was added dropwise at 0°C to a solution of LiAlH<sub>4</sub> (1M in THF, 10 mmol). The reaction mixture was stirred at r.t. for 1h, then quenched with water and NaOH. The resulting lithium salts were filtered and the solution was evaporated.

#### 1-Amino-2-(2-((3-methoxybenzyl)oxy)phenyl)propan-2-ol 40

Compound **40** was synthesised from **37** (1.74 g, 4.88 mmol) following the procedure described in the general procedure. The crude product was directly used in the next step without further purification. **40** (0.83 g, 2.88 mmol, yield 59%): <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.49 (s, 3H, Me); 3.31 (d, 1H, *J* = 12.7 Hz, CH<sub>2</sub>); 3.51 (d, 1H, *J* = 12.7 Hz, CH<sub>2</sub>); 3.83 (s, 3H, OMe); 5.00-5.14 (m, 2H, CH<sub>2</sub>); 6.87-7.05 (m, 4H, Ar); 7.18-7.28 (m, 1H, Ar); 7.35 (d, 2H, *J* = 8.6 Hz, Ar); 7.68 (d, 1H, *J* = 2.0, 8.6 Hz, Ar) ppm.

#### 1-Amino-2-(2-((3,5-dimethoxybenzyl)oxy)phenyl)propan-2-ol 41

Compound **41** was synthesised from **38** (1.91 g, 4.95 mmol) following the procedure described above. The crude product was directly used in the next step without further purification. **41** (1.19 g, 3.76 mmol, yield 76%): <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.54 (s, 3H, Me); 2.77 (d, 1H, *J* = 12.7 Hz, CH<sub>2</sub>); 3.29 (d, 1H, *J* = 12.7 Hz, CH<sub>2</sub>); 3.88 (s, 3H, OMe); 3.89 (s, 3H, OMe); 5.03 (s, 2H, CH<sub>2</sub>); 6.62-7.02 (m, 5H, Ar); 7.20-7.28 (m, 1H, Ar); 7.45-7.50 (m, 1H, Ar) ppm.

#### 1-Amino-2-(2-((3,4,5-trimethoxybenzyl)oxy)phenyl)propan-2-ol 42

Compound **42** was synthesised from **39** (1.91 g, 4.95 mmol) following the procedure previously described. The crude product was directly used in the next step without further purification. **42** (1.21 g, 3.49 mmol, yield 65%): <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.57 (s, 3H, Me); 2.80 (d, 1H, *J* = 12.7 Hz, CH<sub>2</sub>); 3.31 (d, 1H, *J* = 12.7 Hz, CH<sub>2</sub>); 3.85 (s, 9H, OMe); 5.03 (s, 2H, CH<sub>2</sub>); 6.65 (s, 2H, Ar); 6.90-7.02 (m, 2H, Ar); 7.18-7.28 (m, 1H, Ar); 7.48-7.53 (m, 1H, Ar) ppm.

#### General Procedure for Synthesis of Compounds 43-45

A solution of aminoalcohols **40-42** (6.33 mmol) in THF was added dropwise to a solution of *N,N'*-carbonyl diimidazole (CDI) (1.02 g, 6.33 mmol) in THF at 0 °C. The reaction mixtures were stirred r.t. for 5h. The solvent was evaporated under vacuum and the residue diluted with EtOAc and washed with HCl 1N and K<sub>2</sub>CO<sub>3</sub> sat. The organic layers were dried and concentrated under vacuum.

#### 5-(2-((3-Methoxybenzyl)oxy)phenyl)-5-methyloxazolidin-2-one 43

Compound **43** was synthesised from **40** (0.48 g, 1.67 mmol) following the general procedure. The crude product was purified by flash column chromatography eluted by *n*-Hexane/EtOAc (5.5:4.5) to give **43** (0.16 g, 0.5 mmol, yield 30%): <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.73 (s, 3H, Me); 3.62 (d, 1H, *J* = 9.1 Hz, CH<sub>2</sub>); 3.72 (d, 1H, *J* = 9.1 Hz, CH<sub>2</sub>); 3.83 (s, 3H, OMe); 4.99 (s, 2H, CH<sub>2</sub>); 6.93 (d, 2H, *J* = 8.4 Hz, Ar); 6.96-7.04 (m, 2H, Ar); 7.25-7.33 (m, 3H, Ar); 7.59-7.63 (m, 1H, Ar) ppm.

**5-(2-((3,5-Dimethoxybenzyl)oxy)phenyl)-5-methyloxazolidin-2-one44**

Compound **44** was synthesised from **41** (0.53 g, 1.67 mmol) following the procedure described above. The crude product was directly used in the next step without further purification. **44** (0.45 g, 1.30 mmol, yield 78%). <sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.75 (s, 3H, Me); 3.63 (d, 1H, *J* = 9.1 Hz, CH<sub>2</sub>); 3.73 (d, 1H, *J* = 9.1 Hz, CH<sub>2</sub>); 3.89 (s, 3H, OMe); 3.90 (s, 3H, OMe); 4.92-5.09 (m, 2H, CH<sub>2</sub>); 6.70-7.04 (m, 5H, Ar); 7.25-7.33 (m, 1H, Ar); 7.58-7.63 (m, 1H, Ar) ppm.

**5-Methyl-5-(2-((3,4,5-trimethoxybenzyl)oxy)phenyl)oxazolidin-2-one45**

Compound **45** was synthesised from **42** (1.21 g, 3.49 mmol) following the general procedure. The crude product was directly used in the next step without further purification. **45** (1.07 g, 2.86 mmol, yield 82%). <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.80 (s, 3H, Me); 3.67-3.93 (m, 11H, CH<sub>2</sub>, OMe); 4.92-5.45 (m, 2H, CH<sub>2</sub>); 6.60 (s, 2H, Ar); 6.92-7.06 (m, 2H, Ar); 7.20-7.30 (m, 1H, Ar); 7.62 (dd, 1H, *J* = 1.6, 7.7 Hz, Ar) ppm.

**General Procedure for Synthesis of Compounds46-48**

To a stirred solution of NaH (0.07 g, 3.00 mmol, 60% dispersion in mineral oil) in dry DMF (10 mL) was added appropriate compounds **43-45** (0.9 g, 4.00 mmol) under N<sub>2</sub> atmosphere. After 30 min, the reaction mixture was cooled at 0 °C and ethyl bromoacetate (0.83 g, 5.00 mmol) was added. The reaction mixture was allowed to warm at 25 °C and stirred for 1 h before being quenched with water and extracted with EtOAc. The combined organic layers were dried, filtered, and concentrated under vacuum.

**Ethyl 2-(5-(2-((3-methoxybenzyl)oxy)phenyl)-5-methyl-2-oxooxazolidin-3-yl)acetate46**

Compound **46** was synthesised from **43** (0.16 g, 0.5 mmol) and ethylbromoacetate (0.08 g, 0.5 mmol) following the procedure described above. The crude product was directly used in the next step without further purification. **46** (0.16 g, 0.4 mmol, yield 81%): <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.18-1.27 (m, 3H, Me); 1.77 (s, 3H, Me); 3.66-3.78 (m, 2H, CH<sub>2</sub>); 3.83 (s, 3H, OMe); 4.02-4.26 (m, 4H, CH<sub>2</sub>); 5.01 (s, 2H, CH<sub>2</sub>); 6.91-7.04 (m, 4H, Ar); 7.26-7.33 (m, 3H, Ar); 7.59-7.63 (m, 1H, Ar) ppm.

**Ethyl 2-(5-(2-((3,5-dimethoxybenzyl)oxy)phenyl)-5-methyl-2-oxooxazolidin-3-yl)acetate47**

Compound **47** was synthesised from **44** (0.17 g, 0.5 mmol) and ethyl bromoacetate (0.08 g, 0.5 mmol) following the general procedure. The crude product was directly used in the next step without further purification. **47** (0.17 g, 0.4 mmol, yield 81%). <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.14-1.33 (m, 3H, Me); 1.79 (s, 3H, Me); 3.65-3.98 (m, 10H, CH<sub>2</sub>, OMe); 4.02-4.25 (m, 2H, CH<sub>2</sub>); 5.01 (s, 2H, CH<sub>2</sub>); 6.70-6.79 (m, 1H, Ar); 6.83-7.03 (m, 4H, Ar); 7.24-7.33 (m, 1H, Ar); 7.58-7.62 (m, 1H, Ar) ppm.

**Ethyl 2-(5-methyl-2-oxo-5-(2-((3,4,5-trimethoxybenzyl)oxy)phenyl)oxazolidin-3-yl)acetate48**

Compound **48** was synthesised from **45** (0.50 g, 1.34 mmol) and ethyl bromoacetate (0.22 g, 1.34 mmol) following the procedure previously described. The crude product was directly used in the next step without further purification. **48** (0.54 g, 1.18 mmol, yield 88%). <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.14-1.33 (m, 3H, Me); 1.84 (s, 3H, Me); 3.70-3.92 (m, 13H, CH<sub>2</sub>, OMe); 4.06-4.22 (m, 2H, CH<sub>2</sub>); 4.94-5.17 (m, 2H, CH<sub>2</sub>); 6.61 (s, 2H, Ar); 6.93-7.05 (m, 2H, Ar); 7.20-7.30 (m, 1H, Ar); 7.63 (dd, 1H, *J* = 1.6, 7.7 Hz, Ar) ppm.

**Biology****Materials and Methods**

Adult Sprague-Dawley albino rats (body weights of 120–140), provided by Harlan Nossan, Italy, were exploited to obtain the target enzymes, ALR2 and ALR1, which were isolated and purified from lens and kidney, respectively, following a previously reported protocol [17 - 19]. Pyridine coenzyme, D,L-glyceraldehyde, and sodium D-glucuronate came from Sigma-Aldrich while Sorbinil was a gift from Pfizer, Groton CT. All other chemicals were of reagent grade.

**Enzymatic Assays**

The activity of the two enzymes was determined spectrophotometrically by monitoring the change in absorbance at 340 nm, which is due to the oxidation of NADPH catalyzed by ALR2 and ALR1. The change in pyridine coenzyme

concentration/min was determined using a Beckman DU-64 kinetics software program (Solf Pack TM Module).

ALR2 activity was determined at 30°C in a reaction mixture containing 10 mM D,L-glyceraldehyde, 0.104 mM NADPH, 0.1 M sodium phosphate buffer (pH 6.2), deionized water and the enzyme extract, in a total volume of 1 mL.

ALR1 activity was assayed at 37°C in a reaction mixture containing 20 mM sodium D-glucuronate, 0.12 mM NADPH, 0.1 M sodium phosphate buffer (pH 7.2) and the enzyme extract, in a total volume of 1 mL.

### **Enzymatic Inhibition**

The new synthesized compounds were tested for the inhibitory activity against ALR2 and ALR1 by adding 0.1 mL of the inhibitor solution to the reaction mixture described above. The inhibitors were solubilized in water, and the pH was adjusted to 7. A reference blank containing all the above assay components except the substrate was prepared, to correct for the non-enzymatic oxidation of NADPH and for absorption by the compounds tested.

The inhibitory effect of the new derivatives was routinely estimated at a concentration of 10<sup>-4</sup> M. Those compounds found to be active were tested at additional concentrations between 10<sup>-5</sup> and 10<sup>-7</sup> M. The IC<sub>50</sub> values were determined through linear regression analysis of the log-dose response curve, generated using at least four concentrations of the inhibitor causing an inhibition between 20% and 80%, with two replicates at each concentration.

### **CONCLUSION**

In this study, we described the design, synthesis and biological evaluation of three small series of spirooxazolidinone, spiromorpholinone and benzyloxy acetic acid derivatives, proposed as novel ALR2 inhibitors.

Moving from our previously developed spirochromanes **1** and **2** [12], and through suitable chemical manipulation of both their substitution pattern and their main heterocyclic structure, we succeeded in obtaining a novel set of analogues which proved to inhibit the target enzyme showing IC<sub>50</sub> values in the micromolar/low micromole range.

According to SARs observed, the presence of a pendant phenyl ring in the 2-position of the central core is essential to confer a good inhibitory activity to the spirooxazolidine and spiromorpholine derivatives bearing a 6-bromine atom.

On the contrary, a *gem*-dimethyl substitution proves to be more fruitful in the case of the 6-unsubstituted analogues. The flexible benzyloxy derivatives benefit from the presence of methoxy groups in suitable positions of the pendant phenyl ring.

Although further substitution patterns are needed to extend and deepen SARs observed for the three sub-series, the novel compounds here proposed represent a good starting point for the obtainment of novel and effective ARIs.

### **CONFLICT OF INTEREST**

The authors confirm that this article content has no conflict of interest.

### **ACKNOWLEDGEMENTS**

Declared none.

### **REFERENCES**

- [1] Roglic, G. WHO Global report on diabetes: A summary. *Int. J. Noncommunicable Dis.*, **2016**, *1*(1), 3. [<http://dx.doi.org/10.4103/2468-8827.184853>]
- [2] Alexiou, P.; Pegklidou, K.; Chatzopoulou, M.; Nicolaou, I.; Demopoulos, V.J. Aldose reductase enzyme and its implication to major health problems of the 21<sup>st</sup> century. *Curr. Med. Chem.*, **2009**, *16*(6), 734-752. [<http://dx.doi.org/10.2174/092986709787458362>] [PMID: 19199934]
- [3] Cousido-Siah, A.; Ruiz, F.X.; Mitschler, A.; Porté, S.; de Lera, Á.R.; Martín, M.J.; Manzanaro, S.; de la Fuente, J.A.; Terwesten, F.; Betz, M.; Klebe, G.; Farrés, J.; Parés, X.; Podjarny, A. Identification of a novel polyfluorinated compound as a lead to inhibit the human enzymes aldose reductase and AKR1B10: structure determination of both ternary complexes and implications for drug design. *Acta Crystallogr. D Biol. Crystallogr.*, **2014**, *70*(Pt 3), 889-903. [<http://dx.doi.org/10.1107/S1399004713033452>] [PMID: 24598757]
- [4] Tang, W.H.; Martin, K.A.; Hwa, J. Aldose reductase, oxidative stress, and diabetic mellitus. *Front. Pharmacol.*, **2012**, *3*, 87. [<http://dx.doi.org/10.3389/fphar.2012.00087>] [PMID: 22582044]
- [5] Da Settimo, F.; Primofiore, G.; La Motta, C.; Sartini, S.; Taliani, S.; Simorini, F.; Marini, A.M.; Lavecchia, A.; Novellino, E.; Boldrini, E. Naphtho[1,2-d]isothiazole acetic acid derivatives as a novel class of selective aldose reductase inhibitors. *J. Med. Chem.*, **2005**, *48*(22),

- 6897-6907.  
[<http://dx.doi.org/10.1021/jm050382p>] [PMID: 16250648]
- [6] El-Kabbani, O.; Carbone, V.; Darmanin, C.; Oka, M.; Mitschler, A.; Podjarny, A.; Schulze-Briese, C.; Chung, R.P. Structure of aldehyde reductase holoenzyme in complex with the potent aldose reductase inhibitor fidarestat: implications for inhibitor binding and selectivity. *J. Med. Chem.*, **2005**, *48*(17), 5536-5542.  
[<http://dx.doi.org/10.1021/jm050412o>] [PMID: 16107153]
- [7] El-Kabbani, O.; Ruiz, F.; Darmanin, C.; Chung, R.P. Aldose reductase structures: implications for mechanism and inhibition. *Cell. Mol. Life Sci.*, **2004**, *61*(7-8), 750-762.  
[<http://dx.doi.org/10.1007/s00018-003-3403-2>] [PMID: 15095000]
- [8] Chen, X.; Zhang, S.; Yang, Y.; Hussain, S.; He, M.; Gui, D.; Ma, B.; Jing, C.; Qiao, Z.; Zhu, C.; Yu, Q. 1,2-Benzothiazine 1,1-dioxide carboxylate derivatives as novel potent inhibitors of aldose reductase. *Bioorg. Med. Chem.*, **2011**, *19*(23), 7262-7269.  
[<http://dx.doi.org/10.1016/j.bmc.2011.07.051>] [PMID: 22036211]
- [9] Ramirez, M.A.; Borja, N.L. Epalrestat: an aldose reductase inhibitor for the treatment of diabetic neuropathy. *Pharmacotherapy*, **2008**, *28*(5), 646-655.  
[<http://dx.doi.org/10.1592/phco.28.5.646>] [PMID: 18447661]
- [10] Feather, M.S.; Flynn, T.G.; Munro, K.A.; Kubiseski, T.J.; Walton, D.J. Catalysis of reduction of carbohydrate 2-oxoaldehydes (osones) by mammalian aldose reductase and aldehyde reductase. *Biochim. Biophys. Acta*, **1995**, *1244*(1), 10-16. [BBA].  
[[http://dx.doi.org/10.1016/0304-4165\(94\)00156-R](http://dx.doi.org/10.1016/0304-4165(94)00156-R)] [PMID: 7766643]
- [11] Ramunno, A.; Cosconati, S.; Sartini, S.; Maglio, V.; Angiuoli, S.; La Pietra, V.; Di Maro, S.; Giustiniano, M.; La Motta, C.; Da Settimo, F.; Marinelli, L.; Novellino, E. Progresses in the pursuit of aldose reductase inhibitors: the structure-based lead optimization step. *Eur. J. Med. Chem.*, **2012**, *51*, 216-226.  
[<http://dx.doi.org/10.1016/j.ejmech.2012.02.045>] [PMID: 22436396]
- [12] Rapposelli, S.; Da Settimo, F.; Digiaco, M.; La Motta, C.; Lapucci, A.; Sartini, S.; Vanni, M. Synthesis and biological evaluation of 2-oxo-2,3-dihydro-3H-spiro[chromene-4,5-[1,3]oxazolidin]-3'yl]acetic acid derivatives as aldose reductase inhibitors. *Arch. Pharm. (Weinheim)*, **2011**, *344*(6), 372-385.  
[<http://dx.doi.org/10.1002/ardp.201000302>] [PMID: 21319207]
- [13] Rapposelli, S.; Breschi, M.C.; Calderone, V.; Digiaco, M.; Martelli, A.; Testai, L.; Vanni, M.; Balsamo, A. Synthesis and biological evaluation of 5-membered spiro heterocycle-benzopyran derivatives against myocardial ischemia. *Eur. J. Med. Chem.*, **2011**, *46*(3), 966-973.  
[<http://dx.doi.org/10.1016/j.ejmech.2011.01.003>] [PMID: 21288603]
- [14] Breschi, M.C.; Calderone, V.; Martelli, A.; Minutolo, F.; Rapposelli, S.; Testai, L.; Tonelli, F.; Balsamo, A. New benzopyran-based openers of the mitochondrial ATP-sensitive potassium channel with potent anti-ischemic properties. *J. Med. Chem.*, **2006**, *49*(26), 7600-7602.  
[<http://dx.doi.org/10.1021/jm061228i>] [PMID: 17181142]
- [15] Breschi, M.C.; Calderone, V.; Digiaco, M.; Manganaro, M.; Martelli, A.; Minutolo, F.; Rapposelli, S.; Testai, L.; Tonelli, F.; Balsamo, A. Spirocyclic benzopyran-based derivatives as new anti-ischemic activators of mitochondrial ATP-sensitive potassium channel. *J. Med. Chem.*, **2008**, *51*(21), 6945-6954.  
[<http://dx.doi.org/10.1021/jm800956g>] [PMID: 18925735]
- [16] Brown, H.C.; Choi, Y.M.; Narasimhan, S. Selective reductions. 29. A simple technique to achieve an enhanced rate of reduction of representative organic compounds by borane-dimethyl sulfide. *J. Org. Chem.*, **1982**, *47*, 3153-3163.  
[<http://dx.doi.org/10.1021/jo00137a025>]
- [17] Sartini, S.; Cosconati, S.; Marinelli, L.; Barresi, E.; Di Maro, S.; Simorini, F.; Taliani, S.; Salerno, S.; Marini, A.M.; Da Settimo, F.; Novellino, E.; La Motta, C. Benzofuroxane derivatives as multi-effective agents for the treatment of cardiovascular diabetic complications. Synthesis, functional evaluation, and molecular modeling studies. *J. Med. Chem.*, **2012**, *55*(23), 10523-10531.  
[<http://dx.doi.org/10.1021/jm301124s>] [PMID: 23134227]
- [18] Ottanà, R.; Maccari, R.; Giglio, M.; Del Corso, A.; Cappiello, M.; Mura, U.; Cosconati, S.; Marinelli, L.; Novellino, E.; Sartini, S.; La Motta, C.; Da Settimo, F. Identification of 5-arylidene-4-thiazolidinone derivatives endowed with dual activity as aldose reductase inhibitors and antioxidant agents for the treatment of diabetic complications. *Eur. J. Med. Chem.*, **2011**, *46*(7), 2797-2806.  
[<http://dx.doi.org/10.1016/j.ejmech.2011.03.068>] [PMID: 21531055]
- [19] La Motta, C.; Sartini, S.; Salerno, S.; Simorini, F.; Taliani, S.; Marini, A.M.; Da Settimo, F.; Marinelli, L.; Limongelli, V.; Novellino, E. Acetic acid aldose reductase inhibitors bearing a five-membered heterocyclic core with potent topical activity in a visual impairment rat model. *J. Med. Chem.*, **2008**, *51*(11), 3182-3193.  
[<http://dx.doi.org/10.1021/jm701613h>] [PMID: 18452283]