

Accepted Manuscript

Hepatitis C virus: life cycle in cells, infection and host response, and analysis of molecular markers influencing the outcome of infection and response to therapy

Lynn B. Dustin, Barbara Bartolini, Maria R. Capobianchi, Mauro Pistello

PII: S1198-743X(16)30364-0

DOI: [10.1016/j.cmi.2016.08.025](https://doi.org/10.1016/j.cmi.2016.08.025)

Reference: CMI 704

To appear in: *Clinical Microbiology and Infection*


Received Date: 7 May 2016

Revised Date: 16 August 2016

Accepted Date: 25 August 2016

Please cite this article as: Dustin LB, Bartolini B, Capobianchi MR, Pistello M, Hepatitis C virus: life cycle in cells, infection and host response, and analysis of molecular markers influencing the outcome of infection and response to therapy, *Clinical Microbiology and Infection* (2016), doi: 10.1016/j.cmi.2016.08.025.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



1 **CLM-16-10263**

2

3 **Hepatitis C virus: life cycle in cells, infection and host response, and analysis of**
4 **molecular markers influencing the outcome of infection and response to therapy**

5

6 Lynn B. Dustin¹, Barbara Bartolini², Maria R. Capobianchi², Mauro Pistello^{3*}

7

8 ¹ *Kennedy Institute for Rheumatology and Peter Medawar Building for Pathogen Research,*
9 *University of Oxford, Oxford, United Kingdom*

10 ² *Laboratory of Virology, National Institute for Infectious Diseases "Lazzaro Spallanzani"*
11 *IRCCS, Rome, Italy.*

12 ³ *Virology Unit, Pisa University Hospital, and Virology Section and Retrovirus Center,*
13 *Department of Translational Research, University of Pisa, Pisa, Italy*

14

15 (Text: 3616 words)

16 * Corresponding author:

17 Mauro Pistello

18 Centro Retrovirus e Sezione Virologia

19 Dipartimento Ricerca Traslazionale e delle Nuove Tecnologie in Medicina e Chirurgia

20 Università di Pisa

21 Via San Zeno 37

22 I-56127, Pisa, Italy

23 Phone: + 39 050 221 3781

24 Fax: + 39 050 221 3524

25 E-mail: mauro.pistello@med.unipi.it

26

1 **Abstract**

2 Hepatitis C virus (HCV) is a major global health burden accounting for around 170
3 million chronic infections worldwide. Since its discovery, which dates back to about 30 years
4 ago, many details of the viral genome organization and the astonishing genetic diversity have
5 been unveiled but, owing to the difficulty of culturing HCV *in vitro* and obtaining fully
6 susceptible yet immunocompetent *in vivo* models, we are still a long way from the full
7 comprehension of viral life cycle, host cell pathways facilitating or counteracting infection,
8 pathogenetic mechanisms *in vivo*, and host defenses. Here, we illustrate the viral life cycle
9 into cells, describe the interplay between immune and genetic host factors shaping the course
10 of infection, and provide details of the molecular approaches currently used to genotype,
11 monitor replication *in vivo*, and studying the emergence of drug-resistant viral variants.

12
13 (137 words)

14
15
16 **Keywords:** Hepatitis C virus; Cell culture-derived HCV; Pathogenesis; Host response;
17 Broadly-reactive neutralizing antibodies; Genotype; Quantitative molecular assays; Drug
18 resistance; Monitoring drug resistance; Direct antiviral agents.

19

1 **HCV: A momentous virus for Virology**

2 The Hepatitis C virus (HCV) was identified in 1989. This year marks a departure from
3 traditional virology, based on isolation, cultivation, and biochemical studies, to modern
4 virology that uses molecular biology and biotechnology to discover, characterize, and
5 monitor viruses. HCV is indeed the first infectious agent discovered thanks to molecular
6 biology techniques that, owing to the difficulty to replicate the virus *in vitro*, have been
7 extensively used to define the molecular aspects of HCV biology.

8 Today HCV is making history again. Recently developed direct-acting antivirals
9 (DAAs) eliminate infection in over 90% treated individuals and are changing the idea that
10 antivirals, in general, can at most block viral replication and slow disease progression. Poorly
11 tolerated interferon (IFN)-based therapeutic regimens are being rapidly replaced with IFN-
12 free DAA regimens, and tissue damage in patients with advanced stages of disease is
13 stabilized and possibly reversed [1]. Further, detailed study of HCV's dependence on host
14 factors has permitted development of host-directed antiviral therapies. This review provides
15 an overview of some key aspects of viral interaction at cellular and host levels and illustrates
16 current methods to monitor viral replication and genotyping with particular emphasis on
17 fundamentals and recent clinical findings important to determine susceptibility or resistance to
18 DAAs.

20 **HCV life cycle and host-cell interactions *in vitro***

21 HCV belongs the Flaviviridae, a large family of enveloped, single-stranded RNA viruses that
22 is organized into the genera Hepacivirus, Flavivirus, Pestivirus, and Pegivirus [2], and that
23 includes many viruses transmitted by arthropods and growing matter of health concern [3].
24 The HCV life cycle is only partly understood; difficulties in establishing an *in vitro* model of
25 replication and the complex network of cell surface molecules used to mediate viral entry

1 have delayed comprehension of various molecular mechanisms [4, 5]. Briefly, as shown in
2 Figure 1, the HCV virion circulates in the bloodstream either as free-particle or surrounded
3 by host low-density lipoproteins [6], attaches onto the target cell membrane by sequential
4 binding of various receptor molecules, and enters into the cell by a clathrin-mediated
5 endocytosis process. Disruption of the viral capsid in the endocytic compartment releases the
6 9.6 kb single-stranded RNA genome of positive polarity into the cytoplasm. The RNA
7 genome is then directly translated at the rough endoplasmic reticulum (ER) in a single
8 polyprotein precursor of about 3,000 amino acid residues that is eventually cleaved by
9 cellular and viral proteases into ten mature products [7, 8]. These proteins, enlisted in the
10 order they are encoded, include the structural core and envelope glycoproteins E1 and E2,
11 and the following nonstructural proteins: p7 viroporin and nonstructural protein 2 (NS2) that
12 participate in virus assembly and release; NS3 and NS4A, the protease complex that, as
13 described in Chapters 3 and 4 of this Theme Section, is actively targeted by the protease
14 inhibitor class of DAAs; NS4B, a membrane-associated protein that mediates virus–host
15 interactions; NS5A, a zinc-binding and proline-rich hydrophilic phosphoprotein involved in
16 HCV RNA replication and targeted by NS5A inhibitor DAAs; and NS5B, the RNA-
17 dependent RNA polymerase targeted by nucleoside and non-nucleoside polymerase inhibitor
18 DAAs. New virions are assembled in an ER-derived compartment and released by exocytosis
19 following a Golgi-dependent secretory pathway. Along this process the virus undergoes
20 maturation and becomes surrounded by endogenous lipoproteins that, as described below, are
21 believed to help immune escape [4, 5]. Binding to host lipoproteins and envelope without
22 clearly discernable surface features confer to HCV virions low buoyant density and broad
23 size range (40–80 nm diameter) [5].

24 Lack of a reliable *in vitro* method to study HCV replication was due to scarce
25 adaptability of primary hepatic cells to *in vitro* propagation, no availability of viral isolates

1 adapted to *in vitro* culture, and large use of cell-to-cell transmission to disseminate infection
2 to neighbor cells. This mechanism has complicated identification of cellular receptors
3 necessary for viral entry and is believed to facilitate immunological escape, virus persistence,
4 and resistance to DAAs [9-12]. Also, receptor usage appears to depend upon cell type and
5 infection via free-particle or cell-to-cell transmission [4, 13].

6 The cell culture derived HCV (HCV_{cc}) and the HCV trans-complemented particles
7 (HCV_{TCP}) are among the most used methods to study replication of HCV replication *in vitro*
8 [4]. HCV_{cc} uses JFH1, a HCV genotype 2a strain isolated from a Japanese patient with
9 fulminant hepatitis and replicates in Huh-7, a human cell line from hepatocellular carcinoma
10 [14, 15]. HCV_{cc} generates infectious virus and, using either native or inter-genotype
11 recombinant JFH1 variants, has allowed identifying some HCV entry factors, defining virion
12 structure and biochemical properties, and testing DAA potency. HCV_{TCP}, described in detail
13 elsewhere [16, 17], employs pseudotyped HCV virions generated in packaging cells
14 transfected with viral proteins provided by different constructs. HCV_{TCP} can be theoretically
15 obtained from any isolate but support only single-round infection and are unable to spread.

16 HCV_{cc}, HCV_{TCP}, and basically all *in vitro* methods use Huh-7 cells that, although
17 permissive to HCV replication, differ from primary hepatocytes for different restriction
18 mechanisms, diverse localization of HCV receptors, and absence of the cell polarity observed
19 in hepatic tissue. As a result, viral entry, assembly, release, and cell-to-cell spread observed *in*
20 *vivo* is not completely reproduced *in vitro* [5]. HepG2 cell clones and hepatoma cells derived
21 from primary hepatocytes are permissive to HCV replication *in vitro* and should allow better
22 understanding of virus-host cell interplay. On this regard, a growing body of evidence shows
23 that host genetics impact disease progression, immune response, and antiviral therapy [18,
24 19]. Nucleotide polymorphism of IFN and IFN-stimulated genes, for instance, strongly
25 correlates with disease and therapy outcome to the point that host-targeting antivirals have

1 become an attractive field of drug research [20].

2

3 **The natural history of infection**

4 An estimated 130-200 million people worldwide are persistently infected with HCV, up
5 to 4 million new infections occur annually, and the majority of infected persons are unaware
6 of their infection status [21-23]. HCV spreads primarily through percutaneous contact with
7 infected blood. Before identification, many people became HCV-infected as a result of unsafe
8 injection practices, blood or blood products. Blood testing greatly reduced the risk of
9 iatrogenic exposure in developed countries; however, unsafe medical procedures remain an
10 important source of new infections particularly in resource-poor settings. In many developed
11 countries, most infections now occur amongst people who use injected or intranasal drugs.
12 Specific populations, notably human immunodeficiency virus (HIV)-positive men who have
13 sex with men, are at increased risk of sexual transmission. Vertical transmission and
14 transmission through piercing or tattooing are also possible [23, 24].

15 HCV causes hepatic inflammation and fibrosis that may progress sub-clinically over
16 decades. Long-term sequelae include cirrhosis, end-stage liver disease, and hepatocellular
17 carcinoma (HCC). In the Western world, chronic HCV infection is the leading indication for
18 liver transplant and the leading cause of end-stage liver disease, HCC, and liver-related death
19 [23, 25, 26]. Individual outcomes are highly variable, with many patients experiencing
20 minimal changes while others progress rapidly [23]. Fibrosis progression is uneven and may
21 accelerate with longer duration of infection; comorbid conditions such as HIV infection
22 accelerate disease progression [22]. A large meta-analysis reported the risk of cirrhosis at 7-
23 18% after 20 years and 41% after 30 years of infection [27]. Cirrhotic patients are at high risk
24 for hepatic decompensation (27.7-39.5% risk over five years) and HCC (2.8-7.4% in the first
25 year, and 8-16.1% over five years) [28]. Factors that increase the risks of fibrosis, cirrhosis,

1 and HCC include male sex, increasing age, alcohol use, and HIV co-infection.

2 Most infected individuals have no symptoms during the acute infection, although 15-
3 30% may experience non-specific symptoms [22]. Because most acute HCV infections are
4 sub-clinical, it is difficult to calculate how many patients clear HCV without treatment [22,
5 23]. This typically occurs in the first six months after exposure and is estimated at between
6 <20% and about 50% in different populations [23]; spontaneous clearance is affected by host
7 genetics, race, age, sex, and comorbidities such as HIV [29-32].

8

9 **The host response and the outcome of HCV infection**

10 *Innate immunity*

11 Soon after establishing infection in hepatic foci, HCV undergoes an exponential “ramp-
12 up” phase of replication [33]; the rate of increase decreases abruptly when cells in the liver
13 express a host of IFN-stimulated genes (ISGs) that limit HCV replication and spread [34].
14 Innate immunity is a first line of defense against HCV infection [35] and stimulates adaptive
15 immunity. HCV RNA binds to retinoic acid-inducible gene I, activating mitochondrial
16 antiviral signaling (MAVS) proteins; double-stranded RNA bound to Toll-like receptor-3
17 induces signaling via TIR domain-containing adaptor inducing IFN- β (TRIF). Both pathways
18 activate NF κ B and IRF3 translocation to the nucleus. Here, they promote expression of IFNs
19 and ISGs to inhibit viral replication, plus proinflammatory cytokines and chemokines to
20 recruit and activate immune cells. HCV’s NS3-4A protease specifically cleaves MAVS and
21 TRIF to dampen IFN induction [36, 37]. HCV may use several additional strategies to reduce
22 innate antiviral responses (reviewed in [36, 37]). ISG expression in HCV-infected
23 hepatocytes, even those with evidence of MAVS cleavage, demonstrates that these
24 mechanisms do not completely abolish innate immunity [38, 39]. Hepatocytes preferentially
25 express IFN- λ following HCV infection [40]. Dendritic cells, Kupffer cells, and other

1 nonparenchymal cells also recognize viral molecular patterns, contributing to IFN and
2 cytokine production and response without themselves harboring replicating HCV [34, 36].

3

4 *Role of adaptive immunity in the outcome of infection*

5 Innate antiviral responses can limit HCV replication and spread, but rarely mediate
6 elimination of infection without action by the adaptive immune response. Indeed, a hallmark
7 of HCV control is the arrival in the liver of T-lymphocytes that produce IFN- γ . Elevated
8 blood levels of transaminases, indicating hepatocyte cell death, are seen as viral loads
9 decline. Chronic infection is defined as infection lasting more than 6 months; spontaneous
10 clearance is rare but not unheard of after this point. Some key immunologic features of
11 spontaneous clearance and chronic infection are summarized in Table 1.

12 In resolving infections, adaptive immunity rapidly mounts a cell-mediated response
13 targeting multiple HCV epitopes and high-titer, broadly-reactive neutralizing antibodies
14 (bNAbs) [41, 42]. By targeting multiple epitopes, T-lymphocytes reduce viral opportunities
15 for immune escape. HCV's error-prone replication strategy permits rapid evolution, and
16 immune responses select for variants that escape recognition. Some immune escape
17 mutations are not tolerated because they impair viral fitness [43]. A second characteristic of
18 effective anti-HCV immunity is the preservation of polyfunctional T-lymphocyte activity.
19 CD8⁺ T-lymphocytes (and other immune cells) are dependent on help from HCV-specific
20 CD4⁺ T-lymphocytes; these are readily detected early after HCV infection regardless of its
21 outcome. In resolving infection, HCV-specific CD4⁺ T-lymphocytes support CD8⁺ T-
22 lymphocyte survival, proliferation, and antiviral activity. Effector T-lymphocytes accumulate
23 in the liver and slow HCV replication through cytokine release (IFN- γ , TNF- α) and killing of
24 infected cells. T-lymphocyte responses persist long after the virus has been cleared, and
25 mediate protection, albeit imperfect, against chronic infection in subsequent exposures. As

1 regards humoral immunity, bNAbs binding viral E2 and, to a lesser extent, E1 glycoproteins
2 are produced at high titer and within the first few months of infection. This results in
3 blocking HCV infection of its target cells. Key viral neutralization targets – i.e. conserved
4 domains required for hepatocyte infection - are hidden behind glycans, lipoproteins, and
5 hypervariable decoy domains. bNAbs bind the essential domains rather than the decoys [9,
6 44]. bNAbs may contribute to clearance because HCV must continuously infect new target
7 cells to maintain even an established infection [45].

8 In contrast to the broad and sustained T-lymphocyte responses in resolving HCV
9 infection, T-lymphocytes in persisting HCV infection may target a more limited set of
10 epitopes; often, an initially broad response narrows. Thus, fewer viral sequence changes are
11 required for immune escape. HCV that persists in chronic infection often shows evidence of
12 immune-mediated selection for variants that avoid recognition by CD8⁺ T-lymphocytes [34].
13 Where viral epitope sequences remain unchanged, HCV-specific T-lymphocyte responses are
14 characterized by progressive loss of function. CD4⁺ T-lymphocyte responses fail through a
15 mechanism independent of epitope escape [34, 42]. CD8⁺ T-lymphocytes, lacking CD4⁺ T-
16 lymphocyte help, lose effector functions, express markers associated with exhaustion, and
17 stop proliferating. Inflammatory T-lymphocytes, frequently not HCV-specific, infiltrate the
18 liver and may mediate tissue damage [46]. Finally, patients with chronic infection also
19 express neutralizing antibodies, but these may arise later and may be isolate-specific, often
20 targeting hypervariable epitopes with high potential for immune escape [9, 34].

21

22 *Role of host genetics in the outcome of infection*

23 HCV infection is more likely to persist in people bearing a set of unfavorable
24 polymorphisms in the IFN- λ locus; the same alleles are associated with failure of IFN- α -
25 based antiviral therapy [40]. High ISG expression in the infected liver is also a poor

1 prognostic indicator for IFN- α -based HCV treatment [47]. The cluster of IFN- λ locus
2 polymorphisms includes a frame-shift in the IFN- λ 4 gene; the favorable allele abolishes IFN-
3 λ 4 protein expression [40]. It has been proposed that IFN- λ 4 might regulate hepatocyte IFN
4 responsiveness, perhaps through negative feedback mechanisms. Alternatively, IFN- λ 4 (or a
5 linked polymorphism) may support prolonged innate immune activation and thereby interfere
6 with maturation of an adaptive immune response [48]. Whether IFN- λ acts strictly via innate
7 immunity, or influences adaptive immunity, is uncertain. Additional important
8 polymorphisms are in the HLA locus. These may influence the outcome of infection through
9 selection of immunodominant epitopes [43], cross-reactive responses [49], and interactions
10 with natural killer cells [50].

11

12 **Importance of viral load and genotyping in the outcome of infection and therapy**

13

14 *HCV viral load*

15 Monitoring of the course of infection and therapeutic response is based on HCV RNA
16 measurement in plasma or serum of patients and, under anti-HCV treatment, is aimed at
17 optimizing therapy duration, and prompting early discontinuation to prevent potential side
18 effects and reduce unnecessary costs. Baseline viral load, extent and sharpness of viremia
19 decay in the early phases of treatment (4 and 12 weeks), and undetectable HCV RNA at the
20 end of treatment represent key parameters guiding IFN-treatment.

21 With DAAs and according to current guidelines, HCV viral load monitoring during
22 therapy remains crucial for patient management with regard to futility rules and assessment
23 of therapy efficacy [51-53]. In addition, due to the rapid decay expected in almost all patients
24 treated with DAAs, HCV RNA monitoring is also used to assess patient adherence.

25 The methods to measure HCV viral load have greatly evolved since their initial

1 establishment. Today automated real-time reverse transcriptase-polymerase chain reaction
2 (RT-PCR) and transcription-mediated amplification (TMA) platforms from different vendors
3 are widely used throughout western countries (Table 2) [54,56]. These systems accurately
4 quantify HCV RNA within a broad linear range, with a lower limit of detection (LLOD)
5 sometimes even below the lower limit of quantification (LLOQ).

6 Despite good inter-assay agreement of linear range, there are important differences
7 concerning LLOD, LLOQ and low HCV RNA concentration, which are crucial for clinical
8 evaluation. Since decision to prolong or stop DDA therapy is often taken with a single
9 measurement, accurate quantification is important. To reduce uncertainty, several strategies
10 have been devised. For instance, with the Roche High Pure System/COBAS TaqMan assay
11 virological response has been set at 25 IU/ml, the LLOQ of this assay. Depending on their
12 LLOQ, other assays use lower levels [52-55]. Thus, the threshold and the number of patients
13 achieving virological response may differ depending on the assay and influence clinical
14 management [56-62]. For instance, patients receiving first generation DAA (Telaprevir)
15 combined with PegIFN and RBV and with HCV RNA undetectable at week 4 were used to
16 identify patients eligible for shortened treatment (24 vs 48 weeks) [63]. Based on this
17 criterion, in the OPTIMIZE study, for instance, 34% or 72% of patients would have been
18 eligible for shorter treatment if HCV RNA was measured with Abbott RealTime or Roche
19 High Pure System/COBAS TaqMan [59]. Thus, current guidelines recommend considering
20 assay performances and performing virological monitoring in the same laboratory for patient
21 management [51, 61].

22 As mentioned, HCV must continuously infect new target cells to maintain even an
23 established infection [45]. It is assumed therefore that clearance from each infected cell
24 occurs when viral replication has been halted for a sufficient time. However, even when HCV
25 RNA is not detected with the most sensitive assay, it is possible that minute, undetected

1 amounts of virus are still present. This “residual viremia”, firstly demonstrated for HIV with
2 highly sensitive tests developed *ad hoc*, with enhanced sensitivity, may indicate suboptimal
3 treatment and enhanced risk of virological failure [64]. To prevent this occurrence in HCV,
4 ultrasensitive versions using increased input sample volume and modified calibration curve
5 have been developed from commercial assays [65]. Preliminary data on patients receiving
6 first generation DAA-based therapy suggest that early achievement of HCV RNA values
7 undetectable with such ultrasensitive tests is predictive of sustained virological response [66].
8 Further studies are warranted to determine whether assays with enhanced sensitivity are
9 necessary for clinical management of patients treated with new DAAs.

10 HCV core antigen (HCV Ag) is also gaining importance for effective monitoring of
11 new generation DAAs. Architect HCV Core Antigen Test, one of most used systems,
12 quantitates down to 0.06 pg/mL HCV Ag, roughly corresponding to 700 IU/mL HCV RNA.
13 This marker proved useful in IFN-based therapy [67, 68] and good predictor of sustained
14 virological response in first generation DAA-based therapy [69, 70]. This analysis is less
15 expensive and time consuming compared to molecular assays, and should be considered
16 whenever high analytical sensitivity is not a must [68, 69].

17

18 *HCV genotyping*

19 HCV shows enormous genomic sequence variability. Currently there are seven
20 confirmed genotypes that are organized in 67 confirmed, 20 provisionally assigned, and 21
21 unassigned subtypes; median variability among genotypes and subtypes is approximately
22 33%, and 10%, respectively [71]. Because type and duration of treatment also depends on the
23 genotype, HCV genotyping is mandatory in patients eligible for antiviral therapy [72].
24 Further, owing to important barriers to resistance, genotyping is also important for DAA-
25 based therapies. As long as highly and broadly effective all oral combination therapies will be

1 available, it is conceivable the importance of HCV genotyping will decline in the future [72,
2 73].

3 Sequencing of conserved genomic regions is the gold standard for HCV genotyping,
4 but this method is limited to high-level laboratories; reverse hybridization or genotype-
5 specific real-time RT-PCR are easier-to-perform and routinely used but, particularly first
6 generation assays that targeted the 5' untranslated region (5'UTR), can misclassify isolates
7 especially at subtype level. A consistent improvement has been achieved by including coding
8 regions in other targets, e.g. NS5B and core protein, possessing non-overlapping sequences
9 amongst genotypes and subtypes [73].

10

11 *Analysis of HCV resistance*

12 Because of high replication rate and no proofreading activity of the viral polymerase,
13 HCV is highly variable and each possible single mutation and combination of mutations may
14 arise every day in a given infected individual. Genome plasticity and drug-driven selection
15 create the conditions for the emergence of resistant variants [33] and, as a consequence, most
16 mutations associated with resistance are located within the drug target regions (Figure 2) [74-
17 76]. Despite the reduced fitness, such variants rapidly overgrow wild-type viruses and during
18 this process they may accumulate additional, fitness-restoring, mutations [76]. Single drug,
19 drug family and genotype/subtype often influence the emerging mutations. In turn, each DAA
20 class displays different genetic barrier to resistance and cross-resistance between drugs
21 inhibiting NS3 protease and NS5A is frequently observed.

22 Direct sequencing of PCR products has been the method of choice for investigating the
23 presence of mutations conferring antiviral resistance. Unfortunately, this approach analyzes
24 the predominant species and, as observed with HIV-1, variants below 20% may escape
25 detection despite being crucial during the initial phases of resistance development [78]. Next

1 generation sequencing (NGS) detects and quantitates variants present at frequencies as low as
2 0.5% [79] and, therefore, permits early detection of resistance mutations, definition of their
3 kinetics, and progressive disappearance after treatment suspension [80-83]. Although there
4 are no commercial kits, many laboratories analyze NS3-4A, NS5A, and NS5B regions with
5 NGS but its use in clinical practice is still limited and requires expert guidance for
6 interpretation.

7 Natural variants carrying resistance-associated mutations have been repeatedly found
8 by conventional and NGS analyses but, with the exception of searching for Q80K mutation in
9 patients infected with genotype 1a and to be enrolled for treatment with Simeprevir, there is
10 no agreement on the usefulness of performing resistance testing before DAA-based therapy
11 [84-86]. Even the use of resistance testing in failing cases is questioned since: *i.* The
12 emergence of resistant variants does not account for all virological failures; *ii.* The resistant
13 variants rapidly decline (but do not always disappear) after stopping treatment [86, 87]; *iii.*
14 The different classes and combination of drugs at disposal enormously increase the genetic
15 barrier to resistance.

16 According to most recent guidelines and since NS5A resistance mutations can persist
17 for over two years post-treatment [87], resistance testing is recommended for patients who
18 need urgent treatment and have failed previous treatment with NS5A inhibitors. In contrast,
19 the utility of routine testing for NS5B mutations is questionable. Resistance to nucleotide
20 analogues targeting NS5B (i.e. Sofosbuvir) is rare and S282T, the only mutation (so far)
21 associated with Sofosbuvir resistance, confers fitness disadvantage. As mentioned, however,
22 unfit variants continue to evolve during treatment and select for compensatory mutations that
23 counterbalance fitness loss.

24

25

1 **Conclusions**

2

3 Understanding the HCV life cycle and host factors that hinder or allow persistence of
4 infection is crucial to develop post-exposure and prophylactic measures for the global control
5 of HCV. Only a fraction of the global patient population has been diagnosed and, in spite of
6 aforementioned remarkable advances in antiviral therapy, less than 1% has been treated [88-
7 90] and those who have been treated remain vulnerable to re-infection [91]. Therapy alone is
8 therefore not sufficient to eliminate the global burden of HCV infection and chronic liver
9 disease in the near future. In contrast, individuals who have mounted a successful immune
10 response against HCV have a substantial degree of protection upon subsequent exposures.
11 This suggests that effective immunological strategies to protect against persistent infection
12 are feasible [42, 89] and likely capable to face HCV's tremendous genetic diversity [71].

13

14 **Transparency declaration**

15

16 The authors declare no conflicts of interest and acknowledge the NIH grants R01 AI060561
17 and R01 AI089957 (awarded to LBD), Fondo Ricerca corrente (MRC), and FAS Salute 2016
18 “UNAVIR: *Malattie Virali Rare: una strategia innovativa per combatterle con un unico*
19 *agente antivirale*”, CUP B32I14001060002 (MP).

20 All authors wrote, revised and commented on the manuscript at all stages.

21

1 **References**

- 2 1. Martin NK, Vickerman P, Dore GJ, Grebely J, Miners A, Cairns J et al. Prioritization of
3 HCV treatment in the direct-acting antiviral era: An economic evaluation. *J Hepatol*
4 2016; 65: 17-25.
- 5 2. Simmonds P. The origin of hepatitis C virus. *Curr Top Microbiol Immunol* 2013; 369:
6 1-15.
- 7 3. Choumet V, Despres P. Dengue and other flavivirus infections. *Rev Sci Tech* 2015; 34:
8 473-8, 467-72.
- 9 4. Maggi F, Focosi D, Pistello M. How current direct-acting antiviral and novel cell
10 culture systems for HCV are shaping therapy and molecular diagnosis of chronic HCV
11 infection. *Curr Drug Targets* 2015 Aug 6.
- 12 5. Paul D, Madan V, Bartenschlager R. Hepatitis C virus RNA replication and assembly:
13 Living on the fat of the land. *Cell Host Microbe* 2014; 16: 569-79.
- 14 6. Andre P, Komurian-Pradel F, Deforges S, Perret M, Berland JL, Sodoyer M et al.
15 Characterization of low- and very-low-density hepatitis C virus RNA-containing
16 particles. *J Virol* 2002; 76: 6919-28.
- 17 7. Lohmann V. Hepatitis C virus RNA replication. *Curr Top Microbiol Immunol* 2013;
18 369: 167-98.
- 19 8. Niepmann M. Hepatitis C virus RNA translation. *Curr Top Microbiol Immunol* 2013;
20 369: 143-66.
- 21 9. Ball JK, Tarr AW, McKeating JA. The past, present and future of neutralizing
22 antibodies for hepatitis C virus. *Antiviral Res* 2014; 105:100-11.
- 23 10. Timpe JM, Stamataki Z, Jennings A, Hu K, Farquhar MJ, Harris HJ et al. Hepatitis C
24 virus cell-cell transmission in hepatoma cells in the presence of neutralizing antibodies.
25 *Hepatology* 2008; 47: 17-24.

- 1 11. Xiao F, Fofana I, Heydmann L, Barth H, Soulier E, Habersetzer F et al. Hepatitis C
2 virus cell-cell transmission and resistance to direct-acting antiviral agents. *PLoS*
3 *Pathogens* 2014; 10: e1004128.
- 4 12. Yoon JC, Lim JB, Park JH, Lee JM. Cell-to-cell contact with hepatitis C virus-infected
5 cells reduces functional capacity of natural killer cells. *J Virol* 2011; 85: 12557-69.
- 6 13. Barretto N, Sainz B, Jr., Hussain S, Uprichard SL. Determining the involvement and
7 therapeutic implications of host cellular factors in hepatitis C virus cell-to-cell spread. *J*
8 *Virol* 2014; 88: 5050-61.
- 9 14. Lindenbach BD, Evans MJ, Syder AJ, Wölk B, Tellinghuisen TL, Liu CC et al.
10 Complete replication of hepatitis C virus in cell culture. *Science* 2005; 309: 623-6.
- 11 15. Wakita T, Pietschmann T, Kato T, Date T, Miyamoto M, Zhao Z et al. Production of
12 infectious hepatitis C virus in tissue culture from a cloned viral genome. *Nat Med* 2005;
13 11: 791-6.
- 14 16. Steinmann E, Brohm C, Kallis S, Bartenschlager R, Pietschmann T. Efficient trans-
15 encapsidation of hepatitis C virus RNAs into infectious virus-like particles. *J Virol*
16 2008; 82: 7034-46.
- 17 17. Suzuki R, Saito K, Kato T, Shirakura M, Akazawa D, Ishii K et al. Trans-
18 complemented hepatitis C virus particles as a versatile tool for study of virus assembly
19 and infection. *Virology* 2012; 432: 29-38.
- 20 18. Mina MM, Luciani F, Cameron B, Bull RA, Beard MR, Booth D et al. Resistance to
21 hepatitis C virus: Potential genetic and immunological determinants. *Lancet Infect Dis*
22 2015; 15: 451-60.
- 23 19. Yang DR, Zhu HZ. Hepatitis C virus and antiviral innate immunity: Who wins at tug-
24 of-war? *World J Gastroenterol* 2015; 21: 3786-3800.
- 25 20. Baugh JM, Garcia-Rivera JA, Gallay PA. Host-targeting agents in the treatment of

- 1 hepatitis C: A beginning and an end? *Antiviral Res* 2013; 100: 555-61.
- 2 21. Mohd Hanafiah K, Groeger J, Flaxman AD, Wiersma ST. Global epidemiology of
3 hepatitis C virus infection: New estimates of age-specific antibody to HCV
4 seroprevalence. *Hepatology* 2013; 57: 1333-42.
- 5 22. Hajarizadeh B, Grebely J, Dore GJ. Epidemiology and natural history of HCV
6 infection. *Nat Rev Gastroenterol Hepatol* 2013; 10: 553-62.
- 7 23. Westbrook RH, Dusheiko G. Natural history of hepatitis C. *J Hepatol* 2014; 61: S58-68.
- 8 24. Webster DP, Klenerman P, Dusheiko GM. Hepatitis C. *Lancet* 2015; 385: 1124-35.
- 9 25. Perz JF, Armstrong GL, Farrington LA, Hutin YJ, Bell BP. The contributions of
10 hepatitis B virus and hepatitis C virus infections to cirrhosis and primary liver cancer
11 worldwide. *J Hepatol* 2006; 45: 529-38.
- 12 26. El-Serag HB. Epidemiology of viral hepatitis and hepatocellular carcinoma.
13 *Gastroenterology* 2012; 142: 1264-73.
- 14 27. Thein HH, Yi Q, Dore GJ, Krahn MD. Estimation of stage-specific fibrosis progression
15 rates in chronic hepatitis C virus infection: A meta-analysis and meta-regression.
16 *Hepatology* 2008; 48: 418-31.
- 17 28. Xu F, Moorman AC, Tong X, Gordon SC, Rupp LB, Lu M et al. All-cause mortality
18 and progression risks to hepatic decompensation and hepatocellular carcinoma in
19 patients infected with hepatitis C virus. *Clin Infect Dis* 2016; 62: 289-97.
- 20 29. Thomas DL, Thio CL, Martin MP, Qi Y, Ge D, O'Huigin C et al. Genetic variation in
21 IL28b and spontaneous clearance of hepatitis C virus. *Nature* 2009; 461: 798-801.
- 22 30. Wang CC, Krantz E, Klarquist J, Krows M, McBride L, Scott EP et al. Acute hepatitis
23 C in a contemporary US cohort: Modes of acquisition and factors influencing viral
24 clearance. *J Infect Dis* 2007; 196: 1474-82.
- 25 31. Grebely J, Page K, Sacks-Davis R, van der Loeff MS, Rice TM, Bruneau J et al. The

- 1 effects of female sex, viral genotype, and il28b genotype on spontaneous clearance of
2 acute hepatitis C virus infection. *Hepatology* 2013; 59: 109-20.
- 3 32. Kenny-Walsh E. Clinical outcomes after hepatitis C infection from contaminated anti-D
4 immune globulin. Irish hepatology research group. *N Engl J Med* 1999; 340: 1228-33.
- 5 33. Dustin LB. Too low to measure, infectious nonetheless. *Blood* 2012; 119: 6181-2.
- 6 34. Dustin LB, Cashman SB, Laidlaw SM. Immune control and failure in HCV infection--
7 tipping the balance *J Leukoc Biol* 2014; 96: 535-48.
- 8 35. Schoggins JW, Rice CM. Innate immune responses to hepatitis C virus. *Curr Top*
9 *Microbiol Immunol.* 2013; 369: 219-42.
- 10 36. Horner SM, Gale M, Jr. Regulation of hepatic innate immunity by hepatitis C virus. *Nat*
11 *Med* 2013; 19: 879-88.
- 12 37. Li K, Lemon SM. Innate immune responses in hepatitis C virus infection. *Semin*
13 *Immunopathol* 2013; 35: 53-72.
- 14 38. Sheahan T, Imanaka N, Marukian S, Dorner M, Liu P, Ploss A, Rice CM. Interferon
15 lambda alleles predict innate antiviral immune responses and hepatitis C virus
16 permissiveness. *Cell Host Microbe* 2014; 15: 190-202.
- 17 39. Wieland S, Makowska Z, Campana B, et al. Simultaneous detection of hepatitis C virus
18 and interferon stimulated gene expression in infected human liver. *Hepatology* 2014;
19 59: 2121-30.
- 20 40. Laidlaw SM, Dustin LB. Interferon lambda: Opportunities, risks, and uncertainties in
21 the fight against HCV. *Front Immunology* 2014; 5: 545.
- 22 41. Holz L, Rehermann B. T cell responses in hepatitis C virus infection: Historical
23 overview and goals for future research. *Antiviral Res* 2015; 114: 96-105.
- 24 42. Honegger JR, Zhou Y, Walker CM. Will there be a vaccine to prevent HCV infection?
25 *Semin Liver Dis* 2014; 34: 79-88.

- 1 43. Dazert E, Neumann-Haefelin C, Bressanelli S, Fitzmaurice K, Kort J, Timm J et al.
2 Loss of viral fitness and cross-recognition by CD8+ T cells limit HCV escape from a
3 protective HLA-B27-restricted human immune response. *J Clin Invest* 2009; 119: 376-
4 86.
- 5 44. Cashman SB, Marsden BD, Dustin LB. The humoral immune response to HCV:
6 Understanding is key to vaccine development. *Front Immunol* 2014; 5: 550.
- 7 45. de Jong YP, Dorner M, Mommersteeg MC, Xiao JW, Balazs AB, Robbins JB et al.
8 Broadly neutralizing antibodies abrogate established hepatitis C virus infection. *Sci*
9 *Transl Med* 2014; 6: 254ra129.
- 10 46. Klenerman P, Thimme R. T cell responses in hepatitis C: The good, the bad and the
11 unconventional. *Gut* 2012; 61: 1226-34.
- 12 47. Sarasin-Filipowicz M, Oakeley EJ, Duong FH, Christen V, Terracciano L, Filipowicz
13 W et al. Interferon signaling and treatment outcome in chronic hepatitis C. *Proc Natl*
14 *Acad Sci USA* 2008; 105: 7034-9.
- 15 48. Park SH, Rehermann B. Immune responses to HCV and other hepatitis viruses.
16 *Immunity* 2014; 40: 13-24.
- 17 49. Cornberg M, Wedemeyer H. Hepatitis C virus infection from the perspective of
18 heterologous immunity. *Curr Opin Virol* 2016; 16: 41-8.
- 19 50. Cheent K, Khakoo SI. Natural killer cells and hepatitis C: Action and reaction. *Gut*
20 2011; 60: 268-78.
- 21 51. European Association for the Study of the Liver. EASL Recommendations on
22 Treatment of Hepatitis C 2015. *J Hepatol* 2015; 63: 199-236.
- 23 52. Sarrazin C, Wedemeyer H, Cloherty G, Cohen DE, Chevaliez S, Herman C et al.
24 Importance of very early HCV RNA kinetics for prediction of treatment outcome of
25 highly effective all oral direct acting antiviral combination therapy. *J Virol Methods*

- 1 2015; 214: 29-32.
- 2 53. AASLD/IDSA HCV Guidance Panel. Hepatitis C guidance: AASLD-IDSA
3 recommendations for testing, managing, and treating adults infected with hepatitis C
4 virus. *Hepatology* 2015; 62: 932-54.
- 5 54. Cobb B, Pockros PJ, Vilchez RA, Vierling JM. HCV RNA viral load assessments in
6 the era of direct-acting antivirals. *Am J Gastroenterol* 2013; 108: 471–5.
- 7 55. Schalasta G, Speicher A, Börner A, Enders M. Performance of the new Aptima HCV
8 Quant Dx assay in comparison to the COBAS TaqMan HCV2 assay for use with the
9 High Pure System in the Detection and Quantification of HCV RNA in plasma or
10 serum. *J Clin Microbiol* 2016; 54: 1101-7.
- 11 56. Fevery B, Susser S, Lenz O, Perner D, Picchio G, Sarrazin C et al. HCV RNA
12 quantification with different assays: implications for protease-inhibitor-based response-
13 guided therapy. *Antivir Ther* 2014; 19: 559–67.
- 14 57. Taylor N, Haschke-Becher E, Greil R, Strasser M, Oberkofler H. Performance
15 characteristics of the COBAS Ampliprep/COBAS TaqMan v2.0 and the Abbott
16 RealTime hepatitis C assays – implications for response-guided therapy in genotype 1
17 infections. *Antivir Ther* 2014; 19: 449–54.
- 18 58. Cloherty G, Cohen D, Sarrazin C, Wedemeyer H, Chevaliez S, Herman C et al. HCV
19 RNA assay sensitivity impacts the management of patients treated with direct-acting
20 antivirals. *Antivir Ther* 2015; 20: 177–83.
- 21 59. Sarrazin C, Dierynck I, Cloherty G, Ghys A, Janssen K, Luo D et al. An OPTIMIZE
22 Study retrospective analysis for the management of telaprevir-treated HCV patients
23 using the Abbott RealTime HCV RNA assay. *J Clin Microbiol* 2015; 53: 1264–9.
- 24 60. Vermehren J, Aghemo A, Falconer K, Susser S, Lunghi G, Zeuzem S et al. Clinical
25 significance of residual viremia detected by two real-time PCR assays for response-

- 1 guided therapy of HCV genotype 1 infection. *J Hepatol* 2014; 60: 913–9.
- 2 61. Wiesmann F, Naeth G, Berger A, Hirsch HH, Regenass S, Ross RS et al. Multicentric
3 performance analysis of HCV quantification assays and its potential relevance for HCV
4 treatment. *Med Microbiol Immunol* 2016; 205: 263-8.
- 5 62. Strassl R, Rutter K, Stättermayer AF, Beinhardt S, Kammer M, Hofer H et al. Real-
6 Time PCR Assays for the Quantification of HCV RNA: Concordance, Discrepancies
7 and Implications for Response Guided Therapy. *PLoS One* 2015; 10: e0135963.
- 8 63. Fevery B, Susser S, Lenz O, Cloherty G, Perner D, Gilles L et al. Comparison of three
9 quantitative HCV RNA assays in samples from HCV genotype 1- or 4-infected patients
10 treated with the NS3/4A protease inhibitor simeprevir. *J Clin Virol* 2015; 72: 133-40.
- 11 64. Sahu GK. Potential implication of residual viremia in patients on effective antiretroviral
12 therapy. *AIDS Res Hum Retroviruses* 2015; 31: 25-35.
- 13 65. Amendola A, Bloisi M, Marsella P, Sabatini R, Bibbò A, Angeletti C, Capobianchi MR.
14 Standardization and performance evaluation of "modified" and "ultrasensitive" versions
15 of the Abbott RealTime HIV-1 assay, adapted to quantify minimal residual viremia. *J*
16 *Clin Virol* 2011; 52: 17-22.
- 17 66. Garbuglia AR, Visco-Comandini U, Lionetti R, et al. Ultrasensitive HCV-RNA
18 quantification in PEG-Interferon, Telaprevir and Ribavirin standard treatment protocol:
19 new insights into viral clearance dynamics and treatment outcome predictors.
20 Submitted.
- 21 67. Loggi E, Cursaro C, Scuteri A, Grandini E, Panno AM, Galli S et al. Patterns of HCV-
22 RNA and HCV core antigen in the early monitoring of standard treatment for chronic
23 hepatitis C. *J Clin Virol* 2013; 56: 207–11.
- 24 68. Vermehren J, Susser S, Berger A, Perner D, Peiffer KH, Allwinn R et al. Clinical utility
25 of the ARCHITECT HCV Ag assay for early treatment monitoring in patients with

- 1 chronic hepatitis C genotype 1 infection. *J Clin Virol* 2012; 55: 17–22.
- 2 69. Chevaliez S, Soulier A, Poiteau L, Bouvier-Alias M, Pawlotsky JM. Clinical utility of
3 hepatitis C virus core antigen quantification in patients with chronic hepatitis C. *J Clin*
4 *Virol* 2014; 61: 145–8.
- 5 70. Garbuglia AR, Lionetti R, Lapa D, Taibi C, Visco-Comandini U, Montalbano M et al.
6 The clinical significance of HCV core antigen detection during Telaprevir/Peg-
7 Interferon/Ribavirin therapy in patients with HCV 1 genotype infection. *J Clin Virol*
8 2015; 69: 68-73.
- 9 71. Smith DB, Bukh J, Kuiken C, Muerhoff AS, Rice CM, Stapleton JT et al. Expanded
10 classification of hepatitis C virus into 7 genotypes and 67 subtypes: updated criteria and
11 genotype assignment web resource. *Hepatology* 2014; 59: 318-27.
- 12 72. Lange CM, Jacobson IM, Rice CM, Zeuzem S. Emerging therapies for the treatment of
13 hepatitis C. *EMBO Mol Med* 2014; 6: 4-15.
- 14 73. Bowden DS, Berzsenyi MD. Chronic hepatitis C virus infection: genotyping and its
15 clinical role. *Future Microbiol* 2006; 1: 103-12.
- 16 74. Sarrazin C, Zeuzem S. Resistance to direct antiviral agents in patients with hepatitis C
17 virus infection. *Gastroenterology* 2010; 138: 447–62.
- 18 75. Kieffer TL, Kwong AD, Picchio GR. Viral resistance to specifically targeted antiviral
19 therapies for hepatitis C (STAT-Cs). *J Antimicrob Chemother* 2010; 65: 202–12.
- 20 76. Poveda E, Soriano V. Drug resistance testing in hepatitis C therapy. *Future Virology*
21 2012; 7:309–21.
- 22 77. Buggisch P, Sarrazin C, Mauss S, Hinrichsen H, Simon KG, Vermehren J et al.
23 Sofosbuvir-based treatment under real life conditions in Germany (The SOFGER trial).
24 *J Hepatol* 2015; 62: S622.
- 25 78. Thielen A, Martini N, Thiele B, Däumer M. Validation of HIV-1 drug resistance testing

- 1 by deep sequencing: insights from comparative Sanger sequencing. *Antivir Ther* 2014;
2 19: A78.
- 3 79. Gregori J, Esteban JI, Cubero M, Garcia-Cehic D, Perales C, Casillas R et al. Ultra-
4 deep pyrosequencing (UDPS) data treatment to study amplicon HCV minor variants.
5 *PLoS One* 2013; 8: e83361.
- 6 80. Bartolini B, Selleri M, Garbuglia AR, Giombini E, Taibi C, Lionetti R et al. HCV NS3
7 quasispecies in liver and plasma and dynamics of telaprevir-resistant variants in
8 breakthrough patients assessed by UDPS: A case study. *J Clin Virol* 2015; 72: 60-5.
- 9 81. Bartolini B, Lionetti R, Giombini E, Sias C, Taibi C, Montalbano M et al. Dynamics of
10 HCV genotype 4 resistance-associated variants during virologic escape with
11 pIFN/RBV+daclatasvir: a case study using ultra deep pyrosequencing. *J Clin Virol*
12 2015; 66: 38-43.
- 13 82. Trémeaux P, Caporossi A, Ramière C, Santoni E, Tarbouriech N, Thélu MA et al.
14 Amplification and pyrosequencing of near-full-length hepatitis C virus for typing and
15 monitoring antiviral resistant strains. *Clin Microbiol Infect* 2016; 22: 460.e1-460.e10.
- 16 83. Bergfors A, Leenheer D, Bergqvist A, Ameer A, Lennerstrand J. Analysis of hepatitis C
17 NS5A resistance associated polymorphisms using ultra deep single molecule real time
18 (SMRT) sequencing. *Antiviral Res* 2016; 126: 81-9.
- 19 84. Vermehren J, Sarrazin C. The role of resistance in HCV treatment. *Best Pract Res Clin*
20 *Gastroenterol* 2012; 26: 487–503.
- 21 85. Pawlotsky JM. Hepatitis C treatment: the data flood goes on-an update from the liver
22 meeting 2014. *Gastroenterology* 2015; 148: 468–79.
- 23 86. Sarrazin C. The importance of resistance to direct antiviral drugs in HCV infection in
24 clinical practice. *J Hepatol* 2016; 64: 486-504.
- 25 87. Paolucci S, Fiorina L, Mariani B, Landini V, Gulminetti R, Novati S et al. Development

- 1 and persistence of DAA resistance associated mutations in patients failing HCV
2 treatment. *J Clin Virol* 2015; 72: 114-8.
- 3 88. Rehermann B. HCV in 2015: Advances in hepatitis c research and treatment. *Nat Rev*
4 *Gastroenterol Hepatol* 2016; 13: 70-2.
- 5 89. Cox AL. *Medicine*. Global control of hepatitis C virus. *Science* 2015; 349: 790-1.
- 6 90. Thomas DL. Global control of hepatitis C: Where challenge meets opportunity. *Nat*
7 *Med* 2013; 19: 850-8.
- 8 91. Sulkowski MS, Gardiner DF, Rodriguez-Torres M, et al. Daclatasvir plus sofosbuvir for
9 previously treated or untreated chronic HCV infection. *N Engl J Med* 2014; 370: 211-
10 21.
- 11
- 12

1 **Legend to figures**

2

3 **Figure 1. The HCV replication cycle.** The seven steps of the viral life cycle, indicated in the
4 white boxes, are the following: Attachment, the viral particle, surrounded with lipoproteins,
5 binds the target cells by interacting with several receptors some, most of which shown in
6 figure, considered essential other accessory; Entry: following attachment, the virus enters
7 through clathrin-mediated endocytosis; Uncoating: the cellular and viral membranes fuse and
8 the capsid is disorganized with a process triggered by the low pH of the endosome. After
9 uncoating the positive-strand RNA genome is released into the cytoplasm; Translation: the
10 genomic RNA is directly translated in a polyprotein precursor that is then cleaved into single
11 proteins by both host and viral proteases; Replication: the non-structural proteins and some
12 host factors form a replication complex that synthesized multiple copies of the HCV RNA
13 genome via a minus-strand replicative intermediate; Assembly and maturation: packaging of
14 viral progeny takes place in the endoplasmic reticulum from which the virion acquires the
15 envelope with E1 and E2 glycoproteins. Maturation and association with endogenous
16 lipoproteins to form lipoviral particles immediately follow; Release: virions are released from
17 the cells most likely by exocytosis or transmitted to other cells via a cell-free mechanism.

18

19 **Figure 2. A schematic of the mutations conferring resistance to NS3, NS5A and NS5B**
20 **inhibitors.** Numbers refer to the amino acid positions correlated to resistance in various HCV
21 genotypes.

22


Table 1. Immunologic characteristics of resolving and persisting HCV infections

	<i>Resolving infection</i>	<i>Persisting infection</i>
T-cell responses	Broadly focused: limits viral escape options	Restricted number of epitopes facilitates viral escape
Duration of T-cell response	Rapid, sustained even after clearance	Wanes over time
HCV specific CD4+ T-cells	Persist	Disappear
HCV-specific CD8+ T-cells	<ul style="list-style-type: none"> • Proliferate and expand • Polyfunctional: produce IFNγ, TNFα • Express perforin 	<ul style="list-style-type: none"> • Lose proliferative capacity • Fewer polyfunctional cells • Express markers of exhaustion
Neutralizing antibody	<ul style="list-style-type: none"> • Early • High titer • Broadly-reactive 	<ul style="list-style-type: none"> • Isolate-specific • Viral sequence evolution outpaces antibody reactivity
IFN λ SNPs		
rs12979860	C	T
rs368234815	TT	Δ G
rs8099917	T	G


Table 2. Overview and features of automated assays marketed and licensed to quantitate HCV RNA in plasma or serum samples

<i>Test name</i>	<i>Manufacturer</i>	<i>Technology used</i>	<i>Level of automation</i>	<i>LLOD^a</i> <i>(IU/ml)</i>	<i>LLOQ^b</i> <i>(IU/ml)</i>	<i>Linear range</i> <i>(IU/ml)</i>
Artus Hepatitis C QS-RGQ	Qiagen	Real time RT-PCR	Moderate; extraction and amplification /detection in separate instruments; a few manual steps	15	20	up to 1.77x10 ⁷
CobasAmpliprep/ CobasTaqMan v2.0	Roche	Real time RT-PCR	High; extraction and amplification /detection in separate instruments; sample loading can be fully automated	15	15	up to 1.7x10 ⁸
Abbott RealTime HCV	Abbott	Real time RT-PCR	High; extraction from primary tube; amplification/detection in a separate instrument; a few manual steps	12	12	up to 1.0x10 ⁸
Versant HCV RNA 2.0	Siemens	Real time RT-PCR	High; extraction from primary tube; amplification/detection in a separate instrument	15	15	up to 1.0 x 10 ⁸
Aptima HCV Quant Dx Assay	Hologic	TMA ^c	Fully automated; extraction/amplification /detection all in one instrument, starting from primary tube)	4	12	up to 1.0x10 ⁸

^a Lower limit of detection;^b Lower limit of quantification;^c Transcription-mediated amplification.



ACCEPTED



ACCEPTED