

Elsevier Editorial System(tm) for Food Research International  
Manuscript Draft

Manuscript Number: FOODRES-D-14-01201R1

Title: Evolution of the Anisakis risk management in the European and Italian context

Article Type: Review Article

Keywords: Anisakids, risk management, European legislation, raw fish products.

Corresponding Author: Dr. Andrea Armani,

Corresponding Author's Institution:

First Author: Priscilla D'Amico, Dr.

Order of Authors: Priscilla D'Amico, Dr.; Renato Malandra, Dr.; Francesco Costanzo, Dr.; Lorenzo Castigliano, Dr.; Alessandra Guidi, Prof.; Daniela Gianfaldoni, Professor; Andrea Armani

Abstract: Due to the social and legislative implications, the presence of *Anisakis* spp. larvae in fishery products has become a concern for both the consumers and the official Control Authorities. The issuance of a large number of provisions, aimed at better managing fish products intended to be consumed raw or almost raw and the associated risks, resulted in a very complicate legal framework. In this work, we analyzed the evolution of the normative through an overview on the local and international legislation, focusing on issues that are of practical interest for Food Business Operators (FBOs) in the fishery chain. In addition, we performed a survey across the Departments of Prevention of the Italian Local Health Authorities (LHA) and the main fish markets in Italy to collect the operating procedures and the monitoring plans. Overall, we found many differences, due to the absence of a national reference standard for the management of the *Anisakis* risk. From this examination, it turns clear that only a participation of all the involved Institutions, a strategy of synergistic interventions, as well as a correct training of FBOs, can result in an effective risk management and a proper risk communication, which should overcome states of confusion and unnecessary negative impacts on the economy.

**Dear Editor,**

We would like to submit the following manuscript for possible publication:

***“Evolution of the Anisakis risk management in the European and Italian context”***

The Anisakis risk management has recently been a source of great concern worldwide for both consumers and the Official Authorities in charge for controls. At the international levels, authorities, such as the *Codex Alimentarius* Commission, the EFSA and the U.S. FDA have recognized the need to face this issue through the adoption of preventive measures, capable to reduce, or even eliminate, the risk for public health.

The increased perception of this risk has determined, over the years, the issuance of a large number of international provisions, which have created a legal framework aimed at better managing fish products intended to be consumed raw or almost raw and its associated risks.

In this manuscript we revisited the evolution of the legislation related to the Anisakis risk, with an excursus of all European and Italian amendments issued from the late 60s to date. In particular, we addressed those aspects that, in our opinion, need to be reconsidered. Moreover, thanks to the collaboration of an Official Veterinary who has been working for more than 30 years at the wholesale market of Milan, we took into consideration many practical issues that are of interest for people working in the fishery chain.

Our aim was not only to provide an update on this topic, but to also analyze the data obtained from a survey of operating procedures and the self-monitoring plans, gathered at the Department of Prevention of the Italian Regional Health Authorities (AZ. USL) and at the main fish markets.

Other than to the academic world, this work is addressed to all those, who daily deal with the management of the fishery products as Food Business Operators (manufacturers, vendors, caterers), responsible for self-monitoring plans, and Official Authorities, to better clarify the tasks that they are called to fulfill.

Best regards,

Andrea Armani

Dear Editor,

we revised the manuscript according to the suggestions of the Reviewers. Our answers, comments and rebuttals are listed below.

The modified parts have been written in green to facilitate the revision process.

**Reviewer #1:** Manuscript Number: FOODRES-D-14-01201

Title: Evolution of the Anisakis risk management in the European and Italian context

This work carries out an overview on the legislation about Anisakids which are the most important fish parasites from a sanitary point of view.

The content is of interest; however, some important deficiencies are included:

(AZ. USL) The authors have to explain its meaning.

**Az. USL is the acronym of Azienda Unità Sanitaria Locale. Its name, in English, is Local Health Authority (the name in the text has been modified from Regional Health Authorities to Local Health Authorities). Az. USL has been removed from the text to avoid confusion.**

The authors affirm: "In fact, hypersensitivity reactions may also occur either via or with cooked fish, since allergens are thermostable". There are not evidences about this possibility. Nowadays, we can accept that allergic reactions only occur when live larva are ingested in raw or undercooked parasitized fish.

**The sentence has been modified and, according to the suggestions of reviewer N.3, a short paragraph on the ongoing debate on the necessity of larvae being alive to induce allergy the first time has been added (line 134-154)**

Anisakiasis should be changed by anisakiosis. According to the following paper: T. KASSAI, M. CORDERO DEL CAMPILLO, J. EUZEBY, S. GAAFAR, Th. HIEPE and C.A. HIMONAS Standardized Nomenclature of Animal Parasitic Diseases (SNOAPAD) Veterinary Parasitology, 29 (1988) 299-326 299, "the disease name is constructed solely by the suffix -osis, which is added to the stem of the name of the parasite taxon, formed from the nominative of the taxa". Besides, the reference of Audicana and Kennedy 2008 should be changed by Kassai et al 1988.

**Anisakiasis has been changed in anisakiosis and Kassai et al 1988 (line 128) has been added in the text.**

Hysterothylacium belongs to the family Anisakidae, but to the subfamily Raphidascaridinae Hartwich, 1954. Please include this in the text.

**In this work we used the classification of Fagerholm, 1991, due to the fact that molecular analyses performed to investigate phylogenetic relationships in the superfamily Ascaridoidea support this classification (Murrell, K. Darwin, and Bernard Fried. Food-borne parasitic zoonoses: fish and plant-borne parasites. Vol. 11. Springer, 2007 report two different accepted classification for ascaridoid). For further explanations see answers to the Reviewer 2. Moreover, a new paragraph has been added in the manuscript (line 82-89).**

Infestation should be changed by infection.

**Infestation has been changed in infection in all the manuscript.**

The following reference: De Nicola, P., Napolitano, L., Di Bartolomeo, N., Waku, M., & Innocenti, P. (2005). Su di un caso di Anisachiasi con perforazione del cieco. *Il Giornale di Chirurgia*, 26, 375-377, there is inadequate in order to justify that "infestations constitute a public health problem, especially in Spain".

**The sentence has been modified and the reference has been removed (line 173-176)**

**Reviewer #2:** The topic of this review is interesting. However, in my opinion there is too much generic information. Although this review does not present any new knowledge, after some revisions, it could give contribution in this field.

Detailed reviewer comments:

#### ABSTRACT

The abstract is not sufficiently informative and the objective is not clear (line 35...we analyze the data obtained etc...)

**The abstract has been changed according to the suggestion**

Line 34: operator working in....it could be replaced with food business operators in...

**Done (line 34)**

Line 37: On the basis of or Based on....

The authors should combine some sentences. (lines...37 - 42)

**The sentence has been changed (line 37-42)**

Line 44: European legislation or European Rules

**We prefer European legislation**

#### INTRODUCTION

There is a disconnect in several parts in the section.

**Introduction has been revised**

Line 55: the Authors may delete "significant" in the text.

**Significant has been deleted (line 54-55)**

Lines 106-110: The Authors should explain better these sentences.

**The classification of Hartwich, 1974 was based on the structure of the secretory-excretory system while that of Fagerholm, 1991 was based on the male caudal morphology (see table below from Murrell, K. Darwin, and Bernard Fried. *Food-borne parasitic zoonoses: fish and plant-borne parasites. Vol. 11. Springer, 2007 report two different accepted classification for ascaridoid*). These differences in the morphological characteristics used for phylogenetic**

reconstruction led to an array of contrasting interpretations and hypotheses of relationships, in turn leading to instability of ascaridoid classification, although the classification schemes of Hartwich (1974) and Fagerholm (1991) have been the most commonly used. Molecular data have been used to investigate phylogenetic relationships in the superfamily Ascaridoidea (e.g. Nadler, 1992; Nadler and Hudspeth, 1998, 2000; Zhu et al., 1998). Although these studies have not fully resolved taxonomic uncertainty within the superfamily, they tend to support the classification proposed by Fagerholm (1991). For this reason we decide to report the classification of Fagerholm, 1991, which consider the genus *Hysterothylacium* not belonging to the family Anisakidae, but to the family Raphidascarididae. A new section has been added in the text (Line 82-89).

TABLE 5.1. Contrasting classification schemes of anisakid nematodes by Hartwich (1974) and Fagerholm (1991).

Authority	Family	Subfamily	Genera
Hartwich (1974)	Anisakidae	Anisakinae	<i>Anisakis</i> , <i>Phocanema</i> (= <i>Pseudoterranova</i> ), <i>Terranova</i> , <i>Sulcascaris</i> , <i>Duplicaecum</i> , <i>Galeiceps</i> , <i>Contraecum</i> , <i>Phosascaris</i>
		Geozinae	<i>Goezia</i>
		Raphidascaridinae	<i>Raphidascaris</i> , <i>Raphidascaroides</i> , <i>Thynnascaris</i> (= <i>Hysterothylacium</i> ), <i>Lappetascaris</i> , <i>Aliascaris</i> , <i>Heterotyphlum</i> , <i>Paranisakis</i> , <i>Paranisakiopsis</i>
Fagerholm (1991)	Anisakidae	Anisakinae	<i>Anisakis</i> , <i>Pseudoterranova</i> , <i>Terranova</i> , <i>Sulcascaris</i> , <i>Peritrachelius</i> , <i>Pulchrascaris</i> , <i>Paranisakiopsis</i>
		Contraecinae	<i>Contraecum</i> , <i>Galeiceps</i> , <i>Phosascaris</i>
	Raphidascarididae	<i>Raphidascaris</i> , <i>Raphidascaroides</i> , <i>Hysterothylacium</i> , <i>Lappetascaris</i> , <i>Heterotyphlum</i> , <i>Paranisakis</i> , <i>Goezia</i> , <i>Sprentascaris</i> , <i>Paraheterotyphlum</i>	

Line 117: "the molecular studies carried out have shown" ?

**The sentence has been modified (line 82-85)**

Line 152: (Genchi and Pozio, 2003 before Abe,2008)

**Done Line (169)**

Lines: 164-170 and: these sentences are not clear, Please could you explain these sentences better?  
I think that this sentence is not pertinent for the topic of the review.

**The sentence has been modified and shortened (line 178-184)**

Lines 192 - 193: the authors should eliminate this part.

**The sentence has been modified (line 195-197)**

Line: 202....the reference?

**The reference has been refined (line 211)**

Lines: 215-220 The objectives need to be re-written

**The objectives have been re-written and moved at the beginning of the introduction (line 62-67)**

LEGISLATIVE HYSTORY

Lines: 249-254 The authors should combine this part.

**The sentence has been modified (line 242-249)**

This is not in contrast with the EFSA opinion.

**The reviewer did not report the line. Probably he/she refers to line 414 (original manuscript).  
The sentence has been modified according to the suggestion (line 393-395)**

DISCUSSION

The Authors need to rethink the Discussion.

In my opinion this section is too long and the Authors may cut some parts or combine some sentences.

**Discussion has been revised and shortened**

Lines: 450-452...this part is not clear, the Authors need to rethink the sentence.

**The sentence has been modified (line 427-428)**

There is a disconnect in several parts in the section...please could you organize it better?

In some parts, I have found the same information reported in other sections (Introduction or Legislative Hystory)

**The section has been revised and the repeated information have been deleted**

**Reviewer #3:** The paper gives an comprehensive overview on the European legislation concerning Anisakis in fish as food and describes in detail how the visual inspection was adopted in Italy. It needs some revisions before it is acceptable for publication.

In the introduction you should add a short paragraph on the ongoing debate on the necessity of larvae being alive to induce allergy the first time.

**The section has been modified and some information has been reported on this issue (line 134-154)**

line 130 - 141: the paragraph should be changed because *Anisakis* larvae can be found in most marine species caught in European waters and not only in the species you mentioned, e.g. herring can also be heavily infected.

**According to the suggestions we decided to remove the paragraph from line 137 to line 141 (original manuscript) which is not fundamental for the aim of our review. This to avoid to add an excessive number of references, due to the fact that a multitude of fish species can be infected by *Anisakis*.**

It is correct that according to the RASFF data, the species you mentioned are the most frequently parasitized species, but these species were mainly imported to Spain and Italy, other European countries simply do not check for nematodes in species landed in their harbours. So a general ranking could be different.

**The sentence has been modified highlighting that the data mainly come from Italian and Spanish notification (Line 116-123)**

line 142-148: This paragraph should be moved and included in the explanation given in line 176 - 185.

**In this section we discussed the reason why the *Anisakidosis* is an emerging parasitosis and the factors that have led to an increase of the human cases of parasitosis. Among these factors, there is the decrease of INTERMEDIATE hosts of the larvae. The paragraph previously included in lines 142-148 is about the prevalence of infected species, which is higher where cetaceans are more numerous. This is an issue correlated more to the epidemiology of the infected species than to the increase of parasitosis cases in these last years. Therefore, we think that it should be included in the paragraph 1.1.**

line 208-9: it was also my opinion that farmed *Salmo salar* presents an irrelevant risk of infestation, but Mo et al: *Journal of Fish Diseases* 37, 135-140 (2014) showed the farmed *S.salar* can be infected with *Anisakis*. So your conclusions throughout the paper concerning farmed salmon should be slightly revised according to this new findings.

**The section about farmed salmon has been modified in the light of the results of Mo *et al.*, 2014. Moreover, to avoid repetition, the section has been deleted from introduction and moved to line 450-457).**

line 490 - 495. *Anisakis* free does not mean parasite free (trematodes, cestodes). It would be nice if you could add one sentence on possible consequences the consumption of fresh raw fish may have concerning these parasites.

A new sentence has been added at line 468-469

- The presence of Anisakids larvae in fishery products is an emerging issues
- We carried out a survey on European and local laws on Anisakis risk management
- We analyzed the approach of different control systems to face the Anisakis risk
- The management of the Anisakis risk show differences in local laws
- A reassessment of the Anisakis issue is needed to yield common strategies



1        **Evolution of the Anisakis risk management in the European and Italian context.**

2

3        Priscilla D'amico<sup>1</sup>, Renato Malandra<sup>2</sup>, Francesco Costanzo<sup>1</sup>, Lorenzo Castigliero<sup>1</sup>,

4        Alessandra Guidi<sup>1</sup>, Daniela Gianfaldoni<sup>1</sup>, and Andrea Armani<sup>1\*</sup>

5

6        <sup>1</sup>FishLab, Department of Veterinary Sciences, University of Pisa, Via delle Piagge 2,

7        56124, Pisa (Italy).

8        <sup>2</sup> Wholesale fish market of Milan, ASL of Milan, Viale Molise, 20, 20137 Milan (Italy).

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

\* Author for correspondence. Tel: (+39)0502210207; Fax: (+39)0502210213

Email: [andrea.armani@unipi.it](mailto:andrea.armani@unipi.it)

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

27       **ABSTRACT**

1  
2 28       Due to the social and legislative implications, the presence of *Anisakis* spp. larvae in  
3  
4 29       fishery products has become a concern for both the consumers and the official Control  
5  
6  
7 30       Authorities. The issuance of a large number of provisions, aimed at better managing fish  
8  
9 31       products intended to be consumed raw or almost raw and the associated risks, resulted in a  
10  
11 32       very complicate legal framework. In this work, we analyzed the evolution of the normative  
12  
13 33       through an overview on **the local and international** legislation, focusing on issues that are of  
14  
15 34       practical interest for **Food Business Operators (FBOs)** in the fishery chain. **In addition, we**  
16  
17 35       **performed a survey across the Departments of Prevention of the Italian Local Health**  
18  
19 36       **Authorities (LHA) and the main fish markets in Italy to collect the operating procedures and**  
20  
21 37       **the monitoring plans. Overall, we found many differences, due to the absence of a national**  
22  
23 38       **reference standard for the management of the *Anisakis* risk. From this examination, it turns**  
24  
25 39       **clear that only a participation of all the involved Institutions, a strategy of synergistic**  
26  
27 40       **interventions, as well as a correct training of FBOs, can result in an effective risk management**  
28  
29 41       **and a proper risk communication, which should overcome states of confusion and**  
30  
31 42       **unnecessary negative impacts on the economy.**  
32  
33  
34  
35  
36  
37  
38  
39  
40

41 44       **Keywords: Anisakids, risk management, European legislation, raw fish products.**  
42  
43  
44 45  
45  
46 46  
47  
48 47  
49  
50  
51 48  
52  
53 49  
54  
55 50  
56  
57 51  
58  
59 52  
60  
61  
62  
63  
64  
65

## 1. INTRODUCTION

Parasitic infections associated to the consumption of fish products have always been a concern for the public health and for economy. Recently, the World Health Organization (WHO) has estimated a prevalence of approximately 56 million cases worldwide and a population exposed to the risk of about 400 million individuals (World Health Organization, 2013). The main food-borne zoonoses associated with the consumption of fishery products are mainly attributable to trematodes, cestodes and nematodes. Among the latter, the Anisakids are the most important parasites from a sanitary point of view, since capable to induce pathologies in humans (Chai, Darwin Murrell, & Lymbery, 2005).

In this work, we made an excursus on the European, Italian and regional legislation and analyzed the operating procedures and the monitoring plans for prevention and control of Anisakidosis, the information on which were provided by the Italian Local Health Authorities (LHA) or collected at the main fish markets. We then highlighted the peculiarities in the current normative and the differences in the risk management strategies, implemented by the Official Authorities and by the FBOs.

### *1.1. Anisakids: classification, biological cycle and epidemiology*

The identification of Anisakid species is very difficult due to the limited species-specific differences in morphological characters. Moreover, these differences are only visible in the adult worm and not in the larvae (Fagerholm, 1991; Mattiucci, Abaunza, Damiano, Garcia, Santos, & Nascetti, 2007). For this reason, in the past, only two species were considered responsible for zoonotic forms: *Anisakis simplex*, known as “herring worm”, and *Pseudoterranova decipiens*, known as “cod worm” (Smith and Wootten, 1978a; Murrell and Fried, 2007). However, the molecular studies based on genetic markers have shown that many morphospecies of *Anisakis* and *Pseudoterranova* include a certain number of sibling species with identical morphology, but different genetic make-up and geographical location (Mattiucci and Nascetti, 2008). Currently, 9 species of the *Anisakis* genus, (Mattiucci and

79 Nascetti, 2008; Mattiucci *et al.*, 2011) and 6 of the *Pseudoterranova* genus (Paggi *et al.*,  
80 1991; Mattiucci *et al.*, 1998; George-Nascimento and Urrutia, 2000; McClelland, 2002;  
81 Murrell and Fried, 2007; Mattiucci and Nascetti, 2008) have been described.

82 Even though the taxonomic uncertainty within the superfamily has not been still  
83 completely solved, the molecular analyses used to investigate phylogenetic relationships tend  
84 to support the classification proposed by Fagerholm (1991) based on the male caudal  
85 morphology (Murrell and Fried, 2007). This classification considers the genus  
86 *Hysterothylacium* not to belong to the family Anisakidae, but to the family Raphidascarididae.  
87 In spite of this, there is a tendency to use the term Anisakidosis also for infections provoked  
88 by species belonging to this genus, because likewise hazardous to public health (Audicana, *et*  
89 *al.*, 2002; Genchi and Pozio, 2003; Griglio *et al.*, 2012).

90 The Anisakids are characterized by different stages of larval development, located in  
91 different hosts. In general, the adult forms are located in the stomach of marine mammals,  
92 while the eggs are shed in the feces and require an incubation period before hatching into  
93 second stage larvae (L2). The L2 are ingested by invertebrates, especially crustaceans, such as  
94 Euphausiacea and Copepoda, where they grow inside the hemocoel (McClelland, 2002;  
95 Murrell and Fried, 2007; Smith and Snyder, 2005). These larval forms do not seem to be  
96 hazardous to the human health as it is the third stage larvae (L3), which, after being ingested  
97 by a paratenic host, can directly encyst in the edible tissue (Nagasawa, 1990; Anderson and  
98 Anderson, 2000), representing a risk when fishery products are consumed raw or  
99 undercooked. In addition, some studies have shown that L3 larvae are able to migrate from  
100 the viscera to the muscle, even after the death of the fish (Van Thiel, 1962; Smith and  
101 Wootten, 1978a; Adams, Murrell, & Cross, 1997; Abollo, Gestal, & Pascual, 2001; Murrell  
102 and Fried, 2007; Rello, Adroher, Benitez, & Valero, 2009; Pravettoni, Primavesi, &  
103 Piantanida, 2012).

104 At the global level, the *Anisakis* larvae were found in at least 200 different species of fish  
105 and 25 species of cephalopods (Abollo, Gestal, & Pascual, 2001; Klimpel, Palm, Rückert, &  
106 Piantkowsky, 2004; Murrell and Fried, 2007). In the Mediterranean Sea, the L3 larvae were  
107 observed in 31 different species out of 99 species examined belonging to 47 families (about  
108 15.000 specimens), with a prevalence ranging from 1.3 to 100% and with an infection  
109 intensity from 1 to over 300 larvae per specimen (Ponzio, 2004). Moreover, some studies  
110 have found that fishes caught in areas with abundant populations of cetaceans show a greater  
111 degree of infection. For instance, in the Ligurian Sea, which is part of the Marine Mammal  
112 Sanctuary and where the presence of whales and dolphins is stable, the prevalence of infection  
113 is 5 times greater than those found in the South Atlantic or in other Mediterranean areas  
114 (Notarbartolo-di-Sciara, Agardy, Hyrenbach, Scovazzi, & Van Klaveren, 2008; Rello *et al.*,  
115 2009).

116 In the European Union, in the period 2009-2013, the Rapid Alert System for Food and  
117 Feed (RASFF) (Rapid Alert System for Food and Feed, 2014) has reported a total of 333  
118 notifications for the presence of parasites: of these, 262 (78.5%) were due to *Anisakis* spp.  
119 These notifications were mainly forwarded by Italy (147) and Spain (49) and refer to 108  
120 border rejections, 86 information and 68 alerts. The RASFF data, mainly based on Italy and  
121 Spain notification, show that the most frequently parasitized fish, among the species  
122 commercialized in this countries, were: *Merluccius merluccius*, *Scomber scombrus*, *Engraulis*  
123 *engrasicolus*, *Lophius* spp., *Xiphias gladius* and *Lepidopus caudatus*.

## 124 **1.2 Human health issues**

125 In 1988, Kassai *et al.* recommended the use of three different terms to indicate zoonoses  
126 caused by nematodes of the Anisakidae family:

- 127 - Anisakidosis: for zoonoses from species belonging to the Anisakidae family (Anisakids);
- 128 - Anisakiosis: for zoonoses from species belonging to *Anisakis* genus;
- 129 - Pseudoterranovosis: for zoonoses from species belonging to the *Pseudoterranova* genus.

130 Humans, who are accidental hosts of the L3 larvae, may develop diseases, which presents  
131 either non-invasive forms, generally asymptomatic, or invasive forms, characterized by  
132 gastrointestinal symptoms (acute or chronic), nowadays well identified and extensively  
133 described and discussed in the literature.

134 What is less known is the “gastroallergic anisakidosis”, an IgE-mediated allergic reaction  
135 that can be developed after ingestion of Anisakids. It is characterized by clinical  
136 manifestations ranging from urticaria to anaphylactic shock (Daschner, Alonso-Gomez,  
137 Cabanas, Suarez-de-Parga, & Lopez-Serrano, 2000; Dominguez-Ortega *et al.*, 2001). Despite  
138 two episodes of allergy have been described without the involvement of living parasites  
139 (Audicana *et al.*, 1995; Aníbarro *et al.*, 1997), the ingestion of alive larvae is usually required  
140 for sensitization and allergic reactions (Daschner, Alonso-Gomez, Cabanas, Suarez-de-Parga,  
141 & Lopez-Serrano, 2000; Audicana, *et al.*, 2002; Alonso-Gomez *et al.*, 2004), while the  
142 exposure to Anisakis proteins alone may suffice to elicit allergic reactions in sensitized  
143 individuals (Nieuwenhuizen *et al.*, 2006). This possibility makes the “gastroallergic  
144 anisakidosis” a professional pathology in people who have frequent contacts with infected  
145 fish (Aníbarro *et al.*, 1997; Aníbarro and Seoane, 1998; Armentia *et al.*, 1998; Scala *et al.*,  
146 2001). Still, considering that some allergens have been shown to be heat-resistant, cooking  
147 might not provide an effective protection against allergic manifestations (Moneo, Caballero,  
148 GómezOrtega, & Alonso, 2000; Caballero and Moneo, 2004; Moneo *et al.*, 2005). For  
149 instance, it has been reported that in northern Spain the consumption of cooked hake is one of  
150 the main cause of allergies, closely followed by cooked or raw anchovies (Fernández de  
151 Corres *et al.*, 1996; Audicana, Del Pozo, Iglesias, & Ubeira 2003). On the contrary, it has also  
152 been reported that freezing could reduce the risk of allergic reactions (Garcia *et al.*, 2001).  
153 However, the exposure to Anisakid antigens is still a debated issue, due to the many clinical  
154 manifestations that are involved (Audicana and Kennedy, 2008).

155 The 90% of diseases related to the ingestion of Anisakids worldwide is determined by  
156 *Anisakis simplex*, *A. pegreffii* and *Pseudoterranova* spp. (Bouree, Paugam, & Petithory, 1995;  
157 Rosales et al., 1999; Smith, 1999; Audicana, et al., 2002; Murrell and Fried, 2007). In Italy,  
158 the species most frequently associated to health issues is *A. pegreffii* (D'Amelio,  
159 Mathiopoulosv, Brandonisio, Lucarelli, Doronz, & Paggi, 1999; Moschella et al., 2004;  
160 Fumarola et al., 2009; Mattiucci et al., 2011; Bernardi, Gustinelli, Fioravanti, Caffara,  
161 Mattiucci, & Cattaneo, 2011; Mladineo and Poljakb, 2013). *A. physeteris*. *Contracaecum* spp.  
162 and *Hysterothylacium* spp. are other species of *Anisakis* spp. that can be involved in human  
163 infections, even though they are only seldomly involved in pathological forms (Ishikura et al.,  
164 1993; Yagi et al., 1996; Rosales et al., 1999; Smith, 1999; Lima dos Santos and Howgate,  
165 2011).

### 166 ***1.3 The oxymoron of Anisakidosis: a classic but emerging parasitosis.***

167 In the last decade there have been about 20,000 cases of human Anisakidosis, of which more  
168 than 90% located in Japan, where, every year, about 2.000 infections occurs (Genchi and  
169 Pozio, 2003; Abe, 2008). In the United States, a recent survey of patients with generic  
170 gastrointestinal disorders showed that these symptoms were ascribable to parasitic diseases of  
171 fish origin, with such a frequency to require preventive controls throughout the national  
172 territory (Hochberg, Hamer, Hughes, & Wilson, 2010).

173 In Europe, the estimated incidence is about 0.038% (Orphanet, 2009 and Lima dos Santos  
174 and Howgate, 2011). Most of the cases have been registered in Spain, Netherlands and  
175 Germany, (European Food Safety Authority, 2010 and Pravettoni, Primavesi, & Piantanida,  
176 2012). The exact incidence is difficult to establish, but it seems to average 20 cases per  
177 country per year. In France, a report of 2003 estimated an incidence of eight cases per year  
178 (French Agency for Food, Environmental and Occupational Health and Safety, 2011). In  
179 Italy, 54 cases had been reported from 1996 to 2011 (Ponzio, 2004; De Rosa, 2011; Griglio et  
180 al., 2012), most of which occurred in Puglia, Abruzzo, Molise and Sicily, especially in coastal

181 areas. Only few sporadic cases have been reported in other regions as Emilia Romagna,  
182 Liguria, Lombardy, Tuscany, Marche, Lazio and Campania. All cases were associated with the  
183 consumption of traditional dishes made with anchovies, herring and mackerel served raw or  
184 "almost raw" (Ponzio, 2004; De Rosa, 2011; Griglio *et al.*, 2012).

185 The factors that have led to an increase of the Anisakidosis cases over the past 30 years are  
186 many and interdependent. First of all, the food scare crises, such as the "mad cow disease"  
187 and the "avian influenza", which have shifted the orientation of consumers' attention towards  
188 proteins of fish origin, have increased the consumption of fishery products.

189 Another factor to be taken into consideration is the spread of ethnic food on the Western  
190 tables has led to the availability of a variety of Oriental dishes, especially Japanese,  
191 characterized by preparations of raw seafood. In fact, many Japanese restaurants are not really  
192 authentic, but managed by operators of different ethnicity, especially Chinese. These latter  
193 tend to increasingly convert their restaurant activities into Sushi Bars or Sushi Restaurants,  
194 offering cheaper products, often at the expense of quality (Masotti, Amadei, & Lanni, 2010;  
195 Armani, Castigliego, Gianfaldoni, & Guidi, 2011). **However, the lack of a proper knowledge  
196 on the microbiological and parasitic risks associated to dishes based on raw fish could lead to  
197 an inappropriate manipulation and treatment of the raw materials.**

198 Not to be undervalued is also the change, in recent years, of the control responsibility in  
199 the fishery chain. In fact, while once the veterinary inspector was the person in charge for  
200 controls, nowadays, this task is delegated to the FBOs, who are not always adequately trained  
201 in detecting infections.

202 Finally, the unconditioned increase in the supply of wild fishery products has often  
203 determined adverse effects on the availability of natural fisheries, resulting in a depletion of  
204 marine resources. Therefore, the reduction of the species that have "historically" represented  
205 the intermediate hosts, together with the low specificity that these parasites have towards their  
206 intermediate hosts (LyMBERY and Cheah, 2007; Murrell and Fried, 2007), have likely



207 determined the progressive increase of both the number of affected species and the prevalence  
208 of infection.

209 Due to the increasing difficulties in the management of the "Anisakis risk", related to the  
210 involvement of wider geographic areas and to the significant number of individuals affected  
211 by anisakidosis, the European Food Safety Authority (EFSA), in its report (EFSA, Panel on  
212 Biological Hazards (BIOHAZ), 2010), highlighted that no maritime area can be considered  
213 free from Anisakids and that the presence of larvae in fishery products is a natural condition  
214 that claims the "presumption of infection" throughout the supply chain. In conclusion, the  
215 group of EFSA experts stated that to reduce Anisakis infection it is essential to provide to all  
216 the operators involved in the fishery chain and to the final consumers the necessary  
217 information for the proper management of the risks associated to the presence of parasites in  
218 fishery products.

## 2. LEGISLATIVE HISTORY

### 2.1. Before the "Hygiene Package"

221 In Europe, the first regulation related to the management of the *Anisakis* spp. associated  
222 risk was issued by the Dutch Health Authorities in the years 1968/69. This law allowed the  
223 marketing of not-eviscerated herrings only if previously subjected to treatments for  
224 devitalization of the larvae (Panebianco and Lo Schiavo, 1985).

225 In the early 70's, in Germany traders and processors decided to adopt the regulations issued  
226 by the Dutch Authorities (Horst, 2008) but, in the course of the '80s, some companies decided  
227 to no longer adhere to this voluntary agreement. This led to an increase of cases of infection  
228 and to the collapse of sales. Since then, the German government started a series of efforts in  
229 order to minimize the Anisakids risk in the domestic fish products (Horst, 2008). On August  
230 the 8<sup>th</sup> 1988, the Federal Republic of Germany issued an Ordinance (Federal Republic of  
231 Germany, 1988), which ordered the removal of parts containing nematodes (dead or alive)  
232 and their marketing ban.

233 In the same period, both the French and the Dutch Ministry of Agriculture and Fisheries  
1  
2 234 instituted specific rules for herrings processing (Renon and Malandra, 1991). In Italy, the first  
3  
4  
5 235 specific Circular Letters (CL) on *Anisakis* were enacted around the 80's, but already in 1962,  
6  
7 236 with the national Law n.283, it was forbidden to "*use in the preparation of food or drink, sell,*  
8  
9 237 *hold to sell, administer or otherwise distribute for consumption, soiled, invaded by parasites,*  
10  
11 238 *altered or harmful foodstuff*". The parasites considered by the Law n. 283 were not *Anisakids*,  
12  
13  
14 239 but all those that could be found in the products, as a consequence of a poor state of  
15  
16 240 preservation and thus destined to seizure and destruction.

19 241 The first specific legislation on the management of *Anisakis* spp. in fishery products was  
20  
21 242 enacted in 1992, when the presence of the parasite strikingly emerged in bluefish. In order to  
22  
23  
24 243 draw the attention of the industry, consumers and Health Authorities on the need to adopt  
25  
26 244 systems to prevent this zoonosis, the CL No. 10 of 11/03/1992 was issued by the Ministry of  
27  
28  
29 245 Health: *the Italian government recommended the FBOs to proceed with a rapid evisceration*  
30  
31 246 *of the fish at risk of infection (with a size exceeding 18 cm) and not to throw the entrails into*  
32  
33  
34 247 *the sea. On the other hand, the veterinary inspectors were asked to carry out a check of the*  
35  
36 248 *fish through random sampling and visual inspection, after opening the coelomic cavity*  
37  
38  
39 249 *(Circular Letter No. 10/92)*. Finally, if the invasion by the parasites of the edible parts was  
40  
41 250 such as to give a repugnant appearance to the fish, according to the judgment of the veterinary  
42  
43 251 inspector, the Letter No. 10 settled the obligation to destroy the products. In case of slight  
44  
45  
46 252 infection, the veterinary inspector could instead dispose the treatment of products to kill  
47  
48 253 parasites. Such treatment consisted in the freezing at -20 °C for at least 24 hours or,  
49  
50  
51 254 alternatively, in the heat treatment at 60 °C for 10 minutes at least. As for the visual  
52  
53 255 inspection, the aforesaid Letter did not include particular obligations for the retailers, but a  
54  
55  
56 256 verification that the product had been monitored by the Veterinary Service. These provisions  
57  
58 257 were resumed in the Ministerial Decree of 12/05/1992, which stated that preventive  
59  
60  
61  
62  
63  
64  
65

1  
2 258 treatments would have been only carried out in authorized establishments and that they had to  
3 be certified (Ministerial Decree of 12/05/1992).

4  
5 260 Subsequently, the Legislative Decree No. 531 of 30/12/1992 introduced the obligation of  
6  
7 261 the visual inspection and the research of visible parasites, before the introduction on the  
8  
9 262 market. The Decree No. 531 also ordered that the fish or their parts, obviously **infected** with  
10  
11 263 parasites, were not intended for human consumption. Fishes, to be consumed raw or almost  
12  
13 264 raw, cold-smoked (with a core fish temperature of less than 60° C), as well as marinated  
14  
15 265 and/or salted, had to be treated with low temperatures if the process did not guarantee the  
16  
17 266 death of the larvae (Decreto Legislativo No. 531/92).

18  
19  
20  
21 267 In 1993, Commission Decision No. 93/140/EEC defined the concept of "visible parasite"  
22  
23 268 as *"a parasite or a group of parasites which has a dimension, color or texture which is clearly*  
24  
25 269 *distinguishable from fish tissues"* and "visual inspection" as *"a non-destructive examination of*  
26  
27 270 *fish or fishery products without optical means of magnifying and under good light conditions*  
28  
29 271 *for human vision, including, if necessary, candling"*. The recipients of the aforesaid Decision  
30  
31 272 were required to implement controls for the detection of visible parasites in whole products  
32  
33 273 and in fish fillets or slices. They must determine the scale and frequency of the inspections,  
34  
35 274 depending on the nature of the fishery products, their geographical origin and their use. For  
36  
37 275 eviscerated fish, the search for visible parasites must be conducted by professionals on the  
38  
39 276 abdominal cavity, livers and roes, intended for human consumption. According to the type of  
40  
41 277 evisceration, the visual inspection must be performed in a continuous manner at the time of  
42  
43 278 evisceration and washing (manual evisceration) or by sampling a representative number of  
44  
45 279 specimens, being not less than 10 fish per batch (mechanical evisceration). Fish fillets and  
46  
47 280 fish slices must be inspected by qualified personnel during trimming after filleting or slicing.  
48  
49 281 Where an individual examination is not possible, due to the size of the fillets or to the filleting  
50  
51 282 operations, operators must draw up a sampling plan.  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

283 In later years, some Italian regions introduced appropriate measures into their local  
1  
2 284 legislation, in order to protect the health of the consumers. In 1994, the Region of Lombardy  
3  
4  
5 285 defined the procedures for carrying out a statistically significant sampling for the detection of  
6  
7 286 the parasite in fish products (Table 1). The protocol established that, through an initial visual  
8  
9  
10 287 inspection, the Control Authority must ensure that there were no migrating larvae on the outer  
11  
12 288 surface of the fish. Otherwise, the lot of fish should have been withdrawn and subsequently  
13  
14 289 destroyed. As for the anchovies, if no larvae were found on the surface, the Authority would  
15  
16  
17 290 have proceeded with the sampling, on a representative number of specimens, according to the  
18  
19 291 "Table of statistical significance" (Table 1). In 1997, the Liguria Region stated that in all  
20  
21 292 business activities where raw anchovies were commercialized, an informative billboard must  
22  
23  
24 293 have been displayed reporting "*.....the specimens of anchovies (Engraulis encrasicolus) used*  
25  
26 294 *in food preparation as raw, raw marinated "the flagship acciughette" or with lemon etc.,*  
27  
28  
29 295 *must be previously frozen at -20 °C for at least 24 hours"* (Decree of the President of the  
30  
31 296 Regional Council n. 282/49671). Moreover, the Liguria Region defined how the controls must  
32  
33  
34 297 be carried out on lots of anchovies, as well as the appropriate measures to be taken following  
35  
36 298 the outcome of the inspection. In particular, specific procedures were defined for the sampling  
37  
38  
39 299 of specimens (Circular Letter No. 1 of 1997).

40  
41 300 In 1996, an European Regulation took into consideration, for the first time, the relationship  
42  
43 301 between the presence of parasites and the negative effect that they have on the quality of the  
44  
45  
46 302 product, establishing the downgrade of the products despite the excellent organoleptic  
47  
48  
49 303 conditions and the favorable judgment of edibility (Council Regulation (EC) No. 2406/96).

50  
51 304 In 1999, the Norwegian Directorate of Fisheries, on the basis of the European rules,  
52  
53 305 established that a visual control of random samples must be carried out to detect and remove  
54  
55  
56 306 visible parasites. The controls must be performed after gutting and rinsing of the belly cavity  
57  
58  
59 307 and any liver/roe and during filleting and slicing, possibly using the candling. The person in  
60  
61 308 charge of the establishment would set up a plan, as part of the self-monitoring, specifying the  
62  
63  
64  
65

309 scope and frequency of the controls in relation to the type of product, geographical origin and  
1  
2 310 use. Herrings, mackerels, sprats, wild salmons caught in the Atlantic or Pacific Ocean,  
3  
4 311 marinated and/or salted herrings (when the treatment has not been sufficient to destroy  
5  
6 312 nematode larvae) intended to be eaten raw or almost raw, must undergo a freezing treatment  
7  
8  
9 313 at a core temperature of -20° C or lower for at least 24 hour. Moreover, the fish species listed  
10  
11 314 above, if not sold as fresh or frozen products, must be accompanied by a certificate from the  
12  
13 315 producer indicating the type of treatment that they had undergone. Finally, special provisions  
14  
15 316 were required for packed capelin roe, which must, on examination of a minimum of 6  
16  
17 317 samples, contain a maximum average of 1 nematode larva per 100 g in the samples, and a  
18  
19 318 maximum of 3 nematode larvae per 100 g in any single sample (Norwegian Directorate of  
20  
21 319 Fisheries, 1999).

## 26 320 *2.2. After the “Hygiene Package”*

28  
29 321 The new provisions introduced by the “Hygiene Package” led to a radical change in the  
30  
31 322 system of parasites control in fishery products. At the fish market, the figure of the official  
32  
33 323 veterinarian, who was previously responsible for carrying out the sampling and the visual  
34  
35 324 inspection of the fish, has been replaced by the FBO, who has become the first responsible for  
36  
37 325 the risk management. The FBO is required to set, implement and maintain the procedures  
38  
39 326 based on HACCP principles (Hazard Analysis Critical Control Points), through an analysis of  
40  
41 327 risks associated with food production. He should also identify operational techniques to  
42  
43 328 eliminate or reduce the presence of Anisakids in the finished product. Unfortunately, the new  
44  
45 329 regulatory approach does not define the specific methods and procedures that the FBOs have  
46  
47 330 to apply, but it only defines the objectives that they have to attain, possibly assisted by the  
48  
49 331 Competent Authorities. This transposition of tasks, if on one side has streamlined the official  
50  
51 332 inspection practices, on the other implies an adequate training of FBOs. In fact, at present, the  
52  
53 333 FBOs are called to perform a statistically significant and appropriately documented sampling,  
54  
55 334 in order to demonstrate that they have taken all possible precautions to detect parasites. Such  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 335 a sampling should be performed by qualified persons, who are required to submit their  
2 336 products to a control for the detection of visible parasites, without placing on the market those  
3  
4 337 *'obviously contaminated'*.

6  
7 338 The Regulation (EC) No. 853/2004 states that fishery products to be consumed raw, almost  
8  
9 339 raw, marinated, salted and cold smoked with an internal temperature not more than 60° should  
10  
11 340 be frozen at a temperature not higher than -20 °C in all parts of the product for at least 24  
12  
13 341 hours. Subsequently, Regulation (EC) No. 2074/2005 gave further **explanations** about the  
14  
15 342 visual inspections, clarifying the concept of candling: *“Candling means, in respect of flat fish  
16  
17 343 or fish fillets, holding up fish to a light in a darkened room to detect parasites”*.

19  
20 344 In 2006, in Spain, the Royal Decree 1420/2006 was issued, following both the EU policies  
21  
22 345 and the data provided by the Scientific Committee of the Spanish Agency for Food Safety  
23  
24 346 (AESAN) and the National Center for Epidemiology, which showed a higher incidence of the  
25  
26 347 parasite in fish and an increase in Anisakidosis prevalence in the population. With this Decree  
27  
28 348 all the operators dealing with preparation and catering activities were called to accomplish a  
29  
30 349 preventive treatment of fishery products to be consumed as raw or almost raw, cold-smoked,  
31  
32 350 pickled or salted. Afterwards, in 2007, the AESAN provided some recommendations  
33  
34 351 concerning the best criteria to determine if freezing is necessary in marinated or salted fish  
35  
36 352 products (Comité Científico de la Agencia Española de Seguridad Alimentaria y Nutrición,  
37  
38 353 2007).

39  
40 354 In May 2008, the Italian Ministry of Health, following the remarks received by the  
41  
42 355 European Commission about the activation procedure of the national alert system for the  
43  
44 356 detection of *Anisakis* spp. in fishery products coming from Third Countries, issued the CL  
45  
46 357 DGSAN No. 10776-P. This document provided that the alert system must be activated only  
47  
48 358 after detection of viable larvae in the coelomic cavity and in the flesh of fresh fish products or  
49  
50 359 in those products subjected to heat treatment insufficient to kill larvae. Moreover, if during  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 controls, products unsuitable for human consumption were found (regardless of the larvae  
2 were dead or alive), the authorities should proceed with the rejection of the goods.  
3

4  
5 362 Moreover, in 2008, with the amendments introduced by Reg. (EC) No. 1020/2008 to the  
6  
7 363 Reg. (EC) No. 853/2004, the visual inspection become mandatory also for FBOs who operate  
8  
9 364 at the retail level.  
10

11  
12 365 In Italy, following the EFSA opinion of 2010, the Ministerial CL 4379-P and 4380-P of  
13  
14 366 17/02/2011 were enacted. The Letter 4379-P-17/02/2011 stresses the obligation for FBOs to  
15  
16 367 freeze at a temperature not higher than -20 °C and for almost 24 hours, the fish (even fresh  
17  
18 368 water fishes) intended to be administered raw. The type of the treatment must be declared in  
19  
20 369 the self-monitoring plan of the food business activities and, the fishery products always be  
21  
22 370 accompanied, during their placing on the market, by a certificate of the FBO, stating the  
23  
24 371 treatment performed. Moreover, the Letter 4379-P, in order to maintain the correct  
25  
26 372 information on freezing treatment and due to the fact that reporting the term “defrosted” on  
27  
28 373 products previously frozen for health safety purposes might negatively influence the  
29  
30 374 consumer, allowed to report: *"Conforms to Regulation EC No. 853/20054, Annex III, Section  
31  
32 375 VII, Chapter 3, point D, point 3"* on products label.  
33  
34  
35  
36  
37  
38

39 376 The Letter 4380-P1 stated that, if in presence of **infected** products, the operators prove they  
40  
41 377 acted in good faith and in accordance with the law the theory of liability is not "perfectible"  
42  
43 378 against them and therefore the Article 5 of Law 283/62 is not applicable. Before this CL, due  
44  
45 379 to the numerous detections of *Anisakis* spp. in fishery products, the Regional Directorate of  
46  
47 380 Public Health of Piedmont Region issued, in 2010, the Note No. 16294. This note established  
48  
49 381 that retailers *"... have to communicate to the local competent authority the detection of viable  
50  
51 382 larvae found during self-monitoring and selling"*.  
52  
53  
54  
55

56 383 In 2011, in Europe the Reg. (EC) No. 1276/2011 was issued, which amended and  
57  
58 384 expanded the Annex III of Regulation (EC) No. 853/2004. The FBOs were called to respect  
59  
60 385 the requirements also when place on the market *"Fishery products derived from finfish and  
61  
62  
63  
64  
65*

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

386 *cephalopod molluscs*" (in the Reg. (EC) No. 853/2004 generically defined as "fish") at the  
387 retail or catering level. This clarification was necessary, considering that cephalopods, as  
388 already stressed above, may be infected by Anisakids. Moreover, according to this Reg., the  
389 freezing treatment should be aimed at killing the "viable parasite" and not the "nematode  
390 larvae", as defined by Reg. (EC) No. 853/2004. Furthermore, for parasites other than  
391 trematodes, the new provisions introduce the possibility to apply other time/temperature  
392 combinations for the freezing treatment, such as -35° C for at least 15 hours.

393 According to the new provisions, the FBOs can place on the market fresh and not treated  
394 fishery products only if they are intended for cooking or if they come from fishing area and/or  
395 "Anisakis free" farms. However, the FBOs must ensure that the fishing or aquaculture area of  
396 origin are in conformity with the requirements of Reg. (EC) No. 1276/2011, through a check  
397 of the information provided by the accompanying documents. In fact, as previously stated by  
398 the CL 4379-P, if the freezing treatment is carried out in the intermediate steps of the product  
399 chain, the FBO supplier is obliged to provide to the buyer a certificate that states the freezing  
400 method applied. In 2011, both the Piedmont and Puglia Regions, with the Note Circular No.  
401 2998 and 8499, respectively, tried to clarify some aspects of application of Reg. (EC) No.  
402 853/2004 regarding the sale and supply of fishery products. In particular, both notes pointed  
403 out the role of the FBOs and their responsibility for visual inspection, to be carried out for  
404 self-monitoring and during the sales activities, stressing the importance for the operators to  
405 record all the non-conformances detected and the corrective actions taken.

406 In 2012, at the Italian national level, the Ministerial Note DGISAN 0024111-P drew  
407 attention to the official control of parasitic infections, reaffirming that it represents an  
408 effective measure for the risk management during the marketing and administration of fishery  
409 products. Subsequently, the Italian Decree No. 158/2012, known as "Balduzzi" Decree,  
410 ordered that: *"The food business operator, who offers for sale to the final consumer fresh fish*  
411 *and cephalopods as well as products of fresh water, ..... is required to expose a clearly*



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

412 *visible billboard bearing the information specified by the Minister of Health, after*  
413 *consultation with the Minister of Agriculture and Forestry, concerning the correct conditions*  
414 *of employment".* Recently, this provision has been further specified by the Decree of July 17,  
415 2013 of the Italian Ministry of Health, which stated that, at the retail level, a billboard should  
416 be displayed in a prominent position reporting: "*In case of consumption as raw, marinated or*  
417 *not fully cooked, the product must first be frozen for at least 96 hours at - 18 ° C in a domestic*  
418 *freezer marked with three or more stars".*

419 In order to facilitate understanding, the laws related to the standards considered in this  
420 work are listed in chronological order in the Table 2.

### 421 **3. DISCUSSION**

422 The different geographical origin, species and size, method of capture, and post-capture  
423 operations are among the main factors that could influence the chances of fishery products to  
424 be contaminated by Anisakids larvae. On the other hand, good manufacturing practices and  
425 HACCP programs, applied from the primary production to the administration and/or use by  
426 the final consumer, are the two key factors to maximize the safety of fishery products,  
427 reducing the *Anisakis* spp. risk to acceptable levels. **However, in our opinion, some points**  
428 **should be pointed out to better understand the risk management of this zoonoses.**

#### 429 **3.1 Preventive pre-capture measures.**

430 **The preventive measures that FBOs can apply before the capture are very limited and are**  
431 **mainly related to farming. In fact, only sometimes there is the possibility to raise "Anisakis**  
432 **free" fish, through the implementation of management procedures, validated by the**  
433 **Competent Authority and aimed at controlling the environmental and ecological factors that**  
434 **regulate the relationship between hosts and parasites (French Food Safety Agency, 2007).**

435 **One of the factor to be taken into account is the feeding plan.** As reported by **Reg. (EC)**  
436 **No. 1276/2011, after the approval by the Competent Authority, fishery products raised**  
437 **exclusively in free area and fed with a diet without parasites can be considered exempt from**

438 *Anisakis* spp. In particular, it is essential that the farmed fish are not fed with wet fee, but only  
439 with pellet (U.S. Food and Drug Administration, 2013). The geographical location is another  
440 important factor to be considered. In this context, epidemiological studies in marine areas of  
441 interest, aimed at assessing the level of infection of the species are a prerequisite to the  
442 installation of the farm.

443 Despite Peñalver *et al.* (2010) reported the absence of Anisakids larvae in sea bass, EFSA,  
444 based on scientific data examined (Deardorff and Kent, 1989; Angot and Brasseur, 1993; and  
445 Inoue, Oshima, Hirata, & Kimura, 2000; Lunestad, 2003; Wootten, Yoon, & Bron, 2009;  
446 Lima dos Santos and Howgate, 2011), asserted that the risk of parasites can be considered  
447 negligible only for Atlantic salmon (*Salmo salar*) and that the possibility that parasites may  
448 also affect farmed fish grown in open sea must not be underestimated. In fact, it is possible  
449 that invertebrates and fishes could enter into the cages and transfer the parasite to the species  
450 farmed (Marty, 2008; Skov, Kania, Olsen, Lauridsen, & Buchmann, 2009). However, a recent  
451 work of Mo *et al.* 2014, reported the presence of *A. simplex* and *H. aduncum* both in viscera  
452 and in muscle of farmed Atlantic salmon (*Salmo salar*). However, the larvae have been only  
453 found in runts not processed for human consumption. Therefore, even though these  
454 observations should be considered in future risk analyses, currently, the risk of infection for  
455 the Atlantic salmon farmed in floating cages or onshore tanks and fed with feed that does not  
456 contain live parasites can be considered irrelevant (Marty, 2008; European Food Safety  
457 Authority, 2010; Norwegian Food Safety Authority, 2011).

458 In Spain, a research project conducted during 2011-2012 showed that national aquaculture  
459 products, such as sea bass, bream, turbot and Croaker are free of *Anisakis* spp. larvae  
460 (Asociación Empresarial de Productores de Cultivos Marinos, (APROMAR), 2012). It is not  
461 to be rejected the hypothesis that, in this Country, these species could be marketed without the  
462 need to apply the preventive freezing treatment. In Italy, only the Region of Puglia has  
463 indicated cases of exclusion from the obligation of freezing treatment, in relation to the local

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

464 culinary habits and to the frequent consumption of raw or marinated fish products in its own  
465 territory. The possibility to obtain "Anisakis free" products, would not only avoid the freezing  
466 treatment, but also allow the sale of a fresh product, characterized by a better commercial  
467 appeal.

468 However, considering that "Anisakis free" does not mean "parasite free", the presence of  
469 some other kind of parasites, such as trematodes or cestodes, cannot be excluded a priori.

### 470 **3.2 Preventive post-capture measures**

471 **3.2.1 Fish handling on board.** Since no fishing area can be considered free from zoonotic  
472 parasites (European Food Safety Authority, 2010), the only post-capture measure that can  
473 reduce, or rather, not increase the spread of the parasitic infection is related to good  
474 manufacturing practices on vessels. In particular, in order to break the cycle of the parasite, it  
475 is crucial not to dispose the entrails into the sea. With this regard, the Reg. (EC) No.  
476 1069/2009, "laying down health rules as regards animal by-products and derived products  
477 not intended for human consumption...", clarifies specific measures for the disposal of  
478 materials derived from evisceration of fish with signs of disease on board fishing vessels. It  
479 follows that the fish parasitized by *Anisakis* spp. must necessarily be regarded as Category 1  
480 material. In fact, the Article 8, paragraph a), makes clear reference to "wild animals, when  
481 suspected of being infected with diseases communicable to humans or animals".

482 **3.2.2 Inspection control.** During inspection controls, the possible infection in fishery  
483 products can be detected by non-destructive or destructive methods. The first, such as visual  
484 inspection and candling, can be used for gutted and filleted products, while, in case of whole  
485 fish, inspection of the viscera can be performed by opening the coelomic cavity. The  
486 enzymatic digestion (destructive method), despite more sensitive in identifying the parasites,  
487 is not considered an official method, as a consequence of the small number of samples that  
488 can be processed and to the long time requested (Karl and Leinemann, 1997). The visual  
489 inspection method does not suffice to detect all the larvae, but only 50% of the parasites

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

490 detectable by other techniques, such as candling or enzymatic digestion (Llarena-Reino,  
491 González, Vello, Outeiriño, & Pascual, 2012). The U.S. FDA considers candling as a  
492 subjective technique (U.S. Food and Drug Administration, 2001), whose outcome can be  
493 affected by many factors, such as: thickness of the fillet, oil content, presence of skin,  
494 pigmentation, and training level and competence of the FBOs (Smith, 1999; Stormo,  
495 Sivertsen, Heia, Nilsen, & Elvevoll, 2007; Milligan, 2008; Angelucci *et al.*, 2011). However,  
496 candling is considered an official method by the *Codex Alimentarius* (Codex Alimentarius  
497 Commission, 1995).

3.2.3 *Sampling plans*. Currently, no European or Italian legislation defines a statistically  
498 significant sampling system to be used for the collection of samples during the inspection. In  
499 this context, the sampling procedures for the visual search of the parasite often refer to the  
500 internationally accepted protocols, such as those of the *Codex Alimentarius*. Specifically, the  
501 *Codex Alimentarius* (Codex Alimentarius Commission, 1989) establishes that the presence of  
502 two or more parasites per kg of sample unit with a capsular diameter greater than 3 mm, or the  
503 detection of a parasite not encapsulated and greater than 10 mm in length, determine the non-  
504 conformity of the product. In addition, based on the requirements of the *Codex* sampling plans  
505 for prepackaged products (Codex Alimentarius Commission, 1969), Canada has set limits  
506 stating that, for batches of 1 kg or more, the presence of 2 or more parasites per kg of sample  
507 unit entails the rejection of the sample. In case of prepackaged products of less than 1 kg, the  
508 rejection is due to the detection of 1 parasite per kg calculated on the total sample (Canada  
509 Food Inspection Agency, 2013).

51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

In Italy, some Regions have set up specific procedure to look for the *Anisakis* spp.  
parasites in fishery products, especially focusing on anchovies. In 1994, the Lombardy Region  
provided a statistical method to be used for the detection of *Anisakis* larvae, which is still  
used. This protocol, developed for the veterinary inspector, but today also applicable by the  
FBOs, provides criteria for sampling of a statistically significant number of specimens to be

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
516 inspected, given a degree of sensitivity of 10% and confidence of 95%. From the total weight  
517 of the lot it is possible to estimate the number of specimens that compose it, using a  
518 conversion ratio, which differs depending on the species. Then, once the size of the lot has  
519 been determined, and given a sensitivity of 5 or 10%, it is possible to determine the number of  
520 subjects to be examined (Table 1). For the visual inspection, the head and viscera are removed  
521 from the body (edible portion) and left at room temperature for at least 10 minutes. In this  
522 way, the larvae start moving and become visible. The inspection procedure adopted by the  
523 Liguria Region is very similar, but the sampling of the specimens must be performed on the  
524 basis of the number of boxes (Circular Letter No. 1 of 1997). When the lot is composed of  
525 less than 10 boxes, 3 specimens per box (up to 28) were to be sampled, while from 10 to 150  
526 boxes 1 specimens per box (up to 28), and for more than 150 boxes, 1 specimen per box (up  
527 to 40) (Tab3). A basic prerequisite, for a statistically representative sampling control, is that  
528 the lot is uniform with regard to the specimen size and origin. Furthermore, the larger the size,  
529 the greater the risk of infection.

34  
35  
36  
37  
38  
39  
40  
530 Therefore, among the criteria for identifying the lots to be sampled it could be considered:  
531 fish species with higher *Anisakis*-related risk, geographical origin, commercial size and  
532 destination. In addition, it would be appropriate to conform the criteria of sampling by:

- 41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
533
- ensuring the uniformity of the lot;
  - considering the number of crates, carrying out sampling evenly distributed throughout the lot, taking samples at several points;
  - collecting a statistically significant number of specimens;
  - examining the viscera removed from the cavity and the cavity itself after 30/40 minutes after their evisceration and exposure.

56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
539 The experience gained in decades of sampling shows that the degree of infection within a  
540 lot is constant and this allows to state that the visual examination of the viscera has a value  
541 statistically reliable for the purposes of a judgment. Therefore, it is possible to demonstrate a

1  
2 542 directly proportional correlation between the number of specimens parasitized and the number  
3  
4 543 of larvae detectable: in general, to a high number of parasitized subjects corresponds a high  
5  
6 544 number of larvae per subject (and vice versa). Therefore, it is essential to identify the  
7  
8 545 percentage of parasitized subjects that can be tolerated, or rather the percentage that is not  
9  
10 546 perceptible to the observation of the consumer. For instance, on the basis of a confidence level  
11  
12 547 of 95% associated with the sampling method, there is the possibility that in 5% of cases the  
13  
14 548 10% of **infected** subjects (occult infection) may not be detected, a percentage of additional  
15  
16 549 tolerance of 10% could be proposed. This means that, during the inspection, it could be  
17  
18 550 tolerated the presence of up to a maximum of 3 parasitized subjects, up to a maximum of 3  
19  
20 551 larvae each, when the sampling presupposes the collection of the 29 specimens (lot > 600  
21  
22 552 specimens). For batches in which the percentage of parasitized individuals exceeds the level  
23  
24 553 of tolerability, the FBOs should choose the destination depending on the species, size and  
25  
26 554 final use.

30  
31 555 In the other Italian regions, the provisions laid down are more or less those **adopted by the**  
32  
33 556 **Regions of Lombardy and Liguria**. Regarding Sardinia, the sampling is carried out in a  
34  
35 557 different manner depending on the size of the species. For instance, for specimens weighing  
36  
37 558 less than 20 grams, the procedure applied is the same as that applied in the Liguria Region for  
38  
39 559 anchovies. For species weighing more than 20 grams, new criteria have been established  
40  
41 560 (Table 3). Even in the Region of Marche, the Operating Unit (OU) for Fishing Products of the  
42  
43 561 **LHA** has developed a procedure similar to that provided by the **Liguria Region** (Tab3). The  
44  
45 562 procedure laid down by the Veneto Region provides that research for parasites must be  
46  
47 563 carried out only on fish at risk if they are intended for cold processing and if not previously  
48  
49 564 frozen (Tab 3). An interesting aspect to highlight is that, in case of non-compliance, the  
50  
51 565 procedure requires that at least the three successive batches from the same supplier must be  
52  
53 566 checked.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
567 The procedures that must be put in place after the visual inspection are different in the two  
568 Regional laws. In this regard, it may be interesting to consider the weight of a hypothetical  
569 sample of anchovies. Based on the data provided by the CL of the Liguria Region, we can  
570 infer that a sample of 28-30 anchovies may have a weight ranging between 200 and 300  
571 grams depending on the season. Therefore, the approach of the Lombardy Region seems to be  
572 more in line with the provisions of the *Codex* than with those of the Liguria Region, even  
573 though no distinction is made between encysted and free parasites. In fact, for a sample with a  
574 weight less than 1 kg, the entire batch should be rejected following the discovery of a single  
575 parasite. On the other hand, this procedure could be considered overly restrictive, whereas, at  
576 present, only the batches obviously infected should be subjected to restrictions on marketing.  
577 For this reason, the Veterinary Services of the Lombardy Region, and especially those present  
578 at the wholesale fish market of Milan, have adopted an approach that, while bestowing greater  
579 tolerance, allows to identify, at the same time, the lots manifestly contaminated.

31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
580 As for the other Regions, Sardinia provides that batches invaded by clearly visible  
581 parasites should be seized and destroyed. In this regard, the Region of Marche states that the  
582 destruction of the consignment is to be carried out only in case of massive infection and  
583 repulsive appearance of the products, applying the same criteria adopted in Liguria (Table 3).  
584 Unfortunately, the expression “massive infection” is not easily correlatable to “obviously  
585 contaminated”, adopted by the European Commission to indicate a product not suitable for  
586 human consumption.

48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
587 *3.2.4 Obviously contaminated.* The wording "obviously contaminated" introduced by  
588 Council Reg. (EC) No. 853/2004 has been the topic of several discussions in Europe. In fact,  
589 the European Commission has noted that the term "obviously" can be interpreted in different  
590 ways within each Member State. This led the Commission to produce the draft document  
591 SANCO/10137/2013-rev1. In this document, it is reported that a distinction should be made  
592 based on the location of the visible parasite. If during the visual inspection larvae are found in

593 the edible part, such as the muscles, the product is considered to be obviously contaminated.

594 If the visible parasites are found only in the non-edible parts, such as viscera, the raw material  
595 is not to be considered "obviously contaminated" and can be placed on the market.

596 Another concept closely related to the definition of "obviously contaminated" is those of  
597 "visible parasite". On the basis of the definition given by the Decision 93/140/CE and by the  
598 *Codex Alimentarius* (Codex Alimentarius Commission, 1971), the larvae must have a size that  
599 make them detectable and must be clearly differentiable from the tissues of the fish, even in  
600 the absence of optical instruments. In this regard, it must be underlined that the larvae of  
601 *Hysterothylacium* spp. (0.7 mm -2 cm) (Køie, 1993; Borges, Cunha, Santos, Monteiro-Neto,  
602 & Santos 2012) can often co-infect fish together with the larvae of *Anisakis* spp. (10-30 mm  
603 in length and 0.44-0.54 mm in diameter) (Sakanari and McKerrow, 1989; Genchi and Pozio,  
604 2003), from which are not easily distinguishable macroscopically. Thus, even though  
605 *Hysterothylacium* spp. are only rarely responsible for diseases, they fall within the concept of  
606 "visible parasite" and for this reason contribute to determine the total infection degree. On the  
607 basis of these findings, when talking about nematode larvae, it would be more appropriate to  
608 refer to visible larvae, including *Anisakis* spp. and *Hysterothylacium* spp., rather than about  
609 Anisakids larvae.

610 3.2.5 Processing that guarantee the inactivation of the parasites. At present, the freezing  
611 treatment can be also done at the retail level, such as catering establishments, restaurants and  
612 similar administration activities that have been registered under the current legislation.

613 In Italy, if the FBOs want to directly perform the preventive treatment, they have to  
614 provide the necessary guarantees of compliance and effectiveness (Ministerial Circular Letter  
615 4379-P-17/02/2011). First, the FBOs, after setting up a written procedure aimed at controlling  
616 the presence of parasites based on the principles of the HACCP system, must give a prior  
617 communication to the Competent Authority concerning the preventive treatment to be  
618 applied. Then, they are required to be equipped with an appropriate blast chiller, certified for



1 619 the reduction of the temperature at -20° C, which must be exclusively used to perform the  
2 620 freezing treatment and not to preserve fish or other product. About the freezing method, the  
3  
4 621 U.S. FDA stated that the effectiveness of the treatment is related to several factors, the most  
5  
6  
7 622 important of which are the type of parasite involved, the temperature and the duration of the  
8  
9  
10 623 treatment (U.S. Food and Drug Administration, 2013). Also, the time required to reach the  
11  
12 624 intended core temperature and the percentage of fat of the fish can affect the preventive  
13  
14 625 treatment. With regard to the type of parasite involved, it should be underlined that some  
15  
16  
17 626 cestodes, for instance, are more sensitive to freezing than nematodes, which, however, are less  
18  
19 627 resistant than trematodes. According to the U.S. FDA ([U.S. Food and Drug Administration,](#)  
20  
21 [2013](#)), in order to kill the nematode parasite, the product may be subjected to various types of  
22 628  
23  
24 629 preventive treatments, which provide different time/temperatures combination, such as:

- 25  
26 630 - Freezing at -20° C followed by a storage period of 7 days at - 20° C (or lower);
- 27  
28  
29 631 - Freezing at -35° C (or lower) associated with a period of storage at -35° C (or lower) for  
30  
31 632 15 hours;
- 32  
33  
34 633 - Freezing at -35° C (or lower) and at -20° C (or lower) for 24 hours.

35  
36 634 The freezing, as preventive treatment, is a procedure expressly required by law and,  
37  
38  
39 635 according the provisions about the correct information to consumer, the data related to the  
40  
41 636 process have to accompany the product up to the retail sale. However, this information is  
42  
43 637 usually omitted, especially in catering and food service. This is because the defrosted fish  
44  
45  
46 638 creates mistrust among consumers, who consider it of second-rate compared to fresh fish,  
47  
48  
49 639 showing a reluctance to purchase and consume it. In this regard, the term "defrosted" should  
50  
51 640 not appear on fishery and aquaculture products subjected to a preventive treatment for food  
52  
53 641 safety and health purposes ([Regulation \(EC\) No. 404/2011](#)). Beyond this, it would be  
54  
55  
56 642 desirable that, at the retail level, the [FBOs](#) exposed an indication to clearly differentiate the  
57  
58 643 product defrosted from the remaining fish retained. At the same time, it would be also  
59  
60  
61  
62  
63  
64  
65

1  
2 644 desirable that not “Anisakis free” fish were distinguished by an indication, informing the  
3 645 consumer that the product must be “consumed after cooking or after freezing”.

4  
5 646 While the European legislation does not deal with the problem of consumer’s information,  
6  
7 647 in Italy, this issue has been considered many times (Decree of the President of the Regional  
8  
9 648 Council No. 282/49671, Ministerial Circular Letter 4379-P-17/02/2011, "Balduzzi" Decree  
10  
11 649 and Decree of July 17, 2013).

12  
13  
14 650 Some EU countries, such as Spain and France, have specified the technical parameters of  
15  
16 651 salting and pickling to kill the larvae of the parasite, thus excluding the preventive freezing of  
17  
18 652 the products. In Spain, the Scientific Committee of AESAN, following the entry into force of  
19  
20 653 the Royal Decree No. 1420/2006, asserted that freezing is not necessary for those fishery  
21  
22 654 products that reach a concentration of NaCl above 9% for at least six weeks (Comité  
23  
24 655 Científico de la Agencia Española de Seguridad Alimentaria y Nutrición, 2007), between 10  
25  
26 656 and 20% for four/five weeks or more than 20% for at least three weeks. With regard to the  
27  
28 657 salting, the French Food Safety Agency (Afssa), with the opinion No. 2007-SA-0379,  
29  
30 658 reported that in traditional preparations and for small quantities salinity levels of 20% result in  
31  
32 659 the inactivation of the parasite within 21 days, while concentrations of 15% require 28 days  
33  
34 660 (French Food Safety Agency, 2007). Afssa also reported that, according to some authors, fish  
35  
36 661 marinated with 10% acetic acid and 12% salt, maintained for 5 days at 4° C, are not  
37  
38 662 hazardous to health as well as marinated products with 12% salt and 6% of acetic acid for 13  
39  
40 663 days at 4° C. Finally, Afssa affirmed that the freezing treatment can also be replaced by either  
41  
42 664 irradiation or treatments with high pressures. Unfortunately, the irradiation procedure  
43  
44 665 requested to kill the parasite seems to induce changes in the organoleptic characteristic  
45  
46 666 (Farkas, 1998). In addition, in the EU the use of ionizing radiation for fishery products is not  
47  
48 667 approved by most of the Member States, including Italy (Directive 1999/3/EC and Notice No  
49  
50 668 2009/C 283/02). The organoleptic quality of the products is also compromised by too high  
51  
52 669 pressures (French Food Safety Agency, 2007 and U.S. Food and Drug Administration, 2013).  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

670 Twenty-five years ago, this kind of alternative approaches were already implemented by  
671 Germany (Ordinance of Federal Republic of Germany, 1988). In Italy an alternative technique  
672 was never defined and from the very beginning it was established the unconditional  
673 application of preventive treatment to all products to be consumed raw or almost raw.

674 *3.2.7 Training of personnel.* Considering the complexity related to the management of  
675 zoonotic parasites, the training of FBOs is fundamental for a thorough understanding of the  
676 issues associated to *Anisakis* spp.. However, this does not imply that the figure in charge for  
677 carrying out checking of goods ought to be the FBO, since the *Anisakis* risk needs scientific  
678 knowledge to be best managed. Therefore, it is essential that the FBOs are supported by  
679 specific professionals, such as veterinarians, who can assist them during controls. The "know  
680 how" of the veterinarian is also crucial to implement all the necessary preventive and  
681 corrective measures and to define the inspective outcomes. In this regard, the Local  
682 Veterinary Departments could consider the possibility that the responsible of large-scale retail  
683 or wholesale fish markets, where large quantities of fishery products are handled, could be  
684 supported in the management and control of parasites by a veterinarian, as the most suitable  
685 competent figure.

#### 686 **4. CONCLUSIONS**

687 The *Codex Alimentarius* defines food hygiene as "*all conditions and measures necessary*  
688 *to ensure the safety and suitability of food at all stages of the food chain*". In this context,  
689 food safety, defined as "*the assurance that food will not cause harm to the consumer when it*  
690 *is prepared and/or eaten according to its intended use*", represents the priority aspect.  
691 However, food has to be suitable for human consumption, in accordance to the purpose for  
692 which it is intended, beyond its safety. The presence of Anisakids larvae in fishery products is  
693 not only a public health issue, but it also determines a deterioration of the quality product  
694 making it repugnant to the consumer. Therefore, on the one hand it is necessary to avoid that  
695 products containing viable larvae can arrive to the consumer, on the other, it is also

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

696 appropriate to implement corrective measures aimed at reducing the perception of the  
697 problem by the consumers as recently occurred (Anonymus, 2013 and La Pira, 2013). In fact,  
698 even though the larvae had lost their pathogenic power, the products were found not to be  
699 suitable for sale, since repugnant. For this reason, and also in the view of current economic  
700 conditions and limited fish stocks, it would be desirable that alternatives were considered by  
701 the legislation in order to improve the management of the products affected by visible  
702 parasites. For example, the medium/large fish could be eviscerated in an equipped  
703 establishment, while those of small size could be exclusively marketed without viscera and  
704 the retailers could carry out a deferred evisceration at the time of retail.

696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720

Among the emerging critical issues there is also the need to implement harmonized policies during the inspections and controls: with regard to the sampling, the international laws do neither explain in detail how the visual inspection should be carried out nor specify a reference value to identify the "obviously contaminated" products. This uncertainty is largely due to the different perception/acceptance of the risk associated to Anisakidosis that characterizes each Countries, which is also reflected at the level of political bodies. Therefore, in Europe, there is an autonomy on measures to be taken, which are left to the Competent Authorities of each Member States. In Italy, beyond the inevitable local differences, the Competent Authorities have so far adopted a balanced approach between what is expected in terms of prevention by the European and Italian legislation and the need to protect, besides the public health, the production activities related to fishing and their economy. However, Anisakids do no longer involve a risk only for the lesions induced by the viable larvae, but also for the allergic reactions in sensitive subjects, which should be taken into consideration by establishing preventive measures, such as the reporting of Anisakids as allergenic in fishery products, as well as their inclusion in the list of all common allergens by the specific Community regulation (Directive 2003/89/EC).

1 721 Finally, more than 20 years after the CL No. 10 of the Italian Ministry of Health, in which  
2 722 it was stated the importance of the veterinary inspector in the evaluation of an infected  
3  
4 723 product, in a complex framework, such as that related to the *Anisakis* risk management, this  
5  
6  
7 724 professional figure is still essential, especially when a discriminating judgment is required,  
8  
9  
10 725 based on scientific knowledge and professional skill.

11 726

12  
13  
14 727

15  
16  
17 728 **ACKNOWLEDGEMENTS**

18  
19 729 We thank Dr. Patrizia Buratti of ASL 14 Chioggia (Chioggia, Italy), Dr. Pietro Ceddia of  
20  
21 730 ASL Roma H (ROMA, Italy), Dr. Francesco Giacomino of ASL TO3 (Pinerolo, Italy), Dr.  
22  
23  
24 731 Giuliana Giulini of Mare.A (Cattolica, Italy), Dr. Antonio Piccioni of ASL Teramo (Teramo,  
25  
26 732 Italy) and Dr. Salvatore Madeddu of ASL 8 Cagliari (Cagliari, Italy) for providing us useful  
27  
28  
29 733 information and documents.

30  
31 734

32  
33  
34 735

35  
36 736

37  
38  
39 737

40  
41 738

42  
43  
44 739

45  
46 740

47  
48 741

49  
50  
51 742

52  
53 743

54  
55  
56 744

57  
58 745

59  
60 746

61  
62  
63  
64  
65

747 **REFERENCES**

- 1  
2 748
- 3  
4 749 1. Abe, N. (2008). Application of the PCR sequence-specific primers for the discrimination  
5 750 among larval *Anisakis simplex* complex. *Parasitology Research*, 102, 1073-1075.
- 6  
7 751 2. Abollo, E., Gestal, C., & Pascual, S. (2001). *Anisakis* infestation in marine fish and  
8  
9 752 cephalopods from Galician waters: an updated perspective. *Parasitology Research*, 87,  
10  
11 753 492-499.
- 12  
13 754 3. Adams, A.M., Murrell, K.D., & Cross, J.H. (1997). Parasites of fish and risks to public  
14 755 health. *Scientific and Technical Review*, 16, 652-660.
- 16 756 4. Alonso-Gomez, A., Moreno-Ancillo, A., Lopez-Serrano, M.C., Suarez-de-Parga, J.M.,  
17  
18 757 Daschner, A., Caballero MT, Barranco P., & Cabañas R. (2004). *Anisakis simplex*  
19  
20 758 only provokes allergic symptoms when the worm parasitises the gastrointestinal tract.  
21  
22 759 *Parasitology Research*, 93, 378-384.
- 23 760 5. Anderson, R. C., & Anderson, R. C. (2000). Nematode parasites of vertebrates: their  
24  
25 761 development and transmission. (2nd ed.). Oxon: CABI.
- 27 762 6. Angelucci, G., Meloni, M., Merella, P., Sardu, F., Madeddu, S., Marrosu, R., Petza, F.,  
28  
29 763 & Salati, F. (2011). Prevalence of *Anisakis* spp. and *Hysterothylacium* spp. larvae in  
30  
31 764 teleosts and cephalopods sampled from waters of Sardinia. *Journal of Food*  
32  
33 765 *Protection*, 74, 1769-1775.
- 34 766 7. Angot, V., & Brasseur., P. (1993). European farmed Atlantic salmon (*Salmo salar* L.)  
35  
36 767 are safe from anisakid larvae. *Aquaculture*, 118, 339-344.
- 38 768 8. Anibarro, P.C., Carmona, J.B., Gonzalez, F.G., Durantez, M.M., Gil, L.A., Sotillos,  
39  
40 769 M.G., Gimenez, R.P, Picon, S.J., & Ortega. M.C. (1997). Protein contact dermatitis  
41  
42 770 caused by *Anisakis simplex*. *Contact Dermatitis*, 37, 245-247.
- 43 771 9. Anibarro, B., & Seoane, F.J. (1998). Occupational conjunctivitis caused by sensitization  
44  
45 772 to *Anisakis simplex*. *Journal of Allergy and Clinical Immunology*, 102, 331-332.
- 47 773 10. Anonymus (2013). *Anisakis* nei bastoncini di pesce Findus. Cittadini Europei.  
48  
49 774 Available at: [http://www.cittadinieuropei.it/2013/11/08/anisakis-nei-bastoncini-di-](http://www.cittadinieuropei.it/2013/11/08/anisakis-nei-bastoncini-di-pesce-findus/)  
50  
51 775 [pesce-findus/](http://www.cittadinieuropei.it/2013/11/08/anisakis-nei-bastoncini-di-pesce-findus/). Accessed 16 April 2014.
- 52 776 11. Armani, A., Castigliengo, L., Gianfaldoni, D., & Guidi A. (2011). L'insicurezza  
53  
54 777 alimentare della nuova ristorazione cinese. *Industrie Alimentari*, 50, 7-11.
- 56 778 12. Armentia, A., Lombardero M., Callejo A., Martin Santos J.M., Gil F.J., Vega J.,  
57  
58 779 Arranz M.L., & Martinez C. (1998). Occupational asthma by *Anisakis simplex*.  
59  
60 780 Risultati di ricerca. *Journal of Allergy and Clinical Immunology*, 102, 831-834.
- 61  
62  
63  
64  
65

- 781 13. Asociación Empresarial de Productores de Cultivos Marinos (2012). Evaluación de la  
1 782 presencia de nematodos del género *Anisakis* en los pescados de acuicultura marina  
2 783 españoles. Informe Final. Available at: [http://www.apromar.es/Proyecto-](http://www.apromar.es/Proyecto-Anisakis/APROMAR-Informe-ANISAKIS-2012.pdf)  
3 784 [Anisakis/APROMAR-Informe-ANISAKIS-2012.pdf](http://www.apromar.es/Proyecto-Anisakis/APROMAR-Informe-ANISAKIS-2012.pdf). Accessed 16 April 2014.  
4 785  
5 786 14. Audicana, M.T., Fernández de Corres, L., Muñoz, D., Fernández, E., Navarro, J.A., &  
6 787 Del Pozo M.D. (1995). Recurrent anaphylaxis caused by *Anisakis simplex* parasitizing  
7 788 fish. *Journal of Allergy and Clinical Immunology*, 96, 558-560.  
8 789  
9 790 15. Audicana, M.T., Ansotegui, I.J., Corres, L.F., & Kennedy M.W. (2002). *Anisakis*  
10 791 *simplex*: dangerous - dead and alive?. *Trends Parasitology*, 18, 20-25.  
11 792  
12 793 16. Audicana, M.T., Del Pozo, M.D., Iglesias, R., & Ubeira, F. M. (2003). *Anisakis*  
13 794 *simplex* and *Pseudoterranova decipiens*. In R. Learmonth, & M. D. Milliotis (Eds),  
14 795 *International handbook of foodborne pathogens* (pp. 613-636). New York: Marcel  
15 796 Dekker.  
16 797  
17 798 17. Audicana, M.T., & Kennedy, M.W. (2008). *Anisakis simplex*: from obscure infectious  
18 799 worm to inducer of immune hypersensitivity. *Clinical Microbiology Reviews*, 21, 360-  
19 800 379.  
20 801  
21 802 18. Bernardi, C., Gustinelli, A., Fioravanti, M.L., Caffara, M., Mattiucci, S., & Cattaneo,  
22 803 P. (2011). Prevalence and mean intensity of *Anisakis simplex* (sensu stricto) in  
23 804 European sea bass (*Dicentrarchus labrax*) from Northeast Atlantic Ocean.  
24 805 *International Journal of Food Microbiology*, 148, 55-59.  
25 806  
26 807 19. Borges, J.N., Cuhna, L.F.G., Santos, H.L.C., Monteiro-Neto, C., & Santos, C.P.  
27 808 (2012). Morphological and molecular diagnosis of Anisakid nematode larvae from  
28 809 cutlassfish (*Trichiurus lepturus*) off the coast of Rio de Janeiro, Brazil. *PLoS one*, 7,  
29 810 e40447.  
30 811  
31 812 20. Bouree, P., Paugam, A., & Petithory, J.C. (1995). Anisakidosis: report of 25 cases and  
32 813 review of the literature. *Comparative Immunology, Microbiology & Infectious*  
33 814 *Diseases*, 18, 75-84.  
34 815  
35 816 21. Caballero, M.L., & Moneo I. (2004). Several allergens from *Anisakis simplex* are  
36 817 highly resistant to heat and pepsin treatments. *Parasitology Research*, 93, 248-251.  
37 818  
38 819 22. Canada Food Inspection Agency (2013). Fish products. Standards and methods manual  
39 820 Chapter 3, Standard 3. Available at: [http://www.inspection.gc.ca/food/fish-and-](http://www.inspection.gc.ca/food/fish-and-seafood/manuals/standards-and-methods/eng/1348608971859/1348609209602)  
40 821 [seafood/manuals/standards-and-methods/eng/1348608971859/1348609209602](http://www.inspection.gc.ca/food/fish-and-seafood/manuals/standards-and-methods/eng/1348608971859/1348609209602).  
41 822  
42 823 Accessed 16 April 2014.  
43 824  
44 825  
45 826  
46 827  
47 828  
48 829  
49 830  
50 831  
51 832  
52 833  
53 834  
54 835  
55 836  
56 837  
57 838  
58 839  
59 840  
60 841  
61 842  
62 843  
63 844  
64 845  
65 846

- 814 23.Chai, J.Y., Darwin Murrell, K., & Lymbery, A.J. (2005). Fish-borne parasitic  
1 zoonoses: Status and issues. *International Journal for Parasitology*, 35, 1233-1254.  
2 815
- 3 816 24.Circolare No. 1 del 1997 del Consiglio Regionale della Liguria.  
4
- 5 817 25.Circolare Ministeriale No. 10 del 11 Marzo 1992, direttive e raccomandazioni in  
6 merito alla presenza di larve di Anisakis nel pesce. *Gazzetta Ufficiale*, SG62 del 14  
7 818 marzo 1992.  
8 819
- 10 820 26.Circolare Ministeriale DGSAN 10776-P-19/09/08 Riscontro di larve di anisakis in  
11 prodotti della pesca.  
12 821
- 14 822 27.Circolare Ministeriale DGSAN No. 4379-P del 17 Febbraio 2011 chiarimenti  
15 concernenti alcuni aspetti applicativi del Reg. CE No. 853/2004 in materia di vendita e  
16 823 somministrazione di preparazioni gastronomiche contenenti prodotti della pesca  
17 destinati ad essere consumati crudi o praticamente crudi.  
18 824
- 20 825 28.Circolare Ministeriale DGSAN No. 4380-P del 17 Febbraio 2011. Articolo 5 della  
21 legge 283 sulla presenza di Anisakis.  
22 826
- 25 828 29.Circolare Regionale Puglia No. 2998 del 1 marzo 2011 chiarimenti concernenti alcuni  
26 aspetti applicativi del regolamento CE No. 853/2004 in materia di vendita e  
27 829 somministrazione di preparazioni gastronomiche e somministrazione di preparazioni  
28 contenenti prodotti della pesca destinati ad essere consumati crudi o praticamente  
29 830 crudi - Indicazioni Regionali.  
30 831
- 32 832 30.Consiglio Regionale della Lombardia, prot. No. 790/94. Metodi di ispezione igienico  
33 sanitaria applicata a partite di pesce per l'evidenziazione di larve del genere *Anisakis*.  
34 833
- 36 834 31.Codex Alimentarius Commission (1969). Codex Alimentarius Sampling Plans for  
37 Prepackaged Foods. CODEX STAN 233-1969. Available at:  
38 835 [http://down.40777.cn/stardard/10/CODEX%20STAN%20233-  
39 836 1969%20CODEX%20SAMPLING%20PLANS%20FOR%20PREPACKAGED%20FOODS%20%28AQL%206.5%29.pdf](http://down.40777.cn/stardard/10/CODEX%20STAN%20233-1969%20CODEX%20SAMPLING%20PLANS%20FOR%20PREPACKAGED%20FOODS%20%28AQL%206.5%29.pdf). Accessed 14 April 2014.  
40 837  
41 838  
42 839
- 47 840 32.Codex Alimentarius Commission (1971). Report of the eighth session of the joint  
48 FAO/WHO Codex Alimentarius Commision: recommended international standard for  
49 841 quick frozen filet of cod and haddock). CAC/RS-50-1971. Available at:  
50 842 <http://www.fao.org/docrep/meeting/005/c0531e/C0531E09.htm>. Accessed 16 April  
51 843 2014.  
52 844
- 56 845 33.Codex Alimentarius Commission (1989). Codex standard for quick frozen blocks of  
57 fish fillet, minced fish flesh and mixtures of fillets and minced fish flesh. CODEX  
58 846 STAN 165-1989. Available at:  
59  
60 847



- 848 [www.codexalimentarius.org/input/download/standards/111/CXS\\_165s.pdf](http://www.codexalimentarius.org/input/download/standards/111/CXS_165s.pdf). Accessed  
1 849 16 April 2014.
- 3 850 34.Codex Alimentarius Commission (1995). Codex General Standard For Quick Frozen  
4 851 Fish Fillets. CODEX STAN 190 – 1995. Available at:  
5 852 [www.codexalimentarius.org/input/download/standards/115/CXS\\_190e.pdf](http://www.codexalimentarius.org/input/download/standards/115/CXS_190e.pdf). Accessed  
6 853 [16 April 2014](http://www.codexalimentarius.org/input/download/standards/115/CXS_190e.pdf).
- 10 854 35.Comité Científico de la Agencia Española de Seguridad Alimentaria y Nutrición  
11 855 (2007). Informe del Comité Científico de la Agencia Española de Seguridad  
12 856 Alimentaria y Nutrición sobre medidas para reducir el riesgo asociado a la presencia  
13 857 de Anisakis. Revista del Comité Científico de la AESAN 6:59-65. Available at:  
14 858 [http://aesan.msssi.gob.es/AESAN/docs/docs/publicaciones\\_estudios/revistas/comite\\_c](http://aesan.msssi.gob.es/AESAN/docs/docs/publicaciones_estudios/revistas/comite_cientifico_6.pdf)  
15 859 [ientifico\\_6.pdf](http://aesan.msssi.gob.es/AESAN/docs/docs/publicaciones_estudios/revistas/comite_cientifico_6.pdf). Accessed 16 April 2014.
- 21 860 36.Commission Decision 93/140/EEC of 19 January 1993 laying down the detailed rules  
22 861 relating to the visual inspection for the purpose of detecting parasites in fishery  
23 862 products. *Official Journal of European Union*, L56, 50-52.
- 27 863 37.Commission Implementing Regulation (EU) No. 404/2011 laying down detailed rules  
28 864 for the implementation of Council Regulation (EC) No. 1224/2009 establishing a  
29 865 Community control system, for ensuring compliance with the rules of the Common  
30 866 Fisheries Policy. *Official Journal of European Union*, L112, 1-153.
- 34 867 38.Commission Regulation (EC) No. 2074/2005 of 05 December 2004 laying down  
35 868 implementing measures for certain products under Regulation (EC) No. 853/2004 of  
36 869 the European Parliament and of the Council and for the organisation of official  
37 870 controls under Regulation (EC) No. 854/2004 of the European Parliament and of the  
38 871 Council and Regulation (EC) No. 882/2004 of the European Parliament and of the  
39 872 Council, derogating from Regulation (EC) No. 852/2004 of the European Parliament  
40 873 and of the Council and amending Regulations (EC) No. 853/2004 and (EC) No.  
41 874 854/2004. *Official Journal of European Union*, L338, 27-59.
- 48 875 39.Commission Regulation (EC) No. 1020/2008 of 17 October 2008 amending Annexes  
49 876 II and III to Regulation (EC) No. 853/2004 of the European Parliament and of the  
50 877 Council laying down specific hygiene rules for food of animal origin and Regulation  
51 878 (EC) No. 2076/2005 as regards identification marking, raw milk and dairy products,  
52 879 eggs and egg products and certain fishery products. *Official Journal of European*  
53 880 *Union*, L277, 8-14.
- 54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

- 881 40. Commission Regulation (EC) No. 1069/2009 of the European Parliament and of the  
1 882 Council of 21 October 2009 laying down health rules as regards animal by-products  
2 883 and derived products not intended for human consumption and repealing Regulation  
3 884 (EC) No. 1774/2002 (Animal by-products Regulation). *Official Journal of European*  
4 885 *Union*, L300, 1-33.  
5  
6  
7  
8  
9 886 41. Commission Regulation (EU) No. 1276/2011 of 8 December 2011 amending Annex III  
10 887 to Regulation (EC) No. 853/2004 of the European Parliament and of the Council as  
11 888 regards the treatment to kill viable parasites in fishery products for human  
12 889 consumption. *Official Journal of European Union*, L327, 39-41.  
13  
14  
15  
16 890 42. Council Regulation (EC) No. 2406/96 of 26 November 1996) laying down common  
17 891 marketing standards for certain fishery products. *Official Journal*, L334.  
18  
19  
20 892 43. D'Amelio, S., Mathiopoulovs, K.D., Brandonisiov, O., Lucarelliv, G., Doronzov, F., &  
21 893 Paggi, L. (1999). Diagnosis of a case of gastric Anisakidosis by PCR-based restriction  
22 894 fragment length polymorphism analysis. *Parassitologia*, 41, 591-593.  
23  
24  
25 895 44. Daschner, A., Alonso-Gomez, A., Cabanas, R., Suarez-de-Parga, J.M., & Lopez-  
26 896 Serrano, M.C. (2000). Gastroallergic anisakiasis: borderline between food allergy and  
27 897 parasitic disease: clinical and allergologic evaluation of 20 patients with confirmed  
28 898 acute parasitism by *Anisakis simplex*. *Journal of Allergy and Clinical Immunology*,  
29 899 105, 176-181  
30  
31  
32  
33  
34 900 45. Deardorff, T.L., & Kent, M.L. (1989). Prevalence of larval *Anisakis simplex* in pen-  
35 901 reared and wild-caught salmon (Salmonidae) from Puget Sound, Washington. *Journal*  
36 902 *of Wildlife Diseases*, 25, 416-419.  
37  
38  
39  
40 903 46. Decreto del Presidente della Giunta Regionale della Liguria n. 282/49671 del 2 maggio  
41 904 1997 norme di prevenzione per la vendita di prodotti della pesca da consumarsi crudi  
42 905 o praticamente crudi.  
43  
44  
45 906 47. Decreto Legge No. 158 del 13 settembre 2012 Disposizioni urgenti per promuovere lo  
46 907 sviluppo del Paese mediante un più alto livello di tutela della salute (12G0180).  
47 908 *Gazzetta Ufficiale*. SG 214 del 13 settembre 2012.  
48  
49  
50  
51 909 48. Decreto Legislativo 30 dicembre 1992, n.531 Attuazione della Direttiva 91/493/CEE  
52 910 che stabilisce le norme sanitarie applicabili alla produzione e commercializzazione dei  
53 911 prodotti della pesca, tenuto conto delle modifiche apportate dalla Direttiva 92/48/CEE  
54 912 che stabilisce le norme igieniche minime applicabili ai prodotti della pesca ottenuti a  
55 913 bordo di talune navi. *Gazzetta Ufficiale*, 7.  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

- 914 49. Decreto Legislativo del 17 Luglio 2013 informazioni obbligatorie a tutela del  
1 consumatore di pesce e cefalopodi freschi e di prodotti di acqua dolce, in attuazione  
2 915  
3 dell'articolo 8, comma 4, del decreto-legge 13 settembre 2012 No. 158, convertito,  
4 916  
5 con modificazioni, dalla legge 8 novembre 2012, No. 189. *Gazzetta Ufficiale*, 187 del  
6 917  
7 18 agosto 2013.  
8
- 9 919 50. De Nicola, P., Napolitano, L., Di Bartolomeo, N., Waku, M., & Innocenti, P. (2005).  
10 920 Su di un caso di Anisachiasi con perforazione del cieco. *Il Giornale di Chirurgia*, 26,  
11 921 375- 377.  
12
- 13 922 51. Cristina De Rosa (2011). *Una parassitosi emergente: l'Anisakiasi. Thesis in Medicine*  
14 923 *and Surgery, University of Chieti - Pescara, A.A. 2010-2011.*  
15
- 16 924 52. Directive 1999/3/EC of the European Parliament and of the Council (22 February  
17 925  
18 1999) on the establishment of a Community list of foods and food ingredients treated  
19 926  
20 with ionising radiation. *Official Journal of European Community*, L66, 24-25.  
21
- 22 927 53. Directive 2003/89/EC of the European Parliament and of the Council (10 November  
23 928  
24 2003) amending Directive 2000/13/EC as regards indication of the ingredients present  
25 929  
26 in foodstuffs. *Official Journal of European Union*, L308, 15-18.  
27
- 28 930 54. Dominguez-Ortega, J., Alonso-Llamazares, A., Rodriguez, L., Chamorro, M., Robledo,  
29 931  
30 T., Bartolome, J.M., & Martínez-Cóccera C. (2001). Anaphylaxis due to  
31 932  
32 hypersensitivity to *Anisakis simplex*. *International Archives of Allergy and*  
33 933  
34 *Immunology*, 125, 86-88.  
35
- 36 934 55. European Food Safety Authority, Panel on Biological Hazards (BIOHAZ) (2010).  
37 935  
38 Scientific Opinion on risk assessment of parasites in fishery products. . *EFSA Journal*,  
39 936  
40 8, 1543.  
41
- 42 937 56. Fagerholm, H.P. (1991). Systematic implications of male caudal morphology in  
43 938  
44 ascaridoid nematode parasites. *Systematic Parasitology*, 19, 215-229.  
45
- 46 939 57. Farkas, J. (1998). Irradiation as a method decontaminating food. A review.  
47 940  
48 *International Journal of Food Microbiology*, 44, 189-204.  
49
- 50 941 58. Federal Republic of Germany (1988). Verordnung über gesundheitliche  
51 942  
52 Anforderungen an Fische und Schalentiere (Fisch-Verordnung). Vom 19. *Federal Law*  
53 943  
54 *Gazette BGBl*, I41, 1570.  
55
- 56 944 59. Fernández de Corres, L., Audicana, M., Del Pozo, M. D., Muñoz, D., Fernández, L.,  
57 945  
58 Navarro, J. A., García, M., & Diez J. (1996). *Anisakis simplex* induces not only  
59 946  
60 anisakiasis: report on 28 cases of allergy caused by this nematode. *Journal of*  
61 947  
62 *Investigational Allergology and Clinical Immunology*, 6, 315-319.  
63  
64  
65

- 948 60.Fioravanti, M.L., Gavaudan, S., Vagnini, V., & Tonucci, F. (2004). Indagine sulla  
1 949 diffusione di larve di *Anisakis* e *Hysterothylacium* (Nematoda, Anisakidae) in pesci  
2 del Mar Adriatico centrale. *Webzine Sanità Pubblica Veterinaria*, 23. Available at:  
3 950 [http://spvet.it/arretrati%5Cnumero\\_23%5Canis.html](http://spvet.it/arretrati%5Cnumero_23%5Canis.html). Accessed 15 April 2014.  
4 951  
5 952 61.French Agency for Food, Environmental and Occupational Health and Safety (2011).  
6  
7 953 Data sheet on foodborne biological hazards /*Anisakis* spp., *Pseudoterranova* spp.  
8 Available at: <http://www.anses.fr/Documents/MIC-Fi-AnisakisEN.pdf>. Accessed 23  
9 954 April 2014.  
10 955  
11 956 62.French Food Safety Agency (2007). Opinion of the French Food Safety Agency  
12 (AFSSA) on a risk assessment request concerning the presence of anisakidae in  
13 957 fishery products and the extension of the exemption from the freezing sanitary  
14 958 obligation of fishery products whose feeding is under control and for certain species of  
15 959 wild fish. AFSSA – Request no. 2007-SA.0379. Available at:  
16 960 <http://www.anses.fr/Documents/MIC2007sa0379EN.pdf>. Accessed 16 April 2014.  
17 961  
18 962 63.Fumarola, L., Monno, R., Ierardi, E., Rizzo, G., Giannelli, G., Lalle, M., & Pozio, E.  
19 963 (2009). *Anisakis pegreffii* etiological agent of gastric infections in two Italian women.  
20 964 *Foodborne Pathogens and Disease*, 6, 1157-1159.  
21 965  
22 966 64.Garcia, M., Blanco, J.G., Garces, M., Juste, S., Fuentes, M., & Herrero D. (2001).  
23 967 Freezing protect against allergy to *Anisakis simplex*. *Journal of investigational*  
24 968 *allergology & clinical immunology: official organ of the International Association of*  
25 969 *Asthmology (INTERASMA) and Sociedad Latinoamericana de Alergia e Inmunologia*,  
26 11, 49-52.  
27 970  
28 971 65.Genchi, C., & Pozio E. (2003). *De Carneri. Parassitologia generale e umana*. Milano:  
29 972 Ambrosiana Rozzano.  
30 973  
31 974 66.George-Nascimento, M., & Urrutia X. (2000). *Pseudoterranova cattani* sp. nov.  
32 975 (Ascaridoidea:Anisakidae), a parasite of the South American sea lion *Otaria byronia*  
33 976 De Blainville from Chile. *Revista Chilena de Historia Natural*, 73, 93-98.  
34 977  
35 978 67.Griglio, B., Marro, S., Marotta, V., Testa, A., Sattanino, G., Civera, T., Fazio, G.,  
36 979 Negro, M., Cravero, M., Pairone, B., Raschio, C., Rossignoli M., Biglia, C., &  
37 980 Decastelli, L. (2012). Anisakidae: valutazione del rischio e indicazioni operative per i  
38 981 controlli ufficiali alla luce del quadro normativo. *Associazione Italiana Veterinaria*  
39 982 *Medicina Pubblica*, 4. Available at:  
40 983 <http://www.ceirsa.org/fd.php?path=201402/anisakis.pdf>. Accessed 15 April 2014.  
41 984  
42 985  
43 986  
44 987  
45 988  
46 989  
47 990  
48 991  
49 992  
50 993  
51 994  
52 995  
53 996  
54 997  
55 998  
56 999  
60  
61  
62  
63  
64  
65

- 981 68.Hochberg, N.S., Hamer, D.H., Hughes, J.M., & Wilson, M.E. (2010). Anisakidosis:  
1 982 Perils of the deep. *Clinical Infectious Diseases*, 51, 806-812.
- 3 983 69.Horst, K. (2008). Nematode larvae in fish on the German market 20 years of consumer  
4 related research. *Archiv für Lebensmittelhygiene*, 59, 107-116.
- 5 984  
6  
7 985 70.Inoue, K., Oshima, S.I., Hirata, T., & Kimura, I. (2000). Possibility of anisakidae  
8 larvae infection in farmed salmon. *Fisheries Science*, 66, 1049-1052.
- 9 986  
10 987 71.Ishikura, H., Kikuchi, K., Nagasawa, K., Ooiwa, T., Takamiya, H., Sato, N., & Sugane,  
11 K. (1993). Anisakidae and anisakidosis. In M.D. Tsieh (ed.), *Sun Progress in clinical*  
12 988 *parasitology* (pp. 43-102). New York: Springer.
- 13 989  
14 990 72.Karl, H., Leinemann, M. (1997). A fast and quantitative detection method for  
15 nematodes in fish fillets and fishery products. *Archiv für Lebensmittelhygiene*, 44,  
16 991 105-128.
- 17 992  
18 993 73.Kassai T., Cordero Del Campillo M., Euzeby, J., Gaafar, S., Hiepe Th., & Himonas,  
19 C.A. (1988). Standardized Nomenclature Of Animal Parasitic Diseases (SNOAPAD).  
20 994 *Veterinary Parasitology*, 29, 299-326.
- 21 995  
22 996 74.Klimpel, S., Palm, H.W., Rückert, S., & Piantkowsky, U. (2004). The life cycle of  
23 *Anisakis simplex* in the Norwegian Deep (Northern North Sea). *Journal of*  
24 997 *Parasitology Research*, 94, 1-9.
- 25 998  
26 999 75.Kjøie, M. (1993). Aspects of the cycle and morphology of *Hysterothylacium aduncum*  
27 (Rudolphi, 1802) (Nematoda, Ascaridoidea, Anisakidae). *Canadian Journal of*  
28 1000 *Zoology*, 71, 1289-1296.
- 29 1001  
30 1002 76.La Pira, R. (2013). Sgombri al naturale con *Anisakis* ritirate 80 mila lattine: nessun  
31 pericolo per la salute. Ecco la lista dei prodotti e i lotti. Il Fatto Alimentare. Available  
32 at: [http://www.ilfattoalimentare.it/sgombri-anisakis-lista-prodotti-ritirati-nessun-](http://www.ilfattoalimentare.it/sgombri-anisakis-lista-prodotti-ritirati-nessun-pericolo-salute.html)  
33 1003 [pericolo-salute.html](http://www.ilfattoalimentare.it/sgombri-anisakis-lista-prodotti-ritirati-nessun-pericolo-salute.html). Accessed 16 April 2014.
- 34 1004  
35 1005 77.Legge n. 283 del 30 aprile 1962. Modifica degli articoli 242, 243, 247, 250 e 262 del  
36 testo unico delle leggi sanitarie, approvato con regio decreto 27 luglio 1934, n. 1265:  
37 1006 Disciplina igienica della produzione e della vendita delle sostanze alimentari e delle  
38 1007 bevande. *Gazzetta Ufficiale*, 139 del 4 giugno 1962.
- 39 1008  
40 1009 78.Lima dos Santos, C.A.M., & Howgate, P. (2011). Fishborne zoonotic parasites and  
41 aquaculture: a review. *Aquaculture*, 318:253-261.
- 42 1010  
43 1011 79.Llarena-Reino, M., González, Á.F., Vello, C., Outeiriño, L., & Pascual, S. (2012). The  
44 accuracy of visual inspection for preventing risk of *Anisakis* spp. infection in  
45 unprocessed fish. *Food Control*, 23, 54-58.
- 46 1012  
47 1013  
48 1014  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

- 1015 80.Lunestad, B.T. (2003). Absence of Nematodes in Farmed Atlantic Salmon (*Salmo*  
1 *salar* L.) in Norway. *Journal of Food Protection*, 66, 122-124.  
2 1016
- 3 81.Lymbery, A.J., & Cheah, F.Y. (2007). Anisakid nematodes and Anisakiasis. In S.J.  
4 1017  
5 1018 Black (ed.), *Food-borne parasitic zoonoses fish: plant-borne parasites* (pp. 433). New  
6 1019  
7 1019 York: Springer.  
8 1020
- 9 1020 82.Marty, G.D. (2008). Anisakid larva in the viscera of a farmed Atlantic salmon (*Salmo*  
10 *salar*). *Aquaculture*, 279, 209-210.  
11 1021
- 12 83.Masotti, G., Amadei, P., & Lanni, L. (2010). Pesce crudo: Condizioni igieniche,  
13 1022  
14 1023 operative e strutturali nei ristoranti giapponesi. *SIVEMP Argomenti 2* - Available at:  
15 1024  
16 1024 [http://www.sivemp.it/userfiles/\\_documents/\\_rivista\\_/26\\_58\\_62\\_pesce.pdf](http://www.sivemp.it/userfiles/_documents/_rivista_/26_58_62_pesce.pdf). Accessed  
17 1025  
18 1025 15 April 2014.  
19 1026
- 20 1026 84.Mattiucci, S., Paggi, L., Nascetti, G., Ishikura, H., Kikuchi, K., Sato, N., Cianchi, R., &  
21 1027  
22 1027 Bullini, L. (1998). Allozyme and morphological identification of *Anisakis*,  
23 1028  
24 1028 *Contracaecum* and *Pseudoterranova* (Nematoda: Ascaridoidea) from Japanese waters.  
25 1029  
26 1029 *Systematic Parasitology*, 40, 81-92.
- 27 1030 85.Mattiucci, S., Abaunza, P., Damiano, S., Garcia, A., Santos, M.N., & Nascetti, G.  
28 1031  
29 1031 (2007). Distribution of *Anisakis* larvae, identified by genetic markers, and their use for  
30 1032  
31 1032 stock characterization of demersal and pelagic fish from European waters: an update.  
32 1033  
33 1033 *Journal of Helminthology*, 81, 117-127.
- 34 1034 86.Mattiucci, S., & Nascetti, G. (2008). Advances and trends in the molecular systematics  
35 1035  
36 1035 of anisakid nematodes, with implications for their evolutionary ecology and host-  
37 1036  
38 1036 parasite co-evolutionary processes. *Advances in Parasitology*, 66, 147-148.  
39 1037
- 40 1037 87.Mattiucci, S., Paoletti, M., Borrini, F., Palumbo, M., Macarone-Palmieri, R., Gomes,  
41 1038  
42 1038 V., Casati, A., & Nascetti, G. (2011). First molecular identification of the zoonotic  
43 1039  
44 1039 parasite *Anisakis pegreffii* (Nematoda: Anisakidae) in a paraffin-embedded granuloma  
45 1040  
46 1040 taken from a case of human intestinal anisakiasis in Italy. *BMC Infectious Diseases*,  
47 1041  
48 1041 11, 82.
- 49 1042 88.McClelland, G. (2002). The trouble with sealworms (*Pseudoterranova decipiens*  
50 1043  
51 1043 species complex, Nematoda): a review. *Parasitology*, 124, 183-203.  
52 1044
- 53 1044 89.Milligan, R.J. (2008). The occurrence and behaviour of *Pseudoterranova decipiens* and  
54 1045  
55 1045 *Anisakis simplex* (Nematoda) in *Gadus morhua* and their impacts on commercial  
56 1046  
57 1046 processing. MSc(R) Thesis, University of Glasgow.  
58  
59  
60  
61  
62  
63  
64  
65

- 1047 90.Mladineo, I., & Poljak, V. (2013) Ecology and Genetic Structure of Zoonotic *Anisakis*  
1 1048 spp. from Adriatic Commercial Fish Species. *Applied and Environmental*  
2 1049 *Microbiology*, 80, 1281-1290.
- 3 91.Mo, T.A., Gahr, A., Hansen, H., Hoel E., Oaland, Ø., & Poppe T.T. (2014). Presence  
4 1050 of *Anisakis simplex* (Rudolphi, 1809 det. Krabbe, 1878) and *Hysterothylacium*  
5 1051 *aduncum* (Rudolphi, 1802) (Nematoda; Anisakidae) in runts of farmed Atlantic  
6 1052 salmon, *Salmo salar* L. *Journal of Fish Diseases*, 37, 135–140.
- 7 92.Moneo, I., Caballero, M.L., Gòmez, F., Ortega, E., & Alonso, M.J. (2000). Isolation  
8 1053 and characterization of a major allergen from the fish parasite *Anisakis simplex*.  
9 1054 *Journal of Allergy and Clinical Immunology*, 106, 177-182.
- 10 93.Moneo, I., Caballero, M.L., González-Muñoz, M., Rodríguez-Mahillo, A.I.,  
11 1055 Rodríguez-Perez, R., & Silva A. (2005). Isolation of a heat-resistant allergen from the  
12 1056 fish parasite *Anisakis simplex*. *Parasitology Research*, 96, 285-289.
- 13 94.Moschella, C.M., Mattiucci, S., Mingazzini, P., De Angelis, G., Assenza, M.,  
14 1057 Lombardo, F., Monaco, S., Paggi, L., & Modini, C. (2004). Intestinal Anisakiasis in  
15 1058 Italy: case report. *Journal of Helminthology*, 78, 271-273.
- 16 95.Murrell, K. D., & Fried, B. (2007). *Food-borne parasitic zoonoses: fish and plant-*  
17 1059 *borne parasites*. New York: Springer, (Vol. 11).
- 18 96.Nagasawa, K. (1990). The life cycle of *Anisakis simplex*: a review. In H. Ishikura, & K.  
19 1060 Kikuchi (Eds.), *Intestinal Anisakiasis in Japan* (pp. 31-40). Tokyo: Springer Japan.
- 20 97.Nieuwenhuizen, N., Lopata, A.L., Jeebhay, M.F., De'Broski R. H., Robins, T.G.,  
21 1061 Brombacher F. (2006). Exposure to the fish parasite *Anisakis* causes allergic airway  
22 1062 hyperreactivity and dermatitis. *Journal of Allergy and Clinical Immunology*, 117,  
23 1063 1098–1105.
- 24 98.Norwegian Directorate of Fisheries (1999). Quality regulations relating to fish and  
25 1064 fishery products. Available at: <http://faolex.fao.org/docs/pdf/nor13807E.pdf>. Accessed  
26 1065 23 April 2014.
- 27 99.Norwegian Food Safety Authority (2011). Statement concerning *Anisakis* parasites in  
28 1066 farmed salmonids. Erklæring, anisakis, oppdrettet laksefisk, engelsk 2.2.34-2011-03  
29 1067 Available at:  
30 1068 [http://www.mattilsynet.no/skjema/2234\\_erklaering\\_anisakis\\_oppdrettet\\_laksefisk\\_engelsk\\_2011\\_03\\_specimen.11242/BINARY/2.2.34%20Erkl%C3%A6ring%20anisakis%20oppdrettet%20laksefisk,%20engelsk,%202011%2003%20specimen](http://www.mattilsynet.no/skjema/2234_erklaering_anisakis_oppdrettet_laksefisk_engelsk_2011_03_specimen.11242/BINARY/2.2.34%20Erkl%C3%A6ring%20anisakis%20oppdrettet%20laksefisk,%20engelsk,%202011%2003%20specimen). Accessed 16  
31 1069 April 2014.
- 32 1070  
33 1071  
34 1072  
35 1073  
36 1074  
37 1075  
38 1076  
39 1077  
40 1078  
41 1079  
42 1080  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

- 1081 100.Nota No. 16294 del 25/05/2010 della Direzione Sanità Pubblica della Regione  
1  
2 1082 Piemonte. Chiarimenti concernenti alcuni aspetti applicativi del Regolamento (CE)  
3  
4 1083 No. 853/2004 in materia di vendita e somministrazione di preparazioni gastronomiche  
5  
6 1084 contenenti prodotti della pesca destinati ad essere consumati crudi o praticamente  
7  
8 1085 crudi. Articolo 5 della Legge 283/62 – Presenza di Anisakis
- 9 1086 101.Nota No. 8499 del 22/03/2011 della Regione Piemonte. Chiarimenti concernenti  
10  
11 1087 alcuni aspetti applicativi del Regolamento (CE) No. 853/2004 in materia di vendita e  
12  
13 1088 somministrazione di preparazioni gastronomiche contenenti prodotti della pesca  
14  
15 1089 destinati ad essere consumati crudi o praticamente crudi. Articolo 5 della Legge  
16  
17 1090 283/62 – Presenza di Anisakis.
- 18 1091 102.Nota Ministeriale DGISAN 0024111-P-05/07/2012 sulle preparazioni gastronomiche  
19  
20 1092 contenenti prodotti della pesca destinati ad essere consumati crudi o praticamente  
21  
22 1093 crudi.
- 23  
24 1094 103.Notarbartolo-di-Sciara, G., Agardy, T., Hyrenbach, D., Scovazzi, T., & Van  
25  
26 1095 Klaveren, P. (2008). The Pelagos sanctuary for Mediterranean marine mammals.  
27  
28 1096 *Aquatic Conservation: Marine and Freshwater Ecosystems*, 18, 367-391.
- 29 1097 104.Notice No. 2009/C 283/02 of 24 November 2009 list of Member States’  
30  
31 1098 authorisations of food and food ingredients which may be treated with ionising  
32  
33 1099 radiation (According to Article 4(6) of Directive 1999/2/EC of the European  
34  
35 1100 Parliament and of the Council on the approximation of the laws of the Member States  
36  
37 1101 concerning foods and food ingredients treated with ionising radiation. *Official Journal*  
38  
39 1102 *of European Union*, C283, 5.
- 40 1103 105.Ordinanza Ministeriale 12 maggio 1992 misure urgenti per la prevenzione delle  
41  
42 1104 parassitosi da Anisakis. *Gazzetta Ufficiale* SG 121 del 25 maggio 1992.
- 43  
44 1105 106.Orphanet (2013). Prevalence of rare diseases: Bibliographic data. Orphanet Report  
45  
46 1106 Series, Rare Diseases collection, 1. Available at:  
47  
48 1107 [http://www.orpha.net/orphacom/cahiers/docs/GB/Prevalence\\_of\\_rare\\_diseases\\_by\\_alp](http://www.orpha.net/orphacom/cahiers/docs/GB/Prevalence_of_rare_diseases_by_alphabetical_list.pdf)  
49  
50 1108 [habetical\\_list.pdf](http://www.orpha.net/orphacom/cahiers/docs/GB/Prevalence_of_rare_diseases_by_alphabetical_list.pdf). Accessed 15 April 2014.
- 51 1109 107.Paggi, L., Nascetti, G., Cianchi, R., Orecchia, P., Mattiucci, S., D’Amelio, S.,  
52  
53 1110 Berland, B., Bratney, J., Smith, J.W., & Bullini, L. (1991). Genetic evidence for three  
54  
55 1111 species within *Pseudoterranova decipiens* (Nematoda, Ascaridida, Ascaridoidea) in  
56  
57 1112 the North Atlantic and Norwegian and Barents Seas. *International Journal for*  
58  
59 1113 *Parasitology*, 21, 195-212.
- 60  
61  
62  
63  
64  
65



- 1114 108. Peñalver, J., Dolores, E. M., & Muñoz, P. (2010). Absence of anisakid larvae in  
1 1115 farmed European sea bass (*Dicentrarchus labrax* L.) and gilthead sea bream (*Sparus*  
2 1116 *aurata* L.) in Southeast Spain. *Journal of Food Protection*, 73, 1332-1334.
- 3 1117 109. Panebianco, A., & Lo Schiavo, A. (1985). Ricerca di larve di Anisakidi in aringhe  
4 1118 salate e affumicate del commercio. Considerazioni di carattere ispettivo. *La Clinica*  
5 1119 *Veterinaria*, 108, 180-184.
- 6 1120 110. Ponzio, E. (2004). Zoonosi parassitarie Trasmesse da prodotti ittici. Workshop di  
7 1121 aggiornamento su problematiche emergenti nel settore dei prodotti ittici. Available at:  
8 1122 <http://www.iss.it/binary/publ/publi/05-24.1129716985.pdf> Accessed 15 April 2014.
- 9 1123 111. Pravettoni, V., Primavesi, L., & Piantanida, M. (2012). *Anisakis simplex*: current  
10 1124 knowledge. *European Annals of Allergy and Clinical Immunology*, 44, 150-156.
- 11 1125 112. Rapid Alert System for Food and Feed (2014). RASFF Portal. Available at:  
12 1126 [https://webgate.ec.europa.eu/rasff-](https://webgate.ec.europa.eu/rasff-window/portal/?event=SearchForm&cleanSearch=1)  
13 1127 [window/portal/?event=SearchForm&cleanSearch=1](https://webgate.ec.europa.eu/rasff-window/portal/?event=SearchForm&cleanSearch=1). Accessed 15 April 2014.
- 14 1128 113. Real Decree 1420/2006 of 1 December 2006 on prevention of parasitism by *Anisakis*  
15 1129 in fish products provided by establishments that serve food to final consumers or  
16 1130 Communities. Boletín Oficial de Estado.
- 17 1131 114. Regulation (EC) No. 853/2004 of the European Parliament and of the Council (29  
18 1132 April 2004) laying down specific hygiene rules for the hygiene of foodstuffs. *Official*  
19 1133 *Journal of European Union*, L139, 55-206.
- 20 1134 115. Rello, F.J., Adroher, F.J., Benitez, R., & Valero, A. (2009). The fishing area as a  
21 1135 possible indicator of the infection by Anisakids in anchovies (*Engraulis encrasicolus*)  
22 1136 from southwestern Europe. *International Journal of Food Microbiology*, 129, 277-  
23 1137 281.
- 24 1138 116. Renon, P., & Malandra, R. (1991). Anisakidosi. Indagine sulla presenza di larve di  
25 1139 anisakidi in teleostei marini catturati nei mari italiani. *Obiettivi e Documenti*  
26 1140 *Veterinari*, 12, 73-79.
- 27 1141 117. Risoluzione del Consiglio Regionale Puglia n. 1675 del 17 settembre 2013, indirizzi  
28 1142 per l'adozione di un sistema per la sorveglianza e il controllo delle patologie  
29 1143 dell'uomo causate da *Anisakis spp.* in Puglia. *Bollettino Ufficiale*, 131.
- 30 1144 118. Rodríguez-Mahillo, A.I, González-Muñoz, M., De Las Heras, C., Tejada, M., &  
31 1145 Moneo, I. (2010). Quantification of *Anisakis simplex* allergens in fresh, long-term  
32 1146 frozen, and cooked fish muscle. *Foodborne Pathogens and Disease*, 7, 967-973.
- 33 1147  
34 1148  
35 1149  
36 1150  
37 1151  
38 1152  
39 1153  
40 1154  
41 1155  
42 1156  
43 1157  
44 1158  
45 1159  
46 1160  
47 1161  
48 1162  
49 1163  
50 1164  
51 1165  
52 1166  
53 1167  
54 1168  
55 1169  
56 1170  
57 1171  
58 1172  
59 1173  
60 1174  
61 1175  
62 1176  
63 1177  
64 1178  
65 1179

- 1147 119.Rosales, M.J., Mascaro, C., Fernandez, C., Luque, F., Moreno, M.S., Parras, L.,  
1 1148 Cosano, A., & Muñoz, J.R. (1999). Acute intestinal Anisakiasis in Spain: a fourth-  
2 1149 stage *Anisakis simplex* larva. *Memórias do Instituto Oswaldo Cruz*, 94, 823-826.  
3  
4  
5 1150 120.Sakanari, J.A., & McKerrow, J.H. (1989). Anisakiasis. *Clinical Microbiology*  
6  
7 1151 *Reviews*, 2, 278-284.  
8  
9 1152 121.SANCO/10137/2013-rev1 (2013) Guidance on the term 'obviously contaminated' in  
10  
11 1153 relation to parasites in fishery products. Working document SANCO/10137/2013-  
12  
13 1154 rev1. Available at:  
14 <http://multimedia.food.gov.uk/multimedia/pdfs/obviouslycontaminated>. Accessed 16  
15 1155 April 2014.  
16 1156  
17  
18 1157 122.Scala, E., Giani M., Pirrotta L., Guerra E.C., Cadoni S., Girardelli C.R., De Pita O., &  
19  
20 1158 Puddu P. (2001). Occupational generalised urticaria and allergic airborne asthma due  
21  
22 1159 to *Anisakis simplex*. *European Journal of Dermatology*, 11, 249-250.  
23  
24 1160 123.Schiavo, A. (2005). Anisakidosis: aspetti principali di interesse generale. *Il Pesce* 3.  
25 1161 Available at: <http://max.pubblicitaitalia.com/ilpesce/2005/3/6053.html>. Accessed 15  
26 1162 April 2014.  
27  
28  
29 1163 124.Skov, J., Kania, P.W., Olsen, M.M., Lauridsen, J.H., & Buchmann, K. (2009).  
30  
31 1164 Nematode infections of maricultured and wild fishes in Danish waters: a comparative  
32  
33 1165 study. *Aquaculture*, 298, 24-28.  
34  
35 1166 125.Smith, J.W., & Wootten, R. (1978a). Experimental studies on the migration of  
36 1167 *Anisakis sp.* larvae (Nematoda: ascaridida) into the flesh of herring, *Clupea harengus*  
37  
38 1168 L. *International Journal for Parasitology*, 5, 133-136.  
39  
40 1169 126.Smith, J.W., & Wootten, R. (1978b). *Anisakis* and anisakiasis. *Advances in*  
41  
42 1170 *Parasitology*, 16, 93-163.  
43  
44 1171 127.Smith, J.W. (1999). Ascaridoid nematodes and pathology of the alimentary tract and  
45  
46 1172 its associated organs in vertebrates, including man: a literature review.  
47 1173 *Helminthological Abstracts*, 68, 49-96.  
48  
49 1174 128.Smith, J.W., & Snyder, J.M. (2005). New locality records for third-stage larvae of  
50  
51 1175 *Anisakis simplex* (sensu lato)(Nematoda: Ascaridoidea) in euphausiids *Euphausia*  
52  
53 1176 *pacifica* and *Thysanoessa raschii* from Prince William Sound, Alaska. *Parasitology*  
54  
55 1177 *Research*, 97, 539-542.  
56  
57 1178 129.Stormo, S.K., Sivertsen, A.H., Heia, K., Nilsen, H., & Elvevoll, E. (2007). Effects of  
58 1179 single wavelength selection for Anisakid roundworm larvae detection through  
59  
60 1180 multispectral imaging. *Journal of Food Protection*, 70, 1890-1895.  
61  
62  
63  
64  
65

- 1181 130.U.S. Food and Drug Administration (2001). Fish and Fishery Products Hazards and  
1 1182 Controls Guide. FDA's Guidance 3rd Edition. Available at: <http://archive.is/uBqk>.  
2 1183 Accessed 16 April 2014.  
3  
4  
5 1184 131.U.S. Food and Drug Administration (2013). Processing Parameters Needed to Control  
6 Pathogens in Cold Smoked Fish Chapter V. Potential Hazards in Cold-Smoked Fish:  
7 1185 Parasites. Available at:  
8  
9 1186 [http://www.fda.gov/Food/FoodScienceResearch/SafePracticesforFoodProcesses/ucm0](http://www.fda.gov/Food/FoodScienceResearch/SafePracticesforFoodProcesses/ucm094578.htm)  
10  
11 1187 [94578.htm](http://www.fda.gov/Food/FoodScienceResearch/SafePracticesforFoodProcesses/ucm094578.htm). Accessed 16 April 2014.  
12  
13 1188  
14 1189 132.Van Thiel, P.H. (1962). Anisakiasis. *Parasitology*, 52, 16-17.  
15  
16 1190 133.Wooten, R., Yoon, G.H., & Bron, J. (2009). A Survey of Anisakid Nematodes in  
17 Scottish Farmed Salmon. FSAS Project 14008E. Available at:  
18 1191 [http://www.foodbase.org.uk/results.php?f\\_category\\_id=&f\\_report\\_id=355](http://www.foodbase.org.uk/results.php?f_category_id=&f_report_id=355). Accessed  
19  
20 1192 16 April 2014.  
21 1193  
22  
23 1194 134.World Health Organization (2013). Foodborne trematode infections. Fact sheet N°368  
24 Available at: <http://www.who.int/mediacentre/factsheets/fs368/en/index.html>.  
25 1195 Accessed 15 April 2014.  
26  
27 1196  
28  
29 1197 135.Yagi, K., Nagasawa, K., Ishikura, H., Nagasawa, A., Sato, N., Kikuchi, K., &  
30 Ishikura, H. (1996). Female worm *Hysterothylacium aduncum* excreted from human: a  
31 1198 case report. *Japanese Journal of Parasitology*, 45, 12-23.  
32  
33 1199  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

NUMBER OF SPECIMENS OF THE LOT	NUMBER OF SPECIMENS TO USE FOR VISUAL INSPECTION	
	SENSIBILITY 10 %	SENSIBILITY 5 %
10	All	All
20	16	19
30	19	26
40 – 45	21	31
46 – 50	22	35
51 – 60	23	38
61 – 70	24	40
71 – 80	24	42
81 – 90	25	43
91 – 100	25	45
101 – 120	26	47
121 – 140	26	48
141 – 161	27	49
161 – 180	27	50
181 – 200	27	51
201 – 250	27	53
251 – 350	28	54
351 – 450	28	55
451 – 600	28	56
601 – 1200	29	57
1201 – 4000	29	58
> 4000	29	59

**Table 1.** Sampling protocol for the visual inspection to detect Anisakis larvae in fish batches, according to the Circular Letter VS8/C790/94 of the Lombardy region.

STANDARD OF REFERENCE	DISPOSITIONS
<i>Law n. 293 of 30 April 1962</i>	<ul style="list-style-type: none"> <li>• Prohibition of use in the preparation of food or drink, sell, hold to sell, administer or otherwise distribute for consumption foodstuffs soiled, invaded by parasites, altered or harmful.</li> </ul>
<i>Ordinance of 8 August 1988 of German Federal Republic</i>	<ul style="list-style-type: none"> <li>• Removing parts containing nematodes (dead or alive) and their marketing ban.</li> <li>• Official procedures for frozen, cooked, salted or pickled products.</li> </ul>
<i>Italian Ministry of Health Circular Letter No. 10 of 11 March 1992</i>	<ul style="list-style-type: none"> <li>• Prompt evisceration of fishery products at risk of infestation, longer than 18 cm.</li> <li>• The official veterinarian must perform a check of the fish, through random sampling and visual examination, with the opening of the coelomic cavity.</li> <li>• Freezing treatment at -20 ° C for at least 24h</li> </ul>
<i>Italian Ministry of Health Ministerial Decree of 12 May 1992</i>	<ul style="list-style-type: none"> <li>• The treatment for killing parasites must be certified.</li> </ul>
<i>Legislative Decree No. 531 of 30 December 1992</i>	<ul style="list-style-type: none"> <li>• Obligation to carry out the visual inspection with the purpose of visible parasites detection, prior to the placement on the market of fishery products.</li> </ul>
<i>Decision 93/140/EEC of 19 January 1993</i>	<ul style="list-style-type: none"> <li>• Definition of "<i>visible parasite</i>" and "<i>visual inspection</i>".</li> </ul>
<i>Region of Lombardy Regional Council prot. No. 790/94</i>	<ul style="list-style-type: none"> <li>• Establishment of sampling procedures for <i>Anisakis</i> spp</li> </ul>
<i>Council Regulation (EC) No. 2406/96 of 26 November 1996</i>	<ul style="list-style-type: none"> <li>• Negative impact of parasites on the product quality.</li> </ul>
<i>Region of Liguria Decree of the President of the Regional Council n. 282/49671 of 2</i>	<ul style="list-style-type: none"> <li>• Display of an informative billboard at retail outlets (including points of direct sale from the fisherman to the consumer or retailer) where are marketed and/or sold raw</li> </ul>

<i>May 1997</i>	anchovies.
<i>Region of Liguria</i> <i>Circular Letter No. 1/97 of 24 March 1997</i>	<ul style="list-style-type: none"> <li>• Detailing on sampling procedures for <i>Anisakis</i> spp</li> </ul>
<i>Regulation (EC) No. 852/2004 of the European Parliament and of the Council of 29 April 2004</i>	
<i>Regulation (EC) No. 853/2004 of the European Parliament and of the Council of 29 April 2004</i>	<ul style="list-style-type: none"> <li>• The FBOs must ensure that fishery products are subjected to a visual inspection for the detection of visible endoparasites, before placing on the market.</li> <li>• FBOs must not place on the market, for human consumption, fishery products that are obviously contaminated with parasites.</li> <li>• Prohibition to trade products obviously contaminated with parasites</li> <li>• Obligation of the freezing treatment at -20 ° C (core temperature) for at least 24 hours, to kill the "nematode larvae."</li> </ul>
<i>Commission Regulation (EC) No. 2074/2005 of 5 December 2005</i>	<ul style="list-style-type: none"> <li>• Clarification about the “visible parasite” definition and on how to conduct the visual inspection.</li> </ul>
<i>Spanish Ministry of Health and Consumer Affair</i> <i>Real Decreto No. 1420/2006</i>	<ul style="list-style-type: none"> <li>• Obligation for bars, restaurants, cafes, hotel, hospitals, schools, residences, canteens and schools, catering services and similar, to perform a freezing treatment of the fish, to be consumed raw or almost raw, cold-smoked, marinated or salted.</li> </ul>
<i>Italian Ministry of Health</i> <i>DGSAN 10776-P-26/05/2008</i>	<ul style="list-style-type: none"> <li>• The alert system for the detection of <i>Anisakis</i> in fishery products from Third Countries, should only be activated upon detection of viable larvae of <i>Anisakis</i> in the coelomic cavity or in the flesh of fresh fish products</li> </ul>

<p><i>Commission Regulation (EC) No. 1020/2008 of 17 October 2008</i></p>	<ul style="list-style-type: none"> <li>• The visual inspection becomes mandatory for all the FBOs, including those that operate at the retail level.</li> </ul>
<p><i>Region of Piemont</i> <i>Directorate of Public Health Note No. 16294 of May 25, 2010</i></p>	<ul style="list-style-type: none"> <li>• Obligation for retailers to communicate to the local competent authority the detection of larvae found alive and vital during self-monitoring procedures and during sale.</li> </ul>
<p><i>Italian Ministry of Health</i> <i>DGSAN 4379-P-17/02/2011</i></p>	<ul style="list-style-type: none"> <li>• The preventive treatment carried out must be declared in the operators' self-monitoring plan.</li> <li>• The fishery products, that have been subjected to the treatment, should be always accompanied, up to the placing on the market, by a declaration of the operator responsible of the treatment.</li> <li>• Lettering for products treated "<i>Conforms to Regulation EC no. 853/20054, Annex III, Section VII, Chapter 3, part D, paragraph 3</i>".</li> </ul>
<p><i>Italian Ministry of Health</i> <i>DGSAN 4380-P-17/02/2011</i></p>	<ul style="list-style-type: none"> <li>• Where in the presence of infested products, the operator proves that he acted in good faith and in accordance with the law, implementing all the possible measures to detect parasites, it is not "perfectible" the theory of liability and therefore art. 5 of 283 in '62 is not applicable</li> </ul>
<p><i>Region of Puglia</i> <i>Note Prot. AOO152/1 MAR 2011/2998</i></p>	<ul style="list-style-type: none"> <li>• Recommendations about the timely fulfillment of the Ministerial Circular no. 4379-P and 4380-P in the regional territory.</li> <li>• Clarifying the role of the FBOs and its responsibility for the detection of larvae alive and vital during self-monitoring and during sale.</li> </ul>
<p><i>Region of Piemonte</i></p>	

<i>Note Prot. No. 8499/DB2002 014.140.20</i>	
<i>Commission Regulation (EU) No. 1276/2011</i>	<ul style="list-style-type: none"> <li>• Anisakis risk extended to cephalopods.</li> <li>• The preventive treatments must be targeted to the killing of the "viable parasite".</li> <li>• Possibility to perform the freezing treatment even at -35 ° C for at least 15 hours.</li> </ul>
<i>Italian Ministry of Health</i> <i>DGISAN 0024111-P-05/07/2012</i>	<ul style="list-style-type: none"> <li>• Importance of official control as an effective measure for managing the health risk of parasites, especially during administration and retail.</li> </ul>
<i>“Balduzzi Decree”</i> <i>Decree Law No. 158 September 13, 2012</i>	<ul style="list-style-type: none"> <li>• The FBOs are required to display a visible billboard with the relevant information concerning the correct operating conditions.</li> </ul>
<i>Region of Puglia</i> <i>Regional Council Resolution prot. 1675/2013</i>	<ul style="list-style-type: none"> <li>• Establishment of: <ul style="list-style-type: none"> <li>- Technical Working Group "<i>Prevention and surveillance of Anisakidosis associated to fishery products</i>";</li> <li>- Program "<i>Monitoring and control System for human diseases caused by Anisakis spp. in Puglia</i>";</li> <li>- Allergology Regional Network for the surveillance of the Anisakidosis</li> </ul> </li> </ul>
<i>Italian Ministry of Health</i> <i>Decree 17 July 2013</i>	<ul style="list-style-type: none"> <li>• At the point of sale of fishery products a sign must be displayed in a prominent position with the following information: "In the case of consumption as raw, marinated or not fully cooked, the product must first be frozen for at least 96 hours at - 18 ° C in domestic freezer marked with three or more stars. "</li> </ul>

**Table 2.** Chronological evolution of the European and Italian legislation, with regard to risk management of Anisakis (in white: Italian and Regional disposition; gray: European dispositions; light gray: others Member States' dispositions)



ITALIAN REGION	NORMATIVE REFERENCE	FISH SPECIES AT RISK	DISPOSITIONS	OUTCOMES OF THE VISUAL INSPECTION
<b>Lombardy</b>	Regional Council Circular Sector Health and Hygiene Veterinary Service Protocol No. 201145 / G and N. 47193 of July 20, 1994	Anchovy Sardine Mackerel Melva Bonito/Skipjack Skipjack /Tunny Lanzardo Hake Whiting/Melu Whiting/ Pier Pout /Chaplain Mullet Suro	The sampling must be carried out only for species at risk. <ul style="list-style-type: none"> <li>• Determine the total number of subjects that compose the lot *;</li> <li>• From the total number of subjects that compose the lot, is obtained the statistically significant number of samples and the percentage of desired sensitivity (5% or 10%) ^.</li> </ul> Then the specimens are subjected to visual inspection through the evisceration: <ul style="list-style-type: none"> <li>• Check for the presence of the parasite moving the viscera.</li> </ul>	If only one specimen is found parasitized, the whole lot must be withdrawn.
<b>Liguria</b>	Regional Circular Letter 1/97	Anchovy	1) Visual inspection of the lot to detect migrant parasites on the external surface of the fish; 2) Collection of specimens in a representative number, according to the number of box, composing the lot: <ul style="list-style-type: none"> <li>• &lt; 10 boxes: 3 specimens (MAX 28);</li> <li>• 10 to 150 boxes: 1 specimens per box (Max 28);</li> <li>• &gt; 150 boxes: 1 specimens per box (Max 40).</li> </ul> 3) Opening of the coelomic cavity.	<b>Positive health inspection</b> The lot is withdrawn from the market or destroyed or intended for other uses than human consumption. <b>Visual Inspection</b> <ul style="list-style-type: none"> <li>• Detection of numerous larvae and repulsive appearance of the product: the lot must be withdrawn from the market .</li> <li>• Lot intended for consumption if the n. larvae per anchovy is <math>\leq 3</math>, up to a maximum of 10% of the specimens examined;</li> <li>• Lot intended to preventive treatment if are detected: <ul style="list-style-type: none"> <li>-3 larvae per anchovy in the 10% of specimens examined;</li> <li>- the infested specimens are &lt;10% of the specimens sampled.</li> </ul> </li> </ul>
<b>Lazio</b>	Internal procedure of the Local Health Authority	<b>See Liguria Region</b>		
<b>Emilia Romagna</b>	Self-cmonitoring Procedure	Anchovies Sardines Mackerel Herring Mulletts	Sampling Method From each lot of the species at risk, according to the number of boxes that composes it, are sampled: <ul style="list-style-type: none"> <li>• From 1 to 6 boxes: select at random a total of 10 specimens;</li> </ul>	

		Hake Blue Whiting Scabbardfish Suro	<ul style="list-style-type: none"> <li>• From 7 to 12 boxes: take 1 or 2 specimens per box (minimum of 10 specimens);</li> <li>• Boxes &gt; 12: select at random at least 12 specimens from 12 different boxes.</li> </ul>	
<b>Sardinia</b>	Internal procedure of the Local Health Authority	Fish Sabre Horse Mackerel Blue Whiting Mackerel Hake Mulletts Anchovies	<p>The sampling and the visual inspection are performed for all species, not just those at risk.</p> <p>Fish size of less than 20 grams:</p> <ul style="list-style-type: none"> <li>• 3 specimens per box up to a maximum of 10 boxes;</li> <li>• 1 specimen from 30 boxes, for lot of 150 boxes;</li> <li>• 1 specimen from 40 boxes, for lot of above 150 boxes.</li> </ul> <p>Fish size greater than 20 grams:</p> <ul style="list-style-type: none"> <li>• 1 copy per cassette for a maximum of 10 cassettes</li> <li>• 1 specimen per 15 boxes for batches up to 150 cassettes</li> <li>• 1 specimen per 20 boxes for batches of more than 150 boxes.</li> </ul>	<p>In case of detection of fishery products obviously contaminated, seizure and destruction of the consignment.</p> <p>Condition for which the parasites is obviously contaminated is that the parasite is visible and that for size, color or texture is clearly distinguishable from fish tissues (Reg. 854/2004).</p>
<b>Marche</b>	Internal procedure of the Local Health Authority	Suro Blue Whiting/Whiting Scabbardfish Hake Chaplain/Pout Mackerel Lanzardo Anchovy Sardine Mullet Bonito Squid	<p>Determination of a representative sample:</p> <ul style="list-style-type: none"> <li>• For lot up to 100Kg 3 specimens per box (minimum 28) are sampled</li> <li>• For lot from 110 to 1500Kg 1 specimen per box (minimum 28) is sampled</li> <li>• For lot above 1500kg is sampled 1 specimen per box (minimum 40).</li> </ul>	<p><u>Threshold of acceptability for the products</u></p> <p><b>For Anchovies:</b></p> <ul style="list-style-type: none"> <li>• In case of massive infestation that results in a repulsive appearance of the product, this must be destroyed;</li> <li>• If &gt; 3 parasites/specimen are found in &gt; 10% of the specimens, the lot is sent to preventive treatment;</li> <li>• If &lt;3 parasites/specimen are found in &lt;10% of specimens, the lot is suitable for human consumption.</li> </ul> <p><b>For all other species (including those at risk):</b></p> <ul style="list-style-type: none"> <li>• In case of massive infestation that results in a repulsive appearance of the product, this must be destroyed;</li> <li>• In case of non-massive infestation, the products are subjected to preventive treatment.</li> </ul>
<b>Veneto</b>	Self-monitoring Procedure	Herring Sardines Anchovies Sprat Mackerel Lanzardo Suro Hake	<p>The search for Anisakis spp larvae is performed by sampling the seafood at risk during the exposure, the storage and/or prior to sale, on all specimens intended for cold processing (pickling, smoking, salting) and that have not previously subjected to freezing treatment.</p> <p>The search shall be carried out on a sample of at least:</p> <ul style="list-style-type: none"> <li>• 3 specimens, in case of large species (blue whiting-</li> </ul>	<p>Presence of superficial parasites/modest infestation: presence of parasites in small quantities (max 3 non-viable larvae)</p> <ul style="list-style-type: none"> <li>• Freezing treatment (-20 ° C for 24 hours);</li> <li>• Identification and seizure of the lot and subsequent waste disposal.</li> </ul> <p>Presence of deep parasites/invasive infestation:</p>

		<p>Blue Whiting/Whiting Scabbardfish Monkfish St. Peter's Fish Other species, object of specific controls</p>	<p>whiting, hake, monkfish, St. Peter's fish, Spanish mackerel, scabbard fish) and for lot &lt;50kg</p> <ul style="list-style-type: none"> <li>• 5 specimens in the case of lots between 50 - 500 kg;</li> <li>• 10 specimens in the case of lots &gt; 500 kg;</li> <li>• 150 specimens for small species (anchovies, sardines).</li> </ul>	<p>presence of parasites in large quantities (greater than 3 larvae) found on the surface that even deep in the viscera and penetration into the muscle</p> <ul style="list-style-type: none"> <li>• Preventive treatment before putting it on sale;</li> <li>• Identification and seizure of the lot and subsequent waste disposal with activation of the procedure for recall-retirement.</li> </ul> <p>The detection of Anisakis larvae just in a single specimen inspected (3, 5, 10 depending on the size) determines the non-compliance and therefore the preventive treatment must be performed as described above.</p>
--	--	---	---	--

**Table 3 - Sampling protocols (and related outcomes) adopted in some of the Italian regions to control the Anisakis' risk.**

\* Based on the total weight of the lot, in relation to the average weight of the species or to a conversion index per species, provided in the Circular Letter

^ Based on the table attached to the Circular Letter (for lot with specimens > 600, up to a maximum of 29 specimens are sampled)