

1 **Species diversity and community composition of native arbuscular mycorrhizal fungi in apple roots are affected**
2 **by site and orchard management**

3

4 Alessandra Turrini¹, Monica Agnolucci¹, Michela Palla¹, Elisabetta Tomé², Massimo Tagliavini², Francesca
5 Scandellari², Manuela Giovannetti^{1*}

6

7 ¹Department of Agriculture, Food and Environment, University of Pisa, Via del Borghetto 80, 56124 Pisa, Italy

8 ²Faculty of Science and Technology, Free University of Bolzano, Piazza Università 5, 39100 Bolzano, Italy

9

10 *Corresponding author: M. Giovannetti, Department of Agriculture, Food and Environment, University of Pisa, Via del
11 Borghetto 80, 56124 Pisa, Italy

12 Phone: +39.0502216643, Fax: +39.0502210606, e-mail address: manuela.giovannetti@unipi.it

13

14 **Abstract**

15 Arbuscular mycorrhizal fungi (AMF) are beneficial microorganisms which establish mutualistic symbioses with the roots
16 of most food crops, improving plant performance, nutrient uptake and tolerance to biotic and abiotic stresses. A better
17 understanding of the factors affecting AMF occurrence and diversity is fundamental to implement sustainable
18 agricultural managements effectively profiting from beneficial plant symbionts. Here, we investigated AMF occurrence,
19 diversity and community composition in the roots of apple trees from 21 orchards in South Tyrol, as affected by
20 location, management (organic vs integrated) and altitude, by PCR cloning and sequencing and PCR-DGGE of partial
21 18S rRNA gene. The screening of 448 clones from 21 clone libraries allowed the identification of 6 native AMF at the
22 species level: *Glomus indicum*, *Sclerocystis sinuosa*, *Funneliformis mosseae*, *Rhizoglomus irregulare*, *Septoglomus*
23 *constrictus* and *Claroideoglomus lamellosum*. The most abundant genera were represented by *Glomus* (29.7 % of the
24 sequences), *Paraglomus* (19.4 %), *Claroideoglomus* (17.2 %), *Sclerocystis* (16.1 %) and *Rhizoglomus* (12.3 %).
25 *Septoglomus*, *Diversispora* and *Funneliformis* sequences corresponded to less than 4 % of total sequences. Although the
26 degree of root colonization was unaffected by treatments, ANOSIM analysis of PCR-DGGE clusters revealed
27 significant differences in apple root AMF diversity between sites and agricultural managements. Species richness was
28 significantly higher in organically managed orchards than in integrated ones. Our findings provide insights into
29 important factors affecting native AMF communities of apple trees, which could be exploited in sustainable fruit
30 production systems, where beneficial soil biota boost biogeochemical cycles, energy fluxes and crop productivity.

31

32 **Keywords:** Apple trees; Native AMF communities; Mycorrhizal colonization; Small Ribosomal Subunit (SSU rDNA);
33 Clone libraries; PCR-DGGE

34

35 **1. Introduction**

36 Soil microorganisms are essential elements of soil health, fertility and productivity in sustainable and organic farming
37 systems, playing key roles in the completion of biogeochemical cycles and availability of mineral nutrients, carbon
38 sequestration and soil structure improvement, pest and disease control (Pimentel et al., 1997). Arbuscular mycorrhizal
39 (AM) fungi (AMF, Phylum Glomeromycota) are beneficial microorganisms establishing mutualistic symbioses with the
40 roots of most food crops, including cereals, legumes and fruit plants, and improving plant performance, nutrient uptake
41 and tolerance to biotic and abiotic stresses (Smith and Read, 2008). AMF are obligate biotrophs, obtaining sugars from
42 their host plants while providing soil mineral nutrients - such as P, N, S, K, Ca, Fe, Cu and Zn - absorbed and
43 translocated by means of large and ramified extraradical hyphal networks that spread from plant roots into the
44 surrounding soil (Giovannetti et al., 2015). Moreover, AMF deliver important agroecosystem services, such as soil
45 aggregation and carbon sequestration (Gianinazzi et al., 2010), and are considered important soil biological indicators
46 (Creamer et al., 2016; Stone et al., 2016). AMF enhance plant performance and fitness also through the synergistic
47 action of beneficial mycorrhizosphere bacteria, *i.e.* strictly associated to mycorrhizal hyphae, colonised roots and
48 spores, which display multifunctional activities, from antibiotic, siderophores and indole acetic acid production to P-
49 solubilisation, phytate mineralization and N-fixation (Barea et al., 2002; Philippott et al., 2013; Agnolucci et al., 2015).
50 Thus, AMF represent environmentally-friendly biofertilisers and biostimulants, as they reduce the need of chemical
51 fertilizers and pesticides in sustainable food production systems (Rouphael et al., 2015). In addition, they contribute to
52 the production of safe and high-quality food, positively affecting the synthesis of health-promoting secondary
53 metabolites in food crops (Battini et al., 2016a, b).

54 In spite of all the potential benefits of the symbiosis, AMF role has been often marginalised in agriculture. In
55 field crops, several agronomic practices, such as continuous monocultures, deep ploughing, intensive fertiliser and
56 pesticide use can decrease AMF occurrence, activity and diversity, often leading to a reduction of AMF benefits to crop
57 production and soil quality (Douds and Millner, 1999; Jansa et al., 2003; Oehl et al., 2004, 2005b; Brito et al., 2012;
58 Avio et al., 2013). Moreover, in fruit production the soil is usually replanted with the same tree species short after the
59 preceding trees are removed. In organic farming systems, thanks to the fact that synthetic fertilizers, herbicides and
60 pesticides are not applied, AMF diversity, activity and abundance have been reported to be higher than in conventional
61 systems, supporting the view that AMF play a fundamental functional role in the maintenance of soil fertility and crop

62 production in low-input agroecosystems, compensating for the reduced use of agrochemicals (Mäder et al., 2002; Oehl
63 et al., 2003, 2004; Gosling et al., 2006; Ryan and Tibbet, 2008; Mazzoncini et al., 2010; Verbruggen et al., 2010).

64 Different strategies have been devised to increase the mycorrhizal potential of soils in sustainable agriculture:
65 among them, the inoculation with non-native AMF (Jeffries et al., 2003; Gianinazzi and Vosatka, 2004) has proved
66 effective in improving root colonization, plant biomass production and P uptake (Lekberg and Koide, 2005). However,
67 such an approach involves high costs of production and application when utilised on a large scale, and raises concerns
68 about potential negative impacts of non-native invasive AMF inoculants on the composition and structure of native
69 AMF communities, possibly leading to biodiversity losses (Schwartz et al., 2006). A different approach focuses on the
70 enhancement of native AMF by means of mycotrophic crops able to maintain or increase native mycorrhizal potential
71 of soils and root colonization and growth of the subsequent crops (Kabir and Koide, 2002; Karasawa and Takebe, 2012;
72 Lehman et al., 2012; Njeru et al., 2015, 2014).

73 Little information is available on the impact of cultural practices on AMF occurrence, species richness and
74 composition in apple crop production systems. Different orchard floor managements create different levels of soil
75 disturbance and may therefore promote or depress AMF occurrence. Organic farming guidelines, for example,
76 differently from conventional or integrated fruit production techniques (Kelderer, 2004), do not allow the use of
77 chemical herbicides, but recommend, instead, the use of alternative techniques to manage the orchard floor by either
78 superficial soil tillage, grass mowing or mulching. Similarly, while conventional and integrated management techniques
79 allow the use of synthetic mineral fertilisers, organic farming guidelines recommend increasing soil fertility by
80 enhancing soil organic matter. The use of straw mulches and compost have been reported to increase apple root
81 colonisation, which was positively linked with soil pH and availability of P, K, Zn, Mn and C, suggesting that the
82 relevant organic orchard floor management practices promoted functional AMF associations more effectively than
83 conventional practices (Meyer et al., 2015). A recent work carried out using molecular methods showed that soil
84 characteristics and farming systems affect AMF diversity and community composition in the roots of cultivated apple
85 trees. In particular, a low soil available P content was associated with a higher AMF diversity in organically managed
86 orchards, as compared with conventionally managed ones (Van Geel et al., 2015). In such a study, the native AMF
87 colonizing apple roots were identified by 454-pyrosequencing of small subunit rRNA gene amplicons, with a taxonomic
88 resolution at the family level. So far, the characterization of apple AMF communities at the species level has been
89 performed only by conventional morphological methods, which show several shortfalls, mainly due to the use of spores
90 extracted from the soil, whose integrity and quality may induce species misidentification (Cavallazzi et al., 2007; Miller
91 et al., 1985; Purin et al., 2006).

92 Here we used two molecular methods, PCR-cloning and sequencing and PCR denaturing gradient gel
93 electrophoresis (PCR-DGGE) of partial 18S rRNA gene to characterize AMF communities of apple roots at genus and
94 species levels. We investigated AMF occurrence and diversity in the root systems of apple trees cultivated under
95 organic and integrated management from 21 orchards in the South Tyrol province (Italy), one of the most important
96 apple production districts in Europe, with more than 18,000 ha acreage and approx. 1 million tons apples harvested
97 yearly (Dalla Via and Mantinger, 2012). Our data provide knowledge of the factors which affect AMF colonization and
98 shape the native AMF community composition in apple roots, allowing the implementation of effective sustainable
99 management strategies that take advantage from beneficial plant symbionts.

100

101 **2. Materials and Methods**

102 **2.1. Soil and orchard management parameters**

103 The research was carried out in South Tyrol (Northern Italy). In May 2013, 21 commercial apple (*Malus domestica*
104 Borkh) orchards located either in the municipalities of Terlano (46°31' 59" N, 11°14' 47" E) or in that of Lagundo (46°
105 41' 0" N, 11° 8' 0" E) were randomly selected from a list provided by the local advisory service (The South Tyrolean
106 Advisory Service for Fruit and Grape growing, Lana, Italy). This list included drip-irrigated apple orchards planted
107 between 2007 and 2009, managed since planting following either the integrated fruit production (hereafter referred to as
108 “integrated”) or the organic farming (hereafter referred to as “organic”) guidelines.

109 In the municipality of Lagundo 3 organic and 3 integrated orchards were selected at low (298-334 m a.s.l.) and
110 at high altitude (591-660 m a.s.l.) while in the municipality of Terlano 6 organic and 3 integrated orchards were selected
111 at a low altitude (ranging from 243 to 299 m a.s.l.). Orchards were classified according to the area (Te, Terano; La,
112 Lagundo), management (O, Organic; I, Integrated) and altitude (L, Low altitude; H, High altitude) (Fig. 1).

113 The climate in area is warm temperate according to the Koppen-Geiger classification (Kottek et al., 2006).
114 Average (1993 to 2013) climatic parameters are reported in Table 1. Average annual and maximum temperatures at
115 high elevation were approx. 2 °C lower than at low elevation, while average minimum temperature was approx. 1 °C
116 lower. Soils in the area are mainly sandy loam or loamy sand and are classified as Calcaric Cambisol according to the
117 FAO Soil Taxonomy (IUSS Working Group WRB 2015) (Table 2).

118 The trees were always grafted on the same clonal rootstock (M9, T337 strain) and belonged to one of the
119 following varieties: Gala, Golden Delicious, Pinova, Modì and Red Delicious. Orchards were uniform in terms of
120 training systems (spindle bush) and planting distances (approx. 3 m between rows and 1 m between trees along the
121 row). Orchard floor management always included the presence of grassed alleys between rows (2.0 to 2.3 m wide) that
122 were mowed 3-4 times per year, while in the 0.7-1.0 m soil strip along the tree row, soil management differed between

123 organic and integrated orchards (see below). The organic and integrated orchards differed both for the type of protection
124 against aboveground pests and for soil management. In particular, in the organic orchards weed control was carried out
125 through mechanical removal of weeds (approximately five times per year) and soil fertility was maintained only by the
126 addition of organic fertilisers. In the integrated orchards, weed control in the soil strip under the trees was performed by
127 spring and autumn application of glyphosate; mineral fertilizer supply was carried out according to the nutrient budget,
128 considering expected yields of approximately 60 t ha⁻¹ and soil fertility, in the following ranges, 30-80 kg N, 19-28 kg
129 P, 71-155 kg K.

130

131 **2.2. Root, leaf and soil sampling**

132 In each orchard, root and leaf were samples from 6 randomly chosen trees from different rows, all located in the central
133 part of the plot. Sampling was carried out between May 15th and June 6th, 2013. Six soil cores were collected at a depth
134 of 10-30 cm and at approx. 20 cm distance from the trunk, using a split tube sample of 5 cm-diameter (Eijkelkamp
135 Agrisearch Equipment, BV). Roots were gently cleaned from the soil and approx. 20 g of fine roots (<2 mm diam) per
136 tree were collected and stored at 4 °C in polyethylene bags, to be successively analysed for mycorrhizal colonization. A
137 single root sample per orchard was prepared for molecular analyses (100-125 mg each) pooling together root
138 subsamples from the individual trees. Samples were stored at -20 °C, until processed. Soil samples were oven-dried at
139 65 °C until constant weight and stored at room temperature until analysed. Ten leaves per tree (without petioles) were
140 sampled from the 6th or the 7th nodes of one year-old shoots, oven-dried at 65 °C until constant weight and stored until
141 analysed.

142

143 **2.3. Mycorrhizal colonization**

144 Mycorrhizal colonization was determined on 10 g thoroughly washed root samples. Roots were cleared with 10% KOH
145 in a 80 °C water bath for 15 min and stained with Trypan blue in lactic acid (0.05 %) after 10 min in 2 % aqueous HCl.
146 The percentage of AMF colonization was calculated from root samples from individual trees from each orchard using a
147 dissecting microscope at x25 or x40 magnification and the gridline intersect method (Giovannetti and Mosse, 1980).

148

149 **2.4. Soil and leaf analysis**

150 With the exception of soil N and C concentration (performed by an elemental analyzer, Thermo Scientific), soil
151 parameters were determined according to the methods of the Verband Deutscher Landwirtschaftlicher Untersuchungs-
152 und Forschungsanstalten (VDLUFA, 1991). Leaf N concentration was determined by Elemental Analyser (Flash 2000,
153 Thermo Scientific), while leaf K, Ca, P and magnesium (Mg) concentrations were determined by inductive coupled

154 plasma-optical emission spectrometry (ICP-OES, Spectro Ametek, Arcos and Spectro Ciros CCD) after microwave
155 digestion (Milestone UltraWAVE) with 69 % of ultrapure nitric acid.

156

157 **2.5. Statistical analyses of soil, leaves and mycorrhizal colonization**

158 Data of mycorrhizal colonization (after angular transformation) and leaf mineral nutrient concentrations were analysed
159 by two-way-ANOVA. The sub-dataset from Lagundo was used to assess the interactive effects of orchard management
160 (organic vs. integrated) and orchard altitude (low vs. high). The sub-dataset obtained at low altitude was used to assess
161 the interactive effects of orchard management (organic vs. integrated) and site (Terlano vs. Lagundo). Mycorrhizal
162 colonization data were linearly correlated with soil parameters using the entire dataset. The coefficient of mycorrhizal
163 colonization data variation (CV) within each farm was also calculated. The statistical analyses were conducted using the
164 dedicated software SigmaPlot Centurion XV (StatPoint Technologies, Inc).

165

166 **2.6. Molecular analyses**

167 *2.6.1. DNA extraction from roots*

168 Genomic DNA was isolated from root material by grinding with mortar and pestle in liquid nitrogen and subsequently
169 using DNeasy Plant Mini Kit (Qiagen Milan, Italy), before performing cloning and sequencing analyses. Moreover, in
170 order to obtain DNA containing less inhibitors possibly interfering with the amplification procedure of the PCR-DGGE
171 technique, we extracted DNA from roots using the PowerSoil™ DNA Isolation Kit (MO BIO Laboratories, Solana
172 beach, CA, USA). The isolated DNA was stored at -20 °C for subsequent analyses.

173

174 *2.6.2. Cloning and sequencing*

175 DNA (10-20 ng) extracted from roots was used as template. Partial small subunit (SSU) ribosomal RNA gene fragments
176 were amplified in volumes of 25 µl with 0.125 U of GoTaq Flexi DNA Polymerase (Promega, Milan, Italy), 0.4 µM of
177 each primer (NS31 /AML2, Simon et al., 1992; Lee et al., 2008), 0.2 mM of each dNTP, 1.5 mM of MgCl₂ and 1×
178 manufacturer's reaction buffer. The thermal cycler (Eppendorf Mastercycler personal, Eppendorf, Milan, Italy) was
179 programmed as follows: a manual “hot start” at 94 °C for 3 min, 30 cycles at 94 °C for 30 sec, 58 °C for 40 sec, 72 °C
180 for 55 sec and a final extension step at 72 °C for 10 min. Reactions yields were estimated using a 1 % (w/v) agarose I
181 (Euroclone, Milan, Italy) in TBE 1X buffer (Euroclone) gels stained with ethidium bromide (0.5 µg ml⁻¹).

182 NS31/AML2 amplicons from apple root samples were purified using Wizard® SV Gel and PCR Clean-up
183 system (Promega), then ligated into pGem-T Easy vector (Promega) to transform XL10-Gold Ultracompetent
184 *Escherichia coli* cells (Stratagene, La Jolla, CA, USA).

185 The structure and composition of the AM fungal communities were determined using PCR-RFLP screening of
186 clone libraries. Putative positive clones were amplified by using standard SP6/T7 amplifications, followed by a nested
187 PCR using NS31/AML2 primers as described above. Twenty-five amplicons per clone library were digested by HinfI
188 and Hsp92II restriction enzymes (Promega) and run on 2 % agarose at constant 50 V for 2 h. A 100 bp DNA ladder
189 (Promega) was used as a molecular weight marker. DNA profiles were visualized under UV illumination and captured
190 as TIFF format file by Liscap program for Image Master VDS System (Pharmacia Biotech).

191 A total of 448 clones were examined. Recombinant plasmids of representative clones of each RFLP pattern
192 were purified by Wizard® Plus SV Minipreps (Promega) and sequenced using T7 vector primers at BMR Genomics
193 s.r.l. (University of Padova, Italy).

194 Sixty-nine unique sequences of the clones generated in this study have been deposited in EMBL Nucleotide
195 Sequence Database (www.ebi.ac.uk/embl/) under the accession numbers from LT600783 to LT600851.

196

197 2.6.3. PCR-DGGE analysis

198 To analyse AMF communities by PCR-DGGE, a semi-nested PCR approach was used. Initially a 550 bp fragment of
199 the 18S rRNA gene was amplified using the primer NS31 (Simon et al., 1992) in combination with the primer AM1
200 (Helgason et al., 1998). Amplification reactions were performed in a final volume of 50 µl, using 10-20 ng of DNA, 5
201 µl of 10X *Ex Taq* Buffer (Takara Biotechnology, Milan, Italy), 1.25U of *TaKaRa Ex Taq* (Takara Biotechnology), 0.2
202 mM of each dNTP (Takara Biotechnology), 0.5 µM of each primer (Eurofins genomics, Milan, Italy) and an aliquot of
203 0.3 µg µl⁻¹ acetylated bovine serum albumin (BSA, Promega). PCR amplifications were carried out using an iCycler-iQ
204 Multicolor Real-Time PCR Detection System (Bio-Rad, Milan, Italy) with the following conditions: 94 °C initial
205 denaturation for 1 min; 35 amplification cycles of 30 s at 94 °C, 30 s at 66 °C, 30 s at 72 °C; final extension at 72 °C for
206 5 min. The presence of amplicons was confirmed by electrophoresis in 1.5 % (w/v) agarose gels in TBE 1X buffer
207 (Tris-borate-EDTA, pH 8.0), stained with 0.5 µg mL⁻¹ ethidium bromide.

208 Amplification products from the first PCR reaction were then diluted 1:10 and 1 µl was used as template in a
209 second PCR using the NS31 and the Glo1 (Cornejo et al., 2004) primers. A GC clamp (5'-
210 CGCCCGGGCGCGCCCCGGGCGGGGCGGGGGCACGGGGG -3') was added to the 5' end of the forward primer
211 NS31. PCR amplifications were performed as previously described except for the addition of BSA and the annealing
212 temperature of 52 °C.

213 For the DGGE analysis, 20 µl of the PCR products plus 20 µl of buffer 2X made with 70 % glycerol, 0.05 %
214 xylene cyanol and 0.05 % bromophenol blue, were separated in a 8 % polyacrylamide-bisacrilamide (37.5:1) gel with a
215 35-56 % urea-formamide gradient, using the DCode™ Universal Mutation Detection System (Bio-Rad). A composite

216 mix of AMF 18S rRNA gene fragments from *Funneliformis mossae* AZ225C, *Funneliformis coronatum* IMA3 and
217 *Rhizoglyphus intraradices* IMA5 was added in the middle and at both ends of each gel as DGGE marker (M). Gels were
218 run and profiles were visualized as described in Agnolucci et al. (2013).

219 DGGE bands were excised from polyacrylamide gels and DNA was extracted by eluting in 50 μ l ddH₂O at
220 4°C overnight. One μ L of the supernatant diluted 1:100 was used to re-amplify the 230 bp DNA fragment of the 18S
221 rRNA gene according to the PCR protocol described above, except that the primer NS31 was used without GC clamp.
222 PCR products were then purified by UltraClean PCR CleanUp Kit (MO-BIO Laboratories) according to the
223 manufacturers' protocol, quantified and 5' sequenced at the BMR Genomics (Padova, Italy).

224

225 **2.7. Bioinformatics**

226 Sequences from *E. coli* libraries and DGGE bands were edited in MEGA 6.0 and their similarities were determined
227 using the Basic Local Alignment Search Tool (BLASTn) provided by NCBI. The detection of chimeric sequences was
228 performed using USEARCH 6.0 (http://fungene.cme.msu.edu/FunGenePipeline/chimera_check/form.spr). Sequences
229 were aligned with those corresponding to the closest matches from GenBank as well as with sequences from major
230 clades of Glomeromycota using MUSCLE as implemented in MEGA6. Phylogenetic trees were inferred by Neighbour-
231 joining analysis. The evolutionary distances were computed using the Maximum Composite Likelihood method. The
232 confidence of branching was assessed using 1000 bootstrap resamplings.

233 Richness and composition of AMF communities obtained from clone libraries were evaluated for each
234 treatment. We determined the rarefaction curves with Past software to estimate whether the number of screened
235 sequenced were sufficient to capture AMF diversity of each host. Estimates of community diversity were determined as
236 Richness (S), bias-corrected Chao1 richness, Shannon-Weaver (Hs) and dominance index of Simpson (D). The indices
237 were calculated using PAST 3.0 and 1000 bootstraps were used to determine confidence intervals. Non parametric
238 Kruskal Wallis test was used to determine differences in the diversity indices among AMF communities in the roots of
239 apple plants cultivated in the two sites (Terlano vs Lagundo), under different managements (organic vs integrated) and
240 at different altitudes (low and high). AMF communities were also evaluated by permutational multivariate analysis of
241 variance (PERMANOVA), to test the effect of the management (organic vs integrated), site and altitude. Non
242 parametric tests and PERMANOVA were performed in SPSS version 20 software (IBM Corp., Armon, NY Inc, USA)
243 and PAST 3.0, respectively.

244 DGGE profiles were digitally processed with BioNumerics software version 7.5 (Applied Maths, St-Martens-
245 Latem, Belgium) following the manufacturer's instructions. The lanes were normalized to contain the same amount of
246 total signal after background subtraction and the gel images were straightened and aligned to give a densitometric

247 curve. DGGE markers were used for digital gel normalization to allow comparison between gels. Bands were assigned
248 using the auto search bands option and then checked manually and their positions were converted to Rf % values. All
249 profiles were compared using the band matching tool with a position tolerance and optimization of 0.5 and 0 %
250 respectively.

251 Similarities between DGGE patterns were calculated by determining Dice's similarity coefficients for the total
252 number of lane patterns from the DGGE gel. The similarity coefficients were then used to generate the dendrogram
253 utilizing the clustering method UPGMA (Unweighted Pair Group Method Using Arithmetic Average). DGGE profiles
254 were also analysed using non-metric multidimensional scaling analysis (NMDS) performed from a data matrix based on
255 presence/absence of bands (Bray-Curtis coefficient). The significance of data was assessed by the two way ANOSIM
256 method (analysis of similarities; 999 permutations) (PAST 3.0) with geographical site (Terlano vs Lagundo) and
257 management (organic vs integrated) as variability factors. Note that the stability index R describes the extent of
258 similarity between each pair in the ANOSIM, with values close to unity indicating that the two groups are entirely
259 separate and a value of zero indicating that there is no difference between the groups.

260 DGGE banding data were used to estimate four different indices treating each band as an individual
261 operational taxonomic unit (OTU). Richness (S) indicates the number of OTUs present in a sample and was determined
262 from the number of fragments. The overall diversity index of Shannon-Weaver (Hs) and the dominance index of
263 Simpson (D) were calculated using the equations $H_s = -\sum(P_i \times \ln P_i)$ and $D = \sum P_i^2$ respectively, where the relative
264 importance of each OTU is $P_i = n_i N^{-1}$, and n_i is the peak intensity of a band and N is the sum of all peak intensities in a
265 lane. Evenness index (E), which allows the identification of dominant OTUs, was calculated as $E = H(\ln S)^{-1}$. Two-way
266 ANOVA was applied to diversity indices with site (Terlano vs Lagundo) and management (organic vs integrated) as
267 variability factors, after checking the normal distribution of data. One-way ANOVA was used for Lagundo diversity
268 indices with management as the variability factor. The means were compared by the Tukey's test ($P < 0.05$). Analyses
269 were carried out with the SPSS version 20 software.

270

271 **3. Results**

272 **3.1. Soil fertility and mycorrhizal colonization**

273 Soils texture in the Lagundo area had either a silty loam or a sandy loam texture, while those in the Terlano area were
274 either loamy or sandy loam. In general, as indicated by the standard error values (Table 2), there was a relatively low
275 variability of N concentration among soils belonging to different treatments as well as within each treatment. All soils
276 were well endowed with P and Mg, while soil K availability ranged from moderate-low to normal values. In Lagundo
277 only, soils managed according to organic guidelines had higher pH ($P=0.02$), organic matter (OM) ($P=0.01$) and C

278 concentration (P=0.03) than those managed according to the integrated guidelines (Table 2). Soils at higher elevation
279 also had higher soil OM (P=0.05) and soil C (P=0.06) than those at lower elevation (Table 2). When only data
280 belonging to low elevation were analyzed, no significant differences in soil characteristics between the two orchard
281 managements were detected.

282 No significant effect of altitude and management was recorded on leaf nutrient concentration (Table 3). Leaf P
283 and K were significantly higher (P=0.01 and 0.001, respectively) and leaf Mg significantly lower (P=0.02) in orchards
284 located in Terlano than in Lagundo (Table 3).

285 In all sampled orchards, apple roots were well colonized by AMF (Table 3). The percentage of colonized root
286 length, ranging from 41 to 60 %, was not significantly affected by site, management and altitude. We did not find
287 significant effects of treatments on the degree of mycorrhizal colonization variation among and within orchards (Table
288 S1). The level of mycorrhizal colonization was not significantly correlated with most measured soil parameters, but a
289 negative linear correlation with soil organic matter ($r=-0.49$, $P=0.03$) was found.

290

291 **3.2. Identification of native AMF colonizing apple roots**

292 The DNA extracted from apple roots was successfully amplified using the primer pair NS31/AML2, obtaining a
293 fragment of the expected size (~550 bp). A total of 448 clones from the 21 clone libraries were screened by PCR-RFLP
294 analysis, obtaining 19 different RFLP patterns. For each RFLP group several clones originating from different libraries
295 were sequenced, giving 17 Glomeromycota sequences. All non-redundant sequences from the 21 clone libraries (69 out
296 of 212) and 27 references from GenBank were used for neighbour-joining phylogenetic analyses (Fig. 2). After RFLPs,
297 BLASTn and phylogenetic analyses the sequences were grouped into 17 OTUs supported by a bootstrap value >87 %.

298 Among the 17 OTUs, we retrieved sequences belonging to 8 out of the 19 Glomeromycota genera (Redecker et
299 al., 2013). In particular all the genera of Glomeraceae were found (*Glomus*, *Septoglomus*, *Rhizoglomus*, *Sclerocystis*,
300 *Funneliformis*) together with the genera *Claroideoglomus*, *Diversispora* and *Paraglomus*, belonging to
301 Claroideoglomeraceae, Diversisporaceae and Paraglomeraceae, respectively. The most abundant genera were
302 represented by *Glomus*, accounting for 29.7 % of the sequences, followed by *Paraglomus* (19.4 %), *Claroideoglomus*
303 (17.2 %), *Sclerocystis* (16.1 %) and *Rhizoglomus* (12.3 %). *Septoglomus*, *Diversispora* and *Funneliformis* sequences
304 corresponded to less than 4 % of total sequences. Twelve OTUs were ascribed to Glomeraceae family: within the genus
305 *Glomus* we retrieved seven OTUs, of which only one (Glo7, 5.8 % of total sequences) was identified as *Glomus*
306 *indicum*; the remaining six OTUs (Glo1 to Glo6) represented sequences of uncultured *Glomus* species (Table 4). In the
307 genera *Sclerocystis*, *Funneliformis* and *Rhizoglomus*, OTUs named Scl, Fun and Rhi were identified as *Sclerocystis*
308 *sinuosa*, *Funneliformis mosseae* and *Rhizoglomus irregulare* (synonym *Rhizophagus irregularis*, formerly known as

309 *Glomus irregulare*), respectively. OTUs Sept1 and Sept2 and Cl1, Cl2 and Cl3 assigned to *Septoglomus* and
310 *Claroideoglomus*, respectively, matched to sequences of either known (Sept2, *Septoglomus constrictus* and Cl1,
311 *Claroideoglomus lamellosum*), or unknown species (*Septoglomus* sp., *Claroideoglomus* sp.), while OTUs belonging to
312 *Paraglomus* (Par) and *Diversispora* (Div) matched only unknown Glomeromycota. MaarjAM database
313 (<http://maarjam.botany.ut.ee/>, accessed on March/2016) was used to confirm the assignment of our OTUs to sequences
314 of Glomeromycota (Table 4). Rarefaction analyses indicated that the number of analysed sequences was generally
315 sufficient to capture the AMF diversity in the roots of most orchards, since the curves almost reached the asymptote
316 (Fig. S1).

317 The bands of interest from the DGGE profiles, excised and sequenced to determine their affiliation, identified
318 the same AMF species as those described above, except Claroideoglomeraceae and Paraglomeraceae (data not shown).

319

320 **3.3. AMF community diversity in apple roots as affected by site, management and altitude**

321 **3.3.1. Analysis of clone libraries**

322 The cloning and sequencing method allowed the identification of 14 and 15 OTUs in Terlano and Lagundo orchards,
323 respectively, 12 of which shared between the two sites (97.5 and 88.8 % of the total sequences of the two sites,
324 respectively). Fun (*F. mosseae*) and Glo6 were found only in Terlano apple roots, while Sept2 (*S. constrictus*), Glo2
325 and Cl3 were retrieved only in Lagundo. Two ways PERMANOVA analyses revealed differences in AMF community
326 composition between the two sites (P=0.02) and between organic and integrated managements (P=0.02). Moreover, in
327 Lagundo, where orchards were located at two different altitudes (300 m and 600 m) two ways PERMANOVA analysis
328 showed differences among AMF communities in relation to management (P=0.003) and altitude (P=0.003). In total,
329 orchards under organic management hosted 13 and 6 OTUs at low and high altitude, respectively. In organic orchards
330 growing at higher elevation the most common species (Par, Glo7, Rhi, Scle) present at both sites and under different
331 managements were detected together with Cl1 and Div (species found in all organic orchards at low altitude) (Fig. 3).
332 By contrast, in integrated orchards 10 and 9 OTUs in total were found at low and high altitude, respectively. At high
333 altitude, together with the most common species, a particular OTU (Cl3, 31 % of the sequences) was retrieved (Fig. 3).

334 Among diversity indices, S and Chao-1, indicating observed and estimated AMF richness, were significantly
335 higher in organic orchards (6.33 ± 0.55 and 7.14 ± 0.88 , respectively) than in integrated ones (4.00 ± 0.01 and $4.00 \pm$
336 0.01 , respectively) in Terlano (P=0.027 and P=0.034, respectively). In Terlano 13 OTUs were retrieved in organic
337 management, in contrast with only 6 OTUs found in integrated ones. No differences among diversity indices were
338 detected among orchards in Lagundo.

339

340 3.3.2. Analysis of PCR-DGGE profiles

341 AMF 18S rDNA fragments of approx. 230 bp were successfully amplified from all the samples, by the semi-nested
342 PCR approach. DGGE analyses of PCR products showed profiles characterized by a high number of intense and clearly
343 defined fragments.

344 AMF community composition was assessed by cluster analysis of DGGE profiles (Fig. 4). The dendrogram
345 showed two main subclusters with a low similarity, 20 %. In particular, the first subcluster included all Terlano samples
346 (except one), while the other one included all Lagundo samples (except one). The latter was formed by two subclusters
347 in which organic and integrated samples were separated with a similarity of 36 %. The analysis of DGGE profiles did
348 not separate samples originating from orchards growing at different altitudes (Fig. 4). Terlano samples showed a more
349 complex clustering, with no clearcut separation by management.

350 These findings were confirmed by the NMDS analysis of the DGGE AMF community profiles (Fig. 5).
351 ANOSIM revealed significant differences in AMF community composition of Lagundo samples compared with that of
352 Terlano ($R=0.786$, $P=0.0001$) and between samples from organic and integrated sites ($R=0.227$, $P=0.05$).

353 AMF DGGE profiles were also analysed to assess S, Hs, D and E diversity indices. Richness (S) and Hs
354 indices were significantly different between the two geographical sites, while a significant interaction was observed for
355 Hs and D (Table 5). The relevant one-way ANOVA revealed significant differences between organic and integrated
356 samples in Lagundo, where S and Hs were significantly higher in the organic management ($P=0.049$ and $P=0.029$
357 respectively) (Table 6).

358

359 4. Discussion

360 Our data showed that, in South Tyrol orchards, geographical area, altitude and management shaped AMF species
361 community composition in apple roots, which however always maintained high levels of mycorrhizal colonization.
362 Here, for the first time, we identified at the species and genus level native AMF colonizing apple roots and
363 characterized their diversity and community composition utilizing two molecular methods - PCR cloning and
364 sequencing and PCR-DGGE.

365

366 4.1. Mycorrhizal colonization of apple roots

367 The percentage of colonized root length of apple trees ranged from 41 to 60 %, comparing well with previous data on
368 mycorrhizal colonization of apple seedlings, that ranged from 40 to 60 % after inoculation with *F. mosseae* and *Glomus*
369 *macrocarpum*, respectively (Miller et al., 1989) and with findings obtained by Meyer et al. (2015) in conventionally and
370 organically-managed apple orchards in South Africa. Other authors reported more variable values, ranging from 24 to

371 68 % in micropropagated apple trees, depending on soil pH and the identity of the inoculated AMF species (Cavallazzi
372 et al., 2007). In our work, mycorrhizal colonization of apple roots was uncoupled from soil parameters, except for
373 organic matter, confirming previous findings on the decrease of mycorrhizal colonization levels in an organic apple
374 orchard after a straw mulch floor management treatment (Meyer et al., 2015).

375 Overall, mycorrhizal root length was not significantly affected by site, altitude and management in the apple
376 orchards investigated. As to site, Miller et al. (1985) in a survey of 18 apple rootstock plantings in USA found a high
377 variability in mycorrhizal colonization, ranging from 5 to 75 %, with the lowest levels observed in California and
378 Washington and the highest in Virginia, Pennsylvania, Georgia and Iowa. Such data are expected, given the different
379 range of geographical distances of the apple orchards analysed, entailing the most divergent environmental conditions,
380 at the subcontinental level.

381 Most studies investigating the effects of altitude on mycorrhizal colonization have been carried out at very high
382 elevation gradients (from 1,500 to 5,300 m), in alpine habitats of Tibet, Andes, Rocky Mountains, Alps and Mount Fuji
383 (Read and Haselwandter, 1981; Wu et al., 2007; Lugo et al., 2008; Schmidt et al., 2008; Gai et al., 2012). Overall, these
384 works reported an adverse effect of increasing altitude on AMF colonization across a large altitude gradient. In our
385 study, we assessed the effects of elevation in a single site, Lagundo, from apple orchards cultivated at 300 and 600 m,
386 an altitudinal range probably too low to produce significant effects on AMF ability to colonize apple roots, which,
387 indeed, showed high percentages of mycorrhizal root length, 35-52 %.

388 The type of farming practices has long been known to affect mycorrhizal colonization in different crops, such
389 as wheat (Ryan et al., 1994; Mäder et al., 2000), vetch-rye and grass-clover (Mäder et al., 2000), onion (Galván et al.,
390 2009), maize (Douds et al., 1993; Bedini et al., 2013), soybean (Douds et al., 1993). Very few studies investigated the
391 effects of organic vs. conventional management regimes on the establishment of mycorrhizal symbiosis in apple roots.
392 The absence of differences in the percentage of mycorrhizal root length in our experimental apple orchards may be
393 ascribed to the uniformity of the orchard floor managements, which included the presence of grassed alleys between
394 rows. AMF-host weed species may have contributed to the maintenance of wide and infective extraradical mycelial
395 networks establishing linkages among the roots of cover plants and apple trees (Atkinson, 1983; Giovannetti et al.,
396 2004; Njeru et al., 2014). The application of glyphosate in the small soil strip along the tree rows and that of mineral
397 fertilizer supply in the integrated orchards did not have a significant impact on the ability of AMF to colonize apple
398 roots. This finding is consistent with previous studies reporting variable and unpredictable effects of such chemicals on
399 AMF symbioses in different environments and geographical locations (Gosling et al., 2006).

400

401 **4.2. Identification of native AMF colonizing apple roots**

402 The screening of 448 clones from 21 clone libraries allowed us to affiliate DNA sequences with 8 out of the 19
403 Glomeromycota genera, which encompassed all the genera of the family Glomeraceae described so far: *Glomus*,
404 *Septoglomus*, *Rhizophagus*, *Sclerocystis*, *Funneliformis*. In addition, members of the families Claroideoglomeraceae,
405 Diversisporaceae and Paraglomeraceae were found.

406 Overall, the most common species belonged to the genus *Glomus*, in agreement with previous findings
407 obtained by morphological description of AMF spores in a survey of 18 apple rootstock plantings in USA (Miller et al.,
408 1985). Some of the species retrieved in the quoted study occurred also in our apple roots, such as *F. mosseae* and *S.*
409 *constrictus* (formerly *Glomus mosseae* and *Glomus constrictum*, respectively) and *S. sinuosa*, while no species of the
410 genus *Gigaspora*, common in the USA apple plantings, were found. Our data differ from those obtained in a survey of
411 apple orchards in Santa Catarina, south region of Brazil, where 15 species of the genus *Acaulospora* were retrieved and
412 described after spore sievings from the soil, using morphological methods (Purin et al., 2006). Only few AMF species
413 were in common, belonging to *F. mosseae*, *Claroideoglomus* spp. and *S. sinuosa* (formerly *Glomus sinuosum*). Another
414 interesting work taxonomically characterized AMF species by morphological identification of spores retrieved from soil
415 sieving and trap cultures in Brazil (Cavallazzi et al., 2007): the species with the highest number of spores was
416 *Acaulospora mellea*, followed by *Scutellospora heterogama*, *Gigaspora decipiens* and *Acaulospora spinosa*, none of
417 which occurred in apple roots of South Tyrol orchards. A molecular study, utilising 454-pyrosequencing of small
418 subunit rRNA gene amplicons, identified, at the family level, the AMF colonizing roots of cultivated apple in central
419 Belgium (Van Geel et al., 2015). In the work, 73 % and 19 % of OTUs were affiliated with the Glomeraceae and
420 Claroideoglomeraceae, in agreement with our findings, 70 % and 18 %, respectively. In addition, the authors detected
421 only a few OTUs belonging to the families Gigasporaceae, Diversisporaceae and Acaulosporaceae, while the
422 percentage of OTUs affiliated with Paraglomeraceae was much lower in Belgian apple orchards, compared with South
423 Tyrol ones, 1 % vs. 19 %. A recent work, carried out in an experimental apple orchard near Sint-Truiden, Belgium,
424 confirmed the previous data, with 72 % of OTUs belonging to Glomeraceae, 26 % to Claroideoglomeraceae and 2 % to
425 Paraglomeraceae (Van Geel et al., 2016).

426 The absence or rarity of specific taxa, such as Acaulosporaceae and Gigasporaceae, from the three European
427 apple orchards investigated so far, compared with the frequency of their retrieval from USA and Brazil, may be
428 ascribed to a number of complex and interacting factors - soil, environment, climate, agronomic practices, history of the
429 sites etc. - beyond geographical position. Although the number of experimental data on apple from the two continents is
430 still too small to deduce general trends, recent studies at the global scale revealed that differences in AMF communities
431 diversity do occur among different continents and climatic zones (Kivlin et al., 2011; Öpik et al., 2013; Davison et al.,
432 2015). It is interesting to note that a survey of AMF species distributed all over the world in protected areas reported a

433 general predominance of Acaulosporaceae and Gigasporaceae in tropical or subtropical forests in Brazil and Argentina
434 (Turrini and Giovannetti, 2012).

435 In our study we found sequences (uncultured *Clareidoglomus*-Cl2, *Paraglomus*-Par and *Glomus*-Glo1
436 sequences, corresponding to 18 % of total OTUs), matching those retrieved in apple roots from German orchards
437 (JN644447, unpublished), in roots of other fruit tree plants, such as *Pyrus pyrifolia* (AB695049) (Yoshimura et al.,
438 2013) and *Citrus* rootstocks (JQ350797) (Wang and Wang, 2014). It is tempting to speculate that fruit trees may show a
439 preference for specific fungi, possibly recruiting AMF species in relation to their functional significance.

440

441 **4.3. AMF community diversity in apple roots as affected by site, altitude and management**

442 Present results show that root AMF community composition of apple trees cultivated in two production sites, Terlano
443 and Lagundo, differed significantly and were affected by agricultural management, organic vs. integrated, and altitude.
444 Such findings are supported by consistent data obtained by combining two molecular methods, cloning and sequencing
445 and PCR-DGGE, and utilizing two different primer pairs to amplify the same region of the partial 18S rRNA gene. To
446 the best of our knowledge, this is the first application of such a comprehensive approach for the characterization of root
447 AMF diversity of fruit trees, allowing the differentiation of AMF communities colonizing apple roots in different
448 agricultural conditions.

449 The differences detected in root AMF community composition between the two apple production sites, as
450 revealed by cloning and sequencing analysis, were ascribed to specific taxa occurring only in Terlano (Fun and Glo6) or
451 Lagundo (Sept2, Glo2 and Cl3), to their relative abundance and to the different distribution of shared OTUs between
452 the two sites. Such site differences were consistently detected by PCR-DGGE community profiles analysis, which
453 separated the two relevant clusters with a very high dissimilarity (80 %), and confirmed by NMDS analysis. Our data
454 are in agreement with a previous work, carried out at the regional scale, investigating the distribution of 6 AMF species
455 in 154 agricultural soils across Switzerland, which showed that AMF communities were strongly affected by
456 geographical distance (max. 294 km) (Jansa et al., 2014). Consistent findings were also reported by van der Gast et al.
457 (2011), who investigated AMF community diversity across England (max. 250 km). Although the maximum distances
458 among the sampled apple orchards were much lower than those described above (max. 27 km), our work identified
459 different AMF species and genera in the two sites, Terlano and Lagundo. Such distribution patterns may be the result of
460 biogeographic history, reflecting the dispersal of AMF taxa over time, and the variable climatic and environmental
461 conditions (Morton et al., 1995). Recent molecular works (Hazard et al., 2013; De Beenhouwer et al., 2015; Van Geel et
462 al., 2015) suggested a key role of soil environment in shaping AMF diversity and distribution patterns. Further in-depth
463 and comprehensive studies should be performed in order to separate geographical distance effects from those due to the

464 characteristics of each sampling site, i.e. soil properties, use of pesticides and chemical fertilizers, environs, even when
465 comparing sites at the regional and local scale. In addition, beyond geographical, soil and environmental variables, host
466 plant identity should be taken into account, given its essential role played in the selection of AMF symbionts
467 functionally established in the roots (Helgason et al., 2002; Gollotte et al., 2004; Sýkorová et al., 2007). Actually, at
468 local scale, a marked host preference has been recently found in maize plants, which hosted completely different AMF
469 communities, as compared with those of the preceding cover crops (Turrini et al., 2016).

470 No differences in AMF community composition were found by PCR-DGGE cluster analysis between orchards
471 located at 300 m and 600 m altitude in Lagundo, in contrast with cloning and sequencing, that allowed the detection of
472 a particular OTU (Cl3-*Claroideoglossum* sp., 31 % of the sequences) only in the 600 m high orchards. Such divergent
473 data may be ascribed to the different primers utilized during DNA amplification by the two diverse molecular
474 techniques: indeed, Glo1 primer, used in PCR-DGGE, did not allow the amplification of Claroideoglomeraceae, as
475 shown by band sequencing. The OTU Cl3 is infrequent, probably fitting better to the colder environment and to the
476 higher levels of organic matter and soil C found in the 600 m Lagundo orchards. Indeed, the occurrence of infrequent
477 AMF species at high altitude was previously reported (Liu et al., 2011; Sýkorová et al., 2007), while several new
478 species have been described in soils from Swiss Alps (Oehl and Sieverding, 2004; Oehl et al., 2005a, 2006, 2011).

479 In this work, the relative abundance of single AMF species colonizing apple roots showed significant
480 differences between organic and integrated management, as assessed by PERMANOVA, both in Terlano and in
481 Lagundo orchards. Differences were found also by the analysis of PCR-DGGE AMF community profiles, carried out by
482 NMDS and ANOSIM. Diversity indices showed higher AMF richness in organically managed apples compared with
483 integrated ones, using both molecular approaches. Although scarce information is available on AMF diversity in apple
484 orchards, as affected by management, our findings compare well with those obtained by Purin et al. (2006), who, by a
485 morphological approach, found a higher AMF richness in organic compared with conventional orchards in Brazil. A
486 few other studies, performed on plant species other than apple, reported the effects of organic farming on the
487 composition of AMF communities. For example, Oehl et al. (2004), in the DOC field experiment in Switzerland found
488 that organic management enhanced AMF spore diversity and abundance, while Bedini et al. (2013) showed a
489 progressive increase of AMF richness and composition during the transition from conventional to organic agriculture.
490 Other authors, using molecular techniques, showed that the diversity of AMF species composition in maize and potato
491 roots was higher in organic sites, compared with conventional ones, in agricultural fields throughout the Netherlands
492 (Verbruggen et al., 2010).

493 The experimental data collected so far on the influence of organic and conventional managements on AMF
494 diversity, either in the soil or in the roots, using either morphological or molecular approaches, are limited and do not

495 allow us to infer consistent response patterns. Indeed, the words “organic” and “conventional” may have different
496 meanings in the different experimental studies, as the diverse farming systems often encompass heterogeneous local soil
497 properties, environmental conditions and host plants. Thus, AMF diversity or community composition may be affected
498 by a number of uncontrolled variables, such as the use of weed cover, soil tillage, cover crops, manure, quality and
499 quantity of herbicides, pesticides and fertilizers. Whereas floor management in the orchard alleys was similar in organic
500 and integrated orchards, the two management systems differed mainly by the weed control and fertilizer supply to the
501 soil in the strip centres on the tree row: we speculate that the use of glyphosate and mineral fertilizers in integrated
502 managed orchards may have selectively modulated the abundance and composition of AMF taxa able to tolerate such
503 chemicals.

504

505 **5. Conclusions**

506 In this work, we utilized a multimodal approach to study AMF communities living in symbiosis with apple roots in
507 South Tyrol orchards, under different geographical and environmental conditions. High levels of mycorrhizal
508 colonization were detected across the different variables, while AMF diversity and community composition were
509 affected by geographical area, altitude and farming system management, as detected by PCR cloning and sequencing
510 and PCR-DGGE. In particular, species richness was significantly higher in organically managed orchards than in
511 integrated ones. We identified, for the first time, the native AMF communities of apple roots at the species and genus
512 level, detected infrequent taxa and retrieved some environmental sequences matching those obtained from other fruit
513 plant species. Our findings provide insights into factors affecting native AMF communities of apple trees, which could
514 be exploited to implement sustainable fruit production systems, where beneficial soil biota can boost biogeochemical
515 processes fundamental for energy fluxes, ecosystem functioning and crop productivity.

516

517 **Acknowledgments**

518 This work was supported by the University of Pisa through the project “Molecular and functional biodiversity of plant
519 associated microorganisms” (M.G.) and through Fondi di Ateneo (M.A. and A.T.), and by the Free University of
520 Bolzano-Bozen through the project “Beneficial effects of mycorrhizal fungi to the mineral nutrition and to carbon
521 sequestration in sustainable fruit production systems” (TN5051), granted to F.S. The work represents part of E.T.’s PhD
522 thesis project at UNIBZ and of M.P.’s research fellowship at UNIPI. The authors wish to thank Dr. Wolfgang Drahorad
523 and the owners of the sampling orchards for their precious help in carrying out the field experiment.

524

525 **Authorship and contributorship** M.G., F.S. and M.T. conceived and designed the experiments. F.S. and E.T. carried
526 out mycorrhizal colonization analyses and determined plant and soil parameters, M.A., M.P. and A.T. performed
527 molecular and data analyses. M.A., M.G. and A.T. wrote the paper. F.S., M.T. and E.T. provided agronomic expertise
528 and apple root samples. M.A., M.G., F.S., M.T. and A.T. contributed reagents/ materials/analysis tools.

529

530 **References**

531 Agnolucci, M., Battini, F., Cristani, C., Giovannetti, M., 2015. Diverse bacterial communities are recruited on spores of
532 different arbuscular mycorrhizal fungal isolates. *Biol. Fertil. Soils* 51, 379-389.

533 Agnolucci, M., Cristani, C., Battini, F., Palla, M., Cardelli, R., Saviozzi, A., Nuti, M., 2013. Microbially-enhanced
534 composting of olive mill solid waste (wet husk): bacterial and fungal community dynamics at industrial pilot and
535 farm level. *Bioresour. Technol.* 134, 10-16.

536 Atkinson, D., 1983. The growth activity and distribution of the fruit tree root system. *Plant Soil* 71, 23-35.

537 Avio, L., Castaldini, M., Fabiani, A., Bedini, S., Sbrana, C., Turrini, A., Giovannetti, M., 2013. Impact of nitrogen
538 fertilization and soil tillage on arbuscular mycorrhizal fungal communities in a Mediterranean agroecosystem.
539 *Soil Biol. Biochem.* 67, 285-294.

540 Barea, J.M., Azcón, R., Azcón-Aguilar, C., 2002. Mycorrhizosphere interactions to improve plant fitness and soil
541 quality. *A. Van Leeuw. J. Microb.* 81, 343-351.

542 Battini, F., Bernardi, R., Turrini, A., Agnolucci, M., Giovannetti, M., 2016a. *Rhizophagus intraradices* or its associated
543 bacteria affect gene expression of key enzymes involved in the rosmarinic acid biosynthetic pathway of basil.
544 *Mycorrhiza* 26, 699-707.

545 Battini, F., Turrini, A., Quartacci, M.F., Malorgio, F., Sgherri, C., Picciarelli, P., Pardossi, A., Giovannetti, M.,
546 Agnolucci, M., 2016b. Dual inoculation with AMF and associated bacteria improves nutraceutical value of sweet
547 basil grown under commercial conditions. *Agrochimica* 60, 81-99.

548 Bedini, S., Avio, L., Sbrana, C., Turrini, A., Migliorini, P., Vazzana, C., Giovannetti, M., 2013. Mycorrhizal activity
549 and diversity in a long-term organic Mediterranean agroecosystem. *Biol. Fertil. Soils* 49, 81-790.

550 Brito, I., Goss, M.J., De Carvalho, M., Chatagnier, O., van Tuinen, D., 2012. Impact of tillage system on arbuscular
551 mycorrhiza fungal communities in the soil under Mediterranean conditions. *Soil Tillage Res.* 121, 63-67.

552 Cavallazzi, J.R.P., Filho, O.K., Stürmer, S.L., Rygielwicz, P., Mendonça, M.M., 2007. Screening and selecting
553 arbuscular mycorrhizal fungi for inoculating micropropagated apple rootstocks in acid soils. *Plant Cell Tissue*
554 *Organ Cult.* 90, 117-129.

555 Cornejo, P., Azcón-Aguilar, C., Barea, J.M., Ferrol, N., 2004. Temporal temperature gradient gel electrophoresis
556 (TTGE) as a tool for the characterization of arbuscular mycorrhizal fungi. *FEMS Microbiol. Lett.* 241, 265-270.

557 Creamer, R.E., Hannula, S.E., Van Leeuwen, J.P., Stone, D., Rutgers, M., Schmelz, R.M., De Ruiter, P.C., Hendriksen,
558 N.B., Bolger, T., Bouffaud, M.L., Buee, M., 2016. Ecological network analysis reveals the inter-connection
559 between soil biodiversity and ecosystem function as affected by land use across Europe. *Appl. Soil Ecol.* 97,
560 112-124.

561 Dalla Via, J., Mantinger, H., 2012. Agricultural research in the field of fruit growing in South Tyrol. *Erwerbs-Obstbau*
562 54, 83-115.

563 Davison, J., Moora, M., Öpik, M., Adholeya, A., Ainsaar, L., Bâ, A., Burla, S., Diedhiou, A.G., Hiiesalu, I., Jairus, T.,
564 Johnson, N.C., Kane, A., Koorem, K., Kochar, M., Ndiaye, C., Pärtel, M., Reier, Ü., Saks, Ü., Singh, R., Vasar,
565 M., Zobel, M., 2015. Global assessment of arbuscular mycorrhizal fungus diversity reveals very low endemism.
566 *Science* 349, 970-973.

567 De Beenhouwer, M., Van Geel, M., Ceulemans, T., Muleta, D., Lievens, B., Honnay, O., 2015. Changing soil
568 characteristics alter the arbuscular mycorrhizal fungi communities of Arabica coffee (*Coffea arabica*) in
569 Ethiopia across a management intensity gradient. *Soil Biol. Biochem.* 91, 133-139.

570 Douds, D.D. Jr., Janke, R.R., Peters, S.E., 1993. VAM fungus spore populations and colonization of roots of maize and
571 soybean under conventional and low-input sustainable agriculture. *Agric. Ecosyst. Environ.* 43, 325-335.

572 Douds, D.D. Jr., Millner, P.D., 1999. Biodiversity of arbuscular mycorrhizal fungi in agroecosystems. *Agric. Ecosyst.*
573 *Environ.* 74, 77-93.

574 Gai, J.P., Tian, H., Yang, F.Y., Christie, P., Li, X.L., Klironomos, J.N., 2012. Arbuscular mycorrhizal fungal diversity
575 along a Tibetan elevation gradient. *Pedobiol.* 55, 145-151.

576 Galván, G.A., Parádi, I., Burger, K., Baar, J., Kuyper, T.W., Scholten, O.E., Kik, C., 2009. Molecular diversity of
577 arbuscular mycorrhizal fungi in onion roots from organic and conventional farming systems in the Netherlands.
578 *Mycorrhiza* 19, 317-328.

579 Gianinazzi, S., Gollotte, A., Binet, M.N., van Tuinen, D., Redecker, D., Wipf, D., 2010. Agroecology: the key role of
580 arbuscular mycorrhizas in ecosystem services. *Mycorrhiza* 20, 519-530.

581 Gianinazzi, S., Vosatka, M., 2004. Inoculum of arbuscular mycorrhizal fungi for production systems, science meets
582 business. *Can. J. Bot.* 82, 1264-1271.

583 Giovannetti, M., Avio, L., Sbrana, C., 2015. Functional significance of anastomosis in arbuscular mycorrhizal
584 networks. In: Horton, T.R. (Ed.), *Mycorrhizal networks*. Springer Dordrecht Heidelberg, New York, London, pp.
585 41-67.

586 Giovannetti, M., Mosse, B., 1980. An evaluation of techniques for measuring vesicular arbuscular mycorrhizal infection
587 in roots. *New Phytol.* 84, 489-500.

588 Giovannetti, M., Sbrana, C., Avio, L., Strani, P., 2004. Patterns of below-ground plant interconnections established by
589 means of arbuscular mycorrhizal networks. *New Phytol.* 164, 175-181.

590 Gollotte, A., van Tuinen, D., Atkinson, D., 2004. Diversity of arbuscular mycorrhizal fungi colonising roots of the grass
591 species *Agrostis capillaris* and *Lolium perenne* in a field experiment. *Mycorrhiza* 14, 111-117.

592 Gosling, P., Hodge, A., Goodlass, G., Bending, G.D., 2006. Arbuscular mycorrhizal fungi and organic farming. *Agric.*
593 *Ecosyst. Environ.* 113, 17-35.

594 Hazard, C., Gosling, P., van der Gast, C.J., Mitchell, D.T., Doohan, F.M., Bending, G.D., 2013. The role of local
595 environment and geographical distance in determining community composition of arbuscular mycorrhizal fungi
596 at the landscape scale. *ISME J.* 7, 498-508.

597 Helgason, T., Daniell, T.J., Husband, R., Fitter, A.H., Young, J.P.W., 1998. Ploughing up the wood-wide web? *Nature*
598 394, 431-431.

599 Helgason, T., Merryweather, J.W., Denison, J., Wilson, P., Young, J.P.W., Fitter, A.H., 2002. Selectivity and functional
600 diversity in arbuscular mycorrhizas of co-occurring fungi and plants from a temperate deciduous woodland. *J.*
601 *Ecol.* 90, 371-384.

602 IUSS Working Group W.R.B., 2015. World Reference Base for Soil Resources 2014, update 2015 International soil
603 classification system for naming soils and creating legends for soil maps. *World Soil Res. Rep.* 106. FAO,
604 Rome.

605 Jansa, J., Erb, A., Oberholzer, H.-R., Smilauer, P., Egli, S., 2014. Soil and geography are more important determinants
606 of indigenous arbuscular mycorrhizal communities than management practices in Swiss agricultural soils. *Mol.*
607 *Ecol.* 23, 2118-2135.

608 Jansa, J., Mozafar, A., Kuhn, G., Anken, T., Ruh, R., Sanders, I.R., Frossard, E., 2003. Soil tillage affects the
609 community structures of mycorrhizal fungi in maize roots. *Ecol. Appl.* 13, 1164-1176.

610 Jeffries, P., Gianinazzi, S., Perotto, S., Turnau, K., Barea, J.-M., 2003. The contribution of arbuscular mycorrhizal fungi
611 in sustainable maintenance of plant health and soil fertility. *Biol. Fertil. Soils* 37, 1-16.

612 Kabir, Z., Koide, R.T., 2002. Effect of autumn and winter mycorrhizal cover crops on soil properties, nutrient uptake
613 and yield of sweet corn in Pennsylvania, USA. *Plant Soil* 238, 205-215.

614 Karasawa, T., Takebe, M., 2012. Temporal or spatial arrangements of cover crops to promote arbuscular mycorrhizal
615 colonization and P uptake of upland crops grown after nonmycorrhizal crops. *Plant Soil* 353, 355-366.

616 Kelderer, M., 2004. Organic apple growing: legal roles, production statistics, special features about the cultivation.
617 Compact Fruit Tree 37, 82-85.

618 Kivlin, S.N., Hawkes, C.V., Treseder, K.K., (2011) Global diversity and distribution of arbuscular mycorrhizal fungi.
619 Soil Biol. Biochem. 43, 2294-2303.

620 Kottek, M., Grieser, J., Beck, C., Rudolf, B., Rubel, F., 2006. World Map of the Köppen-Geiger climate classification
621 updated. Meteorol. Z. 15, 259-263.

622 Lee, J., Lee, S., Young, J.P.W., 2008. Improved PCR primers for the detection and identification of arbuscular
623 mycorrhizal fungi. FEMS Microbiol. Ecol. 65, 339-349.

624 Lehman, R.M., Taheri, W.I., Osborne, S.L., Buyer, J.S., Douds, D.D. Jr., 2012. Fall cover cropping can increase
625 arbuscular mycorrhizae in soils supporting intensive agricultural production. Appl. Soil Ecol. 61, 300-304.

626 Lekberg, Y., Koide, R.T., 2005. Is plant performance limited by abundance of arbuscular mycorrhizal fungi? A meta-
627 analysis of studies published between 1988 and 2003. New Phytol. 168, 189-204.

628 Liu, Y., He, J., Shi, G., An, L., Öpik, M., Feng, H., 2011. Diverse communities of arbuscular mycorrhizal fungi inhabit
629 sites with very high altitude in Tibet Plateau. FEMS Microbiol. Ecol. 78, 355-365.

630 Lugo, M.A., Ferrero, M., Menoyo, E., Estévez, M.C., Siñeriz, F., Anton, A., 2008. Arbuscular mycorrhizal fungi and
631 rhizospheric bacteria diversity along an altitudinal gradient in South American Punagrass land. Microbiol. Ecol.
632 55, 705-713.

633 Mäder, P., Edenhofer, S., Boller, T., Wiemken, A., Niggli, U., 2000. Arbuscular mycorrhizae in a long-term field trial
634 comparing low-input (organic, biological) and high-input (conventional) farming systems in a crop rotation.
635 Biol. Fertil. Soils 31, 150-156.

636 Mäder, P., Fließbach, A., Dubois, D., Gunst, L., Fried, P., Niggli, U., 2002. Soil fertility and biodiversity in organic
637 farming. Science 296, 1694-1697.

638 Mazzoncini, M., Canali, S., Giovannetti, M., Castagnoli, M., Tittarelli, F., Antichi, D., Nannelli, R., Cristani, C.,
639 Bärberi, P., 2010. Comparison of organic and conventional stockless arable systems: A multidisciplinary
640 approach to soil quality evaluation. Appl. Soil Ecol. 44, 124-132.

641 Meyer, A.H., Wooldridge, J., Dames, J.F., 2015. Effect of conventional and organic orchard floor management
642 practices on arbuscular mycorrhizal fungi in a 'Cripp's Pink'/M7 apple orchard soil. Agric. Ecosyst. Environ.
643 213, 114-120.

644 Miller, D.D., Bodmer, M., Schüepp, H., 1989. Spread of endomycorrhizal colonization and effects on growth of apple
645 seedlings. New Phytol. 111, 51-59.

646 Miller, D.D., Domoto, P.A., Walker C., 1985. Mycorrhizal fungi at eighteen apple rootstock plantings in the United
647 States. *New Phytol.* 100, 379-391.

648 Morton, J.B., Bentivenga, S.P., Bever, J.D., 1995. Discovery, measurement, and interpretation of diversity in arbuscular
649 endomycorrhizal fungi (Glomales, Zygomycetes). *Can. J. Bot.* 73, S25-S32.

650 Njeru, E.M., Avio, L., Bocci, G., Sbrana, C., Turrini, A., Bàrberi, P., Giovannetti, M., Oehl, F., 2015. Contrasting
651 effects of cover crops on 'hot spot' arbuscular mycorrhizal fungal communities in organic tomato. *Biol. Fertil.*
652 *Soils* 51, 151-166.

653 Njeru, E.M., Avio, L., Sbrana, C., Turrini, A., Bocci, G., Bàrberi, P., Giovannetti, M., 2014. First evidence for a major
654 cover crop effect on arbuscular mycorrhizal fungi and organic maize growth. *Agron. Sustain. Dev.* 34, 841-848.

655 Oehl, F., Redecker, D., Sieverding, E., 2005a. *Glomus badium*, a new sporocarpic mycorrhizal fungal species from
656 European grasslands with higher soil pH. *J. Appl. Bot. Food Qual.* 79, 38-43.

657 Oehl, F., Sieverding, E., 2004. *Pacispora*, a new vesicular arbuscular mycorrhizal fungal genus in the Glomeromycetes.
658 *J. Appl. Bot. Food Qual.* 78, 72-82.

659 Oehl, F., Sieverding, E., Ineichen, K., Mäder, P., Boller, T., Wiemken, A., 2003. Impact of land use intensity on the
660 species diversity of arbuscular mycorrhizal fungi in agroecosystems of central Europe. *Appl. Environ.*
661 *Microbiol.* 69, 2816-2824.

662 Oehl, F., Sieverding, E., Ineichen, K., Ris, E.-A., Boller, T., Wiemken, A., 2005b. Community structure of arbuscular
663 mycorrhizal fungi at different soil depths in extensively and intensively managed agroecosystems. *New Phytol.*
664 165, 273-283.

665 Oehl, F., Sieverding, E., Mäder, P., Dubois, D., Ineichen, K., Boller, T., Wiemken, A., 2004. Impact of long-term
666 conventional and organic farming on the diversity of arbuscular mycorrhizal fungi. *Oecol.* 138, 574-583.

667 Oehl, F., Silva, G.A., Palenzuela, J., Sánchez-Castro, I., Castillo, C., Sieverding, E., 2011. *Acaulospora punctata*, a new
668 fungal species in the Glomeromycetes from mountainous altitudes of the Swiss Alps and Chilean Andes. *Nova*
669 *Hedwigia* 92, 353-362.

670 Oehl, F., Sýkorová, Z., Redecker, D., Wiemken, A., Sieverding, E., 2006. *Acaulospora alpina*, a new arbuscular
671 mycorrhizal fungal species characteristic for high mountainous and alpine regions of the Swiss Alps. *Mycologia*
672 98, 286-294.

673 Öpik, M., Zobel M., Cantero, J.J., Davison, J., Facelli, J.M., Hiiesalu, I., Jairus, T., Kalwij, J.M., Koorem, K., Leal,
674 M.L., Liira, J., Metsis, M., Neshataeva, V., Paal, J., Phosri, C., Pölme, S., Reier, Ü., Saks, Ü., Schimann, H.,
675 Thiéry, O., Vasar, M., Moora, M., 2013. Global sampling of plant roots expands the described molecular
676 diversity of arbuscular mycorrhizal fungi. *Mycorrhiza* 23, 411-430.

677 Philippott, L., Raaijmakers, J.M., Lemanceau, P., van der Putten, W.H., 2013. Going back to the roots: the microbial
678 ecology of the rhizosphere. *Nat. Rev. Microbiol.* 11, 789-799.

679 Pimentel, D., Wilson, C., McCullum, C., Huang, R., Dwen, P., Flack, J., Tran, Q., Saltman, T., Cliff, B., 1997.
680 Economic and environmental benefits of biodiversity. *Biosci.* 47, 747-757.

681 Purin, S., Klauberg-Filho, O., Sturmer, S.L., 2006. Mycorrhizae activity and diversity in conventional and organic and
682 apple orchards from Brazil. *Soil Biol. Biochem.* 38, 1831-1839.

683 Read, D.J., Haselwandter, K., 1981. Observations on the mycorrhizal status of some alpine plant communities. *New*
684 *Phytol.* 88, 341-352.

685 Redecker, D., Schüßler, A., Stockinger, H., Stürmer, S.R., Morton, J.B., Walker, C., 2013. An evidence-based
686 consensus for the classification of arbuscular mycorrhizal fungi (Glomeromycota). *Mycorrhiza* 23, 515-531.

687 Roupshael, Y., Franken, P., Schneider, C., Schwarz, D., Giovannetti, M., Agnolucci, M., De Pascale, S., Bonini, P.,
688 Colla, G., 2015. Arbuscular mycorrhizal fungi act as biostimulants in horticultural crops. *Sci. Hort.* 196, 91-108.

689 Ryan, M.H., Chilvers, G.A., Dumaresq, D.C., 1994. Colonisation of wheat by VA-mycorrhizal fungi was found to be
690 higher on a farm managed in an organic manner than on a conventional neighbour. *Plant Soil* 160, 33-40.

691 Ryan, M.H., Tibbet, M., 2008. The role of arbuscular mycorrhizas in organic farming. In: Kirchmann, H., Bergström, L.
692 (Eds), *Organic crop production: ambition and limitations*. Springer, Berlin, pp. 189-229.

693 Schmidt, S.K., Sobieniak-Wiseman, L.C., Kageyama, S.A., Halloy, S.R.P., Schadt, C.W., 2008. Mycorrhizal and dark-
694 septate fungi in plant roots above 4270 meters elevation in the Andes and Rocky Mountains. *Arct. Antarc. Alp.*
695 *Res.* 40, 576-583.

696 Schwartz, M.W., Hoeksema, J.D., Gehring, C.A., Johnson, N.C., Klironomos, J.N., Abbott, L.K., Pringle, A., 2006. The
697 promise and the potential consequences of the global transport of mycorrhizal fungal inoculum. *Ecol. Lett.* 9,
698 501-515.

699 Simon, L., Lalonde, M., Bruns, T.D., 1992. Specific amplification of 18S fungal ribosomal genes from vesicular
700 arbuscular endomycorrhizal fungi colonizing roots. *Appl. Environ. Microbiol.* 58, 291-295.

701 Smith, S.E., Read, D.J., 2008. *Mycorrhizal Symbiosis*, third ed. Academic Press, London.

702 Stone, D., Ritz, K., Griffiths, B.G., Orgiazzi, A. and Creamer, R.E., 2016. Selection of biological indicators appropriate
703 for European soil monitoring. *Appl. Soil Ecol.* 97, 12-22.

704 Sýkorová, Z., Wiemken, A., Redecker, A., 2007. Co-occurring *Gentiana verna* and *Gentiana acaulis* and their
705 neighboring plants in two Swiss Upper Montane meadows harbor distinct arbuscular mycorrhizal fungal
706 communities. *Appl. Environ. Microbiol.* 73, 5426-5434.

707 Turrini, A., Giovannetti, M., 2012. Arbuscular mycorrhizal fungi in national parks, nature reserves and protected areas
708 worldwide: a strategic perspective for their in situ conservation. *Mycorrhiza* 22, 81-97.

709 Turrini, A., Sbrana, C., Avio, L., Njeru, E.M., Bocci, G., Bàrberi, P., Giovannetti, M., 2016. Changes in the
710 composition of native root arbuscular mycorrhizal fungal communities during a short-term cover crop-maize
711 succession. *Biol. Fertil. Soils* 52, 643-653.

712 Van der Gast, C.J., Gosling, P., Tiwari, B., Bending, G.D., 2011. Spatial scaling of arbuscular mycorrhizal fungal
713 diversity is affected by farming practice. *Environ. Microbiol.* 13, 241-249.

714 Van Geel, M., Ceustermans, A., Van Hemelrijck, W., Lievens, B., Honnay, O., 2015. Decrease in diversity and changes
715 in community composition of arbuscular mycorrhizal fungi in roots of apple trees with increasing orchard
716 management intensity across a regional scale. *Mol. Ecol.* 24, 941-952.

717 Van Geel, M., De Beenhouwer, M., Ceulemans, T., Caes, T., Ceustermans, A., Bylemans, D., Gomand, A., Lievens, B.,
718 Honnay, O., 2016. Application of slow-release phosphorus fertilizers increases arbuscular mycorrhizal fungal
719 diversity in the roots of apple trees. *Plant Soil* 402, 291-301.

720 Verband Deutscher Landwirtschaftlicher Untersuchungs- und Forschungsanstalten (VDLUFA) (1991) Methodenbuch:
721 Die Untersuchung von Böden. VDLUFA-Verlag.

722 Verbruggen, E., Roling, W.F.M., Gamper, H.A., Kowalchuk, G.A., Verhoef, H.A., van der Heijden, M.G.A., 2010.
723 Positive effects of organic farming on below-ground mutualists: large-scale comparison of mycorrhizal fungal
724 communities in agricultural soils. *New Phytol.* 186, 968-979.

725 Wang, P., Wang, Y. 2014. Community analysis of arbuscular mycorrhizal fungi in roots of *Poncirus trifoliata* and
726 *Citrus reticulata* based on SSU rDNA. *Scientific World J.* 2014, 562797.

727 Wu, B., Hogetsu, T., Isobe, K., Ishii, R., 2007. Community structure of arbuscular mycorrhizal fungi in a primary
728 successional volcanic desert on the southeast slope of Mount Fuji. *Mycorrhiza* 17, 495-506.

729 Yoshimura, Y., Ido, A., Matsumoto, T., Yamato, M., 2013. Communities of arbuscular mycorrhizal fungi in *Pyrus*
730 *pyrifolia* var. *culta* (Japanese pear) and an understory herbaceous plant *Plantago asiatica*. *Microbes Environ.* 28,
731 204-210.

732

733 **Table 1**

734 Main climatic parameters of the studied sites

Parameter	Unit	Altitude	
		Low	High
average annual temperature	°C	11.7	10.3
average minimum temperature (January)	°C	-3.8	- 5.1
average maximum temperature (July)	°C	29.2	27.5
annual rainfall	mm	714	522

735 Data are from the meteorological stations (average 1971-2013) of the Province of Bolzano-Bozen

736

737

738 **Table 2**

739 Main soil characteristics in the selected orchards according to their geographical site, management and altitude

Area	Management	Altitude	Organic matter (g kg ⁻¹)	pH (H ₂ O)	C (g kg ⁻¹)	N (g kg ⁻¹)	P (mg kg ⁻¹)	K (mg kg ⁻¹)	Mg (mg kg ⁻¹)
Lagundo	Organic	Low	58.7 ± 8.7	7.3 ± 0.1	34.1 ± 1.6	3.2 ± 0.1	57 ± 24	43 ± 21	380 ± 45
		High	81.7 ± 10.7	7.2 ± 0.1	34.1 ± 1.0	3.2 ± 0.1	108 ± 19	56 ± 23	293 ± 18
	Integrated	Low	38.7 ± 4.1	6.9 ± 0.2	25.1 ± 1.1	2.8 ± 0.3	79 ± 5	44 ± 13	360 ± 105
		High	48.7 ± 1.8	6.2 ± 0.5	31.1 ± 1.8	3.2 ± 0.1	61 ± 8	103 ± 37	187 ± 26
Terlano	Organic	Low	40.0 ± 6.7	7.3 ± 0.1	32.1 ± 4.7	2.6 ± 0.3	88 ± 16	86 ± 18	293 ± 14
	Integrated	Low	36.0 ± 5.5	6.4 ± 1.0	28.7 ± 2.9	3.0 ± 0.2	43 ± 8	70 ± 17	283 ± 38

740 Data are averages ± s.e. For clarity, statistics is reported only in the text.

741

742 **Table 3**

743 Percentage of root mycorrhizal colonisation and leaf nutrient concentrations in the selected orchards according to their geographical site, management and altitude

Area	Management	Altitude	Mycorrhizal colonization (%)	N (%)	P (%)	K (%)	Mg (%)	Ca (%)
Lagundo	Organic	Low	41.44 ± 12.23	3.64 ± 0.70	0.22 ± 0.01	0.95 ± 0.05	0.34 ± 0.01	1.23 ± 0.16
		High	34.97 ± 15.69	2.94 ± 0.29	0.21 ± 0.01	0.97 ± 0.05	0.29 ± 0.01	1.07 ± 0.04
	Integrated	Low	52.02 ± 14.33	3.21 ± 0.39	0.21 ± 0.01	1.02 ± 0.08	0.36 ± 0.05	1.17 ± 0.12
		High	48.19 ± 11.11	2.42 ± 0.09	0.21 ± 0.01	1.15 ± 0.06	0.25 ± 0.02	1.18 ± 0.10
Terlano	Organic	Low	60.01 ± 13.76	2.82 ± 0.10	0.24 ± 0.01	1.26 ± 0.04	0.3 ± 0.01	1.12 ± 0.06
	Integrated	Low	54.71 ± 23.41	3.01 ± 0.24	0.24 ± 0.01	1.2 ± 0.05	0.27 ± 0.03	0.9 ± 0.15

744 Data are averages ± s.e. For clarity, statistics is reported only in the text.

745

746

747

748

749

750

751

752 **Table4**

753 Sequence types of arbuscular mycorrhizal fungi, identified using NS31-AML2 primers pair, in the roots of apple plants

754 grown in South Tyrol orchards

OTU name	Identity NCBI database (%)	Identity Maarjam database (%)	Maarjam database virtual taxa	Taxonomic affiliation
C11 [†]	AJ276087 (99)	FN869808 (99)	VTX00057	<i>Claroideoglossum lamellosum</i>
C12	JN644447 (99)	HE615004 (99)	VTX00056	<i>Claroideoglossum</i> sp.
C13	KF290671 (99)	KF290671 (99)	VTX00225	<i>Claroideoglossum</i> sp.
Div	FJ831643 (100)	FJ831643 (100)	VTX00062	<i>Diversispora</i> sp.
Fun	AJ306438 (99)	AY635833 (99)	VTX00067	<i>Funneliformis mosseae</i>
Glo1	JQ350797 (99)	KF386274 (99)	VTX00214	<i>Glomus</i> sp.
Glo2	KF467269 (99)	KF467269 (99)	VTX00135	<i>Glomus</i> sp.
Glo3	H4380136 (99)	HG004495	VTX00153	<i>Glomus</i> sp.
Glo4	JX144121 (99)	KC579423	VTX00304	<i>Glomus</i> sp.
Glo5	JN009364 (99)	JN009364 (99)	VTX00151	<i>Glomus</i> sp.
Glo6	JX144133 (99)	HG004465	VTX00301	<i>Glomus</i> sp.
Glo7	GU059539 (99)	GU059539 (99)	VTX00222	<i>Glomus indicum</i>
Par	AB695049 (100)	-	-	Uncultured Glomeromycota sp.
Scle	AJ33706 (99)	AJ33706 (98)	VTX00069	<i>Sclerocystis sinuosa</i>
Sept1	KF386332 (99)	KF386332 (99)	VTX00063	<i>Septoglossum</i> sp.
Sept2	FR750212 (99)	FJ831626 (99)	VTX00064	<i>Septoglossum constrictus</i>
Rhi	FJ009618 (99)	FJ009617 (99)	VTX00114	<i>Rhizoglossum irregulare</i>

755 [†]Names denote the most similar AM fungal species of sequenced clones: C11, *Claroideoglossum* sp.1; C12,756 *Claroideoglossum* sp.2; C13, *Claroideoglossum* sp.3; Div, *Diversispora* sp.; Fun, *Funneliformis mosseae*; Glo1, *Glomus*757 sp.1; Glo2, *Glomus* sp.2; Glo3, *Glomus* sp.3; Glo4, *Glomus* sp.4; Glo5, *Glomus* sp.5; Glo6, *Glomus* sp.6; Glo7, *Glomus*758 sp.7; Par, *Paraglossum* sp.; Scle, *Sclerocystis sinuosa*; Sept1, *Septoglossum* sp.1; Sept2, *Septoglossum* sp.2; Rhi,759 *Rhizoglossum irregulare*.

760 †

761

762 **Table 5**

763 Richness (S), Shannon-Weaver (Hs), Simpson (D) and Evenness (E) indices calculated from AMF DGGE profile
764 associated with apple plant roots growing in orchards cultivated in two geographical sites (Terlano vs Lagundo) and
765 under two managements (organic vs integrated) (mean \pm standard error)

Geographical area	Management	S	Hs	D	E
Lagundo	Organic	9.83 \pm 0.87	2.19 \pm 0.08	0.12 \pm 0.01	0.97 \pm 0.00
	Integrated	6.83 \pm 1.01	1.72 \pm 0.16	0.20 \pm 0.03	0.93 \pm 0.02
Terlano	Organic	10.5 \pm 1.63	2.18 \pm 0.17	0.13 \pm 0.02	0.96 \pm 0.01
	Integrated	12.3 \pm 1.33	2.38 \pm 0.08	0.09 \pm 0.01	0.95 \pm 0.01
Analysis of variance (P values)					
Geographical area		0.033	0.049	0.069	0.734
Management		0.666	0.392	0.252	0.126
Geographical area x Management		0.086	0.044	0.033	0.171

766

767

768

769 **Table 6**
 770 Richness (S), Shannon-Weaver (Hs), Simpson (D) and Evenness (E) indices calculated from AMF DGGE profile
 771 associated with apple plant roots growing in orchards cultivated in Lagundo under two managements (organic vs
 772 integrated) (mean \pm standard error)

773

Geographical area	Management	S	Hs	D	E
Lagundo	Organic	9.83 \pm 0.87	2.19 \pm 0.08	0.12 \pm 0.01	0.97 \pm 0.00
	Integrated	6.83 \pm 1.01	1.72 \pm 0.16	0.20 \pm 0.03	0.93 \pm 0.02
Analysis of variance (P values)					
Management		0.029	0.028	0.049	0.069

774

775

776

777

778 **Captions**

779 **Fig. 1.** a) Maps showing the location of apple orchards in South Tyrol, Italy. ● Integrated managed orchards at high
780 altitude, ● Organically managed orchards at high altitude, ● Integrated managed orchards at low altitude, ●
781 Organically managed orchards at low altitude. Source of the air photo on the right: Google earth V 7.1.2.2041 image©
782 2016 DigitalGlobe <http://www.earth.google.com>. Pictures of organically b) and integrated c) managed apple trees in the
783 study site.

784

785 **Fig. 2.** Neighbor-Joining phylogenetic tree of glomeromycotan sequences derived from apple roots of orchards growing
786 in South Tyrol (Italy). Bootstrap values are shown when they exceed 75 % (1,000 replications). The analysis is based on
787 partial nuclear small subunit ribosomal RNA gene sequences (SSU; ~ 550bp; NS31/AML2 fragment) and involved 96
788 nucleotide sequences. Different sequence types are indicated in brackets and names are reported in Table 1. AMF
789 family are also reported. Sequences obtained in the present study are shown in bold and their accession numbers are
790 prefixed with site/management/altitude clone identifiers (Te, Terlano; La, Lagundo; O, Organic; I, Integrated; L, Low
791 altitude; H, High altitude). The tree is rooted with a reference sequence of *Corallochytrium lymacisporum* (L42528).

792

793 **Fig. 3.** Relative abundance (%) of AMF phylotypes detected in the roots of the different orchards cultivated in two
794 geographical sites (Lagundo and Terlano), under two managements (organic and integrated) and at two altitudes (Low,
795 300 m and High, 600m).

796

797 **Fig. 4.** Dendrogram obtained by UPGMA (Unweighted Pair Group Method Using Arithmetic Average) based on AMF
798 DGGE profiles obtained from apple plant roots growing in orchards cultivated in two geographical sites (Terlano and
799 Lagundo) and under two managements (■ Organic, ■ Integrated) (Te, Terlano; La, Lagundo; O, Organic; I, Integrated;
800 L, Low altitude; H, High altitude).

801

802 **Fig. 5.** Non-metric multidimensional scaling plot of DGGE analysis. Each point on the plot represents the AMF
803 community composition associated with apple plant roots growing in orchards cultivated in Terlano area (blue symbols)
804 or Lagundo area (orange symbols). The stress value is 0.2, the ANOSIM values (R) indicates significant differences
805 between Lagundo and Terlano (0.786, P=0.0001) and between organic and integrated (0.227, P=0.05) (Te, Terlano; La,
806 Lagundo; O, Organic; I, Integrated; L, Low altitude; H, High altitude).




Fig. 1




Fig. 3




Fig.4




Fig. 5




Fig. 2