

SIMPLE LINEAR COMPACTIFICATIONS OF ODD ORTHOGONAL GROUPS

JACOPO GANDINI

ABSTRACT. We classify the simple linear compactifications of $\mathrm{SO}(2r+1)$, namely those compactifications with a unique closed orbit which are obtained by taking the closure of the $\mathrm{SO}(2r+1) \times \mathrm{SO}(2r+1)$ -orbit of the identity in a projective space $\mathbb{P}(\mathrm{End}(V))$, where V is a finite dimensional rational $\mathrm{SO}(2r+1)$ -module.

INTRODUCTION

Let G be a semisimple and simply-connected algebraic group defined over an algebraically closed field \mathbb{k} of characteristic zero. Fix a maximal torus $T \subset G$ and a Borel subgroup $B \supset T$, denote Φ the associated root system of G and $\Delta \subset \Phi$ the associated basis. Denote Λ the weight lattice of G and $\Lambda^+ \subset \Lambda$ the semigroup of dominant weights. For $\lambda \in \Lambda^+$, denote $V(\lambda)$ the simple G -module of highest weight λ .

A $G \times G$ -variety X is called *linear* if it admits an equivariant embedding in the projective space of a finite dimensional $G \times G$ -module, while is called *simple* if it possesses a unique closed $G \times G$ -orbit. If $\Pi \subset \Lambda^+$ is a finite subset, consider the $G \times G$ -variety

$$X_\Pi = \overline{(G \times G)[\mathrm{Id}]} \subset \mathbb{P}\left(\bigoplus_{\lambda \in \Pi} \mathrm{End}(V(\lambda))\right) :$$

it is a linear compactification of a quotient of G , and conversely every linear compactification of a quotient of G arise in such a way for some $\Pi \subset \Lambda^+$. We say that X_Π is *adjoint* if it is a compactification of a quotient of the adjoint group G_{ad} .

The variety X_Π was studied by Timashev in [12]: there are studied the local structure and the $G \times G$ -orbit structure, and normality and smoothness are characterized as well. The conditions of normality in particular rely on some properties of the tensor product, and together with the conditions of smoothness they were remarkably simplified by Bravi, Gandini, Maffei, Ruzzi in [3] in case X_Π is simple and adjoint, and by Gandini, Ruzzi in [6] in case X_Π is simple. In particular, in [3] it was shown that every simple adjoint linear compactification is normal if G is simply laced, whereas several examples of non-normal simple adjoint linear compactifications arise in the non-simply laced case.

By a theorem of Sumihiro (see [10]), every simple normal $G \times G$ -variety is linear. Hence if we restrict to simple normal adjoint X_Π 's, a classification follows by the general Luna-Vust theory of spherical embeddings (see [9]): they are classified by their closed orbits, i.e. by non-empty subsets of Δ . However, as far as we know, no explicit classification is known in the general spherical context without assuming normality: this paper stems from the attempt to understand this classification in some explicit case. More precisely, the aim of this work is to classify the

simple linear compactifications of $\mathrm{SO}(2r+1)$: this will be done by classifying the subsets $\Pi \subset \Lambda^+$ which give rise to isomorphic simple compactifications.

Consider the *dominance order* on Λ , defined by $\mu \leq \lambda$ if $\lambda - \mu \in \mathbb{N}\Delta$, and the *rational dominance order* $\leq_{\mathbb{Q}}$, defined by $\mu \leq_{\mathbb{Q}} \lambda$ if $\lambda - \mu \in \mathbb{Q}_{\geq 0}\Delta$. If $\Pi \subset \Lambda^+$ is finite, then the closed orbits of X_{Π} correspond to some maximal elements of Π w.r.t. $\leq_{\mathbb{Q}}$, and X_{Π} is simple if and only if Π contains a unique maximal element w.r.t. $\leq_{\mathbb{Q}}$. If this is the case, we say that $\Pi \subset \Lambda^+$ is a *simple subset*. On the other hand, X_{Π} is an adjoint compactification if and only if Π is contained in a coset of $\Lambda/\mathbb{Z}\Delta$, in which case we say that $\Pi \subset \Lambda^+$ is an *adjoint subset*. Therefore X_{Π} is a simple adjoint variety if and only if Π contains a unique maximal element w.r.t. \leq .

For simplicity, in case $\Pi = \{\lambda\}$, we denote X_{Π} by X_{λ} , while in case $\Pi = \{\lambda, \mu\}$, we denote X_{Π} by $X_{\lambda, \mu}$. Let $\Pi \subset \Lambda^+$ be a simple adjoint subset with maximal element λ , denote \tilde{X}_{λ} the normalization of X_{λ} and $\Pi^+(\lambda) = \{\mu \in \Lambda^+ : \mu \leq \lambda\}$. Then $\Pi \subset \Pi^+(\lambda)$ and we get natural projections

$$X_{\Pi^+(\lambda)} \longrightarrow X_{\Pi} \longrightarrow X_{\lambda}$$

While Kannan shown in [8] that $X_{\Pi^+(\lambda)}$ is projectively normal, De Concini proved in [4] that $X_{\Pi^+(\lambda)} \simeq \tilde{X}_{\lambda}$. In particular, if Π is adjoint and simple with maximal element λ , it follows that $\tilde{X}_{\lambda} \rightarrow X_{\Pi}$ is the normalization.

If $\lambda \in \Lambda^+$, we say that a weight $\mu \in \Pi^+(\lambda)$ is *trivial* if $X_{\lambda, \mu}$ is equivariantly isomorphic to X_{λ} . We denote by $\Pi_{\mathrm{tr}}^+(\lambda) \subset \Pi^+(\lambda)$ the subset of trivial weights: if G is simply laced, then by [3] we have $\Pi_{\mathrm{tr}}^+(\lambda) = \Pi^+(\lambda)$. If $\mathrm{Supp}(\lambda) \subset \Delta$ denotes the set of simple roots non-orthogonal to λ , then the variety X_{λ} depends only on $\mathrm{Supp}(\lambda)$: therefore a first step to classify the simple linear compactifications X_{Π} such that $\tilde{X}_{\lambda} \rightarrow X_{\Pi} \rightarrow X_{\lambda}$ is to characterize the set $\Pi_{\mathrm{tr}}^+(n\lambda)$ for $n \in \mathbb{N}$.

In the case $G = \mathrm{Spin}(2r+1)$, we will give the following combinatorial description of trivial weights. Denote $\Delta = \{\alpha_1, \dots, \alpha_r\}$, where the numbering is the usual one as in [2], and denote $\omega_1, \dots, \omega_r$ the associated fundamental weights.

Theorem 1 (see Theorem 2.1). *Let $G = \mathrm{Spin}(2r+1)$. Let $\lambda \in \Lambda^+$ and $\mu \in \Pi^+(\lambda)$, denote q and l the maximal integers such that $\alpha_q \in \mathrm{Supp}(\lambda)$ and $\alpha_l \in \mathrm{Supp}(\mu)$ and write $\lambda - \mu = \sum_{i=1}^r a_i \alpha_i$. Then $\mu \in \Pi_{\mathrm{tr}}^+(\lambda)$ if and only if a_r is even or $a_r > 2 \min\{r-l, r-q\}$.*

Previous theorem essentially expresses some properties of the tensor product. A main motivation to explain the combinatorial condition in the previous theorem arises by considering the case of the first fundamental weight, where it can be deduced by the Schur-Weyl duality (Proposition 2.5).

To reduce the classification of the simple linear compactifications of $\mathrm{SO}(2r+1)$ to the classification of the trivial weights, it is possible to define a partial order relation \leq^{λ} on Λ with the following geometrical meaning: if $\mu, \nu \in \Pi^+(\lambda) \setminus \Pi_{\mathrm{tr}}^+(\lambda)$, then

$$\nu \leq^{\lambda} \mu \quad \text{if and only if} \quad \begin{array}{l} \text{there exists a } G \times G\text{-morphism} \\ X_{\lambda, \mu} \longrightarrow X_{\lambda, \nu} \end{array}$$

From a combinatorial point of view, \leq^{λ} is the degeneration of the dominance order associated to the set $\Phi^+(\lambda)$ of the positive roots of Φ which are non-orthogonal to λ : if $\mu, \nu \in \Lambda$, then $\nu \leq^{\lambda} \mu$ if and only if $\mu - \nu \in \mathbb{N}\Phi^+(\lambda)$.

In case λ is regular, then \leq^{λ} coincides with the usual dominance order \leq , while if $\lambda = 0$ then \leq^{λ} is the trivial order. In the general case of a (possibly non-adjoint) simple subset Π , the partial order \leq^{λ} was used in [6] to characterize combinatorially the normality of the variety X_{Π} .

If $\Pi \subset \Lambda^+$ is an adjoint simple subset with maximal element λ , denote

$$\Pi_{\text{red}} = \{\mu \in \Pi \setminus \Pi_{\text{tr}}^+(\lambda) : \mu \text{ is maximal w.r.t. } \leq^\lambda\} \cup \{\lambda\}.$$

In case $\Pi = \Pi_{\text{red}}$ we say that Π is a *reduced adjoint subset*. If Π' is another adjoint simple subset with maximal element λ' , then we say that Π and Π' are *equivalent* (and we write $\Pi \sim \Pi'$) if $\text{Supp}(\lambda') = \text{Supp}(\lambda)$ and $\Pi' - \lambda' = \Pi - \lambda$.

Theorem 2 (see Corollary 3.6). *Let $G = \text{Spin}(2r+1)$ and let $\Pi, \Pi' \subset \Lambda^+$ be adjoint simple subsets with maximal elements resp. λ and λ' .*

- i) *Suppose that $\text{Supp}(\lambda) = \text{Supp}(\lambda')$. There exists an equivariant morphism $X_\Pi \rightarrow X_{\Pi'}$ if and only if for every $\mu' \in \Pi'_{\text{red}}$ there exists $\mu \in \Pi_{\text{red}}$ such that $\mu' - \lambda' \leq^\lambda \mu - \lambda$.*
- ii) *The varieties X_Π and $X_{\Pi'}$ are equivariantly isomorphic if and only if $\Pi_{\text{red}} \sim \Pi'_{\text{red}}$.*

As a corollary it follows that the simple linear compactifications of $\text{SO}(2r+1)$ are classified by simple reduced subsets up to equivalence.

The paper is organized as follows. In Section 1, we study the compactifications X_λ and X_Π in full generality: throughout this section (and only in this section) G will denote an arbitrary simply connected semisimple algebraic group. In Section 2, we describe combinatorially the set of trivial weights $\Pi_{\text{tr}}^+(\lambda)$, where λ is a dominant weight for $\text{Spin}(2r+1)$, and we prove Theorem 1. In Section 3, we introduce the reduction of a simple subset and we characterize combinatorially the existence of an equivariant morphism between two simple linear compactifications of $\text{SO}(2r+1)$ possessing isomorphic closed orbits and we prove Theorem 2. In Section 4, we give examples by means of tables in the case of the simple linear compactifications of $\text{SO}(7)$ and of $\text{SO}(9)$.

Differently from the introduction, since we will only deal with adjoint compactifications, we will refer to *simple adjoint sets* of dominant weights just as *simple sets*. It will be also convenient to adopt a “dual viewpoint” in the definition of the variety X_Π : the simple modules $V(\mu)$ occurring in its definition will be substituted with their duals.

Acknowledgments. I would like to thank A. Maffei, who suggested the problem, for his advice and for his continuous support. As well, I thank the referee for his careful reading and useful comments.

1. THE VARIETIES X_λ AND X_Π

Let G be a semisimple simply connected algebraic group over an algebraically closed field \mathbb{k} of characteristic zero and denote \mathfrak{g} its Lie algebra. Fix a maximal torus T and a Borel subgroup $B \supset T$, denote B^- the opposite Borel subgroup of B w.r.t. T and denote $U \subset B$ and $U^- \subset B^-$ the associated maximal unipotent subgroups. Correspondingly to the choice of T and B , we fix $T \times T$ as a maximal torus and $B \times B^-$ as a Borel subgroup in $G \times G$. Denote Φ the root system associated to T and W the Weyl group of Φ . Denote Δ the basis of Φ associated to B and Φ^+ the associated set of positive roots. In case Φ is irreducible and non-simply laced, then we write $\Phi^+ = \Phi_s^+ \cup \Phi_l^+$, where Φ_s^+ and Φ_l^+ denote respectively the set of the positive short roots and that of the positive long roots.

Denote Λ the weight lattice of Φ and $\Lambda^+ \subset \Lambda$ the semigroup of the dominant weights associated to Δ and set $\Lambda_{\mathbb{Q}} = \Lambda \otimes \mathbb{Q}$. If $\lambda \in \Lambda^+$ then we denote by $V(\lambda)$ the simple G -module of highest weight λ , however, if we deal with different groups, we will use also the notation $V_G(\lambda)$. Let $\lambda \mapsto \lambda^*$ be the linear involution of Λ defined by $V(\lambda)^* \simeq V(\lambda^*)$ for any dominant weight λ . If

$\alpha \in \Delta$, we denote by ω_α the associated fundamental weight and by $\{e_\alpha, \alpha^\vee, f_\alpha\}$ an $\mathfrak{sl}(2)$ -triple of T -weights $\alpha, 0, -\alpha$. Given a weight $\lambda = \sum_{\alpha \in \Delta} n_\alpha \omega_\alpha \in \Lambda$, denote $\lambda^+ = \sum_{n_\alpha > 0} n_\alpha \omega_\alpha$ and $\lambda^- = \sum_{n_\alpha < 0} |n_\alpha| \omega_\alpha$. When we deal with explicit root systems, we use the numbering of simple roots and fundamental weights of Bourbaki [2].

If $\lambda \in \Lambda^+$, denote $\Pi(\lambda) \subset \Lambda$ the set of weights occurring in $V(\lambda)$, $\mathcal{P}(\lambda) \subset \Lambda_{\mathbb{Q}}$ the convex hull of $\Pi(\lambda)$ and $\Pi_{\mathbb{Q}}(\lambda) = \mathcal{P}(\lambda) \cap \Lambda$. Denote \leq and $\leq_{\mathbb{Q}}$ resp. the *dominance order* and the *rational dominance order* on Λ , defined by $\mu \leq \lambda$ (resp. $\mu \leq_{\mathbb{Q}} \lambda$) if and only if $\lambda - \mu \in \mathbb{N}\Delta$ (resp. $\mathbb{Q}_{\geq 0}\Delta$). Then we have $\Pi(\lambda) = W\Pi^+(\lambda)$ and $\Pi_{\mathbb{Q}}(\lambda) = W\Pi_{\mathbb{Q}}^+(\lambda)$, where we denote

$$\begin{aligned}\Pi^+(\lambda) &= \Pi(\lambda) \cap \Lambda^+ = \{\mu \in \Lambda^+ : \mu \leq \lambda\}, \\ \Pi_{\mathbb{Q}}^+(\lambda) &= \mathcal{P}(\lambda) \cap \Lambda^+ = \{\mu \in \Lambda^+ : \mu \leq_{\mathbb{Q}} \lambda\}.\end{aligned}$$

If $\lambda \in \Lambda$, define its *support* as $\text{Supp}(\lambda) = \{\alpha \in \Delta : \langle \lambda, \alpha^\vee \rangle \neq 0\}$, while if $\theta = \sum_{\alpha \in \Delta} n_\alpha \alpha \in \mathbb{Z}\Delta$, define its *support over Δ* as $\text{Supp}_\Delta(\theta) = \{\alpha \in \Delta : n_\alpha \neq 0\}$. If $\lambda \in \Lambda$, denote $\Phi^+(\lambda) \subset \Phi^+$ the subset of the positive roots which are non-orthogonal to λ :

$$\Phi^+(\lambda) = \{\theta \in \Phi^+ : \text{Supp}_\Delta(\theta) \cap \text{Supp}(\lambda) \neq \emptyset\}.$$

In case Φ is irreducible and non-simply laced, we also set $\Phi_s^+(\lambda) = \Phi_s^+ \cap \Phi^+(\lambda)$ and $\Phi_l^+(\lambda) = \Phi_l^+ \cap \Phi^+(\lambda)$.

If λ is a non-zero dominant weight, consider the $G \times G$ -variety

$$X_\lambda = \overline{(G \times G)[\text{Id}]} \subset \mathbb{P}(\text{End}(V(\lambda)^*)),$$

Since $\mathbb{P}(\text{End}(V(\lambda)^*))$ possesses a unique closed $G \times G$ orbit and since the diagonal of G fixes the identity, it follows that X_λ is a simple compactification of a quotient of the adjoint group G_{ad} .

Proposition 1.1 ([3, Prop. 1.2]). *Let $\lambda, \mu \in \Lambda^+$. Then $X_\lambda \simeq X_\mu$ as $G \times G$ -varieties if and only if $\text{Supp}(\lambda) = \text{Supp}(\mu)$.*

If $\lambda \in \Lambda^+$ is regular (i.e. if $\text{Supp}(\lambda) = \Delta$), then X_λ coincides with the wonderful compactification of G_{ad} introduced by De Concini and Procesi in [5]. We will denote this variety by M : it is smooth and the complement of its open orbit is the union of smooth prime divisors with normal crossings whose intersection is the closed orbit $G/B \times G/B$.

Since G is semisimple and simply connected, we may identify the Picard group $\text{Pic}(G/B)$ with the weight lattice Λ : we identify a weight $\lambda \in \Lambda$ with the line bundle on G/B whose T -weight in the B -fixed point is $-\lambda$. The restriction of line bundles to the closed orbit induces an homomorphism

$$\omega : \text{Pic}(M) \longrightarrow \Lambda \times \Lambda$$

which is injective and which identifies $\text{Pic}(M)$ with the sublattice $\{(\lambda, \lambda^*) : \lambda \in \Lambda\} \subset \Lambda \times \Lambda$. Therefore $\text{Pic}(M)$ is identified with Λ and we will still denote by $\mathcal{L}_\lambda \in \text{Pic}(M)$ the line bundle whose image is (λ, λ^*) . A line bundle $\mathcal{L}_\lambda \in \text{Pic}(M)$ is generated by its sections if and only if $\lambda \in \Lambda^+$, in which case, as a $G \times G$ -module, the following decomposition holds ([5, Theorem 8.3]):

$$\Gamma(M, \mathcal{L}_\lambda) \simeq \bigoplus_{\mu \in \Pi^+(\lambda)} \text{End}(V(\mu)).$$

Fix now $\lambda \in \Lambda^+$ (possibly non-regular). Then the map $G_{\text{ad}} \rightarrow \mathbb{P}(\text{End}(V(\lambda)^*))$ extends to a map $M \rightarrow \mathbb{P}(\text{End}(V(\lambda)^*))$ whose image is X_λ and such that \mathcal{L}_λ is the pullback of the hyperplane bundle on $\mathbb{P}(\text{End}(V(\lambda)^*))$. If we pull back the homogeneous coordinates of $\mathbb{P}(\text{End}(V(\lambda)^*))$ to

M , we get a submodule of $\Gamma(M, \mathcal{L}_\lambda)$ which is isomorphic to $\text{End}(V(\lambda))$; by abuse of notation we still denote this submodule by $\text{End}(V(\lambda))$.

Consider the algebra

$$\tilde{A}(\lambda) = \bigoplus_{n \in \mathbb{N}} \Gamma(M, \mathcal{L}_{n\lambda})$$

and denote $A(\lambda) \subset \tilde{A}(\lambda)$ the subalgebra generated by $\text{End}(V(\lambda)) \subset \Gamma(M, \mathcal{L}_\lambda)$; consider the natural gradings on $\tilde{A}(\lambda)$ and $A(\lambda)$ respectively defined by $\tilde{A}_n(\lambda) = \Gamma(M, \mathcal{L}_{n\lambda})$ and $A_n(\lambda) = \tilde{A}_n(\lambda) \cap A(\lambda)$. Since $A(\lambda)$ is the projective coordinate ring of X_λ , if we set $\tilde{X}_\lambda = \text{Proj } \tilde{A}(\lambda)$ then we get a commutative diagram as follows:

$$\begin{array}{ccc} M & \xrightarrow{\quad} & \tilde{X}_\lambda \\ & \searrow & \downarrow \\ & & X_\lambda \end{array}$$

In [8], Kannan shown that $\tilde{A}(\lambda)$ is generated in degree 1, while in [4] De Concini shown that $\tilde{X}_\lambda \rightarrow X_\lambda$ is the normalization.

If $\lambda \in \Lambda^+$, denote $\text{Id}_\lambda \in \text{End}(V(\lambda)^*)$ the identity. Similarly, if $\Pi \subset \Lambda^+$ is a finite subset, denote Id_Π the identity vector $(\text{Id}_\mu)_{\mu \in \Pi} \in \bigoplus_{\mu \in \Pi} \text{End}(V(\mu))$. Given such a subset Π , consider the $G \times G$ -variety

$$X_\Pi = \overline{(G \times G)[\text{Id}_\Pi]} \subset \mathbb{P}\left(\bigoplus_{\mu \in \Pi} \text{End}(V(\mu)^*)\right).$$

If $\Pi = \{\lambda\}$, then we get the variety X_λ , while if $\Pi = \Pi^+(\lambda)$ we get its normalization \tilde{X}_λ . If $\Pi = \{\mu_1, \dots, \mu_m\}$, for simplicity we will denote X_Π also by X_{μ_1, \dots, μ_m} . We say that the variety X_Π is *simple* if it contains a unique closed $G \times G$ orbit, while we say that X_Π is *adjoint* if it is a compactification of a quotient of G_{ad} .

Proposition 1.2 ([12, §8]). *Let $\Pi \subset \Lambda^+$ be a finite subset and denote $\mathcal{P}(\Pi) \subset \mathbb{Q}\Delta$ the polytope generated by the T -weights occurring in the G -module $\bigoplus_{\mu \in \Pi} V(\mu)$.*

- i) X_Π is *adjoint* if and only if Π is contained in a coset of $\Lambda/\mathbb{Z}\Delta$.
- ii) Let $\mu \in \Pi$. Then X_Π contains the closed orbit of $\mathbb{P}(\text{End}(V(\mu)^*))$ if and only if μ is an extremal vertex of $\mathcal{P}(\Pi)$.

It follows by previous proposition that X_Π is simple if and only if Π possesses a unique maximal element w.r.t. $\leq_{\mathbb{Q}}$, whereas it is simple and adjoint if and only if Π contains a unique maximal element w.r.t. \leq . Correspondingly, we will say that Π is *adjoint* if it is contained in a coset of $\Lambda/\mathbb{Z}\Delta$, and we say that an adjoint subset is *simple* if it possesses a unique maximal element w.r.t. \leq . Since we will deal only with adjoint subsets, for simplicity we will refer to adjoint simple subsets just as *simple subsets*. For a general treatment on the case of a possibly non-adjoint simple linear group compactification see [6].

Suppose that $\Pi \subset \Lambda^+$ is simple with maximal element λ and consider the line bundle $\mathcal{L}_\lambda \in \text{Pic}(M)$. By its decomposition, it follows that $\Gamma(M, \mathcal{L}_\lambda)$ possesses a $G \times G$ submodule $A_1(\Pi)$ isomorphic to $\bigoplus_{\mu \in \Pi} \text{End}(V(\mu))$, which is base point free since $\lambda \in \Pi$. On the other hand $\Gamma(M, \mathcal{L}_\lambda) = A_1(\Pi^+(\lambda))$ and $X_{\Pi^+(\lambda)} \simeq \tilde{X}_\lambda$, hence we get morphisms

$$M \longrightarrow \tilde{X}_\lambda \longrightarrow X_\Pi \longrightarrow X_\lambda$$

and it follows that $\tilde{X}_\lambda \rightarrow X_\Pi$ is the normalization. Denote $A(\Pi)$ the projective coordinate ring of X_Π , namely the subalgebra of $\tilde{A}(\lambda)$ generated by $\bigoplus_{\mu \in \Pi} \text{End}(V(\mu))$, and denote $A_n(\Pi) = \tilde{A}_n(\lambda) \cap A(\Pi)$.

If $\mu \in \Pi$, denote $\phi_\mu \in \text{End}(V(\mu))$ a highest weight vector. Consider the $B \times B^-$ -stable affine open subsets $X_\lambda^\circ \subset X_\lambda$ and $X_\Pi^\circ \subset X_\Pi$ defined by the non-vanishing of ϕ_λ : then we get

$$\mathbb{k}[X_\Pi^\circ] = \left\{ \frac{\phi}{\phi_\lambda^n} : \phi \in A_n(\Pi), n \in \mathbb{N} \right\} \supset \left\{ \frac{\phi}{\phi_\lambda^n} : \phi \in A_n(\lambda), n \in \mathbb{N} \right\} = \mathbb{k}[X_\lambda^\circ].$$

Previous rings are not $G \times G$ -module, however they are $\mathfrak{g} \oplus \mathfrak{g}$ -modules.

Lemma 1.3. *Let $\Pi \subset \Lambda^+$ be simple with maximal element λ . As a $\mathfrak{g} \oplus \mathfrak{g}$ -algebra, $\mathbb{k}[X_\Pi^\circ]$ is generated by $\mathbb{k}[X_\lambda^\circ]$ together with the set $\{\phi_\mu/\phi_\lambda\}_{\mu \in \Pi}$.*

Proof. Since the projective coordinate ring $A(\Pi)$ is generated by its degree one component $A_1(\Pi) = \bigoplus_{\mu \in \Pi} \text{End}(V(\mu))$, it follows that $\mathbb{k}[X_\Pi^\circ]$ is generated as an algebra by its subset $B(\Pi) = \{\phi/\phi_\lambda : \phi \in A_1(\Pi)\}$. Using the action of $\mathfrak{g} \oplus \mathfrak{g}$, let's show that $B(\Pi)$ is contained in the $\mathfrak{g} \oplus \mathfrak{g}$ -subalgebra $B'(\Pi) \subset \mathbb{k}[X_\Pi^\circ]$ generated by $\mathbb{k}[X_\lambda^\circ]$ together with $\{\phi_\mu/\phi_\lambda\}_{\mu \in \Pi}$. Suppose indeed that α is a simple root and that $\phi/\phi_\lambda \in B'(\Pi)$: then $f_\alpha(\phi)/\phi_\lambda \in B'(\Pi)$ as well since

$$\frac{f_\alpha(\phi)}{\phi_\lambda} = f_\alpha \left(\frac{\phi}{\phi_\lambda} \right) + \frac{\phi}{\phi_\lambda} \cdot \frac{f_\alpha(\phi_\lambda)}{\phi_\lambda}. \quad \square$$

Given $\lambda, \mu \in \Lambda^+$, consider the multiplication map

$$m_{\lambda, \mu} : \Gamma(M, \mathcal{L}_\lambda) \times \Gamma(M, \mathcal{L}_\mu) \rightarrow \Gamma(M, \mathcal{L}_{\lambda+\mu}),$$

which is surjective by [8]. In order to describe combinatorially $m_{\lambda, \mu}$, as in [8] or in [4] it is possible to identify sections of a line bundle on M with functions on G and use the description of the multiplication of matrix coefficients. Recall that as a $G \times G$ -module it holds the decomposition

$$\mathbb{k}[G] = \bigoplus_{\lambda \in \Lambda^+} \text{End}(V(\lambda)) \simeq \bigoplus_{\lambda \in \Lambda^+} V(\lambda)^* \otimes V(\lambda).$$

More explicitly if V is a G -module, define the matrix coefficient $c_V : V^* \otimes V \rightarrow \mathbb{k}[G]$ by $c_V(\psi \otimes v)(g) = \langle \psi, gv \rangle$. If we multiply functions in $\mathbb{k}[G]$ of this type then we get

$$c_V(\psi \otimes v) \cdot c_W(\chi \otimes w) = c_{V \otimes W}((\psi \otimes \chi) \otimes (v \otimes w)) :$$

in particular we get that the image of the multiplication $\text{End}(V(\lambda)) \otimes \text{End}(V(\mu)) \rightarrow \mathbb{k}[G]$ is the sum of all $\text{End}(V(\nu))$ with $V(\nu) \subset V(\lambda) \otimes V(\mu)$. We get then the following combinatorial description of $m_{\lambda, \mu}$.

Lemma 1.4 ([4, Lemma 3.4], [8, Lemma 3.1]). *Let $\lambda, \mu \in \Lambda^+$, $\lambda' \in \Pi^+(\lambda)$ and $\mu' \in \Pi^+(\mu)$. Then the image of $\text{End}(V(\lambda')) \times \text{End}(V(\mu')) \subset \Gamma(M, \mathcal{L}_\lambda) \times \Gamma(M, \mathcal{L}_\mu)$ in $\Gamma(M, \mathcal{L}_{\lambda+\mu})$ via $m_{\lambda, \mu}$ is*

$$\bigoplus_{V(\nu) \subset V(\lambda') \otimes V(\mu')} \text{End}(V(\nu))$$

If $\Pi \subset \Lambda^+$ is simple with maximal element λ , define

$$\Omega(\Pi) = \left\{ \nu - n\lambda : V(\nu) \subset \left(\bigoplus_{\mu \in \Pi} V(\mu) \right)^{\otimes n} \right\}.$$

Notice that if $\nu \in \Pi$ then $\nu - \lambda \in \Omega(\Pi)$. Notice also that, if $\Pi_1 \subset \Pi_2$ are simple subsets with the same maximal element, then $\Omega(\Pi_1) \subset \Omega(\Pi_2)$. If $\Pi = \{\mu_1, \dots, \mu_m\}$, for simplicity sometimes we will denote $\Omega(\Pi)$ also by $\Omega(\mu_1, \dots, \mu_m)$.

Remark 1.5. If $V_\Pi \doteq \bigoplus_{\mu \in \Pi} V(\mu)$, then by Lemma 1.4 we have $A_n(\Pi) = \bigoplus_{V(\nu) \subset V_\Pi^{\otimes n}} \text{End}(V(\nu))$. Hence by the description of $\mathbb{k}[X_\Pi^\circ]$ we get the following identification

$$\Omega(\Pi) \simeq \{(\theta, \theta^*) : \theta \in \Omega(\Pi)\} = \{B \times B^- \text{-weights in } \mathbb{k}[X_\Pi^\circ]\},$$

and in particular $\Omega(\Pi)$ is a semigroup of $\mathbb{Z}\Delta$ respect to the addition.

Since $BB^- \subset G$ is an open subset, it follows that G is a spherical $G \times G$ variety and every non-zero $B \times B^-$ -semiinvariant function $\phi \in \mathbb{k}(G)$ is uniquely determined by its weight up to a scalar factor. Following the general theory of spherical varieties (see for instance [9]), the semigroup $\Omega(\Pi)$ encodes a lot of information on the geometry of X_Π . In particular, we may characterize the existence of an equivariant morphism $X_\Pi \rightarrow X_{\Pi'}$ in terms of the semigroups $\Omega(\Pi)$ and $\Omega(\Pi')$ and of the isotypic decomposition of the tensor powers of $\bigoplus_{\mu \in \Pi} V(\mu)$.

Proposition 1.6. *Let Π, Π' be simple subsets with resp. maximal elements λ, λ' and suppose that $\text{Supp}(\lambda) = \text{Supp}(\lambda')$. There exists a $G \times G$ -equivariant morphism $X_\Pi \rightarrow X_{\Pi'}$ if and only if, for every $\mu' \in \Pi'$, they exist $\mu_1, \dots, \mu_n \in \Pi$ such that $V(\mu' - \lambda' + n\lambda) \subset V(\mu_1) \otimes \dots \otimes V(\mu_n)$.*

Proof. Identify the open orbits $(G \times G)[\text{Id}_\Pi] \subset X_\Pi$ and $(G \times G)[\text{Id}_{\Pi'}] \subset X_{\Pi'}$ with the same quotient of G_{ad} , say G_1 . Since $\text{Supp}(\lambda) = \text{Supp}(\lambda')$, it follows that $G_1 \cap X_\Pi^\circ$ and $G_1 \cap X_{\Pi'}^\circ$ are both identified with the same open subset $G_1^\circ \subset G_1$. On the other hand, since $X_\Pi^\circ \subset X_\Pi$ and $X_{\Pi'}^\circ \subset X_{\Pi'}$ intersect the respective closed orbits, they intersect every orbit, hence $X_\Pi = (G \times G)X_\Pi^\circ$ and $X_{\Pi'} = (G \times G)X_{\Pi'}^\circ$. Therefore the identity on G_1 extends to an (equivariant) morphism $X_\Pi \rightarrow X_{\Pi'}$ if and only if the identity on G_1° extends to a morphism $X_\Pi^\circ \rightarrow X_{\Pi'}^\circ$ if and only if $\mathbb{k}[X_{\Pi'}^\circ] \subset \mathbb{k}[X_\Pi^\circ]$.

By Lemma 1.3, the coordinate ring $\mathbb{k}[X_\Pi^\circ]$ is generated as $\mathfrak{g} \oplus \mathfrak{g}$ -algebra by $\mathbb{k}[X_\lambda^\circ]$ together with the set $\{\phi_\mu/\phi_\lambda\}_{\mu \in \Pi}$, while $\mathbb{k}[X_{\Pi'}^\circ]$ is generated by $\mathbb{k}[X_{\lambda'}^\circ]$ together with the set $\{\phi_{\mu'}/\phi_{\lambda'}\}_{\mu' \in \Pi'}$. On the other hand, by Proposition 1.1 it follows that $\mathbb{k}[X_{\lambda'}^\circ] \simeq \mathbb{k}[X_\lambda^\circ]$, therefore there exists an equivariant morphism $X_\Pi \rightarrow X_{\Pi'}$ if and only if $\phi_{\mu'}/\phi_{\lambda'} \in \mathbb{k}[X_\Pi^\circ]$ for every $\mu' \in \Pi'$. By the description of $\mathbb{k}[X_\Pi^\circ]$ in terms of the projective coordinates of X_Π , it follows that $\phi_{\mu'}/\phi_{\lambda'} \in \mathbb{k}[X_\Pi^\circ]$ if and only if there exists $n \in \mathbb{N}$ and $\phi \in A_n(\Pi) = \left(\bigoplus_{\mu \in \Pi} \text{End}(V(\mu))\right)^n$ such that $\phi_{\mu'}/\phi_{\lambda'} = \phi/\phi_\lambda^n$. On the other hand, ϕ has to be $B \times B^-$ -semiinvariant, hence the claim follows by Lemma 1.4. \square

In terms of the semigroup $\Omega(\Pi)$, we may reformulate previous proposition as follows.

Corollary 1.7. *Let Π, Π' be simple subsets with resp. maximal elements λ, λ' and assume that $\text{Supp}(\lambda) = \text{Supp}(\lambda')$. There exists a $G \times G$ -equivariant morphism $X_\Pi \rightarrow X_{\Pi'}$ if and only if $\Omega(\Pi') \subset \Omega(\Pi)$ if and only if $\mu' - \lambda' \in \Omega(\Pi)$ for all $\mu' \in \Pi'$.*

Proof. If $X_\Pi \rightarrow X_{\Pi'}$, then in particular we have $X_\Pi^\circ \rightarrow X_{\Pi'}^\circ$: hence by Remark 1.5 it follows that $\Omega(\Pi') \subset \Omega(\Pi)$ and we get $\mu' - \lambda' \in \Omega(\Pi)$ for all $\mu' \in \Pi'$. Suppose conversely that $\mu' - \lambda' \in \Omega(\Pi)$ for all $\mu' \in \Pi'$: then X_Π dominates $X_{\Pi'}$ by Proposition 1.6. \square

Definition 1.8. i) Suppose that $\Pi, \Pi' \subset \Lambda^+$ are simple with resp. maximal elements λ, λ' . Then Π and Π' are *equivalent* (and we write $\Pi \sim \Pi'$) if $\text{Supp}(\lambda') = \text{Supp}(\lambda)$ and $\Pi' - \lambda' = \Pi - \lambda$.

- ii) A weight $\mu \in \Pi^+(\lambda)$ is *trivial* if $X_{\lambda, \mu} \simeq X_\lambda$ as $G \times G$ -varieties. We will denote the subset of the trivial weights in $\Pi^+(\lambda)$ by $\Pi_{\text{tr}}^+(\lambda)$.

Remark 1.9. i) By Proposition 1.6, a weight $\mu \in \Pi^+(\lambda)$ is trivial if and only if there exists $n \in \mathbb{N}$ such that $V(\mu + (n-1)\lambda) \subset V(\lambda)^{\otimes n}$. Equivalently, $\mu \in \Pi^+(\lambda)$ is trivial if and only if $\Omega(\lambda, \mu) = \Omega(\lambda)$, if and only if $\mu - \lambda \in \Omega(\lambda)$.

- ii) We may describe the semigroup $\Omega(\lambda)$ in terms of trivial weights as follows:

$$\Omega(\lambda) = \{\mu - n\lambda : \mu \in \Pi_{\text{tr}}^+(n\lambda)\}.$$

Suppose indeed that $\mu \in \Pi_{\text{tr}}^+(n\lambda)$: then by Proposition 1.1 we have $X_{n\lambda} \simeq X_\lambda$, and it follows $\mu - n\lambda \in \Omega(n\lambda) = \Omega(\lambda)$. Conversely, if $\theta \in \Omega(\lambda)$, then $V(n\lambda + \theta) \subset V(\lambda)^{\otimes n}$, hence $n\lambda + \theta \in \Pi_{\text{tr}}^+(n\lambda)$.

- iii) Consider the semigroup

$$\tilde{\Omega}(\lambda) = \{\mu - n\lambda : \mu \in \Pi^+(n\lambda)\}.$$

Since $X_{n\lambda} \simeq X_\lambda$ for all $n > 0$, considering the normalizations $\tilde{X}_{n\lambda} \simeq \tilde{X}_\lambda$ it follows also $X_{\Pi^+(n\lambda)} \simeq X_{\Pi^+(\lambda)}$, hence $\tilde{\Omega}(\lambda) = \Omega(\Pi^+(\lambda))$. On the other hand, for every $\mu \in \Pi^+(\lambda)$ there exists $n \in \mathbb{N}$ such that $V(n\mu) \subset V(\lambda)^{\otimes n}$ (see [1, Lemma 4.9] or [12, Lemma 1]), therefore $\tilde{\Omega}(\lambda) = \Omega(\lambda)_{\mathbb{Q}} \cap \mathbb{Z}\Delta$ is the *saturation* of $\Omega(\lambda)$ in $\mathbb{Z}\Delta$ (where $\Omega(\lambda)_{\mathbb{Q}}$ denotes the cone generated by $\Omega(\lambda)$ in $\mathbb{Q}\Delta$). We say that $\Omega(\Pi)$ is *saturated* in $\mathbb{Z}\Delta$ if $\Omega(\Pi) = \tilde{\Omega}(\lambda)$.

- iv) Suppose that Π is simple with maximal element λ and let $\pi \in \tilde{\Omega}(\lambda)$, then by iii) it exists $n \in \mathbb{N}$ such that $n\pi \in \Omega(\Pi)$. Since $B \times B^-$ -semiinvariant functions in $\mathbb{k}(G)$ are uniquely determined by their weights up to scalar multiples, if $f_\pi, f_{n\pi} \in \mathbb{k}(G_{\text{ad}})^{(B \times B^-)}$ are $B \times B^-$ -semiinvariant functions of weights (π, π^*) and $(n\pi, n\pi^*)$, then by Remark 1.5 we have $f_\pi \in \mathbb{k}[\tilde{X}_\lambda^\circ]^{(B \times B^-)}$ and $f_{n\pi} \in \mathbb{k}[X_\Pi^\circ]^{(B \times B^-)}$. Since they have the same weight, $f_{n\pi}$ and f_π^n are proportional, hence the normality of X_Π implies that $f_\pi \in \mathbb{k}[X_\Pi^\circ]$, i.e. $\pi \in \Omega(\Pi)$. Together with iii), it follows that the normality of X_Π is equivalent to the saturation of $\Omega(\Pi)$ in $\mathbb{Z}\Delta$.

Corollary 1.10. *Let $\Pi, \Pi' \subset \Lambda^+$ be simple subsets with resp. maximal elements λ and λ' .*

- i) *If $\Pi \sim \Pi'$, then $X_\Pi \simeq X_{\Pi'}$ as $G \times G$ -varieties. In particular $\Omega(\Pi) = \Omega(\Pi')$.*
ii) *Assume that Π, Π' both have cardinality 2 and that $X_\Pi \not\simeq X_\lambda$. If $X_{\Pi'} \simeq X_\lambda$ as $G \times G$ -varieties, then $\Pi' \sim \Pi$.*

Proof. i) If $\mu \in \Pi$, let $\mu' \in \Pi'$ be such that $\mu - \lambda = \mu' - \lambda'$. Since $B \times B^-$ eigenfunctions in $\mathbb{k}(G)$ are uniquely determined by their weight up to scalar factors, we have that ϕ_μ/ϕ_λ and $\phi_{\mu'}/\phi_{\lambda'}$ are proportional. If $\Pi \sim \Pi'$ it follows then by Lemma 1.3 that $\mathbb{k}[X_\Pi^\circ] \simeq \mathbb{k}[X_{\Pi'}^\circ]$, and reasoning as in Proposition 1.6 we get an isomorphism of $G \times G$ -varieties $X_\Pi \simeq X_{\Pi'}$. The last claim follows by Corollary 1.7.

ii) Denote $\Pi = \{\lambda, \mu\}$ and $\Pi' = \{\lambda', \mu'\}$ and suppose that $X_{\Pi'} \simeq X_\lambda$. By Proposition 1.2 ii) it follows that $\text{Supp}(\lambda) = \text{Supp}(\lambda')$, since otherwise X_Π and $X_{\Pi'}$ would have non-isomorphic closed orbits. By Proposition 1.6 it follows that $V(\mu' - \lambda' + n\lambda) \subset V(\mu)^{\otimes k} \otimes V(\lambda)^{\otimes n-k}$ for some $n \in \mathbb{N}$ and $k \leq n$, so comparing highest weights on the right and on the left we get that $\mu' - \lambda' \leq k(\mu - \lambda)$. Since $X_{\Pi'} \not\simeq X_\lambda$, by Proposition 1.6 it must be $k > 0$, hence $\mu' - \lambda' \leq \mu - \lambda$. An analogous argument shows that $\mu - \lambda \leq \mu' - \lambda'$, and the claim follows. \square


Remark 1.11. Together with Remark 1.5, previous corollary shows that the set

$$\tilde{\Omega}(\lambda) \setminus \Omega(\lambda) = \{\mu - n\lambda : \mu \in \Pi^+(n\lambda) \setminus \Pi_{\text{tr}}^+(n\lambda)\}$$

classifies the simple linear compactifications X_{Π} such that $\tilde{X}_{\lambda} \rightarrow X_{\Pi} \rightarrow X_{\lambda}$ and $\text{card}(\Pi) = 2$ up to equivariant isomorphism. If indeed X_{Π} is such a compactification and if $X_{\Pi} \not\cong X_{\lambda}$, then it must be $\Pi = \{\lambda', \mu'\}$ for some $\lambda' \in \Lambda^+$ with $\text{Supp}(\lambda') = \text{Supp}(\lambda)$ and some $\mu' \in \Pi^+(\lambda') \setminus \Pi_{\text{tr}}^+(\lambda')$, and by Corollary 1.10 X_{Π} is uniquely determined by the difference $\mu' - \lambda'$. On the other hand up to consider an equivalent simple subset we may assume that $\lambda' = n\lambda$ for some $n \in \mathbb{N}$, therefore $\mu' - \lambda' \in \tilde{\Omega}(\lambda) \setminus \Omega(\lambda)$.

Suppose that $\Pi \subset \Lambda^+$ is simple with maximal element λ . Then, by the isomorphism $\tilde{X}_{\lambda} \simeq X_{\Pi^+(\lambda)}$, Proposition 1.6 yields as well a tensorial criterion of normality for X_{Π} : X_{Π} is normal if and only if, for every $\nu \in \Pi^+(\lambda)$, they exist $\mu_1, \dots, \mu_n \in \Pi$ such that $V(\nu + (n-1)\lambda) \subset V(\mu_1) \otimes \dots \otimes V(\mu_n)$. As shown in [3], this characterization turns out to be equivalent to a combinatorial property of Π .

Definition 1.12 ([3, Def. 2.7]). If $\Delta' \subset \Delta$ is a non-simply laced connected component, order the simple roots in $\Delta' = \{\alpha_1, \dots, \alpha_r\}$ starting from the extreme of the Dynkin diagram of Δ' which contains a long root and denote α_q the first short root in Δ' . If $\lambda \in \Lambda^+$ is such that $\alpha_q \notin \text{Supp}(\lambda)$ and such that $\text{Supp}(\lambda) \cap \Delta'$ contains a long root, denote α_p the last long root which occurs in $\text{Supp}(\lambda) \cap \Delta'$; for instance, if Δ' is not of type G_2 , then the numbering is as follows:



The *little brother* of λ with respect to Δ' is the dominant weight

$$\lambda_{\Delta'}^{\text{lb}} = \lambda - \sum_{i=p}^q \alpha_i = \begin{cases} \lambda - \omega_1 + \omega_2 & \text{if } G \text{ is of type } G_2 \\ \lambda + \omega_{p-1} - \omega_p + \omega_{q+1} & \text{otherwise} \end{cases}$$

where ω_i is the fundamental weight associated to α_i if $1 \leq i \leq r$, while we set $\omega_0 = \omega_{r+1} = 0$. We will denote the set of the little brothers of λ by $\text{LB}(\lambda)$. Notice that $\text{LB}(\lambda)$ is empty if and only if λ satisfies the following condition:

- (\star) For every non-simply laced connected component $\Delta' \subset \Delta$, if $\text{Supp}(\lambda) \cap \Delta'$ contains a long root, then it contains also the short root which is adjacent to a long simple root.

In case Δ is connected and λ does not satisfy (\star), then we set $\lambda^{\text{lb}} = \lambda_{\Delta}^{\text{lb}}$.

Theorem 1.13 ([3, Thm. 2.10]). *Let G be a semisimple group and let $\Pi \subset \Lambda^+$ be simple with maximal element λ . Then the variety X_{Π} is normal if and only if $\text{LB}(\lambda) \subset \Pi$. In particular, X_{λ} is normal if and only if λ satisfies (\star).*

Corollary 1.14. *Suppose that $\lambda \in \Lambda^+$ satisfies (\star). Then $X_{\Pi} \simeq X_{\lambda}$ for every simple subset Π with maximal element λ . In particular $\Pi_{\text{tr}}^+(\lambda) = \Pi^+(\lambda)$.*

Proof. Let $\Pi \subset \Lambda^+$ be simple with maximal element λ . Then the normalization of X_{λ} factors through X_{Π} , so the claim follows by the normality of X_{λ} . \square

1.1. Some remarks on tensor product decompositions. We conclude this section with some explicit results on tensor products that will be needed in the following.

Lemma 1.15. *Let $\lambda_1, \dots, \lambda_n \in \Lambda^+$ and let L be a Levi subgroup of G . Denote $\Lambda_L^+ \subset \Lambda$ the semigroup of dominant weights for L and, for $\pi \in \Lambda_L^+$, denote by $V_L(\pi)$ the simple L -module of highest weight π . If $\mu \in \Lambda_L^+$ is such that $V_L(\mu) \subset V_L(\lambda_1) \otimes \dots \otimes V_L(\lambda_n)$, then $\mu \in \Lambda^+$ and $V(\mu) \subset V(\lambda_1) \otimes \dots \otimes V(\lambda_n)$.*

Proof. Denote $\Delta' \subset \Delta$ the set of simple roots associated to L . Since $\mu \in \Lambda_L^+$, we have $\langle \mu, \alpha^\vee \rangle \geq 0$ for every $\alpha \in \Delta'$. On the other hand, being $V_L(\mu) \subset V_L(\lambda_1) \otimes \dots \otimes V_L(\lambda_n)$, we have $\sum_i \lambda_i - \mu \in \mathbb{N}\Delta'$: hence we get $\langle \mu, \alpha^\vee \rangle \geq \langle \sum_i \lambda_i, \alpha^\vee \rangle$ for every $\alpha \in \Delta \setminus \Delta'$, and it follows that $\mu \in \Lambda^+$ since $\sum_i \lambda_i$ is so.

We now prove the second claim by induction on n , the basis being the case $n = 2$ (see [3, Lemma 2.4]). If \mathfrak{a} is any Lie algebra, denote $\mathfrak{U}(\mathfrak{a})$ the corresponding universal enveloping algebra. Assume $n = 2$ and regard $V_L(\lambda_1) \otimes V_L(\lambda_2) \subset V(\lambda_1) \otimes V(\lambda_2)$. Fix maximal vectors $v_1 \in V(\lambda_1)$ and $v_2 \in V(\lambda_2)$ for B and let $p \in \mathfrak{U}(\mathfrak{l} \cap \mathfrak{u}^-) \otimes \mathfrak{U}(\mathfrak{l} \cap \mathfrak{u}^-)$ be such that $p(v_1 \otimes v_2) \in V_L(\lambda_1) \otimes V_L(\lambda_2)$ is a maximal vector of weight μ for the Borel $B \cap L \subset L$: to prove the claim we only need to show that $p(v_1 \otimes v_2)$ is a maximal vector for B too. If $\alpha \in \Delta'$ then we have $e_\alpha p(v_1 \otimes v_2) = 0$ by hypothesis. On the other hand, if $\alpha \in \Delta \setminus \Delta'$, then e_α commutes with p , since by its definition p is supported only on the f_α 's with $\alpha \in \Delta'$. Since $v_1 \otimes v_2$ is a maximal vector for B , we get then $e_\alpha p(v_1 \otimes v_2) = p e_\alpha(v_1 \otimes v_2) = 0$, therefore $p(v_1 \otimes v_2)$ generates a simple G -module of highest weight μ .

Suppose now $n > 2$ and let $\mu' \in \Lambda_L^+$ be such that $V_L(\mu') \subset V_L(\lambda_1) \otimes \dots \otimes V_L(\lambda_{n-1})$ and $V_L(\mu) \subset V_L(\mu') \otimes V_L(\lambda_n)$. Then $\mu' \in \Lambda^+$ by the first part of the proof, while by the inductive hypothesis we get $V(\mu') \subset V(\lambda_1) \otimes \dots \otimes V(\lambda_{n-1})$ and $V(\mu) \subset V(\mu') \otimes V(\lambda_n)$, so the claim follows. \square

Corollary 1.16. *Let $\lambda \in \Lambda^+$ and let $\mu \in \Pi^+(\lambda)$ be such that $\text{Supp}_\Delta(\lambda - \mu)$ is simply laced regarded as a subset of the vertices of the Dynkin diagram of G . Then $\mu \in \Pi_{\text{tr}}^+(\lambda)$.*

Proof. Denote L the Levi subgroup associated to $\Delta' = \text{Supp}_\Delta(\lambda - \mu)$. By Corollary 1.14 applied to the semisimple part of L it follows that $\mu \in \Pi^+(\lambda)$ is trivial w.r.t. L , hence by Remark 1.9 it follows that there exists $n \in \mathbb{N}$ such that $V_L(\mu + (n-1)\lambda) \subset V_L(\lambda)^{\otimes n}$. Therefore by Lemma 1.15 we get $V(\mu + (n-1)\lambda) \subset V(\lambda)^{\otimes n}$, and by Remark 1.9 i) it follows that $\mu \in \Pi_{\text{tr}}^+(\lambda)$. \square

Let $n \in \mathbb{N}$ and consider the set

$$\text{Tens}_n(G) = \{(\lambda_0, \dots, \lambda_n) \in (\Lambda^+)^{n+1} : V(\lambda_0) \subset V(\lambda_1) \otimes \dots \otimes V(\lambda_n)\}.$$

Following lemma has been proved in several references, usually in the case $n = 2$. Since we will need that, we claim it in a slightly more general form, which is easily reduced to the case $n = 2$ proceeding by induction on n .

Lemma 1.17 ([11, Lemma 3.9]). *The set $\text{Tens}_n(G)$ is a semigroup with respect to the addition.*

An easy application of previous lemma which will be very useful for us is the following.

Corollary 1.18. *Let $\lambda_0, \dots, \lambda_n \in \Lambda^+$ be such that $V(\lambda_0) \subset V(\lambda_1) \otimes \dots \otimes V(\lambda_n)$. Then, for any $\mu \in \Lambda^+$, it also holds $V(\lambda_0 + \mu) \subset V(\lambda_1 + \mu) \otimes V(\lambda_2) \otimes \dots \otimes V(\lambda_n)$.*

Corollary 1.19. *Let $\lambda, \mu \in \Lambda^+$ and let $\nu \in \Pi^+(\mu)$ be such that $\text{Supp}_\Delta(\mu - \nu) \cap \text{Supp}(\lambda) \neq \emptyset$, suppose moreover that $\text{Supp}_\Delta(\mu - \nu)$ is connected and that $\mu - \nu$ is the highest root of the root subsystem generated by $\text{Supp}_\Delta(\mu - \nu)$. Then $V(\lambda + \nu) \subset V(\mu) \otimes V(\lambda)$.*

Proof. Denote L the Levi subgroup associated to $\text{Supp}_\Delta(\mu - \nu)$ and denote \mathfrak{l} its Lie algebra. Consider $\mu - \nu$: by the assumption on $\mu - \nu$, we have an isomorphism of \mathfrak{l} -modules $V_L(\mu - \nu) \simeq \mathfrak{l}$. Since $\text{Supp}_\Delta(\mu - \nu) \cap \text{Supp}(\lambda) \neq \emptyset$, the \mathfrak{l} -action induces a non-zero morphism $V_L(\mu - \nu) \otimes V_L(\lambda) \rightarrow V_L(\lambda)$ which is surjective by irreducibility, hence we get $V_L(\lambda) \subset V_L(\mu - \nu) \otimes V_L(\lambda)$. By Corollary 1.18 this implies $V_L(\lambda + \nu) \subset V_L(\mu) \otimes V_L(\lambda)$, and the claim follows applying Lemma 1.15. \square

We now describe an explicit result which we will need in the special case $G = \text{Spin}(2r + 1)$, which we will treat in the rest of the paper. Set $\Delta = \{\alpha_1, \dots, \alpha_r\}$ and denote $\omega_1, \dots, \omega_r$ the fundamental weights. For convenience, we also denote $\varpi_k = \sum_{j=1}^{k-1} j\alpha_j + k \sum_{j=k}^r \alpha_j$: therefore if $0 < k < r$ we have $\varpi_k = \omega_k$, whereas $\varpi_0 = 0$ and $\varpi_r = 2\omega_r$.

Lemma 1.20. *Let $G = \text{Spin}(2r + 1)$. Let $\lambda, \mu \in \Lambda^+$ and $\nu \in \Pi^+(\mu)$.*

- i) *If $\mu - \nu \in \Phi_l^+(\lambda)$, then $V(\lambda + \nu) \subset V(\mu) \otimes V(\lambda)$.*
- ii) *If $\alpha_r \in \text{Supp}(\lambda + \nu)$ and if $\mu - \nu \in \Phi^+(\lambda)$, then $V(\lambda + \nu) \subset V(\mu) \otimes V(\lambda)$.*
- iii) *If $\text{Supp}(\lambda) \neq \{\alpha_r\}$ and if $\mu - \nu \in 2\Phi_s^+(\lambda)$, then $V(\lambda + \nu) \subset V(\mu) \otimes V(\lambda)$.*

Proof. Denote $\theta = \mu - \nu$ and set $\text{Supp}_\Delta(\theta) = \{\alpha_{p+1}, \dots, \alpha_q\}$, where $0 \leq p < q \leq r$.

i) Notice that it holds one of the followings:

- $q < r$ and $\theta = \sum_{i=p+1}^q \alpha_i$;
- $q = r$ and $\theta = \sum_{i=p+1}^k \alpha_i + 2 \sum_{i=k+1}^r \alpha_i$ for some k with $p < k < r$.

Suppose that we are in the first case: then θ is the highest root of the subsystem generated by $\text{Supp}_\Delta(\theta)$ and the claim follows by Corollary 1.19. Suppose that we are in the second case: then we have $\mu = \nu - \varpi_p + \varpi_{p+1} - \varpi_k + \varpi_{k+1}$. Since μ is dominant, it must be $\alpha_p, \alpha_k \in \text{Supp}(\nu)$. Notice also that by Lemma 1.15 we may assume that $\text{Supp}_\Delta(\theta) = \Delta$, i.e. $p = 0$. Let $\alpha_j \in \text{Supp}(\lambda)$: then by applying Corollary 1.18 twice (first with $\nu - \omega_k$ and then with $\lambda - \omega_j$) we are reduced to the following inclusion, which can be checked directly:

- If $1 \leq j \leq r$ and $1 \leq k \leq r - 1$, then $V(\varpi_k + \varpi_j) \subset V(\varpi_j) \otimes V(\varpi_1 + \varpi_{k+1})$.

ii) By part i), we only need to consider the case where θ is a positive short root. Notice that $\theta = \sum_{i=p+1}^r \alpha_i$ for some p with $0 \leq p < r$. By Lemma 1.15 we may assume $p = 0$, so we have that $\mu = \nu + \omega_1$. Suppose that $\alpha_r \in \text{Supp}(\lambda)$: then by applying Corollary 1.18 twice (first with ν and then with $\lambda - \omega_r$) we are reduced to the following inclusion, which can be checked directly:

- $V(\omega_r) \subset V(\omega_1) \otimes V(\omega_r)$;

Suppose now that $\alpha_r \in \text{Supp}(\nu)$ and let $\alpha_j \in \text{Supp}(\lambda)$: then by applying Corollary 1.18 twice (first with $\nu - \omega_r$ and then with $\lambda - \omega_j$) we are reduced to the following inclusion, which can be checked directly:

- If $1 \leq j \leq r$, then $V(\omega_j + \omega_r) \subset V(\omega_j) \otimes V(\omega_1 + \omega_r)$.

iii) Notice that $\theta = 2 \sum_{i=p+1}^r \alpha_i$ for some p with $0 \leq p < r$. By Lemma 1.15 we may assume $p = 0$, so we have that $\mu = \nu + 2\omega_1$. Let $\alpha_j \in \text{Supp}(\lambda)$ with $j < r$: then by applying Corollary 1.18 twice (first with ν and then with $\lambda - \omega_j$) we are reduced to the following inclusion, which can be checked directly:

- If $1 \leq j < r$, then $V(\omega_j) \subset V(2\omega_1) \otimes V(\omega_j)$. \square

2. TRIVIAL WEIGHTS IN THE ODD ORTHOGONAL CASE

From now on we will suppose $G = \text{Spin}(2r + 1)$. Set $\Delta = \{\alpha_1, \dots, \alpha_r\}$ and denote $\omega_1, \dots, \omega_r$ the fundamental weights. For convenience, if $0 \leq k \leq r$ we also denote $\varpi_k = \sum_{j=1}^{k-1} j\alpha_j + k \sum_{j=k}^r \alpha_j$: therefore $\varpi_k = \omega_k$ if $0 < k < r$, whereas $\varpi_0 = 0$ and $\varpi_r = 2\omega_r$. If $\lambda \in \Lambda^+$ is non-zero we denote by $q(\lambda)$ the maximum such that $\langle \lambda, \alpha_{q(\lambda)}^\vee \rangle \neq 0$, while we set $q(\lambda) = 0$ if $\lambda = 0$. We are going to prove the following combinatorial characterization of trivial weights, the rest of the section will be devoted to its proof.

Theorem 2.1. *Suppose that $G = \text{Spin}(2r + 1)$. Let $\lambda \in \Lambda^+$, $\mu \in \Pi^+(\lambda)$ and denote $\lambda - \mu = \sum_{i=1}^r a_i \alpha_i$. Then $\mu \in \Pi_{\text{tr}}^+(\lambda)$ if and only if a_r is even or $a_r > 2 \min\{r - q(\lambda), r - q(\mu)\}$.*

Remark 2.2. Let $\lambda \in \Lambda^+$ and $\mu \in \Pi^+(\lambda)$. Set $\theta = \lambda - \mu \in \mathbb{N}\Delta$, say $\theta = \sum_{i=1}^r a_i \alpha_i$, and denote $l(\theta) \leq r$ the minimum such that $a_i = a_r$ for every $i \geq l(\theta)$. Since μ is dominant, it follows that $\text{Supp}(\theta^+) \subset \text{Supp}(\lambda)$. If $q(\lambda) < i < r$, then we have $a_{i-1} - 2a_i + a_{i+1} = \langle \mu, \alpha_i^\vee \rangle \geq 0$, whereas if $q(\lambda) < r$ then $2a_{r-1} - 2a_r = \langle \mu, \alpha_r^\vee \rangle \geq 0$. In particular this implies $a_{q(\lambda)} \geq a_{q(\lambda)+1} \geq \dots \geq a_r$ and it follows that $\max\{q(\lambda), q(\mu)\} = \max\{l(\theta), q(\lambda)\}$.

By Corollary 1.10, the triviality of μ depends only on the equivalence class of the simple subset $\{\lambda, \mu\}$. Therefore we may restate previous theorem as follows, not dealing with λ but just with the semigroup $\Omega(\lambda)$, which depends only on $\text{Supp}(\lambda)$.

Corollary 2.3. *Suppose that $G = \text{Spin}(2r + 1)$ and let $\lambda \in \Lambda^+$. Then*

$$-\Omega(\lambda) = \left\{ \theta \doteq \sum_{i=1}^r a_i \alpha_i \in \mathbb{N}\Delta : \begin{array}{l} \text{Supp}(\theta^+) \subset \text{Supp}(\lambda) \text{ and} \\ a_r \text{ is even or } a_r > 2 \min\{r - l(\theta), r - q(\lambda)\} \end{array} \right\}$$

where $l(\theta) \leq r$ denotes the minimum such that $a_i = a_r$ for every $i \geq l(\theta)$.

Proof. By Remark 1.9 ii) we have $-\Omega(\lambda) = \{n\lambda - \mu : \mu \in \Pi_{\text{tr}}^+(n\lambda)\}$. Let $\theta = \sum_{i=1}^r a_i \alpha_i \in \mathbb{N}\Delta$. If $\theta \in -\Omega(\lambda)$, then by previous theorem together with Remark 2.2 we get that a_r is even or that $a_r > 2 \min\{r - l(\theta), r - q(\lambda)\}$. Conversely if $\text{Supp}(\theta^+) \subset \text{Supp}(\lambda)$ then $n\lambda - \theta$ is dominant for some $n \in \mathbb{N}$, and if moreover a_r is even or $a_r > 2 \min\{r - l(\theta), r - q(\lambda)\} = 2 \min\{r - q(n\lambda), r - q(n\lambda - \theta)\}$, then by previous theorem we have $n\lambda - \theta \in \Pi_{\text{tr}}^+(n\lambda)$, hence $\theta \in -\Omega(\lambda)$. \square

Remark 2.4. Suppose that $\text{Supp}(\lambda) = \{\alpha_{r-1}\}$. Then previous corollary implies that $\text{SO}(2r + 1)$ admits a unique non-normal linear compactification X such that $\tilde{X}_\lambda \rightarrow X \rightarrow X_\lambda$, namely X_λ . If indeed $\mu = \lambda - \sum_{i=1}^r a_i \alpha_i \in \Pi^+(\lambda) \setminus \Pi_{\text{tr}}^+(\lambda)$, then by Corollary 2.3 it must be $a_{r-1} = a_r = 1$, and it follows $a_1 = \dots = a_{r-2} = 0$.

We now prove Theorem 2.1, the proof will be split in several lemmas. If $\alpha_r \in \text{Supp}(\lambda)$, then by Corollary 1.14 we have $\Pi_{\text{tr}}^+(\lambda) = \Pi^+(\lambda)$. Therefore throughout this section we will assume that $\alpha_r \notin \text{Supp}(\lambda)$. First we will prove that the conditions are necessary (Corollary 2.7). A basic case is that of the first fundamental weight, treated in the following proposition, where we deduce the isotypic decomposition of the tensor powers of the the standard representation by the Schur-Weyl duality (see for instance [7, Appendix F]).

Proposition 2.5. *Suppose that $G = \text{Spin}(2r + 1)$ and let $n \in \mathbb{N}$. If $\mu \in \Pi^+(n\omega_1)$, denote $n\omega_1 - \mu = \sum_{i=1}^r a_i \alpha_i$. Then $V(\mu) \subset V(\omega_1)^{\otimes n}$ if and only if a_r is even or $a_r > 2(r - q(\mu))$.*

Proof. Embed standardly $\mathrm{SO}(2r+1)$ in $\mathrm{GL}(2r+1)$ and denote $\mathfrak{h} \subset \tilde{\mathfrak{h}}$ the respective Cartan subalgebras of diagonal matrices. Denote $\varepsilon_1, \dots, \varepsilon_{2r+1}$ the basis of $\tilde{\mathfrak{h}}^*$ defined by $\varepsilon_i(\mathrm{diag}(a_1, \dots, a_{2r+1})) = a_i$ and, for any weight $\lambda = \sum_{i=1}^r \lambda_i \varepsilon_i$, denote $|\lambda| = \sum_{i=1}^r \lambda_i$. With respect to this basis μ is expressed as follows

$$\mu = (n - a_1)\varepsilon_1 + \sum_{i=2}^r (a_{i-1} - a_i)\varepsilon_i.$$

By the Schur-Weyl duality it follows that $V(\mu) \subset V(\omega_1)^{\otimes n}$ if and only if μ extends to a dominant weight $\tilde{\mu} = \sum_{i=1}^{2r+1} \tilde{\mu}_i \varepsilon_i \in \tilde{\mathfrak{h}}^*$ such that

$$\begin{cases} |\tilde{\mu}| \leq n \\ |\tilde{\mu}| \equiv n \pmod{2} \\ \tilde{\mu}_1^t + \tilde{\mu}_2^t \leq 2r + 1 \end{cases}$$

where $\tilde{\mu}^t = (\tilde{\mu}_1^t, \dots, \tilde{\mu}_{\tilde{\mu}_1}^t)$ denotes the transposed of $\tilde{\mu} = (\tilde{\mu}_1, \dots, \tilde{\mu}_{2r+1})$ regarded as a partition of $|\tilde{\mu}|$.

Suppose that $V(\mu) \subset V(\omega_1)^{\otimes n}$ and let $\tilde{\mu} \in \tilde{\mathfrak{h}}^*$ be an extension of μ as above. Then either $\tilde{\mu}_i = 0$ for every $i > r$ or

$$\tilde{\mu} = \sum_{i=1}^{\tilde{\mu}_2^t} \tilde{\mu}_i \varepsilon_i + \sum_{i=\tilde{\mu}_2^t+1}^{\tilde{\mu}_1^t} \varepsilon_i$$

with $\tilde{\mu}_i \geq 2$ for $i \leq \tilde{\mu}_2^t$. Suppose that $\tilde{\mu}$ is of the first type: then a_r is even since $a_r = n - |\mu| = n - |\tilde{\mu}| \equiv 0 \pmod{2}$. If instead $\tilde{\mu}$ is of the second type, then $\tilde{\mu}_1^t + \tilde{\mu}_2^t \leq 2r + 1$ implies $q(\mu) = 2r + 1 - \tilde{\mu}_1^t$ and we get $a_r > 2(r - q(\mu))$ since

$$a_r = n - |\mu| = n - |\tilde{\mu}| + 2(\tilde{\mu}_1^t - r - 1) + 1 = n - |\tilde{\mu}| + 2(r - q(\mu)) + 1.$$

Suppose conversely that a_r is even or that $a_r > 2(r - q(\mu))$, let's show that $V(\mu) \subset V(\omega_1)^{\otimes n}$. Define the weight $\tilde{\mu} \in \tilde{\mathfrak{h}}^*$ as follows:

$$\tilde{\mu} = \begin{cases} \sum_{i=1}^{q(\mu)} \mu_i \varepsilon_i & \text{if } a_r \text{ is even} \\ \sum_{i=1}^{q(\mu)} \mu_i \varepsilon_i + \sum_{i=q(\mu)+1}^{2r-q(\mu)+1} \varepsilon_i & \text{if } a_r > 2(r - q(\mu)) \text{ is odd} \end{cases}$$

Then $\tilde{\mu}$ satisfies the conditions of the Schur-Weyl duality and it follows $V(\mu) \subset V(\omega_1)^{\otimes n}$. \square

Following lemma will allow us to deduce the necessity of the conditions in Theorem 2.1 from the case $\mathrm{Supp}(\lambda) = \{\alpha_1\}$.

Lemma 2.6. *Let $\lambda \in \Lambda^+$ and $\mu \in \Pi_{\mathrm{tr}}^+(\lambda)$. Then $\mathbb{k}[X_\mu^\circ] \subset \mathbb{k}[X_\lambda^\circ]_{(\phi_\mu/\phi_\lambda)}$. In particular $\Omega(\mu) \subset \Omega(\lambda)_{\lambda-\mu}$, where the latter denotes the semigroup generated in $\mathbb{Z}\Delta$ by $\Omega(\lambda)$ together with $\lambda - \mu$.*

Proof. Since $X_\lambda \simeq X_{\lambda, \mu}$, it follows that X_λ is endowed with a linearized ample line bundle \mathcal{L} possessing a $B \times B^-$ -semiinvariant section s_μ of weight (μ, μ^*) which generates a submodule of $\Gamma(X_\lambda, \mathcal{L})$ isomorphic to $\mathrm{End}(V(\mu))$. Correspondingly we get a rational application $X_\lambda \dashrightarrow X_\mu$ which is regular in the affine set $(X_\lambda^\circ)_{(\phi_\mu/\phi_\lambda)} \subset X_\lambda^\circ$ defined by the non-vanishing of $\phi_\mu/\phi_\lambda \in \mathbb{k}[X_\lambda^\circ]$, and it follows that $\mathbb{k}[X_\mu^\circ] \subset \mathbb{k}[X_\lambda^\circ]_{(\phi_\mu/\phi_\lambda)}$. The second claim follows by the first one applying Remark 1.5. \square

Corollary 2.7. *Suppose that $G = \mathrm{Spin}(2r + 1)$ and let $\lambda \in \Lambda^+$ be such that $\alpha_r \notin \mathrm{Supp}(\lambda)$. Let $\mu \in \Pi_{\mathrm{tr}}^+(\lambda)$ and denote $\lambda - \mu = \sum_{i=1}^r a_i \alpha_i$, then either a_r is even or $a_r > 2 \min\{r - q(\lambda), r - q(\mu)\}$.*

Proof. Since $\alpha_r \notin \text{Supp}(\lambda)$, we have $\lambda \in \mathbb{Z}\Delta$ and there exists $n > 0$ such that $\lambda \leq n\omega_1$ with $\text{Supp}_\Delta(n\omega_1 - \lambda) \subset \{\alpha_1, \dots, \alpha_{q(\lambda)-1}\}$. Since $\alpha_r \notin \text{Supp}_\Delta(n\omega_1 - \lambda)$, we have $V(\lambda) \subset V(\omega_1)^{\otimes n}$, hence $\lambda - n\omega_1 \in \Omega(\omega_1)$ by Remark 1.9 ii). On the other hand by Corollary 1.7 we have $\Omega(\omega_1) = \Omega(n\omega_1)$, hence $\lambda \in \Pi_{\text{tr}}^+(n\omega_1)$ by Remark 1.9 i). By Lemma 2.6 we get then $\mu - \lambda \in \Omega(\lambda) \subset \Omega(\omega_1)_{n\omega_1 - \lambda}$, hence by Remark 1.9 ii) they exist $k, m \in \mathbb{N}$ and $\mu' \in \Pi_{\text{tr}}^+(m\omega_1)$ such that $\mu - \lambda = \mu' - m\omega_1 + k(n\omega_1 - \lambda)$.

Denote $m\omega_1 - \mu' = \sum a'_i \alpha_i$. By the definition of n it follows that $a_i = a'_i$ for all $i \geq q(\lambda)$: hence by Proposition 2.5 either a_r is even or $a_r > 2 \min\{r-1, r-q(\mu')\}$. If $\mu' = 0$, then $q(\mu') = 0$, hence $a_r > 2(r-1) \geq 2(r-q(\lambda))$. Suppose instead $\mu' \neq 0$ and assume that $q(\mu') > q(\lambda)$: then it must be $q(\mu') = q(\mu)$, therefore we have $2(r-q(\mu')) \geq 2 \min\{r-q(\lambda), r-q(\mu)\}$. \square

We now show that the conditions of Theorem 2.1 are sufficient. We distinguish three different cases:

- i) $a_{r-1} \neq a_r$, i.e. $\alpha_r \in \text{Supp}(\mu)$ (Lemma 2.8).
- ii) $a_{r-1} = a_r$ is even (Lemma 2.9).
- iii) $a_{r-1} = a_r > 2 \min\{r-q(\lambda), r-q(\mu)\}$ is odd (Lemma 2.10).

Lemma 2.8. *Let $\lambda \in \Lambda^+$ be such that $\alpha_r \notin \text{Supp}(\lambda)$ and let $\mu \in \Pi^+(\lambda)$. If $\alpha_r \in \text{Supp}(\mu)$, then $\mu \in \Pi_{\text{tr}}^+(\lambda)$.*

Proof. We proceed by induction on $a_{r-1}a_r$. Suppose that either $a_{r-1} = 0$ or $a_r = 0$: then $\text{Supp}_\Delta(\lambda - \mu)$ has all components of type A and the claim follows by Corollary 1.16. Suppose now that a_{r-1} and a_r are both non-zero and denote $p < r-1$ the maximum such that $a_p = 0$, or set $p = 0$ if $a_i \neq 0$ for all i . Define

$$\mu' = \mu + \sum_{i=p+1}^r \alpha_i = \mu - \varpi_p + \varpi_{p+1}.$$

Notice that μ' is dominant: if indeed $p > 0$, then $\alpha_p \in \text{Supp}(\mu)$ since $\langle \mu, \alpha_p^\vee \rangle \geq \langle \lambda, \alpha_p^\vee \rangle + a_{p+1} > 0$. Therefore $\mu' \in \Pi^+(\lambda)$ and by construction we have $\mu < \mu'$. Hence by Lemma 1.20 ii) it follows $V(\lambda + \mu) \subset V(\lambda) \otimes V(\mu')$ and we get $\mu - \lambda \in \Omega(\lambda, \mu')$.

Consider now μ' and denote $\lambda - \mu' = \sum a'_i \alpha_i$: then $\alpha_r \in \text{Supp}(\mu')$ and $a'_{r-1}a'_r < a_{r-1}a_r$, so by the inductive hypothesis it follows that $\mu' \in \Pi_{\text{tr}}^+(\lambda)$ and by Remark 1.9 we get $\Omega(\lambda, \mu') = \Omega(\lambda)$. It follows then $\mu - \lambda \in \Omega(\lambda)$, i.e. $\mu \in \Pi_{\text{tr}}^+(\lambda)$. \square

Lemma 2.9. *Let $\lambda \in \Lambda^+$ be such that $\alpha_r \notin \text{Supp}(\lambda)$, let $\mu \in \Pi^+(\lambda)$ and denote $\lambda - \mu = \sum_{i=1}^r a_i \alpha_i$. If $a_{r-1} = a_r$ is even, then $\mu \in \Pi_{\text{tr}}^+(\lambda)$.*

Proof. By Corollary 1.10, the triviality of μ depends only on the equivalence class of the simple set $\{\lambda, \mu\}$. Hence we may replace the simple subset $\{\lambda, \mu\}$ with the equivalent simple subset $\{\lambda + \omega_{q(\lambda)}, \mu + \omega_{q(\lambda)}\}$, in particular we may assume that $\alpha_{q(\lambda)} \in \text{Supp}(\mu)$.

We proceed by induction on a_r . Suppose that $a_r = 0$: then $\text{Supp}_\Delta(\lambda - \mu)$ has all components of type A and the claim follows by Corollary 1.16. Suppose now that $a_{r-1} = a_r \geq 2$. Since $q(\lambda) < r$ and since μ is dominant, it must be $a_{q(\lambda)} \geq a_{q(\lambda)+1} \geq \dots \geq a_{r-1} = a_r \geq 2$. Denote p the maximum such that $a_p = 0$, or set $p = 0$ if $a_i \neq 0$ for all i , and define

$$\mu' = \mu + \sum_{i=p+1}^{q(\lambda)} \alpha_i + \sum_{i=q(\lambda)+1}^r 2\alpha_i = \mu - \varpi_p + \varpi_{p+1} - \varpi_{q(\lambda)} + \varpi_{q(\lambda)+1}.$$

Notice that μ' is dominant: while $\alpha_{q(\lambda)} \in \text{Supp}(\mu)$ by the assumption at beginning of the proof, if $p > 0$ we have also $\alpha_p \in \text{Supp}(\mu)$ since $\langle \mu, \alpha_p^\vee \rangle \geq \langle \lambda, \alpha_p^\vee \rangle + a_{p+1} > 0$. Therefore $\mu' \in \Pi^+(\lambda)$ and by construction we have $\mu < \mu'$. Hence by Lemma 1.20 i) and iii) we get $V(\lambda + \mu) \subset V(\lambda) \otimes V(\mu')$, which implies $\mu - \lambda \in \Omega(\lambda, \mu')$.

Consider now μ' and denote $\lambda - \mu' = \sum a'_i \alpha_i$: then either $q(\lambda) = r - 1$ and $\alpha_r \in \text{Supp}(\mu')$ or $a'_{r-1} = a'_r = a_r - 2$. It follows that $\mu' \in \Pi_{\text{tr}}^+(\lambda)$, in the first case by Lemma 2.8 and in the second case by inductive hypothesis. Therefore by Remark 1.9 we get $\Omega(\lambda, \mu') = \Omega(\lambda)$ and it follows $\mu - \lambda \in \Omega(\lambda)$, i.e. $\mu \in \Pi_{\text{tr}}^+(\lambda)$. \square

Lemma 2.10. *Let $\lambda \in \Lambda^+$ be such that $\alpha_r \notin \text{Supp}(\lambda)$, let $\mu \in \Pi^+(\lambda)$ and denote $\lambda - \mu = \sum_{i=1}^r a_i \alpha_i$. If $a_r > 2 \min\{r - q(\lambda), r - q(\mu)\}$, then $\mu \in \Pi_{\text{tr}}^+(\lambda)$.*

Proof. By Corollary 1.10, the triviality of μ depends only on the equivalence class of the simple set $\{\lambda, \mu\}$. Hence we may replace the simple subset $\{\lambda, \mu\}$ with the equivalent simple subset $\{\lambda + \omega_{q(\lambda)}, \mu + \omega_{q(\lambda)}\}$, in particular we may assume that $q(\lambda) \leq q(\mu)$. Moreover by Lemma 2.9 we may assume that a_r is odd.

We proceed by induction on $r - q(\mu)$, the basis being the case $q(\mu) = r$ (Lemma 2.8). Suppose that $q(\mu) < r$: since μ is dominant and since $a_r > 2(r - q(\mu))$, it must be

$$a_{q(\lambda)} \geq a_{q(\lambda)+1} \geq \dots \geq a_{q(\mu)} = a_{q(\mu)+1} = \dots = a_r \geq 3.$$

Denote p the maximum such that $a_p = 0$, or set $p = 0$ otherwise, and define

$$\mu' = \mu + \sum_{i=p+1}^{q(\mu)} \alpha_i + \sum_{i=q(\mu)+1}^r 2\alpha_i = \mu - \varpi_p + \varpi_{p+1} - \varpi_{q(\mu)} + \varpi_{q(\mu)+1}.$$

Notice that μ' is dominant: while $\alpha_{q(\mu)} \in \text{Supp}(\mu)$ by definition, if $p > 0$ we have also $\alpha_p \in \text{Supp}(\mu)$ since $\langle \mu, \alpha_p^\vee \rangle \geq \langle \lambda, \alpha_p^\vee \rangle + a_{p+1} > 0$. Therefore $\mu' \in \Pi^+(\lambda)$ and by construction we have $\mu < \mu'$. Hence by Lemma 1.20 i), iii) we get $V(\lambda + \mu) \subset V(\lambda) \otimes V(\mu')$, which implies $\mu - \lambda \in \Omega(\lambda, \mu')$.

Consider now μ' and denote $\lambda - \mu' = \sum a'_i \alpha_i$. Suppose that $a'_r = 1$. Then $a_r = 3$ and we get $q(\mu) = r - 1$, since by hypothesis we have $a_r > 2(r - q(\mu))$ and $q(\mu) < r$: therefore $\alpha_r \in \text{Supp}(\mu')$ and by Lemma 2.8 it follows $\mu' \in \Pi_{\text{tr}}^+(\lambda)$. Otherwise we have $q(\mu') = q(\mu) + 1 < r$ and $a'_r = a_r - 2 > 2(r - q(\mu'))$: therefore $\mu' \in \Pi^+(\lambda)$ still satisfies the hypothesis of the lemma and by the inductive hypothesis we get $\mu' \in \Pi_{\text{tr}}^+(\lambda)$. Therefore by Remark 1.9 it follows $\Omega(\lambda, \mu') = \Omega(\lambda)$ and we get $\mu - \lambda \in \Omega(\lambda)$, i.e. $\mu \in \Pi_{\text{tr}}^+(\lambda)$. \square

3. SIMPLE REDUCED SUBSETS IN THE ODD ORTHOGONAL CASE

Let $\Pi \subset \Lambda^+$ be a simple subset. In this section we will define the *reduction* of Π , which is the minimal simple subset $\Pi_{\text{red}} \subset \Pi$ such that X_Π and $X_{\Pi_{\text{red}}}$ are equivariantly isomorphic. This subset is canonical, in the sense that if $\lambda \in \Pi$ is the maximal element, then the set of differences $\Pi_{\text{red}} - \lambda$ depends only on $\Pi - \lambda$. If moreover $\Pi' \subset \Lambda^+$ is another simple subset such that

$$\begin{array}{ccc} \tilde{X}_\lambda & \longrightarrow & X_{\Pi'} \\ \downarrow & & \downarrow \\ X_\Pi & \longrightarrow & X_\lambda \end{array}$$

the reductions of Π and Π' will allow to characterize combinatorially the existence of an equivariant morphism $X_\Pi \rightarrow X_{\Pi'}$ which makes the diagram commute. In particular, it will follow a combinatorial criterion to establish whether two simple subsets give rise to isomorphic compactifications.

If $\lambda \in \Lambda^+$, denote $\Phi^+(\lambda) \subset \Phi^+$ the set of the positive roots non-orthogonal to λ and consider the following degeneration of the dominance order:

$$\nu \leq^\lambda \mu \quad \text{if and only if} \quad \mu - \nu \in \mathbb{N}\Phi^+(\lambda).$$

Notice that \leq^λ depends only on $\text{Supp}(\lambda)$ and that it coincides with the usual dominance order if λ is a regular weight. The partial order \leq^λ was studied in the general semisimple case by Gandini and Ruzzi in [6], where it is used to characterize the normality of a simple linear compactification of a semisimple group. In particular, there are proved the following properties.

Proposition 3.1 (see [6, Prop. 2.1 and Cor. 2.2]). *Let $\lambda \in \Lambda^+$.*

- i) *If $\mu \in \Pi(\lambda)$, then $\mu \leq^\lambda \lambda$.*
- ii) *Let $\lambda_1, \dots, \lambda_n \in \Pi^+(\lambda)$. If $\mu, \nu \in \Lambda^+$ are such that $V(\nu) \subset V(\mu) \otimes V(\lambda_1) \otimes \dots \otimes V(\lambda_n)$, then $\nu \leq^\lambda \mu + n\lambda$.*

Suppose that $\Pi \subset \Lambda^+$ is simple with maximal element λ and define the *reduction* of Π as follows:

$$\Pi_{\text{red}} \doteq \{\mu \in \Pi \setminus \Pi_{\text{tr}}^+(\lambda) : \mu \text{ is maximal w.r.t. } \leq^\lambda\} \cup \{\lambda\}.$$

If $\Pi = \Pi_{\text{red}}$, then we say that Π is *reduced*. For instance, if $\alpha_r \notin \text{Supp}(\lambda)$ and $\Pi = \Pi^+(\lambda)$, then we have $\Pi_{\text{red}} = \{\lambda, \lambda^{\text{lb}}\}$: this is a consequence of Proposition 3.1 together with Theorem 1.13.

Let $\lambda \in \Lambda^+$ and denote $\Xi(\lambda) \subset \mathbb{N}\Delta$ the semigroup of the elements $\tau = \sum a_i \alpha_i$ which satisfy the following inequalities:

- (λ -C1) If $p \geq 1$ is the minimum such that $\alpha_p \in \text{Supp}(\lambda)$, then $a_1 \leq a_2 \leq \dots \leq a_p$.
- (λ -C2) If $s < t$ are such that $\alpha_s, \alpha_t \in \text{Supp}(\lambda)$ and $\alpha_i \notin \text{Supp}(\lambda)$ for every $s < i < t$, then $\sum_{i=s}^{t-1} |a_{i+1} - a_i| \leq a_s + a_t$.
- (λ -C3) Let $q \leq r$ be the maximum such that $\alpha_q \in \text{Supp}(\lambda)$. If $q < r$, then a_r is even and $2 \sum_{i \in I_q} (a_{i+1} - a_i) \leq a_r$, where $I_q = \{i < r : i \geq q \text{ and } a_i < a_{i+1}\}$.

Remark 3.2. If $\theta \in \Phi_l^+$ is a long root, then they exist $j, k \in \mathbb{N}$ with $j < k < r$ such that $\theta = \sum_{i=j+1}^k \alpha_i$ or $\theta = \sum_{i=j+1}^k \alpha_i + 2 \sum_{i=k+1}^r \alpha_i$, while if $\theta \in \Phi_s^+$ is a short root, then there exists $j < r$ such that $\theta = \sum_{i=j+1}^r \alpha_i$. Let now $\lambda \in \Lambda^+$: then by the description above it follows that $\Phi_l^+(\lambda) \cup 2\Phi_s^+(\lambda) \subset \Xi(\lambda)$, while $\Phi_s^+(\lambda) \subset \Xi(\lambda)$ if and only if $\alpha_r \in \text{Supp}(\lambda)$. Notice that we always have $\Phi_s^+(\lambda) + \Phi_s^+(\lambda) \subset \Xi(\lambda)$: indeed $\Phi_s^+(\lambda) + \Phi_s^+(\lambda) \subset \Phi_l^+(\lambda) \cup 2\Phi_s^+(\lambda)$.

Lemma 3.3. *Let $\lambda \in \Lambda^+$ and $\tau \in \Xi(\lambda)$ be non-zero. There exists $\theta \in \Phi^+(\lambda)$ such that $\tau - \theta \in \Xi(\lambda)$ and $\lambda + \tau^- + \theta \in \Lambda^+$. If moreover $\alpha_r \notin \text{Supp}(\lambda)$, then it is possible to choose $\theta \in \Phi_l^+(\lambda)$.*

Proof. Denote p the minimum such that $\alpha_p \in \text{Supp}(\lambda)$ and q the maximum such that $\alpha_q \in \text{Supp}(\lambda)$ and let $\tau = \tau^+ - \tau^- = \sum a_i \alpha_i \in \Xi(\lambda)$. Denote $s_0 \geq 0$ the minimum such that $\alpha_{s_0} \in \text{Supp}(\lambda) \cap \text{Supp}_\Delta(\tau)$ and notice that $a_i \leq a_{i+1}$ for every $i < s_0$. This follows by (λ -C1) if $s_0 = p$, whereas if $s_0 > p$ then (λ -C2) implies $\sum_{i=1}^{s_0-1} |a_{i+1} - a_i| = a_{s_0} = \sum_{i=1}^{s_0-1} (a_{i+1} - a_i)$, hence $a_{i+1} - a_i \geq 0$ for every $i < s_0$. Denote $j < s_0$ the maximum such that $a_{j+1} \neq 0$: then $0 < a_{j+1} \leq \dots \leq a_{s_0}$. Notice that, if $j > 0$, then $\alpha_j \in \text{Supp}(\tau^-)$: indeed $\langle \tau^-, \alpha_j^\vee \rangle = \langle \tau^+, \alpha_j^\vee \rangle + a_{j+1} > 0$. In order to construct the root θ , we distinguish the following cases:

Case 1. $s_0 < q$;

Case 2. $s_0 = q = r$;

Case 3. $s_0 = q$ and $a_r = 0$;

Case 4. $s_0 = q < r$ and $a_r \neq 0$.

Case 1. Suppose that $s_0 < q$. Denote $t_0 > s_0$ the minimum such that $\alpha_{t_0} \in \text{Supp}(\lambda)$ and define k as follows:

$$k = \begin{cases} t_0 - 1 & \text{if } a_{s_0} \leq a_{s_0+1} \leq \dots \leq a_{t_0} \\ \max\{i < t_0 : a_i > a_{i+1}\} & \text{otherwise} \end{cases}$$

Therefore $j < s_0 \leq k < t_0$. Set $\theta = \sum_{i=j+1}^k \alpha_i$ and denote $\tau' = \tau - \theta$: then by construction $\tau' \in \mathbb{N}\Delta$ and $\theta \in \Phi_l^+(\lambda)$, we claim that $\tau' \in \Xi(\lambda)$. Notice that τ' satisfies $(\lambda\text{-C1})$ and $(\lambda\text{-C3})$ since τ is so, therefore we only need to show that τ' satisfies $(\lambda\text{-C2})$. Denote $\tau' = \sum_{i=1}^r a'_i \alpha_i$ and suppose that $s < t$ are such that $\alpha_s, \alpha_t \in \text{Supp}(\lambda)$ and $\alpha_i \notin \text{Supp}(\lambda)$ for every $s < i < t$. Since otherwise $a'_i = a_i$ for every i with $s \leq i \leq t$, we may assume that either $s \leq j < t$ or $s \leq k < t$: then $a'_s + a'_t = a_s + a_t - 1$ and

$$|a'_{i+1} - a'_i| = \begin{cases} |a_{i+1} - a_i| & \text{if } i \in \{s, \dots, t-1\} \setminus \{j, k\} \\ |a_{i+1} - a_i| - 1 & \text{if } i = j \text{ or } i = k \end{cases}$$

By construction, $\{s, \dots, t-1\}$ cannot contain both j and k . Therefore we get

$$\sum_{i=s}^{t-1} |a'_{i+1} - a'_i| = \sum_{i=s}^{t-1} |a_{i+1} - a_i| - 1 \leq a_s + a_t - 1 = a'_s + a'_t$$

and τ' satisfies $(\lambda\text{-C2})$.

To show that $\lambda + \tau^- + \theta \in \Lambda^+$, notice that $\lambda + \tau^- + \theta = \lambda + \tau^- - \varpi_j + \varpi_{j+1} + \varpi_k - \varpi_{k+1}$. Suppose that $\alpha_{k+1} \notin \text{Supp}(\lambda)$: then by the definition of k we have $a_k > a_{k+1} \leq a_{k+2}$ and it follows $\alpha_{k+1} \in \text{Supp}(\tau^-)$ since

$$\langle \tau^-, \alpha_{k+1}^\vee \rangle = \langle \tau^+, \alpha_{k+1}^\vee \rangle + a_k - 2a_{k+1} + a_{k+2} > 0.$$

Case 2. Suppose that $s_0 = q = r$, denote $\theta = \sum_{i=j+1}^r \alpha_i$ and set $\tau' = \tau - \theta$. Then $\theta \in \Phi_s^+(\lambda)$ and by the definition of j we have $\text{Supp}_\Delta(\tau) = \{\alpha_{j+1}, \dots, \alpha_r\}$, $\text{Supp}(\lambda) \cap \text{Supp}_\Delta(\tau) = \{\alpha_r\}$ and $0 < a_{j+1} \leq \dots \leq a_r$. If moreover $\tau' = \sum a'_i \alpha_i$, then we still have $\text{Supp}_\Delta(\tau') \subset \{\alpha_{j+1}, \dots, \alpha_r\}$, $\text{Supp}(\lambda) \cap \text{Supp}_\Delta(\tau) = \{\alpha_r\}$ and $0 \leq a'_{j+1} \leq \dots \leq a'_r$: therefore $\tau' \in \Xi(\lambda)$ and $\tau^- + \theta = \tau^- - \varpi_j + \varpi_{j+1} \in \Lambda^+$.

Case 3. Suppose that $s_0 = q$ and that $a_r = 0$. Since $a_{s_0} > 0$, it must be $q < r$, hence $(\lambda\text{-C3})$ implies $a_q \geq a_{q+1} \geq \dots \geq a_r = 0$. Denote $k \geq q$ the maximum such that $a_k > 0$: then by the definition of j we get $\text{Supp}_\Delta(\tau) = \{\alpha_{j+1}, \dots, \alpha_k\}$, $\text{Supp}(\lambda) \cap \text{Supp}_\Delta(\tau) = \{\alpha_q\}$ and

$$0 < a_{j+1} \leq \dots \leq a_q \geq \dots \geq a_k > 0.$$

Set $\theta = \sum_{i=j+1}^k \alpha_i$ and $\tau' = \tau - \theta$: then by construction $\theta \in \Phi_l^+(\lambda)$ and $\tau' \in \mathbb{N}\Delta$. If moreover $\tau' = \sum a'_i \alpha_i$, then we still have $\text{Supp}_\Delta(\tau') \subset \{\alpha_{j+1}, \dots, \alpha_k\}$, $\text{Supp}(\lambda) \cap \text{Supp}_\Delta(\tau') = \{\alpha_q\}$ and $0 \leq a'_{j+1} \leq \dots \leq a'_q \geq \dots \geq a'_k \geq 0$: therefore $\tau' \in \Xi(\lambda)$. Consider now $\tau^- + \theta = \tau^- - \varpi_j + \varpi_{j+1} + \varpi_k - \varpi_{k+1}$. Then we have $\langle \tau^-, \alpha_{k+1}^\vee \rangle = \langle \tau^+, \alpha_{k+1}^\vee \rangle + a_k > 0$, therefore $\tau^- + \theta \in \Lambda^+$.

Case 4. Suppose that $s_0 = q < r$ and that $a_r \neq 0$. If $a_i = 0$ for some $i > q$, then $(\lambda\text{-C3})$ implies that $a_r = 0$, hence it must be $a_i \neq 0$ for every $q \leq i \leq r$. Therefore we have $\text{Supp}_\Delta(\tau) =$

$\{\alpha_{j+1}, \dots, \alpha_r\}$ and $\text{Supp}(\lambda) \cap \text{Supp}_\Delta(\tau) = \{\alpha_q\}$. Denote $I_q = \{i < r : i \geq q \text{ and } a_i < a_{i+1}\}$ and define k as follows:

$$k = \begin{cases} q & \text{if } I_q = \emptyset \\ \min\{i \geq q : a_i < a_{i+1}\} & \text{if } I_q \neq \emptyset \end{cases}$$

Therefore we have $j < q \leq k < r$. Notice that $a_i \geq 2$ for every $k < i \leq r$: indeed $a_r \geq 2$ by the definition of $\Xi(\lambda)$, whereas if $I_q \neq \emptyset$ and $a_i = 1$ for some $i > q$ then $(\lambda\text{-C3})$ implies $a_r = 2$ and $I_q = \{k\}$. Therefore, if we set $\theta = \sum_{i=j+1}^k \alpha_i + 2 \sum_{i=k+1}^r \alpha_i$ and $\tau' = \tau - \theta$, then we have $\theta \in \Phi_l^+(\lambda)$ and $\tau' \in \mathbb{N}\Delta$. We claim that $\tau' \in \Xi(\lambda)$. Since $a_{j+1} \leq \dots \leq a_q$, we have that τ' satisfies $(\lambda\text{-C1})$ and $(\lambda\text{-C2})$ as a direct consequence of the fact that these conditions are satisfied by τ . To show that τ' satisfies $(\lambda\text{-C3})$, denote $\tau' = \sum_{i=1}^r a'_i \alpha_i$ and set $I'_q = \{i \geq q : a'_i < a'_{i+1}\}$. If $i \geq q$, notice that we have $a'_{i+1} - a'_i = a_{i+1} - a_i$ unless $i = k$, in which case $a'_{k+1} - a'_k = a_{k+1} - a_k - 1$. Hence we get that

$$2 \sum_{i \in I'_q} (a'_{i+1} - a'_i) = 2 \sum_{i \in I_q} (a_{i+1} - a_i) - 2 \leq a_r - 2 = a'_r,$$

therefore $\tau' \in \Xi(\lambda)$. To show that $\lambda + \tau^- + \theta \in \Lambda^+$, since $a_r \neq 0$, notice that we have $\tau^- + \theta = \tau^- - \varpi_j + \varpi_{j+1} - \varpi_k + \varpi_{k+1}$. If $k \neq q$, then by its definition we have $a_{k-1} \geq a_k < a_{k+1}$. Therefore if $\alpha_k \notin \text{Supp}(\lambda)$ it follows that

$$\langle \tau^-, \alpha_k^\vee \rangle = \langle \tau^+, \alpha_k^\vee \rangle + a_{k-1} + a_{k+1} - 2a_k > 0 :$$

hence $\alpha_k \in \text{Supp}(\tau^-)$ and $\lambda + \tau^- + \theta \in \Lambda^+$. \square

Corollary 3.4. *Let $\lambda \in \Lambda^+$.*

- i) *If $\alpha_r \in \text{Supp}(\lambda)$, then $\Xi(\lambda) = \mathbb{N}\Phi^+(\lambda)$.*
- ii) *If $\alpha_r \notin \text{Supp}(\lambda)$, then $\Xi(\lambda) = \mathbb{N}\Phi_l^+(\lambda)$.*

Proof. By Remark 3.2 we have $\Phi_l^+(\lambda) \subset \Xi(\lambda)$, whereas $\Phi_s^+(\lambda) \subset \Xi(\lambda)$ if and only if $\alpha_r \in \text{Supp}(\lambda)$. Let $\tau \in \Xi(\lambda)$ and denote $\tau = \sum_{i=1}^r a_i \alpha_i$. If $\tau \neq 0$, then by Lemma 3.3 there exists $\tau' \in \Xi(\lambda)$ with $\tau - \tau' \in \Phi^+(\lambda)$, and if $\alpha_r \notin \text{Supp}(\lambda)$ we may assume $\tau - \tau' \in \Phi_l^+(\lambda)$. Therefore the claim follows proceeding by induction on $\sum_{i=1}^r a_i$. \square

Theorem 3.5. *Let $\mu, \nu \in \Pi^+(\lambda) \setminus \Pi_{\text{tr}}^+(\lambda)$. Then $\Omega(\lambda, \nu) \subset \Omega(\lambda, \mu)$ if and only if $\nu \leq^\lambda \mu$.*

Proof. Since otherwise $\Pi_{\text{tr}}^+(\lambda) = \Pi^+(\lambda)$, it must be $\alpha_r \notin \text{Supp}(\lambda)$. Suppose that $\Omega(\lambda, \nu) \subset \Omega(\lambda, \mu)$. In particular we have $\nu - \lambda \in \Omega(\lambda, \mu)$, so it exists $k \leq n$ such that $V((n-1)\lambda + \nu) \subset V(\lambda)^{\otimes k} \otimes V(\mu)^{\otimes n-k}$. On the other hand $\nu \notin \Pi_{\text{tr}}^+(\lambda)$ so by Remark 1.9 i) it must be $k < n$ and by Proposition 3.1 ii) it follows $\nu \leq^\lambda \mu$.

Suppose conversely that $\nu \leq^\lambda \mu$, we will show that $\Omega(\lambda, \nu) \subset \Omega(\lambda, \mu)$ proceeding by induction on the difference $\mu - \nu$. Denote $\lambda - \mu = \sum_{i=1}^r m_i \alpha_i$, $\lambda - \nu = \sum_{i=1}^r n_i \alpha_i$ and set $a_i = m_i - n_i$. By Theorem 2.1 m_r and n_r are both odd integers, so that a_r is even. Denote $\mu - \nu = \sum a_i \alpha_i = \theta_1 + \dots + \theta_n$ with $\theta_i \in \Phi^+(\lambda)$ and let k be the number of short roots which occur in the set $\{\theta_1, \dots, \theta_n\}$. Denote $\theta_i = \sum b_j^i \alpha_j$ and notice that θ_i is short if and only if b_r^i is odd. Since a_r is even, k is even as well: by Remark 3.2 together with Corollary 3.4 it follows then $\mu - \nu \in \mathbb{N}\Phi_l^+(\lambda) = \Xi(\lambda)$. Hence by Lemma 3.3 together with Corollary 3.4 there exists $\theta \in \Phi_l^+(\lambda)$ such that $\mu - \nu - \theta \in \mathbb{N}\Phi_l^+(\lambda)$ and $\lambda + \nu + \theta \in \Lambda^+$.

Denote $\mu' = \lambda + \mu$, $\nu' = \lambda + \nu$ and $\pi = \lambda + \nu + \theta$. Then $\{\lambda, \mu\} \sim \{2\lambda, \mu'\}$ and $\{\lambda, \nu\} \sim \{2\lambda, \nu'\}$ are equivalent simple subsets, so that by Corollary 1.10 i) it follows $\Omega(2\lambda, \mu') = \Omega(\lambda, \mu)$ and

$\Omega(\lambda, \nu) = \Omega(2\lambda, \nu')$. Moreover by Lemma 1.20 i) it follows that $V(2\lambda + \nu') \subset V(2\lambda) \otimes V(\pi)$, hence $\nu' - 2\lambda \in \Omega(2\lambda, \pi)$ and by Corollary 1.7 we get $\Omega(2\lambda, \nu') \subset \Omega(2\lambda, \pi)$. Consider now the weights $\mu', \pi \in \Pi^+(2\lambda)$: then we have $\pi \leq^\lambda \mu'$ and $\mu' - \pi < \mu - \nu$, hence by the inductive hypothesis it follows $\Omega(2\lambda, \pi) \subset \Omega(2\lambda, \mu')$. Therefore we get

$$\Omega(\lambda, \nu) = \Omega(2\lambda, \nu') \subset \Omega(2\lambda, \pi) \subset \Omega(2\lambda, \mu') = \Omega(\lambda, \mu). \quad \square$$

Corollary 3.6. *Let $\Pi, \Pi' \subset \Lambda^+$ be simple with maximal elements resp. λ, λ' and assume that $\text{Supp}(\lambda) = \text{Supp}(\lambda')$.*

- i) *There exists an equivariant morphism $X_\Pi \rightarrow X_{\Pi'}$ if and only if for every $\mu' \in \Pi'_{\text{red}}$ there exists $\mu \in \Pi_{\text{red}}$ such that $\mu' - \lambda' \leq^\lambda \mu - \lambda$.*
- ii) *The varieties X_Π and $X_{\Pi'}$ are equivariantly isomorphic if and only if $\Pi_{\text{red}} \sim \Pi'_{\text{red}}$.*

Proof. i) Suppose that X_Π dominates $X_{\Pi'}$ and let $\mu' \in \Pi'_{\text{red}} \setminus \{\lambda'\}$. By Corollary 1.10 i) we have $\Omega(\lambda) = \Omega(\lambda')$, while by Proposition 1.6 they exist $\mu_1, \dots, \mu_n \in \Pi$ such that $V(\mu' - \lambda' + n\lambda) \subset V(\mu_1) \otimes \dots \otimes V(\mu_n)$. Since $\mu' \in \Pi' \setminus \Pi_{\text{tr}}^+(\lambda')$, we have $\mu' - \lambda' \notin \Omega(\lambda') = \Omega(\lambda)$, hence some μ_i is not trivial, say μ_1 , and Proposition 3.1 ii) implies $\mu' - \lambda' \leq^\lambda \mu_1 - \lambda$. Therefore if $\mu \in \Pi_{\text{red}}$ is any weight such that $\mu_1 \leq^\lambda \mu$, we get $\mu' - \lambda' \leq^\lambda \mu - \lambda$.

Suppose conversely that for every $\mu' \in \Pi'_{\text{red}}$ there exists $\mu \in \Pi_{\text{red}}$ such that $\mu' - \lambda' \leq^\lambda \mu - \lambda$ and let $\nu' \in \Pi'$. If $\nu' \in \Pi_{\text{tr}}^+(\lambda')$, then $\nu' - \lambda' \in \Omega(\lambda') = \Omega(\lambda)$ by Proposition 1.1 together with Remark 1.9. Suppose that $\nu' \in \Pi' \setminus \Pi_{\text{tr}}^+(\lambda')$ and let $\mu' \in \Pi'_{\text{red}}$ be such that $\nu' \leq^\lambda \mu'$: then by Theorem 3.5 it follows that $\Omega(\lambda', \nu') \subset \Omega(\lambda', \mu')$. Let $\mu \in \Pi_{\text{red}}$ be such that $\mu' - \lambda' \leq^\lambda \mu - \lambda$ and denote $\lambda'' = \lambda + \lambda'$: since μ and μ' are non-trivial, it follows that $\lambda + \mu', \lambda' + \mu \in \Pi^+(\lambda'') \setminus \Pi_{\text{tr}}^+(\lambda'')$. Then $\{\lambda, \mu\} \sim \{\lambda'', \lambda' + \mu\}$ and $\{\lambda', \mu'\} \sim \{\lambda'', \lambda + \mu'\}$ are equivalent simple sets, so by Corollary 1.10 i) we have $\Omega(\lambda, \mu) = \Omega(\lambda'', \lambda' + \mu)$ and $\Omega(\lambda', \mu') = \Omega(\lambda'', \lambda + \mu')$. Moreover by construction we have $\lambda + \mu' \leq^\lambda \lambda' + \mu$, hence $\Omega(\lambda'', \lambda + \mu') \subset \Omega(\lambda'', \lambda' + \mu)$ by Theorem 3.5. Finally by the definition of $\Omega(\Pi)$ we have $\Omega(\lambda, \mu) \subset \Omega(\Pi)$, so it follows

$$\Omega(\lambda', \nu') \subset \Omega(\lambda', \mu') = \Omega(\lambda'', \lambda + \mu') \subset \Omega(\lambda'', \lambda' + \mu) = \Omega(\lambda, \mu) \subset \Omega(\Pi).$$

Therefore we have shown that $\nu' - \lambda' \in \Omega(\Pi)$ for every $\nu' \in \Pi'$ and the claim follows by Corollary 1.7.

ii) If $\Pi_{\text{red}} \sim \Pi'_{\text{red}}$ then the claim follows by Corollary 1.10 i). Assume that $X_\Pi \simeq X_{\Pi'}$ and let $\mu \in \Pi_{\text{red}} \setminus \{\lambda\}$. Then by i) there exists $\mu' \in \Pi'_{\text{red}}$ such that $\mu - \lambda \leq^\lambda \mu' - \lambda'$ and similarly there exists $\mu_1 \in \Pi_{\text{red}}$ such that $\mu' - \lambda' \leq^\lambda \mu_1 - \lambda$. Therefore $\mu + \lambda' \leq^\lambda \mu' + \lambda \leq^\lambda \mu_1 + \lambda'$ and we get $\mu \leq^\lambda \mu_1$. On the other hand by the definition of Π_{red} we have that $\mu, \mu_1 \in \Pi \setminus \Pi_{\text{tr}}^+$ are maximal w.r.t. \leq^λ , hence it follows that $\mu = \mu_1$ and $\mu - \lambda = \mu' - \lambda'$. Therefore for every $\mu \in \Pi_{\text{red}}$ there exists $\mu' \in \Pi'_{\text{red}}$ such that $\mu - \lambda = \mu' - \lambda'$, and an analogous argument for Π' shows that $\Pi_{\text{red}} - \lambda = \Pi'_{\text{red}} - \lambda$. On the other hand, since the closed orbits of X_Π and $X_{\Pi'}$ are isomorphic, Proposition 1.2 ii) implies that $\text{Supp}(\lambda) = \text{Supp}(\lambda')$, therefore we get $\Pi_{\text{red}} \sim \Pi'_{\text{red}}$. \square

4. EXAMPLES: SIMPLE LINEAR COMPACTIFICATIONS OF $\text{SO}(7)$ AND $\text{SO}(9)$

If $I \subset \Delta$, set $X_I = X_\lambda$ and $\tilde{X}_I \rightarrow X_I$ the normalization, where $\lambda \in \Lambda^+$ is such that $\text{Supp}(\lambda) = I$: by Proposition 1.1 these varieties are well defined and they only depend on I . Consider the set

$$T(I) = \{\text{simple linear compactifications } \text{SO}(2r+1) \hookrightarrow X \text{ such that } \tilde{X}_I \rightarrow X \rightarrow X_I\}.$$

In other words, following the discussion after Proposition 1.2, $T(I)$ is the set of the compactifications X_Π such that $\Pi \subset \Lambda^+$ is a simple subset whose maximal element has support I . We regard $T(I)$ as a partially ordered set, where the order is defined as follows: $X' \leq X$ if there exists an equivariant morphism $X \rightarrow X'$. We also denote

$$T(I, 2) = \{X \in T(I) : X \not\cong X_I \text{ and } X \simeq X_\Pi \text{ for some } \Pi \subset \Lambda^+ \text{ with } \text{card}(\Pi) = 2\}.$$

Denote \leq^I the partial order on $\mathbb{N}\Delta$ defined as follows:

$$\text{If } \theta_1, \theta_2 \in \mathbb{N}\Delta, \text{ then } \theta_1 \leq^I \theta_2 \text{ if and only if } \theta_2 - \theta_1 \in \mathbb{N}\Delta \setminus \mathbb{N}[\Delta \setminus I].$$

This coincides with the partial order \leq^λ defined in Section 3, where λ is any weight such that $\text{Supp}(\lambda) = I$. Following Remark 1.11, Corollary 2.3 and Corollary 3.6, $(T(I, 2), \leq)$ is identified with the partially ordered set $(\mathcal{T}(I, 2), \leq^I)$, where $\mathcal{T}(I, 2) \subset \mathbb{N}\Delta$ is defined as follows:

$$\mathcal{T}(I, 2) = \left\{ \theta = \sum_{i=1}^r a_i \alpha_i \in \mathbb{N}\Delta : \text{Supp}(\theta^+) \subset I, a_r \text{ is odd and } a_r < 2 \min\{r - l(\theta), r - q(I)\} \right\}$$

where $q(I) \leq r$ is the maximum such that $\alpha_{q(I)} \in I$ and where $l(\theta) \leq r$ denotes the minimum such that $a_i = a_r$ for every $i \geq l(\theta)$. Following Theorem 1.13, notice that $\mathcal{T}(I, 2)$ possesses a unique maximal element w.r.t. \leq^I , namely $\theta_I = \sum_{i=q(I)}^r \alpha_i$: this is the element corresponding to the normalization \tilde{X}_I and it coincides with the difference $\lambda - \lambda^{\text{lb}}$, where $\lambda \in \Lambda^+$ is any dominant weight with $\text{Supp}(\lambda) = I$ and where λ^{lb} is the little brother of λ (see Definition 1.12).

Denote $\mathcal{P}(\mathbb{N}\Delta)$ the power set of $\mathbb{N}\Delta$ and extend \leq^I to a partial order relation on $\mathcal{P}(\mathbb{N}\Delta)$ as follows:

$$\text{If } A, A' \subset \mathbb{N}\Delta, \text{ then } A \leq^I A' \text{ if and only if } \forall \theta \in A \quad \exists \theta' \in A' : \theta \leq^I \theta'.$$

Following Corollary 3.6, every element of $T(I)$ is identified with a subset of $\mathcal{T}(I, 2)$ and we may identify $(T(I), \leq)$ with the partially ordered set $(\mathcal{T}(I), \leq^I)$, where $\mathcal{T}(I) \subset \mathcal{P}(\mathcal{T}(I, 2))$ is defined as follows:

$$\mathcal{T}(I) = \{A \subset \mathcal{T}(I, 2) : A \text{ contains no comparable elements w.r.t. } \leq^I\}.$$

In the following tables we represent the poset $(\mathcal{T}(I, 2), \leq^I)$ for $\text{SO}(7)$ and for $\text{SO}(9)$. If $\alpha_r \in I$ then by Theorem 1.13 it follows that $\mathcal{T}(I, 2) = \emptyset$, while if $I = \{\alpha_{r-1}\}$ it follows by Remark 2.4 that $\mathcal{T}(I, 2) = \{\alpha_{r-1} + \alpha_r\}$. Therefore we will assume that $\alpha_r \notin I$ and that $I \neq \{\alpha_{r-1}\}$. We represent an element $\sum a_i \alpha_i \in \mathcal{T}(I, 2)$ as the vector (a_1, \dots, a_r) and we connect two elements $\theta_1, \theta_2 \in \mathcal{T}(I, 2)$ with an arrow $\theta_1 \rightarrow \theta_2$ if and only if $\theta_1 \leq^I \theta_2$ and $\theta_2 \in \mathcal{T}(I, 2)$ is minimal with this property.

TABLE 1. The poset $(\mathcal{T}(I, 2), \leq^I)$ for $G = \text{SO}(7)$, $I = \{\alpha_1, \alpha_2\}$.

$$(0, 1, 1) \longrightarrow (1, 1, 1) \longrightarrow (2, 1, 1) \longrightarrow (3, 1, 1) \longrightarrow (4, 1, 1) \longrightarrow \dots$$

TABLE 2. The poset $(\mathcal{T}(I, 2), \leq^I)$ for $G = \text{SO}(7)$, $I = \{\alpha_1\}$.

$$(1, 1, 1) \longrightarrow (2, 1, 1) \xrightarrow{\nearrow (3, 3, 3)} (3, 1, 1) \longrightarrow (4, 1, 1) \longrightarrow (5, 1, 1) \longrightarrow \dots$$





TABLE 3. The poset $(\mathcal{T}(I, 2), \leq^I)$ for $G = \text{SO}(9)$, $I = \{\alpha_1, \alpha_2, \alpha_3\}$.TABLE 4. The poset $(\mathcal{T}(I, 2), \leq^I)$ for $G = \text{SO}(9)$, $I = \{\alpha_1, \alpha_3\}$.TABLE 5. The poset $(\mathcal{T}(I, 2), \leq^I)$ for $G = \text{SO}(9)$, $I = \{\alpha_2, \alpha_3\}$.TABLE 6. The poset $(\mathcal{T}(I, 2), \leq^I)$ for $G = \text{SO}(9)$, $I = \{\alpha_1, \alpha_2\}$.

TABLE 7. The poset $(\mathcal{T}(I, 2), \leq^I)$ for $G = \text{SO}(9)$, $I = \{\alpha_2\}$.





TABLE 8. The poset $(\mathcal{T}(I, 2), \leq^I)$ for $G = \text{SO}(9)$, $I = \{\alpha_1\}$.



REFERENCES

[1] V. Alexeev and M. Brion, *Stable reductive varieties II: Projective case*, Adv. Math. **184** (2004), no. 2, 380–408.
 [2] N. Bourbaki, *Éléments de mathématique. Fasc. XXXIV. Groupes et algèbres de Lie. Chapitres IV, V, VI*, Actualités Scientifiques et Industrielles **1337**, Hermann Paris 1968.
 [3] P. Bravi, J. Gandini, A. Maffei and A. Ruzzi, *Normality and non-normality of group compactifications in simple projective spaces*, Ann. Inst. Fourier (Grenoble) **61** (2011), no. 6, 2435–2461.
 [4] C. De Concini, *Normality and non normality of certain semigroups and orbit closures*, in: Algebraic transformation groups and algebraic varieties, Encycl. Math. Sci. **132**, Springer, Berlin, 2004, pp. 15–35.
 [5] C. De Concini and C. Procesi, *Complete symmetric varieties*, in: Invariant theory (Montecatini, 1982), Lecture Notes in Math. **996**, Springer, Berlin, 1983, pp. 1–44.
 [6] J. Gandini and A. Ruzzi, *Normality and smoothness of simple linear group compactifications*, Math. Z., to appear.
 [7] R. Goodman and N.R. Wallach, *Symmetry, representations, and invariants*, Graduate Texts in Mathematics **255**, Springer, Dordrecht, 2009. Appendices E,F,G available at <http://www.math.rutgers.edu/~goodman/repbook.html>
 [8] S. S. Kannan, *Projective normality of the wonderful compactification of semisimple adjoint groups*, Math. Z. **239** (2002), no. 4, 673–682.
 [9] F. Knop, *The Luna-Vust theory of spherical embeddings*, in: Proceedings of the Hyderabad Conference on Algebraic Groups (Hyderabad, 1989), Manoj Prakashan, Madras, 1991, pp. 225–249.

- [10] F. Knop, H. Kraft, D. Luna and T. Vust, *Local properties of algebraic group actions*, in: *Algebraische Transformationsgruppen und Invariantentheorie*, DMV Sem. **13**, Birkhäuser, Basel, 1989, pp. 63–75.
- [11] S. Kumar, *Tensor Product Decomposition*, in: *Proceedings of the International Congress of Mathematicians*, Hyderabad, India, 2010, pp. 1226–1261.
- [12] D. A. Timashev, *Equivariant compactifications of reductive groups*, *Sb. Math.* **194** (2003), no. 3–4, 589–616.

Current address: DIPARTIMENTO DI MATEMATICA “GUIDO CASTELNUOVO”, “SAPIENZA” UNIVERSITÀ DI ROMA,
PIAZZALE ALDO MORO 5, 00185 ROMA, ITALY

E-mail address: gandini@mat.uniroma1.it