

1 **Multifunctionality and diversity of culturable bacterial communities strictly associated with**  
2 **spores of the plant beneficial symbiont *Rhizophagus intraradices***

3

4 Fabio Battini, Caterina Cristani, Manuela Giovannetti, Monica Agnolucci\*

5 Department of Agriculture, Food and Environment, University of Pisa, Via del Borghetto 80, 56124

6 Pisa, Italy

7

8 RUNNING HEAD: Multifunctional culturable bacteria from AMF spores

9

10 \*Corresponding author: M. Agnolucci, Department of Agriculture, Food and Environment,

11 University of Pisa, Via del Borghetto 80, 56124 Pisa, Italy

12 Phone: +39.0502216647, Fax: +39.0502220606, e-mail address: monica.agnolucci@unipi.it

13

14

15 **ABSTRACT**

16 Arbuscular Mycorrhizal Fungi (AMF) live in symbiosis with most crop plants and represent  
17 essential elements of soil fertility and plant nutrition and productivity, facilitating soil mineral  
18 nutrient uptake and protecting plants from biotic and abiotic stresses. These beneficial services may  
19 be mediated by the dense and active spore-associated bacterial communities, which sustain diverse  
20 functions, such as the promotion of mycorrhizal activity, biological control of soilborne diseases,  
21 nitrogen fixation, and the supply of nutrients and growth factors. In this work, we utilised culture-  
22 dependent methods to isolate and functionally characterize the microbiota strictly associated to  
23 *Rhizophagus intraradices* spores, and molecularly identified the strains with best potential plant  
24 growth promoting (PGP) activities by 16S rDNA sequence analysis. We isolated in pure culture 374  
25 bacterial strains belonging to different functional groups - actinobacteria, spore-forming,  
26 chitinolytic and N<sub>2</sub>-fixing bacteria - and screened 122 strains for their potential PGP activities. The

27 most common PGP trait was represented by P solubilization from phytate (69.7%), followed by  
28 siderophore production (65.6%), mineral P solubilization (49.2%) and IAA production (42.6%).  
29 About 76% of actinobacteria and 65% of chitinolytic bacteria displayed multiple PGP activities.  
30 Nineteen strains with best potential PGP activities, assigned to *Sinorhizobium meliloti*,  
31 *Streptomyces* spp., *Arthrobacter phenanthrenivorans*, *Nocardioides albus*, *Bacillus* sp. *pumilus*  
32 group, *Fictibacillus barbaricus* and *Lysinibacillus fusiformis*, showed the ability to produce IAA  
33 and siderophores and to solubilize P from mineral phosphate and phytate, representing suitable  
34 candidates as biocontrol agents, biofertilisers and bioenhancers, in the perspective of targeted  
35 management of beneficial symbionts and their associated bacteria in sustainable food production  
36 systems.

37  
38 **Keywords:** Plant growth promoting bacteria; AMF spores; culturable spore-associated bacteria;  
39 *Rhizophagus intraradices*

## 40 41 INTRODUCTION

42 The symbiosis between plants and Arbuscular Mycorrhizal (AM) Fungi (AMF, Glomeromycota) is  
43 the most widespread on Earth. AMF establish beneficial associations with the roots of the large  
44 majority of land plants, including the most important food crops, from cereals to legumes,  
45 vegetables and fruit trees (Smith and Read 2008), and contribute to key agroecosystem processes,  
46 such as nutrient uptake, soil aggregation and carbon sequestration (Gianinazzi et al. 2010). AM  
47 symbionts represent essential elements of soil fertility and plant nutrition and productivity,  
48 facilitating soil mineral nutrient uptake - mainly phosphorus (P), nitrogen (N), sulfur (S), potassium  
49 (K), calcium (Ca), copper (Cu) and zinc (Zn) - by means of an extensive extraradical network of  
50 fungal hyphae spreading from colonized roots into the soil (Giovannetti and Avio 2002). In  
51 addition, AMF protect plants from soilborne fungal pathogens and abiotic stresses, such as drought  
52 and salinity (Augé 2001; Evelin et al. 2009; Sikes et al. 2009) and affect the synthesis of beneficial

53 phytochemicals, contributing to the sustainable production of high-quality food (Giovannetti et al.  
54 2012).

55 Such beneficial services may be affected by diverse factors, including agronomic practices (Njeru et  
56 al. 2014) and further mediated by a third component of the symbiosis, the mycorrhizospheric  
57 microbiota, represented by the dense and active bacterial communities living tightly associated with  
58 AMF (Rambelli 1973). Mycorrhizospheric bacteria sustain diverse functions, driving plant growth  
59 and health and nutrient acquisition, *i.e.* the promotion of mycorrhizal activity (Mayo et al. 1986;  
60 Xavier and Germida 2003; Giovannetti et al. 2010; Hori and Ishii 2006), biological control of  
61 soilborne diseases (Citernesi et al. 1996; Budi et al. 1999; Li et al. 2007; Bharadwaj et al. 2008a),  
62 nitrogen fixation, and the supply of nutrients and growth factors (Barea et al. 2002; Xavier and  
63 Germida 2003; Bharadwaj et al. 2008b).

64 A number of studies investigated spore-associated microbiota, as AMF spores have long been  
65 known to harbor a wide diversity of bacterial species, living either intracellularly (Mosse 1970;  
66 MacDonald and Chandler 1981; MacDonald et al. 1982; Bianciotto et al. 1996) or intimately  
67 associated with the spore walls. Indeed, spore walls represent a privileged habitat where bacteria  
68 can thrive on exudates and by hydrolyzing the relevant wall biopolymers, such as proteins and  
69 chitin (Walley and Germida 1996; Filippi et al. 1998; Roesti et al. 2005) and survive before  
70 colonizing the surface of germlings and extraradical hyphae (Lecomte et al. 2011). For this reason,  
71 AMF spores are the preferential source of AMF-associated bacteria to be studied for their  
72 prospective use as biocontrol agents, biofertilisers and bioenhancers, in order to develop strategies  
73 able to minimize anthropogenic energy inputs and promote plant productivity and health, and soil  
74 fertility. The isolation and functional characterization of Spore-Associated Bacteria (SAB) are of  
75 key agronomical importance and represent a prerequisite for understanding entirely how the  
76 complex network of microbial interactions in the mycorrhizosphere affects plant performance and  
77 can be managed in sustainable plant production systems.

78 So far, a few works reported the isolation and characterization of SAB from a small number of  
79 AMF. For example, ten bacterial species were recovered from *Glomus clarum* spores (Xavier and  
80 Germida 2003), seven species from *Glomus irregulare* spores harvested from the field (Lecomte et  
81 al. 2011), while a total of 36 species were obtained from *Glomus intraradices* and *Glomus mosseae*  
82 spores extracted from the rhizosphere of *Festuca ovina* and *Leucanthemum vulgare* (Bharadwaj et  
83 al. 2008a). Despite these studies, little information is currently available on the functional  
84 significance of culturable bacterial communities associated with AMF spores, and on their possible  
85 synergistic interactions. Some bacteria showed antagonistic activity against plant pathogens (Budi  
86 et al. 1999; Bharadwaj et al. 2008b), phosphate-solubilizing and nitrogenase activity (Cruz et al.  
87 2008; Cruz and Ishii 2011), and indole acetic acid production (Bharadwaj et al. 2008b), suggesting  
88 their possible role as Plant Growth Promoting (PGP) bacteria.

89 Using a molecular approach, we recently reported the occurrence of specific and diverse microbial  
90 communities tightly associated with spores of six different AMF isolates. In particular, sequencing  
91 of DGGE bands lead to the identification of bacteria belonging to *Actinomycetales*, *Bacillales*,  
92 *Rhizobiales*, *Pseudomonadales*, *Burkholderiales*, and *Mollicutes* related endobacteria (Mre).  
93 Several strains belonged to species known to play important roles in promoting plant growth either  
94 directly, by affecting nutrient availability - for example solubilizing phosphate and other nutrients  
95 from insoluble sources, fixing nitrogen and producing phytohormones, mainly Indol Acetic Acid  
96 (IAA) - or indirectly by protecting plants against pathogens through the production of siderophores,  
97 antibiotics and extracellular hydrolytic enzymes.

98 As the different combinations of AMF and bacterial activities may result complementary and/or  
99 synergistic and represent the basis of the differential symbiotic performance of AMF isolates, it is  
100 crucial to gain knowledge on the functional significance of bacteria associated with AMF spores, in  
101 order to exploit the potential of such multipartite association in sustainable food production  
102 systems. In this work, we investigated the diversity and functional PGP potential of 122 culturable  
103 bacterial strains strictly associated to spores of the AMF species *Rhizophagus intraradices*. To this

104 aim, we i) utilised culture-dependent methods to isolate the microbiota associated with *R.*  
105 *intraradices* spores, ii) functionally characterized bacterial strains in order to detect their beneficial  
106 role in plant growth and health, nitrogen fixation, P solubilization, siderophore and IAA production,  
107 iii) identified the strains with best potential PGP activities by 16S rDNA sequence analysis.

108

## 109 **MATERIALS AND METHODS**

### 110 *Fungal material*

111 The AM fungus used was *Rhizophagus intraradices* (N.C. Schenck & G.S. Sm.) C. Walker &  
112 Schuessler isolate IMA6, obtained from pot-cultures maintained in the collection of Microbiology  
113 Labs (International Microbial Archives, IMA) of the Department of Agriculture, Food and  
114 Environment, University of Pisa, Italy. The isolate was maintained in sterilized calcinated clay  
115 (OILDRI, Chicago, IL, USA) and field soil (1:1, v/v) pot cultures, using *Trifolium alexandrinum* L.  
116 and *Medicago sativa* L. as host plants.

117

### 118 *Spore collection and bacterial isolation*

119 *R. intraradices* spores were extracted from three pot cultures using the wet-sieving and decanting  
120 technique, down to a mesh size of 50 µm (Gerdermann and Nicolson 1963). Spores retained on  
121 sieves were flushed into Petri dishes and manually collected with a capillary pipette under a  
122 dissecting microscope (Leica MS5, Milan, Italy). For each pot 200 intact and healthy spores were  
123 suspended in 1 mL sterile physiological solution (9 g L<sup>-1</sup> NaCl) in a 1.5 mL Eppendorf tube and  
124 vigorously washed using a vortex mixer at 1500 rpm for 1 min. Spores were aseptically washed 15  
125 times in succession, then crushed aseptically with a pestle and suspended in 3.5 mL of sterile  
126 physiological solution. Spores were not washed further, as previous experiments showed that 15  
127 washings were effective in spore surface decontamination (Agnolucci et al. 2015).

128 An aliquot (1 mL) of spore suspension was heat-treated for spore forming bacteria isolation (80 °C  
129 for 10 min). 100 µL suspension for each sample were plated in triplicate onto different culture

130 media. Culturable heterotrophic and spore-forming bacteria were isolated on TSA (30 g L<sup>-1</sup> tryptic  
131 soy broth, 20 g L<sup>-1</sup> bacteriological agar, Oxoid, Milan, Italy), a non-selective medium which allows  
132 the recovery of a wide range of aerobic and facultative anaerobic gram-negative and gram-positive  
133 bacteria. The media were supplemented with 100 mg L<sup>-1</sup> of cyclohexymide and 500 UI L<sup>-1</sup> of  
134 nystatin (Sigma-Aldrich, Milan, Italy) to inhibit possible fungal development.

135 Selective media were used to isolate specific functional bacterial groups, *i.e.* actinobacteria,  
136 chitinolytic and nitrogen-fixing bacteria. Actinobacteria were isolated using Waksman's agar  
137 medium (10 g L<sup>-1</sup> dextrose, 5 g L<sup>-1</sup> sodium chloride, 5 g L<sup>-1</sup> bacteriological peptone, 3 g L<sup>-1</sup> lab-  
138 lemco powder, 20 g L<sup>-1</sup> bacteriological agar; Oxoid, Milan, Italy) supplemented with 5 mg L<sup>-1</sup> of  
139 polymyxin (Sigma-Aldrich, Milan, Italy) and with 100 mg L<sup>-1</sup> of cyclohexymide and 500 UI L<sup>-1</sup> of  
140 nystatin (Sigma-Aldrich, Milan, Italy) to inhibit the growth of gram-negative bacteria and fungi,  
141 respectively. Chitinolytic bacteria were isolated after plating samples on minimal medium  
142 containing chitin as sole carbon source (Souza et al. 2009), while bacteria able to grow on N-free  
143 medium were isolated from Winogradsky culture agar (Tchan 1984). Both media were  
144 supplemented with 100 mg L<sup>-1</sup> of cyclohexymide and 500 UI L<sup>-1</sup> of nystatin (Sigma-Aldrich, Milan,  
145 Italy) to inhibit fungal growth.

146 The number of Colony Forming Units (CFU) was assessed after 2 and 7 days of incubation at 28 °C  
147 for TSA and the other media, respectively. From each isolation medium, representative bacterial  
148 strains were randomly selected on the basis of phenotypic colony characteristics, *i.e.* shape, size,  
149 edge morphology, surface and pigment, in order to include the most diverse strains, and purified by  
150 streaking four times onto the same medium used for isolation. Each strain was named on the basis  
151 of the acronym of the isolation medium, followed by a progressive number. Spore-forming bacteria  
152 isolated on TSA were named as TSAT. Purified strains were maintained at -80 °C in cryovials with  
153 20% (v/v) of glycerol.

154

155 *Screening for PGP traits*

156 *IAA production*

157 The production of IAA was investigated using Luria-Bertani Broth (LBB, 10 g L<sup>-1</sup> bacto-tryptone, 5  
158 g L<sup>-1</sup> yeast extract, 5 g L<sup>-1</sup> sodium chloride, pH 7.5, Oxoid, Milan, Italy) (Bharadwaj et al. 2008b).  
159 The strains were inoculated in 15 mL tubes containing 4 mL of LBB amended with 1 mg mL<sup>-1</sup> of L-  
160 tryptophan (Sigma-Aldrich, Milan, Italy), incubated at 20 °C with continuous shaking (200 rpm)  
161 until they reached exponential growth phase and then centrifuged at 7500 rpm for 10 min. After  
162 centrifugation, 1 mL of supernatant was transferred in a 24-well plate, mixed with 2 mL of  
163 Salkowski reagent (1.2% FeCl<sub>3</sub> in 37% sulphuric acid) and incubated in the dark for 30 min. The  
164 non-inoculated medium was used as negative control, while the medium amended with pure IAA  
165 was used as positive control. Development of red-purple color indicated positive strains for IAA  
166 production. Strains were classified using a rating scale as follows: - = no production (no color  
167 development), +/- = low production (pale pink), + = production (light purple), ++ = moderate  
168 production (bright purple), +++ = high production (dark purple).

169

170 *Siderophore production*

171 Siderophore-producing strains were detected using the overlay Chrome Azurol S assay (CAS)  
172 described by Pérez-Miranda et al. (2007). CAS agar was prepared following the procedure provided  
173 by Loudon et al. (2011) using 72.9 mg L<sup>-1</sup> hexadecyltrimethyl ammonium bromide (HDTMA), 30.24  
174 g L<sup>-1</sup> piperazine-1,4-bis(2-ethanesulfonic acid) (PIPES), 1 mM FeCl<sub>3</sub> 6H<sub>2</sub>O in 10 mM HCl 10 mL  
175 and 0.9 g L<sup>-1</sup> bacteriological agar. Four isolates per plate were inoculated on TSA and incubated 2-7  
176 days at 28 °C. After incubation, 10 mL of CAS agar were spread as an overlay on the  
177 microorganisms and incubated at room temperature. Siderophore-producing strains showed a  
178 change in color, from-blue to yellow or from blue to orange, in the overlaid medium around the  
179 colonies. After 7 days the radius of the halo was measured (mm) from the colony edge to the edge  
180 of the colored halo. Strains were classified using a rating scale as follows: no production (halo = 0

181 mm), +/- = low production (halo < 2 mm), + = production (2 mm ≤ halo ≤ 8 mm), ++ = moderate  
182 production (8 mm < halo < 14 mm), +++ = high production (halo > 15 mm).

183

#### 184 *P solubilisation from mineral phosphate and phytate*

185 The ability of isolated bacteria to solubilize inorganic and organic phosphate was assayed on  
186 National Botanical Research Institute's Phosphate growth medium (NBRIP, 10 g L<sup>-1</sup> D-glucose, 5 g  
187 L<sup>-1</sup> Ca<sub>3</sub>(PO<sub>4</sub>)<sub>2</sub>, 5 g L<sup>-1</sup> MgCl<sub>2</sub> 6H<sub>2</sub>O, 0.25 g L<sup>-1</sup> MgSO<sub>4</sub>·7H<sub>2</sub>O, 0.2 g L<sup>-1</sup> KCl, 0.1 g L<sup>-1</sup> (NH<sub>4</sub>)<sub>2</sub> SO<sub>4</sub>,  
188 15 g L<sup>-1</sup> agar, pH 7) (Nautiyal et al. 1999), and Phytate Screening Medium (PSM, 10 g L<sup>-1</sup> D-  
189 glucose, 4 g L<sup>-1</sup> C<sub>6</sub>H<sub>18</sub>P<sub>6</sub>O<sub>24</sub>·12Na·xH<sub>2</sub>O, 2 g L<sup>-1</sup> CaCl<sub>2</sub>, 5 g L<sup>-1</sup> NH<sub>4</sub>NO<sub>3</sub>, 0.5 g L<sup>-1</sup> KCl, 0.5 g L<sup>-1</sup>  
190 MgSO<sub>4</sub>·7H<sub>2</sub>O, 0.01 g L<sup>-1</sup> FeSO<sub>4</sub>·7H<sub>2</sub>O, 0.01 g L<sup>-1</sup> MnSO<sub>4</sub>·H<sub>2</sub>O, 15 g L<sup>-1</sup> agar, pH 7) respectively  
191 (Jorquera et al. 2008). In both assays, four strains were spot inoculated on each agar plate and  
192 incubated at 28 °C for 7 days. The formation of halo zones around bacterial colonies indicated  
193 phytate and phosphate solubilization capacity of the strains. After incubation, colony diameter and  
194 halo zones were recorded. The ability of bacteria to solubilize insoluble phosphate was evaluated as  
195 phosphate Solubilization Efficiency (SE), as described by Rokhbakhsh-Zamin et al. (2011):  
196  $SE = [\text{Halo zone (z)}/\text{Diameter of colonies (n)}] \times 100$ . The Phosphate Solubilization Index (PSI) was  
197 also calculated according to the following formula:  $PSI = \frac{\text{total diameter (colony + halo zone)}}{\text{the colony diameter}}$  (Islam et al. 2007).

199

#### 200 *PCR amplification of NifH gene*

201 Bacteria isolated from N-free medium were tested for the presence of nifH gene by PCR  
202 amplification. Genomic DNA was extracted from bacterial liquid cultures grown overnight at 28 °C  
203 using “MasterPure™ Yeast DNA Purification Kit” (Epicentre®) according to the manufacturer’s  
204 protocols. The amplification of nifH gene 390 bp fragment was carried out using the specific  
205 primers 19F (5’-GCIWTYTAYGGIAARGGIGG-3’) and 407R (5’-AAICCRCCRCIAIACIACRTC-  
206 3’) (Ueda et al. 1995). Amplification reaction was prepared in a final volume of 25 µl, with 10-20



207 ng of DNA, 1X Reaction buffer (Takara), 1.25 U of Takara ex Taq DNA polymerase, 0.2 mM of  
208 each dNTPs (Takara) and 0.5  $\mu$ M of each primers (Primm). The reaction was carried out using an  
209 iCycler-iQ Multicolor Real-Time PCR Detection System (Biorad) with the following denaturation,  
210 amplification and extension procedure: 94 °C 1 min; 94 °C 30 sec, 56 °C 30 sec, 72 °C 30 sec for  
211 35 cycles; 72 °C 5 min. The presence of amplicons was confirmed by electrophoresis in 1.5% (w/v)  
212 Agarose I (Euroclone®) in TBE 1 x buffer (Euroclone®) gels stained with ethidium bromide 0.5  $\mu$ g  
213 mL<sup>-1</sup>. All gels were visualized and captured as TIFF format files by the Liscap program for Image  
214 Master VDS system (Pharmacia Biotech).

215

#### 216 *Identification of bacterial strains with best potential PGP activities*

217 Bacterial isolates showing multiple PGP activities were identified based on 16S rDNA sequencing.  
218 Genomic DNA was extracted from bacterial liquid cultures grown overnight at 28 °C using  
219 “MasterPure™ Yeast DNA Purification Kit” (Epicentre®) according to the manufacturer’s  
220 protocols. The amplification of 16S rDNA was carried out using the primers 27f (5’-  
221 GAGAGTTTGACTCTGGCTCAG-3’) and 1495r (5’-CTACGGCTACCTTGTTACGA-3’) (Lane  
222 1991; Weisburg et al. 1991). Amplification reaction was carried out in a final volume of 50  $\mu$ l,  
223 using 10-20 ng of DNA, 1X Reaction buffer (EuroClone®), 2 mM MgCl<sub>2</sub> (EuroClone®), 1.25 U  
224 EuroTaq DNA polymerase (EuroClone®), 0.2 mM of each dNTPs (GeneAmp dNTP Mix, Applied  
225 Biosystem) and 0.2  $\mu$ M of each primers (Primm). The reaction was carried out using an iCycler-iQ  
226 Multicolor Real-Time PCR Detection System (Biorad) with the following denaturation,  
227 amplification and extension procedure: 95 °C 2 min; 94 °C 1 min and 20 sec, 54 °C 1 min, 72 °C 1  
228 min and 30 sec for 35 cycles; 72 °C 5 min. PCR amplicons were analysed by 1.5% agarose gel  
229 electrophoresis, stained with ethidium bromide and visualized under UV light as describe above.  
230 The amplification products were then purified with the EuroGold Cycle Pure Kit (EuroClone®)  
231 according to the manufacturer’s protocol, quantified and 5’ sequenced by BMR Genomics (Padova,  
232 Italy). Sequences were analyzed using BLAST on the NCBI web

233 (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>). The related sequences were collected and aligned using  
234 MUSCLE (Edgar 2004a; 2004b), phylogenetic trees constructed using the Neighbor-Joining  
235 method and the evolutionary distances computed using the Kimura 2-parameter method (Kimura  
236 1980) in Mega 6.0 software (<http://www.megasoftware.net/>) (Tamura et al. 2013) with 1,000  
237 bootstraps replicates.

238 The sequences of 16S rRNA genes were submitted to the European Nucleotide Archive (ENA)  
239 under the accession numbers from LN871737 to LN871755.

240

## 241 RESULTS

### 242 *Culturable bacteria strictly associated with R. intraradices spores*

243 Heterotrophic bacteria strictly associated with *R. intraradices* spores ranged from  $5.4 \pm 0.9$  to  $22.9$   
244  $\pm 0.7$  CFU spore<sup>-1</sup>, spore forming culturable bacteria from  $0.8 \pm 0.4$  to  $2.9 \pm 0.7$  CFU spore<sup>-1</sup>,  
245 actinobacteria from  $3.7 \pm 0.4$  to  $23.0 \pm 1.5$  CFU spore<sup>-1</sup>, bacteria growing on N-free medium from  
246  $0.7 \pm 0.5$  to  $1.0 \pm 0.6$  CFU spore<sup>-1</sup>, and chitinolytic bacteria from  $0.2 \pm 0.1$  to  $1.1 \pm 0.1$  CFU spore<sup>-1</sup>  
247 (Table 1). A total of 374 bacterial strains were isolated and purified. Among the isolated bacteria  
248 50.5% was obtained from TSA, 36.1% from Waksman's agar, 7.2% from N-free Winogradsky  
249 medium and 6.2% developed on the selective chitin containing medium (Figure 1a).

250 Bacterial strains from TSA were morphologically examined and grouped into 16 morphotypes,  
251 according to their colony characteristics. Actinobacteria were categorized into 13 morphotypes,  
252 according to the color of the mature sporulated aerial mycelium and of the substrate mycelium (Fig.  
253 S1, supplementary electronic material). With the aim of detecting bacteria showing PGP traits, from  
254 the pool of the 374 isolated SAB a total of 122 strains were selected for further *in vitro* screening.  
255 Such strains included all the isolated chitinolytic and putative nitrogen fixers (23 and 27 strains,  
256 respectively), along with 38 heterotrophic bacteria and 34 actinobacteria, selected from the different  
257 morphotypes previously identified and chosen taking into account the number of isolates grouped in  
258 each morphotype.

259

260 *Screening for PGP traits*

261 The 122 selected bacterial strains were screened *in vitro* for the presence of PGP traits, *i.e.* the  
262 ability to produce IAA and siderophores, and to solubilize mineral phosphate and phytate. The most  
263 common PGP trait was represented by P solubilization from phytate (69.7%), followed by  
264 siderophore production (65.6%), phosphate solubilization (49.2%) and IAA production (42.6%).

265 The different bacterial strains showed a large variability in their PGP abilities: for example, the halo  
266 zone produced by phytate solubilizing bacteria ranged from 0.1 to 1.80 cm, the SE of phosphate  
267 solubilizers from 10 to 150%, while for siderophore production the radius of color change ranged  
268 from about 0.2 cm to more than 1.5 cm (Tables 2-5).

269 The four functional bacterial groups (heterotrophs, actinobacteria, chitinolytic and putative N<sub>2</sub>-  
270 fixing bacteria) showed different percentages of strains expressing single PGP activities. For  
271 example, the percentage of strains showing IAA production ranged from 70% of actinobacteria to  
272 14% of putative N-fixers, while that of phosphate solubilizing bacteria (PSB) varied from less than  
273 20% of heterotrophic isolates to 74% of actinobacteria and chitinolytic bacteria. Phytate  
274 solubilizing activity was detected in 50% of heterotrophic bacteria and in 100% of actinobacteria.  
275 Finally, siderophores were produced by 96% of bacteria growing on N-free medium, 69% of  
276 chitinolytic bacteria, and only 25% of the heterotrophic bacteria (Figure 1b). Interestingly, only 7  
277 out of 38 heterotrophic strains showed high siderophore production activity *vs.* 19 out of 34  
278 actinobacterial strains.

279 Interestingly, a large number of strains were able to express multiple PGP activities. Indeed, the  
280 percentage of strains displaying three to four PGP traits was about 76% in actinobacteria, 65% in  
281 chitinolytic bacteria, and 48% among bacteria growing on N-free medium (Figure 1c). Conversely,  
282 among heterotrophic bacteria no strain showed four PGP traits, and only 16% displayed three.

283 Venn diagrams relevant to each functional group detected 17 actinobacterial strains (W1,  
284 W22,W39,W43N, W47, W54, W56, W58, W60B, W60R, W64, W68, W77, W90, W94, W115,

285 W132) and 8 chitinolytic strains (CH5, CH6G, CH10, CH11, CH14, CH15, CH17, CH20)  
286 displaying the four PGP traits investigated. Among bacteria growing on N-free medium, only one  
287 isolate (N24) showed the four activities, while 12 strains (N14, N18, N19, N20, N21, N22, N23,  
288 N25, N26, N27, N28, N29) displayed three. Six heterotrophic bacteria (TSA3, TSA10, TSA26,  
289 TSA41, TSA46, TSAT102B) showed three PGP activities (Figure 2).

290

#### 291 *Amplification of NifH genes*

292 Bacterial strains isolated from N-free medium were further tested for the presence of nifH genes by  
293 PCR amplification, using the specific primers 19F and 407R (Ueda et al. 1995). The expected 390  
294 bp DNA fragment was obtained from 10 strains (N18, N19, N20, N21, N22, N23, N24, N25, N28,  
295 and N29).

296

#### 297 *16 S rDNA identification of selected PGP bacterial strains*

298 Representative strains from each functional group were selected on the basis of relevant PGP traits,  
299 and identified by 16S rDNA sequencing. A total of 19 strains were sequenced and affiliated to  
300 bacterial species using BLAST and phylogenetic trees analyses. Blast nucleotide searches of the  
301 16S rDNA sequences showed at least 98% similarities to database entries (Table 6). Figure 3 shows  
302 the related phylogenetic trees with the affiliation of sequences to bacterial species. Sequences were  
303 affiliated with *Actinomycetales* (*Arthrobacter*, *Streptomyces* and *Nocardiodes*), *Bacillales* (*Bacillus*,  
304 *Fictibacillus* and *Lysinibacillus*) and *Rhizobiales* (*Sinorhizobium*). Among the sequenced strains the  
305 majority could be assigned to *Sinorhizobium meliloti* (47.4%), followed by *Streptomyces* spp.  
306 (26.3%), while *Arthrobacter phenethrenivorans*, *Nocardiodes albus*, *Bacillus* sp. *pumilus* group.,  
307 *Fictibacillus barbaricus* and *Lysinibacillus fusiformis* were represented by only one isolate each.

308

309 DISCUSSION

310 In this work we showed the multifunctional traits of culturable bacteria strictly associated with  
311 spores of the beneficial plant symbiont *R. intraradices* IMA6. We isolated 374 spore-associated  
312 bacterial strains belonging to different functional groups known to possess PGP activities -  
313 actinobacteria, spore-forming, chitinolytic and N<sub>2</sub>-fixing bacteria. Strains with best potential PGP  
314 activities measured in pure culture, identified by 16S rDNA sequencing, showed the ability to  
315 produce IAA and siderophores and to solubilize P from mineral phosphate and phytate, representing  
316 suitable candidates as biofertilizers and bioenhancers.

317

### 318 *Spore-associated culturable bacterial communities*

319 Among the 374 bacterial strains strictly associated with *R. intraradices* spores isolated in this work,  
320 189 were recovered from TSA medium, a general growth substrate for heterotrophic bacteria and  
321 135 from Waksman's agar, a selective growth medium for actinobacteria. Quantitative analyses  
322 allowed us to confirm that heterotrophs and actinobacteria represented the majority of SAB, as  
323 bacterial counts reached a maximum of 23 and 22 CFU per spore, respectively. Such high numbers  
324 indicate that AMF spores represent a preferential ecological niche supporting bacterial metabolic  
325 activity as a result of their high levels of nutrients and exudates. Specific physiological groups of  
326 bacteria may be selectively favored, for their ability to hydrolyze spore wall biopolymers (Filippi et  
327 al. 1998; Roesti et al. 2005) or to form biofilms by producing exopolysaccharides, which allow  
328 them to adhere to spore walls (Toljander et al. 2006). Present results confirm our recent molecular  
329 findings on the occurrence of large bacterial communities intimately associated with spores of 6  
330 different AMF isolates, including the isolate IMA6 of the species *R. intraradices* studied here  
331 (Agnolucci et al. 2015). Other authors have previously shown that the microbiota associated with  
332 AMF spores is a dynamic and complex community, whose structure has been molecularly dissected  
333 (Roesti et al. 2005; Long et al. 2008). However, isolation in pure culture of putatively beneficial  
334 microbiota represents an indispensable step in order to integrate knowledge of spore-associated  
335 bacterial communities obtained through molecular studies with data on their functional properties,

336 in the perspective of opening new avenues for targeted management of beneficial symbionts and  
337 their associated bacteria in sustainable food production systems.

338 Actinobacteria represented 36.1% of the total culturable bacteria recovered from *R. intraradices*  
339 spores, consistently with previous data on their widespread occurrence in the mycorrhizosphere and  
340 sporosphere (Ames et al. 1989; Filippi et al. 1998; Bharadwaj et al. 2008a). Recent findings,  
341 obtained using a culture-independent approach - PCR-DGGE analysis of the 16S rRNA gene -  
342 showed that DNA sequences affiliated with the order *Actinomycetales* were found in spore  
343 homogenates of six different AMF isolates originating from diverse geographical areas. Such AMF  
344 spores harbored different actinobacterial species, which were affiliated to the genera *Streptomyces*,  
345 *Arthrobacter*, *Amycolatopsis* and *Propionibacterium* (Agnolucci et al. 2015). Similar results were  
346 obtained by Long et al. (2008) who found *Streptomyces*, *Amycolatopsis*, and *Pseudonocardia*  
347 species associated with *G. margarita* spores. Their physiological characteristics, such as the ability  
348 to produce a vast array of enzymes that break down insoluble organic polymers, including chitin  
349 and chitosan, the major components of spore walls, may explain their intimate association with  
350 AMF spores. Indeed, 23 chitinolytic strains were isolated from *R. intraradices* spores, confirming  
351 previous findings on spore wall degrading activity of culturable bacteria embedded in spore walls of  
352 *Funneliformis mosseae*, where chitinolytic strains represented 72% of all the isolated  
353 microorganisms (Filippi et al. 1998). Other authors, using culture-independent methods, reported  
354 that most DNA sequences obtained from *Glomus geosporum* and *Glomus constrictum* spores were  
355 affiliated with bacterial strains able to hydrolyze biopolymers (Roesti et al. 2005). Such  
356 physiological traits may play important functional roles in the promotion of spore germination and  
357 germling growth, thus positively affecting AMF root colonisation and functioning (Mayo et al.  
358 1986; Xavier and Germida 2003; Bharadwaj et al. 2008b; Giovannetti et al. 2010).

359 A very interesting result is represented by the isolation of 27 bacterial strains from N-free medium,  
360 10 (37%) of which produced the expected *nifH* gene 390 bp amplicon, suggesting their possible role  
361 as biofertilizers by mediating the acquisition of nitrogen, a major plant nutrient. Such data confirm

362 our previous molecular findings, showing that nitrogen fixing bacteria, *Sinorhizobium meliloti*,  
363 *Agrobacterium radiobacter*, *Rhizobium giardinii* and *Rhizobium rhizogenes*, were associated with  
364 spores of different AMF isolates (Agnolucci et al. 2015). Bacteria belonging to *Rhizobiales* have  
365 been rarely isolated from AMF spores (Bharadwaj et al. 2008a), though their co-inoculation with  
366 AMF has been shown to positively affect nutrient uptake and photosynthetic rate in diverse plant  
367 species and to promote mycorrhizal functioning, improving spore germination, mycelial growth and  
368 mycorrhizal colonization (Gopal et al. 2012).

369

#### 370 *Functional diversity of bacterial isolates*

371 Among the 374 bacterial strains isolated, 122 were selected for functional significance studies: 38  
372 heterotrophs (including 15 spore forming bacteria), 34 actinobacteria, 23 chitinolytic and 27  
373 putative N<sub>2</sub>-fixing bacteria were further screened in pure culture for PGP traits, such as the ability to  
374 produce IAA and siderophores, and to solubilize P from mineral phosphate and phytate.

375 We obtained a high percentage of IAA producing strains (42.6% ), with 76% of positive isolates  
376 among actinobacteria. As IAA produced by different bacterial strains is able to stimulate the  
377 development of plant root systems (Glick et al. 1995; Patten et al. 2002), while displaying no  
378 functions in bacterial cells, IAA-producing strains may play a beneficial role in the performance of  
379 the complex plant-AMF interaction.

380 Siderophore-producing strains represented 65.6% of the total isolates, although their activity was  
381 differentially expressed in the different bacterial groups. Such a trait may be functional to AMF  
382 potential biocontrol activity against soilborne diseases, by means of bacterial siderophore-mediated  
383 competition for iron (Thomashow et al. 1990; Glick 1995; Whipps 2001), thus indirectly  
384 stimulating plant growth by inhibiting the development of deleterious plant pathogens (Davison  
385 1988; Arora et al. 2001).

386 The ability to solubilize P was shown by 69.7% and 49.2% of SAB, when tested in the presence of  
387 phytate and inorganic P, respectively. The majority of P-solubilizing strains were recovered from

388 Waksman's agar medium and minimal medium containing chitin as sole carbon source. Other  
389 authors investigated the occurrence of P-solubilizing bacteria associated with *F. mosseae* spores in  
390 the mycorrhizosphere of *Psidium guajava* L. and found that all the isolates showing high P-  
391 solubilizing activity belonged to the genera *Streptomyces* and *Leifsonia* (Mohandas et al. 2013).  
392 Since P is a key plant nutrient usually applied as fertilizer in the form of inorganic phosphates,  
393 which are rapidly immobilized in the soil, thus becoming unavailable to plant roots, P-mobilizing  
394 bacteria could act in synergy with AMF symbionts to enhance P availability to host plants  
395 (Rodriguez and Fraga 1999). AMF are increasingly reported to play a fundamental role in P  
396 biogeochemical cycle, improving P availability in the soil solution by means of the large  
397 extraradical mycelial networks spreading from mycorrhizal roots into the soil environment and able  
398 to absorb P far from the depletion zone surrounding the roots. As phosphate transporter genes are  
399 differentially expressed on such hyphae, we could hypothesize that the higher P concentrations  
400 found in AMF inoculated plants may be ascribed also to the activity of large communities of  
401 mineral phosphate and phytate solubilizing bacteria (Harrison and van Buuren 1995; Fiorilli et al.  
402 2013).

403 Finally, the highest number of high siderophore producing strains (halo zone > 15mm) (TSA20,  
404 TSA44, TSA49, TSA108B and TSA120) and the two best phosphate solubilizing bacteria (TSA41  
405 with SE =150 and TSA3 with SE =115.38), along with one out of the three best IAA producers  
406 strains (TSA50, CH8, CH17) were found among heterotrophic bacteria (Tables 2 and 4).

407

#### 408 *Identification of bacterial isolates*

409 Nineteen strains with best potential PGP activities were selected and identified by 16S rDNA  
410 sequencing. Interestingly, 9 out of 19 strains (47,4%) were identified as *Sinorhizobium meliloti*:  
411 three strains - TSA3, TSA26 and TSA41 - were able to produce IAA, six strains - TSA3, TSA26  
412 and TSA41, N23, N28, N29 - were able to solubilize mineral and organic P *in vitro* and the three  
413 chitinolytic strains - CH5, CH8, CH17 - showed siderophore and IAA production activity and



414 solubilized mineral and organic P *in vitro* (Tables 2, 4 and 5). The strict association of members of  
415 the order *Rhizobiales* with AMF spores may be ascribed to their ability to form biofilms by  
416 producing exopolysaccharides, allowing an efficient colonization of roots and mycorrhizal hyphae  
417 (Bianciotto et al. 1996; Toljander et al. 2006). In addition to the beneficial effect in terms of  
418 biological nitrogen fixation, rhizobia promote plant growth by producing phytohormones,  
419 improving plant nutritional status and biocontrolling phytopathogens (Chandra et al. 2007). Here,  
420 three strains of *S. meliloti* were isolated from a medium containing chitin as sole carbon source,  
421 confirming previous data on the ability of several rhizobial strains isolated from root nodules to  
422 produce chitinolytic enzymes (Sridevi et al. 2008). Such data highlight the multifunctionality of our  
423 isolates, which could be further investigated in order to select the most efficient strains able to  
424 degrade fungal cell walls, thus acting as biocontrol agents against fungal pathogens (Mazen et al.  
425 2008). Indeed, two chitinase producing *Bradyrhizobium* strains were reported to inhibit mycelial  
426 growth, sclerotia formation and germination of *Macrophomina phaseolina* (Tassi) Goid., a major  
427 pathogen of more than 500 plant hosts (Dubey et al. 2012).

428 Five strains (26.3%) were assigned to *Streptomyces* (W43N, W64, W77, W94 and W115), a genus  
429 representing a predominant component of the soil microbial population, capable of producing a vast  
430 array of complex and biologically active secondary metabolites, including antibacterial, antifungal,  
431 antiparasitic, anticancer and immunosuppressant drugs. These strains showed high growth  
432 promoting potential exhibiting all the PGP traits tested. Among the species affiliated to our strains,  
433 *Streptomyces phaeochromogenes* was reported to produce chloramphenicol, and one strain (LL-  
434 P018) to produce phaeochromycins, a novel anti-inflammatory polyketides inhibitors of the  
435 biosynthesis of a tumor necrosis factor (TNF- $\alpha$ ) (van Pée and Lingens 1985; Ritacco and Eveleigh  
436 2008), *Streptomyces collinus* strain Tü 365 to produce the antibiotic kirromycin (Wolf and Zähler  
437 1972), *Streptomyces iakyrus* DSM 41873 actinomycin G (Qin et al. 2014) and *Streptomyces*  
438 *viridochromogenes* Tü57 avilamycin A (Weitnauer et al. 2001). In addition, *Streptomyces* spp. are  
439 considered promising taxa of PGP and mycorrhizal helper bacteria, due to their ability to solubilize

440 phosphates, produce chitinase and growth regulators (Mohandas et al. 2013; Hamedi and  
441 Mohammadipannah 2015) and stimulate AMF spore germination and hyphal growth (Mugnier and  
442 Mosse 1987; Tylka et al. 1991; Carpenter-Boggs et al. 1995). Interestingly, sequences affiliated to  
443 *Streptomyces flavogriseus* and *S. phaeochromogenes*, with a similarity of 99%, were recovered  
444 from *F. mosseae* and *R. intraradices* IMA6, respectively (Agnolucci et al. 2015). The other two  
445 isolates affiliated with *Actinomycetales* were represented by *Arthrobacter phenanthrenivorans*  
446 (N17) and *Nocardioides albus* (N13). Both isolates were able to grow on the N-free selective  
447 medium. For a long time it was believed that the ability to fix atmospheric nitrogen was limited to  
448 the actinomycetes *Frankia*, but lately, nitrogen fixation (*nifH*) genes were found in other non-  
449 *Frankia* actinomycetes including *Streptomyces*, *Arthrobacter* (Gtari et al. 2012; Sharon and Daniel  
450 2013) and *Nocardioides*.

451 Actinobacteria of the genus *Arthrobacter* are ubiquitous in all soil types and able to utilize a wide  
452 range of natural as well as xenobiotic compounds. In particular, *A. phenanthrenivorans* strain Sphe3  
453 showed the ability to grow on phenanthrene as the sole carbon and energy source (Vandera et al.  
454 2015). *Arthrobacter* species have been reported to be associated with AMF spores and hyphae of *F.*  
455 *mosseae* and *R. intraradices* (Bharadwaj et al. 2008a), although Andrade et al. (1997) found that  
456 they were most frequent in the hyphosphere, the zone of soil surrounding individual AMF hyphae  
457 (Artursson et al. 2006). It is interesting to note that *A. phenanthrenivorans* sequences were  
458 previously recovered from *F. coronatum*, *F. mosseae* and *R. intraradices*, including the isolate  
459 IMA6 analysed in the present study (Agnolucci et al. 2015). The strict association of *Arthrobacter*  
460 spp. with AMF spores may be functional to their role as PGP and mycorrhizal helper bacteria:  
461 indeed, some *Arthrobacter* strains have been reported to improve mycorrhizal colonization and root  
462 length, and to display antagonistic activity against plant pathogens (Bharadwaj et al. 2008b).

463 Three isolates were affiliated to *Bacillales*, namely to a species of *Bacillus pumilus* group (CH10),  
464 *Lysinibacillus fusiformis* (CH19) and *Fictibacillus barbaricus* (TSA50). *Bacillus* isolates closely  
465 related to *B. pumilus* are not easily distinguished from each other. Definitely, the *B. pumilus* group

466 contains 5 species, *B. pumilus*, *B. safensis*, *B. stratosphericus*, *B. altitudinis* and *B. aerophilus*,  
467 which are nearly identical in 16S rRNA gene sequence, sharing similarity over 99.5% (Liu et al.  
468 2013). *Lysinibacillus fusiformis* and *Fictibacillus barbaricus* originally assigned to the genus  
469 *Bacillus* were reclassified respectively in 2007 (Trivedi et al. 2011) and in 2013 (Glaeser et al.  
470 2013). *Bacillus* strains are among the major chitin decomposers, producing and secreting many  
471 kinds of chitinase (Heravi et al. 2013). For example, *Bacillus pumilus* strain SG2 produces two  
472 chitinases, namely ChiS and ChiL (Heravi et al. 2013), and *Lysinibacillus fusiformis* strain B-CM18  
473 a purified chitinase of 20 Kd, possessing an *in vitro* strong antifungal activity (Singh et al. 2013).  
474 Accordingly, our isolates assigned to *Bacillus pumilus* group and to *Lysinibacillus fusiformis* were  
475 retrieved from the selective media for the isolation of chitinolytic bacteria. Interestingly, *Bacillus*  
476 *pumilus* strains have a wide range of attributed applications, as PGP rhizobacteria and animal and  
477 human probiotic (Branquinho et al. 2014). An interesting example of such application is represented  
478 by the endophytic strain *Bacillus pumilus* INR7, which has been commercialized as a biological  
479 control product, active by direct antagonism and induction of systemic resistance, against soilborne  
480 pathogens as well as foliar pathogens (Jeong et al. 2014).

481 Interestingly, the isolation in pure culture of bacterial strains belonging to *Bacillales* support our  
482 previous data on the occurrence of DNA sequences affiliated with *Paenibacillus castaneae* and  
483 *Bacillus firmus* in spore homogenates of *F. mosseae* and *R. intraradices*, respectively (Agnolucci et  
484 al. 2015). The role of members of *Bacillales* as PGP has long been known and their activity as  
485 mycorrhizal helper bacteria, increasing mycorrhizal establishment and promoting plant growth, has  
486 been recently reported (Budi et al. 2013; Zhao et al. 2014; Pérez-Montaña et al. 2014).

487 In conclusion, our work demonstrates that AMF spores are a privileged source of bacteria with  
488 potential PGP activities, whose isolation and functional characterization represent a prerequisite for  
489 their use as biocontrol agents, biofertilisers and bioenhancers. The emerging picture of  
490 AMF/bacteria interactions suggests that different partners of tripartite associations - host plants,  
491 AMF and bacteria - may act in synergy and provide new multifunctional benefits, improving plant

492 and fungal performances. SAB can be transferred from spores to soil-based hyphae, where they may  
493 enhance nutrient availability (phosphate solubilizing, nitrogen fixing and chitinolytic bacteria),  
494 control plant pathogens (siderophore producing bacteria) and promote plant growth (IAA producing  
495 bacteria).

496 Further investigations are in progress aimed at selecting the best performing AMF/bacteria  
497 combinations in the perspective of exploiting the potential of such multipartite association in a new  
498 food production system, aimed at maintaining and increasing soil biological fertility and protecting  
499 natural processes that are at the basis of energy flows and matter cycles in sustainable  
500 agroecosystems.

501

## 502 **Acknowledgments**

503 This work was funded by a University of Pisa grant (Fondi di Ateneo).

504

## 505 **References**

- 506 Agnolucci M, Battini F, Cristani C, Giovannetti M. Diverse bacterial communities are recruited on  
507 spores of different arbuscular mycorrhizal fungal isolates. *Biol Fertil Soils* 2015;51:379–389.
- 508 Ames RN, Mihara KL, Bayne HG. Chitin-decomposing actinomycetes associated with a vesicular-  
509 arbuscular mycorrhizal fungus from a calcareous soil. *New Phytol* 1989;111:67–71.
- 510 Andrade G, Mihara KL, Linderman RG, Bethlenfalvay GJ. Bacteria from rhizosphere and  
511 hyphosphere soils of different arbuscular-mycorrhizal fungi. *Plant Soil* 1997;192:71–79.
- 512 Arora NK, Kang SC, Maheshwari DK. Isolation of siderophore-producing strains of *Rhizobium*  
513 *meliloti* and their biocontrol potential against *Macrophomina phaseolina* that causes charcoal rot of  
514 groundnut. *Curr Sci* 2001;81:673–677.
- 515 Artursson V, Finlay RD, Jansson JK. Interactions between arbuscular mycorrhizal fungi and  
516 bacteria and their potential for stimulating plant growth. *Environ Microbiology* 2006;8:1–10.

517 Augé RM. Water relations, drought and vesicular-arbuscular mycorrhizal symbiosis. *Mycorrhiza*  
518 2001;11:3–42.

519 Barea JM, Azcon R, Azcon-Aguilar C. Mycorrhizosphere interactions to improve plant fitness and  
520 soil quality. *Antonie Van Leeuwenhoek* 2002;81:343–351.

521 Bharadwaj DP, Lundquist PO, Persson P, Alström S. Evidence for specificity of cultivable bacteria  
522 associated with arbuscular mycorrhizal fungal spores. *FEMS Microbiol Ecol* 2008a;65:310–322.

523 Bharadwaj DP, Lundquist PO, Alström S. Arbuscular mycorrhizal fungal spore-associated bacteria  
524 affect mycorrhizal colonization, plant growth and potato pathogens. *Soil Biol Biochem*  
525 2008b;40:2494–2501.

526 Bharadwaj D, Ansari MW, Sahoo RK, Tuteja N. Biofertilizers function as key player in sustainable  
527 agriculture by improving soil fertility, plant tolerance and crop productivity. *Microb Cell Fact*  
528 2014;13:66.

529 Bhattacharyya PN, Jha DK. Plant growth-promoting rhizobacteria (PGPR): emergence in  
530 agriculture. *World J Microbiol Biotechnol* 2012;28:1327–1350.

531 Bianciotto V, Bandi CD, Minerdi M, Sironi H, Tichy V, Bonfante P. An obligately endosymbiotic  
532 mycorrhizal fungus itself harbors obligately intracellular bacteria. *Appl Environ Microbiol* 1996;62:  
533 3005–3010.

534 Branquinho R, Sousa C, Osório H, Meirinhos-Soares L, Lopes J, Carriço JA Busse HJ,  
535 Abdulmawjood A, Klein G, Kämpfer P, Pintado ME, Peixe LV. *Bacillus invictae* sp. nov., isolated  
536 from a health product. *Int J Syst Evol Microbiol* 2014;64:3867–3876.

537 Budi SW, Van Tuinen D, Martinotti G, Gianinazzi S. Isolation from *Sorghum bicolor*  
538 mycorrhizosphere of a bacterium compatible with arbuscular mycorrhiza development and  
539 antagonistic towards soil-borne fungal pathogens. *Appl Environ Microbiol* 1999;65:5148–5150.

540 Budi SW, Bakhtiar Y, May NL. Bacteria associated with arbuscula mycorrhizal spores *Gigaspora*  
541 *margarita* and their potential for stimulating root mycorrhizal colonization and neem (*Melia*  
542 *azedarach* Linn) seedling growth. *Microbiol Indones* 2013;6:180–188.

543 Carpenter-Boggs L, Loynachan TE, Stahl PD. Spore germination of *Gigaspora margarita*  
544 stimulated by volatiles of soil-isolated actinomycetes. *Soil Boil Biochem* 1995;27:1445–1451.

545 Chandra S, Choure K, Dubey RC, Maheshwari DK. Rhizosphere competent *Mesorhizobium loti*  
546 MP6 induces root hair curling, inhibits *Sclerotinia sclerotiorum* and enhances growth of Indian  
547 mustard (*Brassica campestris*). *Braz J Microbiol* 2007;38:124–130.

548 Citernesi AS, Fortuna P, Filippi C, Bagnoli G, Giovannetti M. The occurrence of antagonistic  
549 bacteria in *Glomus mosseae* pot cultures. *Agronomie* 1996;16:671–677.

550 Cruz AF, Horii S, Ochiai S, Yasuda A, Ishii T. Isolation and analysis of bacteria associated with  
551 spores of *Gigaspora margarita*. *J Appl Microbiol* 2008;104:1711–1717.

552 Cruz AF, Ishii T. Arbuscular mycorrhizal fungal spores host bacteria that affect nutrient  
553 biodynamics and biocontrol of soil- borne plant pathogens. *Biol Open* 2011;1:52–57.

554 Davison J. Plant beneficial bacteria. *Nat Biotechnol* 1088;6:282–286.

555 Dubey D, Sahu MC, Rath S, Paty BP, Debata NK, Padhy RN. Antimicrobial activity of medicinal  
556 plants used by aborigines of Kalahandi, Orissa, India against multidrug resistant bacteria. *Asian Pac*  
557 *J Trop Biomed* 2012;2:S846–S854.

558 Edgar RC. MUSCLE: a multiple sequence alignment method with reduced time and space  
559 complexity. *BMC Bioinformatics* 2004a;5:113.

560 Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high throughput.  
561 *Nucleic Acids Res* 2004b;32:1792–1797.

562 Evelin H, Kapoor R, Giri B. Arbuscular mycorrhizal fungi in alleviation of salt stress: a review.  
563 *Ann Bot* 2009;104:1263–1280.

564 Filippi C, Bagnoli G, Citernesi AS, Giovannetti M. Ultrastructural spatial distribution of bacteria  
565 associated with sporocarps of *Glomus mosseae*. *Symbiosis* 1998;24:1–12.

566 Fiorilli V, Lanfranco L, Bonfante P. The expression of *GintPT*, the phosphate transporter of  
567 *Rhizophagus irregularis*, depends on the symbiotic status and phosphate availability. *Planta*  
568 2013;237:1267–1277.

569 Gerdermann JW, Nicolson TH. Spores of mycorrhizal *Endogone* species extracted from soil by wet  
570 sieving and decanting. *Trans Br Mycol Soc* 1963;46:235–244.

571 Gianinazzi S, Gollotte A, Binet MN, van Tuinen D, Redecker D, Wipf D. Agroecology: the key role  
572 of arbuscular mycorrhizas in ecosystem services. *Mycorrhiza* 2010;20:519–530.

573 Giovannetti M, Avio L. Biotechnology of arbuscular mycorrhizas. In: Khachatourians GG, Arora  
574 DK, editors. *Applied mycology and biotechnology*, vol. 2. Agriculture and food production, 2002.  
575 p. 275–310.

576 Giovannetti M, Avio L, Sbrana C. Fungal spore germination and pre-symbiotic mycelial growth–  
577 physiological and genetic aspects. In: Koltai H, Kapulnik Y, editors. *Arbuscular mycorrhizas:*  
578 *physiology and function*. Dordrecht: Springer; 2010. p. 3–32.

579 Giovannetti M, Avio L, Barale R, Ceccarelli N, Cristofani R, Iezzi A, Mignolli F, Picciarelli P,  
580 Pinto B, Reali D, Sbrana C, Scarpato R. Nutraceutical value and safety of tomato fruits produced by  
581 mycorrhizal plants. *Br J Nutr* 2012;107:242–251.

582 Glaeser SP, Wolfgang D, Busse H-J, Kampf P. *Fictibacillus phosphorivorans* gen. nov., sp. nov.  
583 and proposal to reclassify *Bacillus arsenicus*, *Bacillus barbaricus*, *Bacillus macauensis*, *Bacillus*  
584 *nanhaiensis*, *Bacillus rigui*, *Bacillus solisalsi* and *Bacillus gelatini* in the genus *Fictibacillus*. *Int J*  
585 *Syst Evol Microbiol* 2013;63:2934–2944.

586 Glick BR. The enhancement of plant growth by free- living bacteria. *Can J Microbiol* 1995;41:109–  
587 117.

588 Gopal S, Chandrasekaran M, Shagol C, Kim K, Sa T. Spore associated bacteria (SAB) of arbuscular  
589 mycorrhizal fungi (AMF) and plant growth promoting rhizobacteria (PGPR) increase nutrient  
590 uptake and plant growth under stress conditions. *Korean J Soil Sci Fert* 2012;45:582–592.

591 Gtari M, Ghodhbane-Gtari F, Nouioui I, Beauchemin N, Tisa LS. Phylogenetic perspectives of  
592 nitrogen-fixing actinobacteria. *Arch Microbiol* 2012;194:3–11.

593 Hamedi J, Mohammadipanah F. Biotechnological application and taxonomical distribution of plant  
594 growth promoting actinobacteria. *J Ind Microbiol Biotechnol* 2015;42: 157–171.

595 Harrison MJ, van Buuren ML. A phosphate transporter from the mycorrhizal fungus *Glomus*  
596 *versiforme*. *Nature* 1995;378:626–629.

597 Heravi KM, Shali A, Naghibzadeh N, Ahmadian G. Characterization of *cis*-acting elements residing  
598 in the chitinase promoter of *Bacillus pumilus* SG2. *World J Microbiol Biotechnol* 2014;30:1491–  
599 1499.

600 Horii S, Ishii T. Identification and function of *Gigaspora margarita* growth-promoting  
601 microorganisms. *Symbiosis* 2006;41:135–141.

602 Islam MT, Deora, A, Hashidoko Y, Rahman A, Ito T, Tahara S. Isolation and identification of  
603 potential phosphate solubilizing bacteria from the rhizoplane of *Oryza sativa* L. cv. BR29 of  
604 Bangladesh. *Z Naturforsch C* 2007;62:103–110.

605 Jeong H, Choi SK, Kloepper JW, Ryu CM. Genome sequence of the plant endophyte *Bacillus*  
606 *pumilus* INR7, triggering induced systemic resistance in field crops. *Genome Announc* 2014;doi:  
607 10.1128/genomeA.01093-14.

608 Jorquera MA, Hernández MT, Rengel Z, Marschner P, De la Luz Mora M. Isolation of culturable  
609 phosphobacteria with both phytate-mineralization and phosphate-solubilization activity from the  
610 rhizosphere of plants grown in a volcanic soil. *Biol Fertil Soils* 2008;44:1025–1034.

611 Kimura M. A simple method for estimating evolutionary rate of base substitutions through  
612 comparative studies of nucleotide sequences. *J Mol Evol* 1980;16:111–120.

613 Lane DJ. 16S/23S rRNA sequencing. In: Stackebrandt E, Goodfellow M. editors. *Nucleic acid*  
614 *techniques in bacterial systematics*. Chichester: Wiley; 1991. p. 115–175.

615 Lecomte J, St-Arnaud M, Hijri M. Isolation and identification of soil bacteria growing at the  
616 expense of arbuscular mycorrhizal fungi. *FEMS Microbiol Lett* 2011;317:43–51.

617 Li B, Ravnskov S, Xie G, Larsen J. Biocontrol of *Pythium* damping-off in cucumber by arbuscular  
618 mycorrhiza-associated bacteria from the genus *Paenibacillus*. *Biocontrol* 2007;52:863–875.



619 Liu Y, Lai Q, Dong C, Sun F, Wang L, Li G, Shao Z. Phylogenetic diversity of the *Bacillus pumilus*  
620 group and the marine ecotype revealed by multilocus sequence analysis. PLoS ONE 8:e80097.  
621 2013;doi: 10.1371/journal.pone.0080097.

622 Long L, Zhu H, Yao Q, Ai Y. Analysis of bacterial communities associated with spores of  
623 *Gigaspora margarita* and *Gigaspora rosea*. Plant Soil 2008;310:1–9.

624 Louden BC, Haarmann D, Lynne AM. Use of blue agar CAS assay for siderophore detection. J  
625 Microbiol Biol Educ 2011;12:51–53.

626 MacDonald RM, Chandler MR. Bacterium-like organelles invesicular-arbuscular mycorrhizal fungus  
627 *Glomus caledonium*. New Phytol 1981;89:241–246.

628 MacDonald RM, Chandler MR, Mosse B. The occurrence of bacterium-like organelles in vesicular-  
629 arbuscular mycorrhizal fungi. New Phytol 1982;90:659–663.

630 Mayo K, Davis RE, Motta J. Stimulation of germination of spores of *Glomus versiforme* by spore-  
631 associated bacteria. Mycologia 1986;78:426–431.

632 Mazen MM, El-Batanony NH, Abd El-Monium MM, Massoud ON. Cultural filtrate of *Rhizobium*  
633 spp. and arbuscular mycorrhiza are potential biological control agents against root rot fungal  
634 diseases of faba bean. Global J Biotechnol Biochem 2008;3:32–41.

635 Mohandas S, Poovarasan S, Panneerselvam P, Saritha B, Upreti KK, Kamal R, Sita T. Guava  
636 (*Psidium guajava* L.) rhizosphere *Glomus mosseae* spores harbor actinomycetes with growth  
637 promoting and antifungal attributes. Sci Hortic Amsterdam 2013;150:371-376.

638 Mosse B. Honey-coloured sessile *Endogone* spores. II. Changes in fine structure during spore  
639 development. Arch Mikrobiol 1970;74:146–159.

640 Mugnier J, Mosse B. Spore germination and viability of a vesicular arbuscular mycorrhizal fungus,  
641 *Glomus mosseae*. Trans Br Mycol Soc 1987;88:411–413.

642 Nautiyal CS. An efficient microbiological growth medium for screening phosphate solubilizing  
643 microorganisms. FEMS Microbiol Lett 1999;170: 265–270.

644 Njeru EM, Avio L, Bocci G, Sbrana C, Turrini A, Bàrberi P, Giovannetti M, Fritz O. Contrasting  
645 effects of cover crops on ‘hot spot’ arbuscular mycorrhizal fungal communities in organic tomato.  
646 Biol Fertil Soils 2014;51:151–166.

647 Patten CL, Glick BR. Role of *Pseudomonas putida* indoleacetic acid in development of the host  
648 plant root system. Appl Environ Microbiol 2002;68: 3795–3801.

649 Pérez-Miranda S, Cabirol N, George-Téllez R, Zamudio-Rivera LS, Fernández FJ. O-CAS, a fast  
650 and universal method for siderophore detection. J Microbiol Meth 2007;70:127–131.

651 Pérez-Montaña F, Alías-Villegas C, Bellogín RA, Del Cerro P, Espuny MR, Jiménez-Guerrero I,  
652 López-Baena FJ, Ollero FJ, Cubo T. Plant growth promotion in cereal and leguminous agricultural  
653 important plants: from microorganism capacities to crop production. Microbiol Res 2014;169:325-  
654 336.

655 Qin Z, Wang X, Rateb ME, Ass'ad LA, Jaspars M, Deng Z, Yu Y, Deng H. Disruption of a  
656 methyltransferase gene in actinomycin G gene cluster in *Streptomyces iakyrus* increases the  
657 production of phenazinomycin. FEMS Microbiol Lett 2014;352: 62–68.

658 Rambelli A. The rhizosphere of mycorrhizae. In: Marks GL, Koslowski TT. editors.  
659 Ectomycorrhizae. New York: Academic; 1973. p. 299–343.

660 Ritacco FV, Eveleigh DE. Molecular and phenotypic comparison of phaeochromycin-producing  
661 strains of *Streptomyces phaeochromogenes* and *Streptomyces ederensis*. J Ind Microbiol Biotechnol  
662 2008;35:931–945.

663 Rodriguez H, Fraga R. Phosphate solubilizing bacteria and their role in plant growth promotion.  
664 Biotechnol Adv 1999;17: 319–339.

665 Roesti D, Ineichen K, Braissant O, Redecker D, Wiemken A, Aragno M. Bacteria associated with  
666 spores of the arbuscular mycorrhizal fungi *Glomus geosporum* and *Glomus constrictum*. Appl  
667 Environ Microbiol 2005;71:6673–6679.

668 Rokhbakhsh-Zamin F, Sachdev D, Kazemi-Pour N, Engineer A, Pardesi KR, Zinjarde S Chopade  
669 BA. Characterization of plant-growth-promoting traits of *Acinetobacter* species isolated from  
670 rhizosphere of *Pennisetum glaucum*. J Microbiol Biotechnol 2011;21:556–566.

671 Sharon FB, Daniel RR. Homology Modeling of Nitrogenase Iron Protein of Nitrogen Fixing  
672 Actinomycete *Arthrobacter* sp. 2013;Int J Comput Appl 61:13–19.

673 Singh RK, Kumar DP, Solanki MK, Singh P, Srivastva AK, Kumar S, Kashyap PL, Saxena AK,  
674 Singhal PK, Arora DK. Optimization of media components for chitinase production by chickpea  
675 rhizosphere associated *Lysinibacillus fusiformis* B-CM18. J Basic Microbiol 2013;53:451–460.

676 Sikes BA, Kottenie K, Klironomos JN. Plant and fungal identity determines pathogen protection of  
677 plant roots by arbuscular mycorrhizas. J Ecol 2009;97:1274–1280.

678 Smith SE, Read DJ. Mycorrhizal symbiosis. Cambridge: Academic; 2008.

679 Souza CP, Burbano-Rosero EM, Almeida BC, Martins GG, Albertini LS, Rivera ING. Culture  
680 medium for isolating chitinolytic bacteria from seawater and plankton. World J Microbiol  
681 Biotechnol 2009;25:2079–2082.

682 Sridevi M, Veera Mallaiah K. Bioproduction of indole acetic acid by *Rhizobium* strains isolated  
683 from root nodules of green manure crop, *Sesbania sesban* (L.) Merr. Iranian J Biotech 2007;5:178–  
684 182.

685 Tamura K, Stecher G, Peterson D, Filipinski A, Kumar S. MEGA6: molecular evolutionary genetics  
686 analysis version 6.0. Mol Biol Evol 2013;30:2725–2729.

687 Tchan YT. Azotobacteraceae. In: Krieg N, Holt JG. editors. Bergey’s manual of systematic  
688 bacteriology, vol. 1. London: Williams and Wikins; 1984. p. 219–225.

689 Thomashow LS, Weller DM, Bonsall RF, Pierson LS. Production of the antibiotic phenazine-1-  
690 carbox-yllic acid by fluorescent *Pseudomonas* species in the rhizosphere of wheat. Appl Environ  
691 Microbiol 1990;56:908–912.

692 Toljander JF, Artursson V, Paul LR, Jansson JK, Finlay RD. Attachment of different soil bacteria to  
693 arbuscular mycorrhizal fungal extraradical hyphae is determined by hyphal vitality and fungal  
694 species. *FEMS Microbiol Lett* 2006;254:34–40.

695 Trivedi P, Spann T, Wang N. Isolation and characterization of beneficial bacteria associated with  
696 citrus roots in Florida. *Microb Ecol* 2011;62:324–336.

697 Tylka GL, Hussey RS, Roncadori RW. Axenic germination of vesicular-arbuscular mycorrhizal  
698 fungi: effects of selected *Streptomyces* species. *Phytopathology* 1991;81:754–759.

699 Ueda T, Suga Y, Yahiro N, Matsuguchi T. Phylogeny of Sym plasmids of rhizobia by PCR-based  
700 sequencing of a *nodC* segment. *J Bacteriol* 1995;177:468–472.

701 van Pée KH, Lingens F. Detection of a bromoperoxidase in *Streptomyces phaeochromogenes*.  
702 *FEBS Lett* 1984;173:5–8.

703 Vandera E, Samiotaki M, Parapouli M, Panayotou G, Koukkou AI. Comparative proteomic analysis  
704 of *Arthrobacter phenanthrenivorans* Sphe3 on phenanthrene, phthalate and glucose. *J Proteomics*  
705 2015;113:73-89.

706 Walley FL, Germida JJ. Failure to decontaminate *Glomus clarum* NT4 spores is due to spore wall-  
707 associated bacteria. *Mycorrhiza* 1996;6:43–49.

708 Weisburg WG, Barns SM, Pelletier DA, Lane DJ. 16S ribosomal DNA amplification for  
709 phylogenetic study. *J Bacteriol* 1991;173:697–703.

710 Weitnauer G, Mühlenweg A, Trefzer A, Hoffmeister D, Süssmuth RD, Jung G, Welzel K, Vente A,  
711 Girreser U, Bechthold A. Biosynthesis of the orthosomycin antibiotic avilamycin A: deductions  
712 from the molecular analysis of the *avi* biosynthetic gene cluster of *Streptomyces viridochromogenes*  
713 Tü57 and production of new antibiotics. *Chem Biol* 2001;8:569–581.

714 Whipps JM. Microbial interactions and biocontrol in the rhizosphere. *J Exp Bot* 2001;52:487–511.

715 Wolf H, Zähner H. Metabolic products of microorganism. *Arch Mikrobiol* 1972;83:147–154.

716 Xavier LJC, Germida JJ. Bacteria associated with *Glomus clarum* spores influence mycorrhizal  
717 activity. *Soil Biol Biochem* 2003;35:471–478.

718 Zhao L, Wu XQ, Ye JR, Li H, Li GE. Isolation and characterization of a mycorrhiza helper  
719 bacterium from rhizosphere soils of poplar stands. *Biol Fertil Soils* 2014;50:593–601.  
720

721 **Table 1** Abundance of culturable bacteria isolated from three batches (A, B, C) of 200 spores of  
722 *Rhizophagus intraradices* IMA6 on different media (mean CFU spore<sup>-1</sup> ± SE). TSA, Tryptic soil  
723 agar; TSAT, TSA plus thermic treatment; W, Waksman agar; N, nitrogen-free medium; CH, chitin  
724 agar.

Media	A	B	C
TSA	22.9±0.7	7.9±1.6	5.4±0.9
TSAT	2.9±0.7	1.9±0.4	0.8±0.4
W	23.0±1.5	6.1±0.5	3.7±0.4
N	nd	1.0±0.6	0.7±0.5
CH	nd	0.2±0.1	1.1±0.1

726 nd = not determined.

727

728

729

730

731

732

733

734

735

736

737

738

739

740

741

742

743 **Table 2** Plant growth promoting traits of heterotrophic bacteria isolated from spores of  
 744 *Rhizophagus intraradices* IMA6.

Isolate	IAA	Siderophore	Phosphate		Phytate
	production	production	solubilization	solubilization	solubilization
		Activity (after 7 days)	SE(%)	PSI	Halo zone (cm)
TSA2	+	-	-	-	-
TSA3	++	-	115.38	2.15	0.85
TSA10	-	+	35.71	1.36	0.85
TSA13	-	-	-	-	-
TSA20	-	+++	-	-	0.40
TSA25	-	-	-	-	0.20
TSA26	+	-	81.82	1.82	0.90
TSA32	-	-	-	-	-
TSA39B	-	-	-	-	-
TSA39G	-	++	-	-	0.40
TSA41	++	-	150	2.50	0.70
TSA43B	-	-	-	-	0.45
TSA44	-	+++	-	-	0.35
TSA46	-	+	11.76	1.12	0.25
TSA47	-	-	-	-	-
TSA49	-	+++	-	-	0.40
TSA50	+++	-	-	-	-
TSA58	-	+	-	-	-
TSA60	+	-	-	-	-
TSA108B	-	+++	-	-	0.65
TSA113	-	-	-	-	-
TSA136	-	-	-	-	-
TSA142	-	-	-	-	0.35
TSAT5G	-	-	-	-	-
TSAT7	-	-	-	-	-
TSAT11	-	-	-	-	0.15
TSAT14	-	+	-	-	-
TSAT28	-	-	-	-	-
TSAT38	+	-	-	-	-
TSAT50B	+	-	-	-	-
TSAT50T	+	-	-	-	-
TSAT51	-	-	-	-	0.25
TSAT60	-	-	-	-	-
TSAT92	+	-	-	-	0.30
TSAT101	+	-	-	-	0.20
TSAT102B	-	+++	16.67	1.17	0.35
TSAT113	-	-	-	-	-
TSAT115	+	-	-	-	-

745 IAA production: - = no production, +/- = low production, + = production, ++ = moderate production, +++ = high  
 746 production.

747 Siderophore production: - = no production (halo = 0 cm), +/- = low production (halo  $\leq$  0.2 cm), + = production (0.3  
748 cm  $\leq$  halo  $\leq$  0.8 cm), ++ = moderate production (0.8 cm < halo < 1.4 cm), +++ = high production (halo > 1.5 cm).  
749 Phosphate solubilization: SE = solubilization efficiency, PSI = phosphate solubilization index, - = absence of  
750 solubilization.  
751 Phytate solubilization: - = absence of solubilization.  
752



753 **Table 3** Plant growth promoting traits of actinobacteria isolated from spores of *Rhizopagus*  
 754 *intraradices* IMA6.

Isolate	IAA production	Siderophore production	Phosphate solubilization		Phytate solubilization
		Activity (after 7 days)	SE(%)	PSI	Halo zone (cm)
W1	++	++	14.29	1.14	0.60
W2	-	+++	10.53	1.11	0.50
W19	-	++	11.11	1.11	0.20
W22	++	++	20	1.2	0.85
W27	++	++	-	-	1.10
W31	++	-	16.67	1.17	0.15
W39	++	++	28.57	1.29	0.75
W40	-	-	-	-	0.50
W41	-	++	-	-	0.05
W43N	++	++	63.64	1.64	0.80
W47	++	++	17.65	1.18	1.50
W49	++	+/-	-	-	0.45
W54	++	++	13.64	1.14	0.25
W56	++	++	18.18	1.18	0.25
W58	+	++	11.11	1.11	0.75
W60B	+	+++	11.11	1.11	0.85
W60R	+	+/-	57.14	1.57	0.35
W64	++	++	45.45	1.45	1.20
W65	-	-	23.53	1.24	0.95
W66	-	-	23.08	1.23	0.15
W68	++	++	13.33	1.13	0.70
W69	-	+	-	-	1.15
W71	-	++	25	1.25	0.90
W77	++	++	36.36	1.36	0.90
W85	-	-	-	-	1.80
W87	-	++	-	-	0.35
W90	+	++	20	1.2	0.80
W92	++	-	15.38	1.15	1.00
W94	++	+	54.55	1.55	1.15
W105	++	++	-	-	1.05
W115	++	++	38.46	1.38	0.50
W129	++	-	33.33	1.33	0.75
W132	++	+	35.71	1.36	0.80
W133	++	++	-	-	0.95

755 IAA production: - = no production, +/- = low production, + = production, ++ = moderate production, +++ = high  
 756 production.

757 Siderophore production: - = no production (halo = 0 cm), +/- = low production (halo ≤ 0.2 cm), + = production (0.3  
 758 cm ≤ halo ≤ 0.8 cm), ++ = moderate production (0.8 cm < halo < 1.4 cm), +++ = high production (halo > 1.5 cm).

759 cm ≤ halo ≤ 0.8 cm), ++ = moderate production (0.8 cm < halo < 1.4 cm), +++ = high production (halo > 1.5 cm).

760 Phosphate solubilization: SE = solubilization efficiency, PSI = phosphate solubilization index, - = absence of  
761 solubilization.

762 Phytate solubilization: - = absence of solubilization.

763

764 **Table 4** Plant growth promoting traits of putative chitinolytic bacteria isolated from spores of  
 765 *Rhizophagus intraradices* IMA 6.

Isolate	IAA production	Siderophore production	Phosphate solubilization	Phytate solubilization	
		Activity (after 7 days)	SE(%)	PSI	Halo zone (cm)
CH1	-	+	-	-	-
CH2	-	+	-	-	-
CH3	-	-	63.16	1.63	0.50
CH4	-	+	-	-	0.20
CH5	++	+	31.25	1.31	0.25
CH6B	-	+	30.77	1.31	0.30
CH6G	+	+	25	1.25	0.30
CH7	++	-	21.43	1.21	0.62
CH8	+++	+	-	-	0.15
CH9	-	-	15.38	1.15	0.15
CH10	+	+	69.23	1.69	0.25
CH11	+/-	+	63.64	1.64	0.35
CH12	-	+	36.36	1.36	0.25
CH13	-	+	46.15	1.46	0.20
CH14	+	+	50	1.5	0.35
CH15	+	+	44.44	1.44	0.30
CH16L	++	-	50	1.5	0.35
CH16B	-	-	20.83	1.21	-
CH17	+++	+	50	1.5	0.30
CH18	-	+	-	-	0.70
CH19	+	-	86.67	1.87	0.45
CH20	+	+	30	1.3	0.35
CH21	-	-	-	-	-

766 IAA production: - = no production, +/- = low production, + = production, ++ = moderate production, +++ = high  
 767 production.

768 Siderophore production: - = no production (halo = 0 cm), +/- = low production (halo  $\leq$  0.2 cm), + = production (0.3  
 769 cm  $\leq$  halo  $\leq$  0.8 cm), ++ = moderate production (0.8 cm < halo < 1.4 cm), +++ = high production (halo > 1.5 cm).

770 Phosphate solubilization: SE = solubilization efficiency, PSI = phosphate solubilization index, - = absence of  
 771 solubilization.

772 Phytate solubilization: - = absence of solubilization.

773

774

775

776

777

778 **Table 5** Plant growth promoting traits of putative N<sub>2</sub>-fixing bacteria isolated from spores of  
 779 *Rhizophagus intraradices* IMA6.

Isolate	IAA production	Siderophore production	Phosphate solubilization		Phytate solubilization
		Activity (after 7 days)	SE(%)	PSI	Halo zone (cm)
N1	-	+	-	-	-
N3	-	+	-	-	-
N4	-	+	-	-	-
N5	-	+	-	-	-
N6	-	+	-	-	-
N7	+/-	+	-	-	-
N8	-	+	-	-	-
N9	-	+	-	-	-
N10	-	+	-	-	-
N11	-	+	-	-	-
N12	-	+	10	1.1	-
N13	-	++	-	-	0.10
N14	+/-	+	-	-	0.10
N16	-	-	-	-	-
N17	-	++	-	-	-
N18	+	+	-	-	0.45
N19	-	+	41.67	1.42	0.50
N20	-	+	61.54	1.62	0.60
N21	-	+	81.25	1.81	0.55
N22	-	+	30.77	1.31	0.55
N23	-	+	71.43	1.71	0.65
N24	+/-	+	33.33	1.33	0.65
N25	-	+	58.33	1.58	0.70
N26	-	+	72.73	1.73	0.50
N27	-	+	64.29	1.64	0.60
N28	-	+	91.67	1.92	0.10
N29	-	+	84.62	1.85	0.60

780 IAA production: - = no production, +/- = low production, + = production, ++ = moderate production, +++ = high  
 781 production.

782 Siderophore production: - = no production (halo = 0 cm), +/- = low production (halo ≤ 0.2 cm), + = production (0.3  
 783 cm ≤ halo ≤ 0.8 cm), ++ = moderate production (0.8 cm < halo < 1.4 cm), +++ = high production (halo > 1.5 cm).

784 Phosphate solubilization: SE = solubilization efficiency, PSI = phosphate solubilization index, - = absence of  
 785 solubilization.

786 Phytate solubilization: - = absence of solubilization.

787

788

789 **Table 6** Phylogenetic identification of the 19 best performing plant growth promoting bacterial  
790 strains.  
791

Strain	Identification	Identity (%)	Most closely related GeneBank sequence
CH5	<i>Sinorhizobium meliloti</i>	100%	NR113670.1
CH8	<i>Sinorhizobium meliloti</i>	99%	NR113670.1
CH10	<i>Bacillus</i> sp. <i>pumilus</i> group	99%	KM087337.1
CH17	<i>Sinorhizobium meliloti</i>	99%	GU129568.1
CH19	<i>Lysinibacillus fusiformis</i>	99%	NR112569.1
N13	<i>Nocardioides albus</i>	99%	AF005004.1
N17	<i>Arthrobacter phenanthrenivorans</i>	100%	NR042469.2
N23	<i>Sinorhizobium meliloti</i>	100%	AB535689
N28	<i>Sinorhizobium meliloti</i>	99%	AB535689
N29	<i>Sinorhizobium meliloti</i>	99%	AB535689
TSA3	<i>Sinorhizobium meliloti</i>	99%	GU129568.1
TSA26	<i>Sinorhizobium meliloti</i>	99%	GU129568.1
TSA41	<i>Sinorhizobium meliloti</i>	99%	GU129568.1
TSA50	<i>Fictibacillus barbaricus</i>	99%	KJ831620.1
W43N	<i>Streptomyces</i> sp.	99%	NR041063.1
W64	<i>Streptomyces</i> sp.	98%	NR114792.1
W77	<i>Streptomyces</i> sp.	99%	NR112526.1
W94	<i>Streptomyces</i> sp.	99%	NR041231.1
W115	<i>Streptomyces</i> sp.	99%	JN969025.1

792

793

794

795

796

797

798

799

800

801

802 **FIGURE LEGENDS**

803 **Fig. 1.** (a) Number of bacterial strains isolated from spores of *Rhizophagus intraradices* IMA6  
804 within each functional group (heterotrophs, actinobacteria, chitinolytic and putative N<sub>2</sub>-fixing  
805 bacteria). (b) Percentage of selected isolates displaying plant growth promoting (PGP) traits within  
806 each functional group. (c) Percentage of selected isolates showing increasing numbers of PGP  
807 activities within each functional group.

808

809 **Fig. 2.** Venn diagram showing, within each functional group, the number of spore associated  
810 bacteria displaying plant growth promoting traits - IAA production, siderophore production, mineral  
811 phosphate solubilization and phytate solubilization.

812

813 **Fig. 3.** (a,b,c) Affiliation of the sequences of *Actinomycetales*, *Bacillales* and *Rhizobiales* isolated  
814 from spores of *Rhizophagus intraradices* IMA6 with the existing 16S rRNA gene sequences, using  
815 Neighbor-Joining trees. Bootstrap values below 70 are not shown. The sequences from the database  
816 are indicated by their accession numbers. The DNA sequences retrieved in this work are shown in  
817 boldface.

818

Figure






Fig. 1


### Heterotrophs



### Actinobacteria



### Chitinolytic Bacteria



### Putative N<sub>2</sub>-fixing bacteria




Fig. 2






Fig. 3