

1 **Identification and characterization of lactic acid bacteria and yeasts of PDO Tuscan bread**
2 **sourdough by culture dependent and independent methods**

3

4 Michela Palla, Caterina Cristani, Manuela Giovannetti, Monica Agnolucci*

5 Department of Agriculture, Food and Environment, University of Pisa, Via del Borghetto 80, 56124

6 Pisa, Italy

7

8 RUNNING HEAD: Lactic acid bacteria and yeasts of PDO Tuscan bread sourdough

9

10 *Corresponding author: M. Agnolucci, Department of Agriculture, Food and Environment,
11 University of Pisa, Via del Borghetto 80, 56124 Pisa, Italy

12 Phone: +39.0502216647, Fax: +39.0502220606, e-mail address: monica.agnolucci@unipi.it

13

14 **Highlights**

- 15 • The microbiota of PDO Tuscan bread sourdough was isolated and molecularly identified.
- 16 • A distinctive tripartite microbial association characterized the PDO Tuscan bread sourdough.
- 17 • A large number of *Lactobacillus sanfranciscensis* (98) and *Candida milleri* (65) strains were
18 isolated.
- 19 • Among yeasts, *Candida milleri* dominated over *Saccharomyces cerevisiae*.

20

21 **Abstract**

22 Sourdough fermentation has been increasingly used worldwide, in accordance with the demand of
23 consumers for tasty, natural and healthy food. The high diversity of lactic acid bacteria (LAB) and
24 yeast species, detected in sourdoughs all over the world, may affect nutritional, organoleptic and
25 technological traits of leavened baked goods. A wide regional variety of traditional sourdough
26 breads, over 200 types, has been recorded in Italy, including special types selected as worthy of

27 either Protected Geographical Indication (PGI) or Protected Designation of Origin (PDO), whose
28 sourdough microbiota has been functionally and molecularly characterized. As, due to the very
29 recent designation, the microbiota of Tuscan bread sourdough has not been investigated so far, the
30 aim of the present work was to isolate and characterize the species composition of LAB and yeasts
31 of PDO Tuscan bread sourdough by culture-independent and dependent methods. A total of 130
32 yeasts from WLN medium and 193 LAB from both mMRS and SDB media were isolated and
33 maintained to constitute the germplasm bank of PDO Tuscan bread. Ninety six LAB from mMRS
34 medium and 68 yeasts from WLN medium were randomly selected and molecularly identified by
35 ARDRA (Amplified Ribosomal DNA Restriction Analysis) and PCR-RFLP analysis of the ITS
36 region, respectively, and sequencing. The yeast identity was confirmed by 26S D1/D2 sequencing.
37 All bacterial isolates showed 99% identity with *Lactobacillus sanfranciscensis*, 65 yeast isolates
38 were identified as *Candida milleri*, and 3 as *Saccharomyces cerevisiae*. Molecular characterization
39 of PDO Tuscan bread sourdough by PCR-DGGE confirmed such data. The distinctive tripartite
40 species association, detected as the microbiota characterizing the sourdough used to produce PDO
41 Tuscan bread, encompassed a large number of *L. sanfranciscensis* and *C. milleri* strains, along with
42 a few of *S. cerevisiae*. The relative composition and specific physiological characteristics of such
43 microbiota could potentially affect the nutritional features of PDO Tuscan bread, as suggested by
44 the qualitative functional characterization of the isolates. Investigations on the differential
45 functional traits of such LAB and yeast isolates could lead to the selection of the most effective
46 single strains and of the best performing strain combinations to be used as starters for the
47 production of baked goods.

48

49 **Keywords:** ARDRA, ITS-RFLP, PCR-DGGE, *Lactobacillus sanfranciscensis*, *Candida milleri*,
50 *Saccharomyces cerevisiae*.

51

52 **1. Introduction**

53 Sourdough fermentation represents one of the oldest biotechnologies used by humans to ferment
54 cereals for bread production. In recent times, it has been increasingly used worldwide, in
55 accordance with the demand of consumers for tasty, natural and healthy foods (Gobbetti and
56 Gänzle, 2013). Indeed, sourdough fermentation improves bread sensory, structural and nutritional
57 properties, and prolongs its shelf-life (Arendt et al., 2007; Katina et al., 2005; Minervini et al.,
58 2014). In addition, microbial metabolism during sourdough fermentation positively affects several
59 bread functional features, producing bioactive compounds, such as peptides, beta-glucans and other
60 exopolisaccharides (Gobbetti et al., 2014).

61 A large number of lactic acid bacteria (LAB) and yeast species, establishing positive
62 interactions and often stable associations, has been isolated from sourdoughs all over the world, *i.e.*
63 about 80 bacterial and 20 yeast species (De Vuyst and Neisens, 2005; Gänzle and Ripari, 2016).
64 Notwithstanding, only a few species characterize single batches of sourdough, where no more than
65 6 different species have been usually identified. The typical most commonly detected bacterial
66 species are *Lactobacillus brevis*, *Lactobacillus pontis*, *Lactobacillus reuteri*, *Lactobacillus*
67 *plantarum* and *Lactobacillus sanfranciscensis*, the latter representing the dominant member of the
68 microbiota, as it has been isolated in more than 75% of sourdoughs globally (Ganzle and Ripari,
69 2016). The most common sourdough yeast is *Saccharomyces cerevisiae*, although, as reviewed by
70 De Vuyst et al. (2016) other species may occur in spontaneously developed stable sourdoughs (in
71 decreasing order of abundance): *Candida humilis/Candida milleri*, *Wickerhamomyces anomalus*,
72 *Torulaspora delbrueckii*, *Kazachstania exigua*, *Pichia kudriavzevii* and *Candida glabrata*. Some of
73 these species, such as *K. exigua*, *C. humilis* and *C. milleri*, are maltose-negative yeasts reported to
74 form stable mutualistic associations with *L. sanfranciscensis*, which is able to hydrolyze maltose
75 (De Vuyst et al., 2014).

76 Several studies investigating the microbiota of sourdough from different countries, *e.g.*
77 Belgium (Scheirlinck et al., 2007), France (Ferchichi et al., 2008; Vera et al., 2012), Turkey (Dertli
78 et al., 2016), China (Liu et al., 2016; Zhang et al., 2011), showed that the diversity of microbial

79 communities depends on process technologies, types of flour and other ingredients traditionally
80 associated with local culture and origin. Such diversity is at the basis of differential metabolic
81 products, affecting nutritional, organoleptic and technological traits of leavened baked goods. In
82 particular, traditional or type I sourdough is characterized by a spontaneous fermentative process,
83 based on continuous backslopping, carried out by LAB and yeasts originating from the flour, other
84 dough ingredients and the environment. Type I sourdough is utilized to produce various leavened
85 baked products, such as San Francisco bread, French bread, rye bread, Altamura bread (Corsetti,
86 2013).

87 A wide regional variety of traditional sourdough breads, over 200 types, has been recorded
88 in Italy (INSOR 2000), including special types selected as worthy of either Protected Geographical
89 Indication (PGI) (*Coppia Ferrarese*, *Pane Casareccio di Genzano*, *Cornetto di Matera*) or
90 Protected Designation of Origin (PDO) (*Pagnotta del Dittaino* and *Pane di Altamura*). The
91 sourdoughs of such traditional regional breads have been functionally and molecularly
92 characterized, *i.e.* *Cornetto di Matera* (Zotta et al., 2008), *Pane di Altamura* (Ricciardi et al., 2005),
93 along with breads from Abruzzo (Valmorri et al., 2006; 2010), Marche (Osimani et al., 2009),
94 Molise (Reale et al., 2005), Sicily (Pulvirenti et al., 2001) and Sardinia (Catzeddu et al., 2006).

95 Recently, *Pane Toscano* has obtained the PDO status from the European Community
96 (Commission implementing regulation (EU) 2016/303 of 1 March 2016, Official Journal of the
97 European Union L 58 of 04 March 2016). PDO Tuscan bread has to be manufactured by a typical
98 method generally adopted in Tuscan bakeries. It requires the exclusive use of sourdough starters,
99 water, the absence of added salt, and type '0' soft-wheat flour from wheat varieties grown in
100 Tuscany, according to the production guideline described in the EU Regulation for PDO Tuscan
101 bread (Official Journal of the European Union C 235 of 14 August 2013). Due to the very recent
102 designation, the microbiota of Tuscan bread sourdough has not been investigated so far. The aim of
103 the present work was to isolate and characterize the species composition of lactic acid bacteria and
104 yeasts of PDO Tuscan bread sourdough. To this aim, we utilized *i)* a culture-independent approach,

105 Polymerase Chain Reaction (PCR) Denaturing Gradient Gel Electrophoresis (DGGE), a
106 molecular technique able to avoid underestimates deriving from the constraints of culture conditions
107 and from the presence of microorganisms in Viable But Non-Culturable (VBNC) state; *ii*) a culture-
108 dependent approach to isolate and molecularly identify LAB and yeast species; *iii*) a preliminary
109 qualitative screening to characterize LAB and yeasts with functionally important traits.

110

111 **2. Materials and methods**

112 *2.1. Sourdough sampling*

113 The sourdough analyzed in this study was collected from the Consortium of Promotion and
114 Protection of Tuscan Sourdough Bread - *Consorzio Pane Toscano a Lievitazione Naturale* (CPT).

115

116 *2.2. Microbiological analysis and isolation of LAB and yeasts*

117 Three samples of about 10 g of PDO Tuscan bread sourdough were homogenized in a sterile
118 stomacher bag containing 90 mL of saline-peptone water (9 g/L NaCl, 1 g/L bacteriological
119 peptone, Oxoid, Milan, Italy) for 2 min at 260 rpm, using a Stomacher (Stomacher 400, Laboratory
120 Blender). Further, a tenfold serial dilution (10^{-1} to 10^{-6}) was carried out in the same solution and
121 aliquots (100 μ L) were added in triplicate into a Petri dish containing the agar media listed below.
122 LAB were counted on Sourdough Bacteria (SDB) (Kline and Sugihara, 1971) medium and on
123 mMRS agar (de Man, Rogosa, Sharpe, 1960) modified by adding 20 g/L maltose and 50 mL/L fresh
124 yeast extract and adjusted to pH 5.6. To inhibit yeast growth, media were supplemented with 100
125 mg/L cycloheximide. Inoculated plates were incubated for 7 days at 28 °C under anaerobic
126 conditions (AnaeroGen, Oxoid). Yeasts were counted on Wallerstein Laboratory Nutrient (WLN)
127 agar (Oxoid, Basingstoke, UK) and on Yeast Extract Peptone Dextrose (YEPD) agar. Both media
128 were added with 100 mg/L chloramphenicol and incubated at 28 °C for 48 h.

129 LAB were randomly selected picking up at least 20 colonies from each plate of both media
130 and purified by streaking four times onto the same medium used for isolation. About 15 yeast

131 colonies were randomly selected from each plate of WLN medium on the basis of phenotypic
132 colony characteristics and then purified as described above.

133 Each strain was named with the acronym of the Collection of the Department of Agriculture,
134 Food and Environment of the University of Pisa (IMA, International Microbial Archives), followed
135 by a progressive number plus “Y” or “LAB” for yeasts or bacteria, respectively. Purified strains
136 were stored at $-80\text{ }^{\circ}\text{C}$ in the appropriate broth medium, supplemented with 20% (w/v) glycerol.

137

138 *2.3. Molecular identification of LAB and yeast isolates*

139 DNA of isolates and reference strains, listed in Table 1, was extracted from microbial liquid
140 cultures grown at $28\text{ }^{\circ}\text{C}$ using “MasterPureTM Yeast DNA Purification Kit” (Epicentre®) according
141 to the manufacturer’s protocols. LAB strains were identified by Amplified Ribosomal DNA
142 Restriction Analysis (ARDRA) and yeast strains by ITS region amplification and its Restriction
143 Fragment Length Polymorphism (RFLP) analysis. Amplification reactions were carried out in a
144 final volume of $50\text{ }\mu\text{L}$, containing $5\text{ }\mu\text{L}$ of 10X Ex Taq Buffer (Takara Biotechnology), 0.2 mM of
145 each dNTP (Takara Biotechnology), $0.5\text{ }\mu\text{M}$ of each primer (Eurofins), 1.25 U of Takara Ex Taq
146 polymerase (Takara Biotechnology) and $10\text{-}20\text{ ng}$ of DNA. The 16S rRNA gene was amplified
147 using 27f (5’-GAG AGT TTG ATC CTG GCT CAG-3’) and 1495r (5’-CTA CGG CTA CCT TGT
148 TAC GA-3’) primers (Lane 1991; Weisburg et al., 1991) and ITS region was amplified using ITS1
149 (5’-TCC GTA GGT GAA CCT GCG G-3’) and ITS4 (5’-TCC TCC GCT TAT TGA TAT GC-3’)
150 primers (White et al., 1990). PCR amplifications were carried out with an iCycler-iQ Multicolor
151 Real-Time PCR Detection System (Bio-Rad) using the following conditions: $94\text{ }^{\circ}\text{C}$ initial
152 denaturation for 1 min; 35 amplification cycles of 30 s at $94\text{ }^{\circ}\text{C}$, 30 s at annealing temperature, 45 s
153 at $72\text{ }^{\circ}\text{C}$; final extension at $72\text{ }^{\circ}\text{C}$ for 5 min. The annealing temperatures, for bacteria and yeasts,
154 were 60 and $55\text{ }^{\circ}\text{C}$, respectively. The presence of amplicons was confirmed by electrophoresis in
155 1.5% (w/v) agarose gel stained with ethidium bromide ($0.5\text{ }\mu\text{g/mL}$).

156 LAB 16S rRNA gene amplicons were digested at 37 °C overnight with the restriction
157 endonucleases *AluI*, *HinfI* and *HaeIII* (BioLabs), while yeast ITS amplicons were digested with
158 *HaeIII* and *HinfI* (BioLabs) enzymes. The restriction fragments were separated (at 50 V for 2 h) on
159 1.8% (w/v) agarose gels stained with ethidium bromide (0.5 µg/mL) in Tris-borate-EDTA buffer
160 (Sigma-Aldrich). A 100 bp DNA ladder (BioLabs) was used as a molecular weight marker. All gels
161 were visualized and captured as TIFF format files by the Liscap program for Image Master VDS
162 system (Pharmacia Biotech). Isolates and reference strains profiles were digitally processed and
163 analysed with BioNumerics software version 7.5 (Applied Maths, St-Martens-Latem, Belgium).
164 ARDRA and ITS-RFLP profiles from all digestions were respectively combined, for LAB and
165 yeasts, and the similarity calculated on the basis of the Dice's coefficient. For cluster analysis,
166 unpaired group method with arithmetic average (UPGMA) trees with highest resampling support, in
167 a permutation sample of size 200, were constructed.

168 The identification of isolates was confirmed by sequencing LAB 16S and yeast ITS1-5.8S-
169 ITS2 and 26S D1/D2 rRNA gene amplicons. The 26S D1/D2 was amplified using the primers NL1
170 and NL4, as reported by Kurtzman and Robnett (1998).

171 In particular, PCR products were purified with the UltraClean PCR CleanUp kit (CABRU)
172 according to the manufacturer's protocol, quantified and 5' sequenced by BMR Genomics (Padova,
173 Italy). Sequences were analyzed using BLAST on the NCBI web
174 (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>). The related sequences were collected and aligned using
175 MUSCLE (Edgar, 2004a, b), and phylogenetic trees were constructed using the Neighbor-Joining
176 method based on the kimura 2-parameter model (Kimura, 1980) in Mega 6 software ([http://](http://www.megasoftware.net/)
177 www.megasoftware.net/) with 1000 bootstrap replicates. The sequences were submitted to the
178 European Nucleotide Archive under the accession numbers from LT605080 to LT605156 and from
179 LT718652 to LT718656.

180

181 *2.4. Molecular characterization of PDO Tuscan bread sourdough by PCR-DGGE*

182 DNA was extracted from the three PDO Tuscan bread sourdough samples by using “Power Soil
183 DNA Isolation kit” (MO-BIO Laboratories) according to the manufacturer’s protocol and stored at -
184 20 °C until further analyses.

185 For the analysis of LAB communities, the V3-V4 region of 16S rRNA gene was amplified
186 with the Lac1 (5’-AGC AGT AGG GAA TCT TCC A-3’) and Lac2 (5’-ATT YCA CCG CTA
187 CAC ATG-3’) primers (Walter et al., 2001). The primer Lac2 had at its 5’ end an additional 40-
188 nucleotide GC-rich tail (5’-CGC CCG GGG CGC GCC CCG GGC GGC CCG GGG GCA CCG
189 GGG G-3’). For the analysis of yeasts, an approximately 250 bp long fragment of D1/D2 region of
190 the 26S rRNA gene was amplified using NL1 (5’-GCC ATA TCA ATA AGC GGA GGA AAA G-
191 3’) and LS2 (5’-ATT CCC AAA CAA CTC GAC TC-3’) primers (Cocolin et al., 2000). An
192 additionally GC clamp (5’-GCG GGC CGC GCG ACC GCC GGG ACG CGC GAG CCG GCG
193 GCG G-3’) was added to the forward primer NL1. PCR amplifications were performed as
194 previously described, using 10-20 ng of DNA. Amplification conditions were: 94 °C initial
195 denaturation for 1 min; 35 amplification cycles of 30 s at 94 °C, 30 s at annealing temperature, 30 s
196 at 72 °C; final extension at 72 °C for 5 min. The annealing temperatures for bacteria and yeasts,
197 were 58 and 55 °C, respectively. The presence of amplicons was confirmed by electrophoresis in
198 1.5% (w/v) agarose gel stained with ethidium bromide (0.5 µg mL⁻¹).

199 For the DGGE analysis, amplicons were separated in 8% (w/v) polyacrylamide gels with a
200 35–50% and 30–56% urea-formamide gradient, for bacteria and yeasts, respectively, using the
201 DCode™ Universal Mutation Detection System (BioRad). A composite mix of bacterial 16S rRNA
202 gene fragments and fungal 26S rRNA gene fragments from reference strains (Table 1), were added
203 as bacterial and fungal reference DGGE markers (M). Gels were run and visualized as described in
204 Agnolucci et al. (2013). DGGE fragments were cut out from the gels for sequencing. DNA was
205 extracted by eluting for 3 days in 50 µL 10 mM TE at 4 °C. One µL of the supernatant diluted 1:100
206 was used to re-amplify the LAB V3-V4 and the yeast D1/D2 regions of the DNA according to the
207 PCR protocol described above, using Lac2 and NL1 primers without GC clamp. PCR products were

208 than purified by UltraClean PCR CleanUp Kit (MO-BIO Laboratories) according to the
209 manufacturer's protocol, quantified and 5' sequenced at the BMR Genomics (Padova, Italy).
210 Sequences were analyzed using BLAST on the NCBI web and submitted to the European
211 Nucleotide Archive under the accession numbers from LT605157 to LT605165.

212

213 *2.5. Qualitative functional characterization of LAB and yeast isolates*

214 To test phytase activity, yeasts were grown on Phytate Screening Medium (PSM) (Jorquera et al.,
215 2008). The same medium added with 50 mL/L of fresh yeast extract at pH 5.6 was used for LAB.
216 To eliminate false positive results, plates were counterstained as described by Bae et al. (1999).
217 Protease activity of yeasts and LAB was assessed on YEPD without Peptone and on mMRS,
218 containing 2% skim milk, respectively. Isolates were then incubated at 28 °C for 8 days. After
219 incubation the formation of halo zones around microbial colonies indicated phytase and protease
220 capacity. Yeast amylase activity was tested on YEPD without Dextrose containing 10 g/L soluble
221 starch (Fluka), pH 6.46. LAB amylase activity was screened on mMRS agar without maltose,
222 containing 10 g/L soluble starch (Fluka), pH 5.6. Yeasts and LAB were then incubated for 7 days at
223 30 °C and 37 °C, respectively, and starch hydrolysis was revealed by the disappearance of the blue
224 colour of the medium around the microbial colonies after addition of Lugol iodine solution (Sigma-
225 Aldrich).

226

227 **3. Results**

228 *3.1. Microbiological analysis of PDO Tuscan bread sourdough*

229 Lactic acid bacteria counts on mMRS and SDB media were $2.33 \pm 0.54 \cdot 10^9$ CFU/g and
230 $1.03 \pm 0.22 \cdot 10^9$ CFU/g, respectively. Yeast numbers were approximately 100 fold lower than LAB,
231 showing colony counts of $2.14 \pm 0.17 \cdot 10^7$ on WLN and $2.21 \pm 0.37 \cdot 10^7$ CFU·g⁻¹ on YEPD. A total of
232 130 yeasts were isolated from WLN medium as such medium allowed the differentiation among
233 colony morphologies, while 193 LAB, showing only one colony morphology, were isolated from

234 both mMRS and SDB media. All the isolates were maintained to constitute the germplasm bank of
235 PDO Tuscan bread. Ninety six LAB from mMRS medium and 68 yeasts from WLN medium were
236 randomly selected and molecularly characterized at species level.

237

238 3.2. Molecular identification of LAB and yeast isolates

239 The 96 LAB and 68 yeasts selected were molecularly identified by ARDRA and PCR-RFLP
240 analysis of the ITS region, respectively, and sequencing. The expected 16S rRNA gene amplicon of
241 1468 pb was obtained for all the 96 LAB isolates. The amplified product was then subjected to
242 restriction analysis using the enzyme *AluI*, *HaeIII* and *HinfI*. *AluI* produced two different ARDRA
243 profiles consisting of three bands (210, 240 and 620 bp) (52% of isolates) and four bands (210, 240,
244 620 and 870 bp) (48% of isolates). *HinfI* produced three different ARDRA profiles consisting of six
245 bands (70, 90, 120, 320, 400 and 1000 bp) in 51% of isolates, five bands (70, 90, 120, 320 and 1000
246 bp) in 48% of isolates and seven bands (70, 90, 120, 320, 400, 1000 and 1430 bp) in one isolate.
247 *HaeIII* produced only one ARDRA profile of three bands (125, 240, 1330 bp) in all isolates
248 (Supplementary Fig. S1). The dendrogram created by combining the three different ARDRA
249 profiles grouped the isolates into two main clusters with a similarity level of 89.9%. The first one
250 consisted of 49 isolates, which clustered at similarity level of 100% with the reference strain *L.*
251 *sanfranciscensis* DSMZ 20451, and of the isolate IMA 23LAB which separately branched at
252 similarity of 95.2%. The remaining 46 isolates, showing 100% similarity, grouped in the second
253 cluster (Fig. 1). Five isolates from each of the two clusters, along with the isolate IMA 23LAB,
254 were subjected to 16S rRNA gene sequence analysis. All isolates showed 99% identity with *L.*
255 *sanfranciscensis* strain ATCC 27651 (DSMZ 20451) despite the presence of a polymorphism
256 within 16S rRNA gene detected by ARDRA analysis (Table 2).

257 The ITS region of the 68 yeast isolates was successfully amplified. An amplicon of about
258 650 bp, was obtained for 65 isolates and one of about 840 bp for the remaining three isolates (IMA
259 19Y, IMA 36Y and IMA 105Y). Digestion of 650 pb amplicons with *HinfI* produced two fragments

260 of 350 and 300 bp in all the 65 isolates and in the reference strains *C. milleri* DBVPG 6753^T, *C.*
261 *humilis* DBVPG 7219^T and *C. humilis* DBVPG 6754 (Supplementary Fig. S2). Using *Hae*III all the
262 65 isolates produced the same restriction pattern (425 and 230 bp) as the reference strains *C. milleri*
263 DBVPG 6753^T, while *C. humilis* DBVPG 7219^T type strain showed three fragments of 310, 210
264 and 110 bp. Conversely, the reference strain *C. humilis* DBVPG 6754 showed two fragments of 425
265 and 230 bp as obtained for the *C. milleri* DBVPG 6753^T. The other three isolates, producing 840 bp
266 amplicons, showed a profile of 310, 230, 165 and 130 bp with *Hae*III and of 350 and 120 bp with
267 *Hinf*I. The dendrogram, created by combining the size of the ITS amplicons and the different
268 restriction profiles, grouped all the 65 isolates with the reference strains *C. milleri* DBVPG 6753^T
269 and *C. humilis* DBVPG 6754 in a main cluster with a 100% of similarity (Fig. 2). *C. humilis*
270 DBVPG 7219^T branched separately at 73% similarity. In a separate cluster (33.5% similarity) the
271 isolates IMA 19Y, IMA 36Y and IMA 105Y grouped with *S. cerevisiae* ATCC 32167 reference
272 strain at a similarity level of 100%. The 840 bp amplicons were then subjected to restriction
273 analysis by the enzyme *Hpa*II and a profile of 700 and 130 bp, corresponding to that of *S. cerevisiae*
274 (Fernández-Espinar et al., 2000), was obtained. The BLAST analysis of ITS1-5.8S-ITS2 and 26S
275 D1/D2 region sequences confirmed the identity of the three isolates IMA 19Y, IMA 36Y and IMA
276 105Y as *S. cerevisiae* (Table 2).

277 To determine the species affiliation of our 65 *Candida* isolates, all 650 bp ITS amplicons
278 were sequenced and examined using BLAST and phylogenetic trees analyses. The dendrogram
279 obtained by the analysis of the ITS sequences of our isolates, and those of *C. humilis* and *C. milleri*
280 reference strains available in GeneBank, is reported in Fig. 3. Results showed that all our ITS
281 sequences grouped with those of *C. milleri* reference strains with 100% identity. The BLAST
282 analysis of 26S D1/D2 region sequences of two representative isolates (IMA 11Y and IMA 33Y)
283 confirmed the identity of our isolates as *C. milleri* (Table 2).

284

285 *3.3. Molecular characterization of PDO Tuscan bread sourdough by PCR-DGGE*

286 Microbial community diversity of PDO Tuscan bread sourdough was further investigated by PCR-
287 DGGE. The amplification of the V3-V4 region of 16S rRNA gene of LAB and of the partial D1/D2
288 domain of 26S rRNA gene of yeasts from the three sourdough samples produced a DNA fragment
289 of approximately 326 and 250 bp, respectively. DGGE profiles of bacterial PCR products
290 (Supplementary Fig. S3a) showed a main fragment corresponding to that of the reference strain *L.*
291 *sanfranciscensis* DSMZ 20451. DGGE profiles of the yeast community (Supplementary Fig. S3b)
292 showed two main fragments migrating one at the same position of that of *S. cerevisiae* ATCC
293 32167 and one at the same position of both *C. humilis* DBVPG 7219^T and *C. milleri* DBVPG
294 6753^T. The bands from each of the two DGGE gels were excised, sequenced and affiliated to
295 bacterial and yeast species by using BLAST analysis. Results revealed the presence of *L.*
296 *sanfranciscensis* associated with *C. milleri/C. humilis* and *S. cerevisiae* (Supplementary Fig. S3 and
297 Table 2).

298 299 *3.4. Qualitative functional characterization of LAB and yeast isolates*

300 Phytase, amylase and protease proprieties of our isolates were assessed by plate assays. The three
301 strains identified as *S. cerevisiae* were able to solubilize phytate, hydrolyze starch and digest casein.
302 Among *C. milleri* strains, 50% showed protease activity, while only 6% and 2% were able to
303 solubilize phytate and to hydrolyze starch, respectively. Specifically, *C. milleri* IMA 33Y showed
304 all the three activities tested (Table 3). Concerning lactic acid bacteria, the ability to solubilize
305 phytate was detected in 19% of *L. sanfranciscensis* strains, while none was able to digest casein and
306 to hydrolyze starch (Table 3).

307 308 **4. Discussion**

309 Here, a peculiar tripartite species association of *Candida milleri*, *Saccharomyces cerevisiae* and
310 *Lactobacillus sanfranciscensis* was detected as the microbiota characterizing the sourdough of PDO
311 Tuscan bread, by using a multimodal approach.

312 Microbiological analysis showed the occurrence of LAB and yeasts at a level of about 10^9
313 and 10^7 CFU/g respectively, revealing a yeasts/LAB ratio of 1:100, consistently with previous data
314 reported for sourdoughs (Gobbetti, 1998). LAB and yeast counts were not affected by the medium
315 used. Among all the isolates, 96 LAB and 68 yeasts were selected and preliminarily functionally
316 characterized for phytase, amylase and protease activity. In particular, the ability to solubilize
317 phytate was found in 18 out of the 96 *L. sanfranciscensis* isolates, while among yeasts this trait was
318 detected in all the *S. cerevisiae* isolates and in 3 out of the 65 *C. milleri* isolates. Such results are
319 consistent with a study of Nuobariene et al. (2012), where phytase activity was found predominant
320 among *S. cerevisiae* isolates and for the first time detected in one *C. humilis* isolate. Moreover, the
321 three *S. cerevisiae* isolates were found positive to the amylase activity assay, a trait which has been
322 scarcely investigated in *S. cerevisiae* sourdough isolates. Consistently with our results, Osimani et
323 al. (2009), analysing 36 yeast strains isolated from sourdoughs collected in the Marche region,
324 found 21 *S. cerevisiae* isolates able to hydrolyze starch, with different levels of activity. However,
325 further investigations are needed in order to exploit the most important functional properties of
326 yeast sourdough isolates for the production of baked goods.

327 The molecular identification of the selected 96 LAB was carried out by ARDRA and
328 sequence analysis. *HinfI* and *AluI* restriction enzymes revealed a polymorphism within the 16S
329 rRNA gene, allowing us to separate the isolates into three different groups, assigned by sequence
330 analysis to the same species, *L. sanfranciscensis*. Such an intraspecific diversity is consistent with
331 data reported by Foschino et al. (2001) showing the same 16S rRNA gene polymorphism in *L.*
332 *sanfranciscensis* strains isolated from some Italian sourdough samples, when *HinfI* was used.
333 Interestingly, in the sourdough of PDO Tuscan bread the only lactic acid bacterial species identified
334 was *L. sanfranciscensis*, dissimilarly from previous findings on bacterial communities
335 characterizing other Italian sourdoughs, that detected *L. sanfranciscensis* as the predominant LAB
336 species, though associated with several facultatively heterofermentative species, such as
337 *Lactobacillus plantarum* and *Lactobacillus alimentarius* (Minervini et al., 2012; Yazar and

338 Tavman, 2012). Moreover, also in sourdoughs used for other typical Tuscan breads, *i.e.* Bozza
339 Pratese and Pane di Altopascio Tradizionale, *L. sanfranciscensis* was found associated with
340 *Lactobacillus paralimentarius* and *Lactobacillus gallinarum*, respectively (Minervini et al., 2012).

341 The molecular identification of the selected 68 yeasts showed the occurrence of *S. cerevisiae*
342 and *C. milleri* as dominant yeast species in PDO Tuscan bread sourdough, in agreement with
343 previous works reporting that these species are the most frequently found in spontaneously
344 developed stable sourdoughs (De Vuyst et al., 2016). *C. milleri* was the prevalent species in our
345 samples, representing 96% of the isolates identified, whereas *S. cerevisiae* represented the
346 remaining 4%. Such data supplement those obtained from Pagnotta del Dittaino PDO sourdough,
347 where a yeast other than *S. cerevisiae*, *C. humilis*, was the only dominant species recovered (Gullo
348 et al., 2003), which occurred in association with the LAB species *L. sanfranciscensis* and
349 *Enterococcus durans* (Minervini et al., 2012). Similar results were reported also for Pane di
350 Cappelli sourdough, where *C. humilis* and *L. sanfranciscensis* were associated with *L. plantarum*
351 (Minervini et al., 2012). By contrast, other studies on sourdoughs used to produce typical Italian
352 baked goods, detected *S. cerevisiae* as the prevalent species. In particular, Corsetti et al. (2001),
353 analysing 25 different sourdoughs from Apulia region, reported the widespread presence of *S.*
354 *cerevisiae*, which was also the only yeast species isolated from the Altamura bread sourdoughs
355 (Ricciardi et al., 2005). Accordingly, Valmorri et al. (2010), in 20 sourdoughs collected from artisan
356 bakeries throughout Abruzzo region, identified 85% of the isolates as *S. cerevisiae*, associated with
357 *C. milleri* (11%), *Candida krusei* (2.5%), and *Torulaspora delbrueckii* (1%). Moreover, *S.*
358 *cerevisiae* was the only dominant species in three out of four sourdoughs in Marche region
359 (Osimani et al., 2009) and in Northern Italy (Iacumin et al., 2009).

360 It is important to note that the same species association of *C. milleri*, *S. cerevisiae* and *L.*
361 *sanfranciscensis* characterizing the sourdough used to produce PDO Tuscan bread, was previously
362 described only in traditional Italian sweet baked goods sourdoughs, such as Colomba, Legaccio,
363 Panettone and Veneziana (Lattanzi et al., 2013; Venturi et al., 2012; Vernocchi et al., 2004).

364 Although in the sourdough context *C. humilis* is often distinguished from *C. milleri* by ITS-
365 RFLP analysis using the *Hae*III enzyme (Pulvirenti et al., 2001), in our work such analysis was not
366 able to discriminate between *C. milleri* and *C. humilis* since the reference strain *C. humilis* DBVPG
367 6754 showed only one *Hae*III restriction site, corresponding to that expected for *C. milleri*. Similar
368 results were obtained by Vigentini et al. (2014) who found some isolates with only one *Hae*III
369 restriction site, as shown by *C. milleri*, which were positioned closer to *C. humilis* after ITS
370 sequencing. However, ITS region sequence analysis of our 65 *Candida* isolates showed that all our
371 isolates belonged to the species *C. milleri*. The reference strain *C. humilis* DBVPG 6754 grouped in
372 a homogeneous sub-cluster of *C. humilis*, which encompassed other isolates possessing sequences
373 with intermediate traits between the two species. Our data confirm that the ITS sequence analysis is
374 the most reliable method for a correct identification of these two sibling species, as suggested by
375 Vigentini et al. (2014). Interestingly, a phylogenetic analysis of the ITS sequences of *C. milleri* and
376 *C. humilis* collected from GeneBank allowed us to note that all strains affiliated to *C. milleri* were
377 recovered from sourdoughs, whereas those identified as *C. humilis* were isolated also from different
378 fermented foods, such as cacao, bantu beer, tequila etc. (Supplementary Fig. S4). It is tempting to
379 speculate that the species *C. milleri* may represent a key species characteristic of the sourdough
380 environment. Finally the identification of our isolates as *C. milleri* was further confirmed by 26S
381 D1/D2 sequencing. As reported in the literature, *C. humilis* and *C. milleri* differ only by one
382 substitution in the D1/D2 domains of 26SrDNA. Accordingly, our sequences show a C base as *C.*
383 *milleri* CBS 6897^T (U94923.1) instead of a T base found in *C. humilis* CBS 5658^T (U69878.1)
384 (Kurtzman and Robnett, 2003; Lachance et al., 2011). Recently Jacques et al. (2016), using a
385 multigenic analysis based on the comparison of coding genes D1/D2 LSU rRNA, RPB1, RPB2 and
386 EF-1 α , have reconsidered *C. humilis* and *C. milleri* as conspecific and reassigned them to the genus
387 *Kazachastania* proposing the new combination *Kazachastania humilis* (E.E. Nel & Van der Walt)
388 Jacques, Sarilar & Casaregola comb.nov. On the other hand, as the level of intraspecific diversity
389 among the tested *C. milleri* strains was higher than the divergence between *C. humilis* CBS 5658^T

390 and *C. milleri* CBS 6897^T (Vigentini et al., 2014), the same authors suggested that such isolates
391 may be part of a species complex.

392 The microbial community composition of PDO Tuscan bread sourdough was also
393 investigated by PCR-DGGE, a method widely utilized to investigate the microbial diversity of
394 many different fermented foods (Cocolin et al., 2013). The same species composition obtained by
395 microbiological analysis was detected in our sourdough samples, where *L. sanfranciscensis*, *C.*
396 *milleri* and *S. cerevisiae* were the dominant species. However, by PCR-DGGE we could not
397 discriminate between *C. milleri* and *C. humilis* as the only one base, differing between these two
398 species, is located out of the D1/D2 fragment amplified by the primers used.

399 The consistency of PCR-DGGE data with those obtained by culture-dependent methods,
400 was reported also by other authors, who studied the microbiota characterizing the sourdoughs used
401 to produce traditional baked goods from different Italian regions *i.e.* Molise (Gatto and Torriani,
402 2004), Sicily (Randazzo et al., 2005), Abruzzo (Settanni et al., 2006), Northern Italy (Iacumin et al.,
403 2009), Campania (Palomba et al., 2011), confirming that PCR-DGGE is a rapid, economic and
404 efficient tool to investigate yeast and LAB species diversity in the sourdough ecosystem. Moreover,
405 this technique can be conveniently applied to investigate the stability of the microbial communities
406 of specific sourdoughs, particularly those used to produce baked goods protected by the PGI or
407 PDO marks (Palla et al., 2015).

408 In conclusion, here, for the first time, a distinctive tripartite microbial association,
409 represented by yeast and LAB species characterizing the sourdough used to produce PDO Tuscan
410 bread was detected. The association encompassed a large number of *L. sanfranciscensis* and *C.*
411 *milleri* strains, along with a few of *S. cerevisiae*, whose relative composition and specific
412 physiological characteristics could potentially affect the organoleptic, rheological, nutritional and
413 nutraceutical features of Tuscan bread, as suggested by the qualitative functional characterization of
414 the isolates. Further investigations on the differential functional traits of the LAB and yeast strains
415 isolated from PDO Tuscan bread sourdough are the next essential steps, in order to exploit the

416 biotechnological potential of the most effective single strains after assessing their complementary
417 and/or synergistic activities and to select the best performing strain combinations to be used as
418 starters for the production of functional baked goods.

419

420 **Acknowledgments**

421 This work was supported by the Tuscany Region PIF Project INNOVAPANE “Processi innovativi
422 per la produzione del Pane Toscano a lievitazione naturale (DOP)” (D.D. n. 2260 - 12.06.2013).

423 The authors wish to thank Dr. Benedetta Turchetti, curator of the DBVPG Industrial Yeasts
424 Collection (University of Perugia, Italy), for sharing information on DBVPG strains.

425

426 **References**

427 Agnolucci, M., Cristani, C., Battini, F., Palla, M., Cardelli, R., Saviozzi, A., Nuti, M., 2013.
428 Microbially-enhanced composting of olive mill solid waste (wet husk): bacterial and fungal
429 community dynamics at industrial pilot and farm level. *Bioresour. Technol.* 134, 10-16.

430 Arendt, E.K., Ryan, L.A., Dal Bello, F., 2007. Impact of sourdough on the texture of bread. *Food*
431 *Microbiol.* 24, 165-174.

432 Bae, H.D., Yanke, L.J., Cheng, K.J., Selinger, L.B., 1999. A novel staining method for detecting
433 phytase activity. *J. Microbiol. Methods* 39, 17-22.

434 Catzeddu, P., Mura, E., Parente, E., Sanna, M., Farris, G.A., 2006. Molecular characterization of
435 lactic acid bacteria from sourdough breads produced in Sardinia (Italy) and multivariate statistical
436 analyses of results. *Syst. Appl. Microbiol.* 29, 138-144.

437 Cocolin, L., Bisson, L.F., Mills, D.A., 2000. Direct profiling of the yeast dynamics in wine
438 fermentations. *FEMS Microbiol. Lett.* 189, 81-87.

439 Cocolin, L., Alessandria, V., Dolci, P., Gorra, R., Rantsiou, K., 2013. Culture independent
440 methods to assess the diversity and dynamics of microbiota during food fermentations. *Int. J Food*
441 *Microbiol.* 167, 29–43.

442 Corsetti, A., 2013. Technology of sourdough fermentation and sourdough applications. In:
443 Gobbetti, M., Gänzle, M. (Eds.), Handbook on Sourdough Biotechnology. Springer, New York, pp.
444 85-104.

445 Corsetti, A., Lavermicocca, P., Morea, M., Baruzzi, F., Tosti, N., Gobbetti, M., 2001. Phenotypic
446 and molecular identification and clustering of lactic acid bacteria and yeasts from wheat (species
447 *Triticum durum* and *Triticum aestivum*) sourdoughs of Southern Italy. Int. J. Food Microbiol. 64,
448 95-104.

449 De Man, J.C., Rogosa, D., Sharpe, M.E., 1960. A medium for the cultivation of lactobacilli. J.
450 Appl. Microbiol. 23, 130-135.

451 Dertli, E., Mercan, E., Arıcı, M., Yılmaz, M.T., Sağdıç, O., 2016. Characterisation of lactic acid
452 bacteria from Turkish sourdough and determination of their exopolysaccharide (EPS) production
453 characteristics. LWT-Food Sci. Technol. 71, 116-124.

454 De Vuyst, L., Harth, H., Van Kerrebroeck, S., Leroy, F., 2016. Yeast diversity of sourdoughs
455 and associated metabolic properties and functionalities. Int. J. Food Microbiol. 239, 26-34.

456 De Vuyst, L., Neysens, P., 2005. The sourdough microflora: biodiversity and metabolic
457 interactions. Trends Food Sci. Technol. 16, 43-56.

458 De Vuyst, L., Van Kerrebroeck, S., Harth, H., Huys, G., Daniel, H.M., Weckx, S., 2014.
459 Microbial ecology of sourdough fermentations: diverse or uniform? Food Microbiol. 37, 11-29.

460 Edgar, R.C., 2004a. MUSCLE: a multiple sequence alignment method with reduced time and
461 space complexity. BMC bioinformatics 5, 1.

462 Edgar, R.C., 2004b. MUSCLE: multiple sequence alignment with high accuracy and high
463 throughput. Nucleic acids research 32, 1792-1797.

464 Felsenstein J., 1985. Confidence limits on phylogenies: An approach using the bootstrap.
465 Evolution 39, 783-791.

466 Ferchichi, M., Valcheva, R., Oheix, N., Kabadjova, P., Prévost, H., Onno, B., Dousset, X., 2008.
467 Rapid investigation of French sourdough microbiota by restriction fragment length polymorphism

468 of the 16S-23S rRNA gene intergenic spacer region. World J. Microbiol. Biotechnol. 24, 2425-
469 2434.

470 Fernández-Espinar, M.T., Esteve-Zarzoso, B., Querol, A., Barrio, E., 2000. RFLP analysis of the
471 ribosomal internal transcribed spacers and the 5.8S rRNA gene region of the genus *Saccharomyces*:
472 a fast method for species identification and the differentiation of flour yeasts. Antonie van
473 Leeuwenhoek, 78, 87-97.

474 Foschino, R., Arrigoni, C., Mora, D., Galli, A., 2001. Phenotypic and genotypic aspects of
475 *Lactobacillus sanfranciscensis* strains isolated from sourdoughs in Italy. Food Microbiol. 18, 277-
476 285.

477 Gänzle, M., Ripari, V., 2016. Composition and function of sourdough microbiota: From
478 ecological theory to bread quality. Int. J. Food Microbiol. 239, 19-25.

479 Gatto, V., Torriani, S., 2004. Microbial population changes during sourdough fermentation
480 monitored by DGGE analysis of 16S and 26S rRNA gene fragments. Ann. Microbiol. 54, 31-42.

481 Gobbetti, M., 1998. The sourdough microflora: interactions of lactic acid bacteria and yeasts.
482 Trends Food Sci. Technol. 9, 267-274.

483 Gobbetti, M., Gänzle, M., 2013. Handbook on sourdough biotechnology, first ed. Springer, New
484 York.

485 Gobbetti, M., Rizzello, C.G., Di Cagno, R., De Angelis, M., 2014. How the sourdough may
486 affect the functional features of leavened baked goods. Food Microbiol. 37, 30-40.

487 Gullo, M., Romano, A.D., Pulvirenti, A., Giudici, P., 2003. *Candida humilis* - dominant species
488 in sourdoughs for the production of durum wheat bran flour bread. Int. J. Food Microbiol. 80, 55-
489 59.

490 Iacumin, L., Cecchini, F., Manzano, M., Osualdini, M., Boscolo, D., Orlic, S., Comi, G., 2009.
491 Description of the microflora of sourdoughs by culture-dependent and culture-independent
492 methods. Food Microbiol. 26, 128-135.

493 Istituto Nazionale di Sociologia Rurale, 2000. Atlante dei prodotti tipici: il pane (p. 13). Agra-
494 Rai Eri, Rome, Italy.

495 Jacques, N., Sarilar, V., Urien, C., Lopes, M.R., Morais, C.G., Uetanabaro, A.P.T., Tinsley C.R.,
496 Rosa C.A., Sicard D., Casaregola, S., 2016. Three novel ascomycetous yeast species of the
497 *Kazachstania* clade, *Kazachstania saulgeensis* sp. nov., *Kazachstania serrabonitensis* sp. nov. and
498 *Kazachstania australis* sp. nov. Reassignment of *Candida humilis* to *Kazachstania humilis* fa
499 comb nov and *Candida pseudohumilis* to *Kazachstania pseudohumilis* fa comb. nov. *International*
500 *Int. J. Syst. Evol. Microbiol.* 66, 5192-5200.

501 Jorquera, M.A., Hernández, M.T., Rengel, Z., Marschner, P., de la Luz Mora, M., 2008. Isolation
502 of culturable phosphobacteria with both phytate-mineralization and phosphate-solubilization
503 activity from the rhizosphere of plants grown in a volcanic soil. *Biol. Fertil. Soils* 44, 1025-1034.

504 Katina, K., Arendt, E., Liukkonen, K.H., Autio, K., Flander, L., Poutanen, K., 2005. Potential of
505 sourdough for healthier cereal products. *Trends Food Sci. Technol.* 16, 104-112.

506 Kimura, M., 1980. A simple method for estimating evolutionary rates of base substitutions
507 through comparative studies of nucleotide sequences. *J. Mol. Evol.* 16, 111-120.

508 Kline, L., Sugihara, T.F., 1971. Microorganisms of the San Francisco sour dough bread process
509 II. Isolation and characterization of undescribed bacterial species responsible for the souring
510 activity. *Applied Microbiol.* 21, 459-465.

511 Kurtzman, C.P., Robnett, C.J., 1998. Identification and phylogeny of ascomycetous yeasts from
512 analysis of nuclear large subunit (26S) ribosomal DNA partial sequences. *Antonie Van*
513 *Leeuwenhoek* 73, 331-371.

514 Kurtzman, C.P., Robnett, C.J., 2003. Phylogenetic relationships among yeasts of the
515 ‘*Saccharomyces* complex’ determined from multigene sequence analyses. *FEMS Yeast Res.* 3, 417-
516 432.

517 Lachance, M.A., Boekhout, T., Scorzetti, G., Fell, J.W., Kurtzman, C.P., 2011. *Candida*
518 *Berkhout*. In: Kurtzman, C., Fell, J.W., Boekhout, T. (Eds.), *The Yeasts: a Taxonomic Study*, fifth
519 ed., vol. 2. Elsevier, Amsterdam (NL), pp. 987-1278.

520 Lane, DJ., 1991. 16S/23S rRNA sequencing. In: Stackebrandt, E., Goodfellow, M. (Eds),
521 *Nucleic acid techniques in bacterial systematics*. Wiley, New York, pp. 115-175.

522 Lattanzi, A., Minervini, F., Di Cagno, R., Diviccaro, A., Antonielli, L., Cardinali, G., Cappelle,
523 S., De Angelis, M., Gobbetti, M., 2013. The lactic acid bacteria and yeast microbiota of eighteen
524 sourdoughs used for the manufacture of traditional Italian sweet leavened baked goods. *Int. J. Food*
525 *Microbiol.* 163, 71-79.

526 Liu, T., Li, Y., Chen, J., Sadiq, F.A., Zhang, G., Li, Y., He, G., 2016. Prevalence and diversity of
527 lactic acid bacteria in Chinese traditional sourdough revealed by culture dependent and
528 pyrosequencing approaches. *LWT-Food Sci. Technol.* 68, 91-97.

529 Minervini, F., De Angelis, M., Di Cagno, R., Gobbetti, M., 2014. Ecological parameters
530 influencing microbial diversity and stability of traditional sourdough. *Int. J. Food Microbiol.* 171,
531 136-146.

532 Minervini, F., Di Cagno, R., Lattanzi, A., De Angelis, M., Antonielli, L., Cardinali, G., Cappelle,
533 S., Gobbetti, M., 2012. Lactic acid bacterium and yeast microbiotas of 19 sourdoughs used for
534 traditional/typical Italian breads: interactions between ingredients and microbial species diversity.
535 *Appl. Environ. Microbiol.* 78, 1251-1264.

536 Nuobariene, L., Hansen, Å. S., Arneborg, N., 2012. Isolation and identification of phytase-active
537 yeasts from sourdoughs. *LWT-Food Sci. Technol.* 48, 190-196.

538 Osimani, A., Zannini, E., Aquilanti, L., Mannazzu, I.M., Comitini, F., Clementi, F., 2009. Lactic
539 acid bacteria and yeasts from wheat sourdoughs of the Marche region. *Ital. J Food Sci.* 21, 269-286.

540 Palla, M., Cristani, C., Giovannetti, M., Agnolucci, M., 2015. Identificazione molecolare di
541 batteri lattici e lieviti caratterizzanti l'impasto acido per la produzione di Pane Toscano a
542 lievitazione naturale. *Industrie Alimentari* 54, 5-11.

543 Palomba, S., Blaiotta, G., Ventorino, V., Saccone, A., Pepe, O., 2011. Microbial characterization
544 of sourdough for sweet baked products in the Campania region (southern Italy) by a polyphasic
545 approach. *Ann. Microbiol.* 61, 307-314.

546 Pulvirenti, A., Caggia, C., Restuccia, C., Gullo, M., Giudici, P., 2001. DNA fingerprinting
547 methods used for identification of yeasts isolated from Sicilian sourdoughs. *Ann. Microbiol.* 51,
548 107-120.

549 Randazzo, C.L., Heilig, H., Restuccia, C., Giudici, P., Caggia, C., 2005. Bacterial population in
550 traditional sourdough evaluated by molecular methods. *J. Appl. Microbiol.* 99, 251-258.

551 Reale, A., Tremonte, P., Succi, M., Sorrentino, E., Coppola, R., 2005. Exploration of lactic acid
552 bacteria ecosystem of sourdoughs from the Molise region. *Ann. Microbiol.* 55, 17-22.

553 Ricciardi, A., Parente, E., Piraino, P., Paraggio, M., Romano, P., 2005. Phenotypic
554 characterization of lactic acid bacteria from sourdoughs for Altamura bread produced in Apulia
555 (Southern Italy). *Int. J. Food Microbiol.* 98, 63-72.

556 Saitou, N., Nei, M., 1987. The neighbor-joining method: A new method for reconstructing
557 phylogenetic trees. *Mol. Biol. Evol.* 4, 406-425.

558 Scheirlinck, I., Van der Meulen, R., Van Schoor, A., Vancanneyt, M., De Vuyst, L., Vandamme,
559 P., Huys, G., 2007. Influence of geographical origin and flour type on diversity of lactic acid
560 bacteria in traditional Belgian sourdoughs. *Appl. Environ. Microbiol.* 73, 6262-6269.

561 Settanni, L., Valmorri, S., van Sinderen, D., Suzzi, G., Paparella, A., Corsetti, A., 2006.
562 Combination of multiplex PCR and PCR-denaturing gradient gel electrophoresis for monitoring
563 common sourdough-associated *Lactobacillus* species. *Appl. Environ. Microbiol.* 72, 3793-3796.

564 Tamura, K., Stecher, G., Peterson, D., Filipski, A., Kumar, S., 2013. MEGA6: Molecular
565 Evolutionary Genetics Analysis version 6.0. *Mol. Biol. Evol.* 30, 2725-2729.

566 Valmorri, S., Settanni, L., Suzzi, G., Gardini, F., Vernocchi, P., Corsetti, A., 2006. Application
567 of a novel polyphasic approach to study the lactobacilli composition of sourdoughs from the
568 Abruzzo region (central Italy). *Lett. Appl. Microbiol.* 43, 343-349.

569 Valmorri, S., Tofalo, R., Settanni, L., Corsetti, A., Suzzi, G., 2010. Yeast microbiota associated
570 with spontaneous sourdough fermentations in the production of traditional wheat sourdough breads
571 of the Abruzzo region (Italy). *Antonie Van Leeuwenhoek* 97, 119-129.

572 Venturi, M., Guerrini, S., Vincenzini, M., 2012. Stable and non-competitive association of
573 *Saccharomyces cerevisiae*, *Candida milleri* and *Lactobacillus sanfranciscensis* during manufacture
574 of two traditional sourdough baked goods. *Food Microbiol.* 31, 107-115.

575 Vera, A., Ly-Chatain, M.H., Rigobello, V., Demarigny, Y., 2012. Description of a French natural
576 wheat sourdough over 10 consecutive days focussing on the lactobacilli present in the microbiota.
577 *Antonie van Leeuwenhoek* 101, 369-377.

578 Vernocchi, P., Valmorri, S., Gatto, V., Torriani, S., Gianotti, A., Suzzi, G., Guerzoni, M.E.,
579 Gardini, F., 2004. A survey on yeast microbiota associated with an Italian traditional sweet-
580 leavened baked good fermentation. *Food Res. Int.* 37, 469-476.

581 Vigentini, I., Antoniani, D., Roscini, L., Comasio, A., Galafassi, S., Picozzi, C., Corte, L.,
582 Compagno, C., Dal Bello, F., Cardinali, G., Foschino, R., 2014. *Candida milleri* species reveals
583 intraspecific genetic and metabolic polymorphisms. *Food Microbiol.* 42, 72-81.

584 Walter, J., Hertel, C., Tannock, G.W., Lis, C. M., Munro, K., Hammes, W.P., 2001. Detection of
585 *Lactobacillus*, *Pediococcus*, *Leuconostoc*, and *Weissella* species in human feces by using group-
586 specific PCR primers and denaturing gradient gel electrophoresis. *Appl. Environ. Microbiol.* 67,
587 2578-2585.

588 Weisburg, W.G., Barns, S.M., Pelletier, D.A., Lane, D.J., 1991. 16S ribosomal DNA
589 amplification for phylogenetic study. *J. Bacteriol.* 173, 697-703.

590 White, T.J., Bruns, T., Lee, S., Taylor, J.W., 1990. Amplification and direct sequencing of fungal
591 ribosomal RNA genes for phylogenetics. In: Innis, M.A., Gelfand, D.H., Sninsky, J.J., White T.J.
592 (Eds.), *PCR protocols: a guide to methods and applications*. Academic Press Inc., New York, pp.
593 315-322.

594 Yazar, G., Tavman, Ş., 2012. Functional and technological aspects of sourdough fermentation
595 with *Lactobacillus sanfranciscensis*. Food Eng. Rev. 4, 171-190.

596 Zhang, J., Liu, W., Sun, Z., Bao, Q., Wang, F., Yu, J., Chen, W., Zhang, H., 2011. Diversity of
597 lactic acid bacteria and yeasts in traditional sourdoughs collected from western region in Inner
598 Mongolia of China. Food Control 22, 767-774.

599 Zotta, T., Piraino, P., Parente, E., Salzano, G., Ricciardi, A., 2008. Characterization of lactic acid
600 bacteria isolated from sourdoughs for *Cornetto*, a traditional bread produced in Basilicata (Southern
601 Italy). World J. Microbiol. Biotechnol. 24, 1785-1795.

602

603

604

605

606

607

608

609

610

611

612

613

614

615

616

617

618

619

620 **Table 1**

621 Lactic acid bacteria and yeast reference strains used in this study.

<i>Strains^a</i>	<i>Source of isolation</i>
<i>Lactobacillus panis</i> DSMZ 6035 ^T	Sourdough
<i>Lactobacillus sanfranciscensis</i> DSMZ 20451 ^T	San Francisco sourdough
<i>Lactobacillus fermentum</i> DSMZ 20052 ^T	Fermented beets
<i>Lactobacillus brevis</i> DSMZ 20054 ^T	Faeces
<i>Lactobacillus plantarum</i> IMA B23	Boza
<i>Lactobacillus curvatus</i> IMA LB51	Sourdough
<i>Saccharomyces cerevisiae</i> ATCC 32167	Unknow
<i>Dekkera bruxellensis</i> IMA 1L	San Giovese Tuscan wine
<i>Candida milleri</i> DBVPG 6753 ^T	San Francisco sourdough
<i>Candida humilis</i> DBVPG 7219 ^T	Bantu beer
<i>Candida humilis</i> DBVPG 6754	Sourdough, Finland
<i>Kazachstania exigua</i> DBVPG 6956	Wheat sourdough, Italy

622

623 ^TType Strain.624 ^aDSMZ=Deutsche Sammlung von Mikroorganismen und Zellkulturen, Braunschweig, Germany;

625 IMA=International Microbial Archives, Department of Agriculture, Food and Environment,

626 University of Pisa, Pisa, Italy; ATCC=American Type culture Collection, Manassas, Virginia,

627 USA; DBVPG=International Collection of Department of Agricultural, Food and Environmental

628 Science, University of Perugia, Perugia, Italy.

629

630 **Table 2**

631 Lactic acid bacteria and yeasts characterizing the PDO Tuscan bread sourdough: identification by
 632 sequencing of different representative isolate genes and of DNA fragments in DGGE profiles.

<i>Isolates</i>	<i>Genes</i>	<i>Accession number</i>	<i>Species in NCBI database</i>	<i>Sequence identity</i>
IMA 2LAB, 3, 13, 19, 23, 29, 44, 64, 67, 93, 97	16S	From LT605146 to LT605156	<i>L. sanfranciscensis</i> DSMZ 20663 (X76331) <i>L. sanfranciscensis</i> ATCC 27651 ^T (X76327)	99%
IMA 105Y	ITS1-5.8S-ITS2	LT605145	<i>S. cerevisiae</i> ATCC 834 (KU729072)	98%
IMA 19Y, 36, 105	26S D1/D2	From LT718652 to LT718654	<i>S. cerevisiae</i> CBS 2962 (KY109317)	100%
IMA 11Y, 33	26S D1/D2	From LT718655 to LT718656	<i>C. milleri</i> CBS 6897 (KY106585)	100%
<i>DGGE fragments</i>				
5LAB, 14LAB, 29LAB	16S V3-V5	From LT605157 to LT605159	<i>L. sanfranciscensis</i> ATCC 27651 ^T (X76327) <i>L. sanfranciscensis</i> DSMZ 20663 (X76331)	99%
1Y, 9Y, 19Y	partial 26S D1/D2	From LT605160 to LT605162	<i>C. milleri</i> NRRL Y-7245 ^T (U94923) <i>C. humilis</i> NRRL Y-17074 ^T (U69878)	99% 99%
3Y, 15Y, 20Y	partial 26S D1/D2	From LT605163 to LT605165	<i>S. cerevisiae</i> CTBRL121 (JX423567)	100%

633

634

635 **Table 3**

636 Phytase, protease and amylase activities of lactic acid bacteria and yeasts isolated from the PDO

637 Tuscan bread sourdough.

<i>Isolates</i>	<i>Phytase activity</i>	<i>Protease activity</i>	<i>Amylase activity</i>
<i>Lactic acid bacteria</i>			
IMA 1LAB; 3; 5-13; 17-21; 23-26; 28-38; 40-49; 51; 52; 56-58; 60-62; 64; 66-75; 77-82; 84-86; 88-92; 94-96; 99	-	-	-
IMA 2LAB; 22; 39; 53; 63; 76; 87; 93	+	-	-
IMA 14-16LAB; 27; 55; 59; 65; 97	-/+	-	-
IMA 83LAB; 98	++	-	-
<i>Yeasts</i>			
IMA 1Y; 122	+	-/+	-
IMA 2-3Y; 9-10; 12; 17-18; 20; 34; 40; 48-49; 103-104; 106-108	-	-/+	-
IMA 4-8Y; 13-16; 21-26; 28; 30-31; 35; 38; 41; 43; 44-47; 50-51; 109-120	-	-	-
IMA 11Y; 32; 37	-	+	-
IMA 19Y	+	+	+
IMA 27Y; 29	-	++	-
IMA 33Y	+	-/+	++
IMA 36Y	++	-/+	++
IMA 105Y	+	-/+	+

638

639 Lactic acid bacteria. Phytase activity: - = no activity (halo = 0mm), +/- = low activity (halo \leq 1mm),640 + = moderate activity (1mm < halo \leq 4mm), ++ = high activity (halo > 4mm).641 Yeasts. Phytase activity: - = no activity (halo = 0mm), +/- = low activity (halo \leq 1mm), + =642 moderate activity (1mm < halo \leq 15mm), ++ = high activity (halo > 15mm); protease activity: - =643 no activity (halo = 0mm), +/- = low activity (halo \leq 1mm), + = moderate activity (1mm < halo \leq

644 5mm), ++ = high activity (halo > 5mm); amylase activity: - = no activity (halo = 0mm), +/- = low

645 activity (halo \leq 1mm), + = moderate activity (1mm < halo \leq 4mm), ++ = high activity (halo >

646 4mm).

647

648

649

650 **FIGURE CAPTIONS**

651 **Fig. 1.** Dendrogram obtained from UPGMA analysis, using Dice's coefficient, of combined
652 ARDRA profiles of 96 LAB isolates and LAB reference strains. The scale indicates the similarity
653 level.

654

655 **Fig. 2.** Dendrogram obtained from UPGMA analysis, using Dice's coefficient, of combined ITS-
656 RFLP profiles and ITS amplicons size of 68 yeast isolates and yeast reference strains. The scale
657 indicates the similarity level.

658

659 **Fig. 3.** Dendrogram showing multiple sequence alignment of ITS1-5.8S-ITS2 region of *Candida*
660 *milleri* strains isolated from the PDO Tuscan bread sourdough. The evolutionary history was
661 inferred using the Neighbor-Joining method (Saitou and Nei, 1987). The optimal tree with the sum
662 of branch length = 0.54023792 is shown. The percentage of replicate trees in which the associated
663 taxa clustered together in the bootstrap test (1000 replicates) are shown next to the branches
664 (Felsenstein, 1985). The tree is drawn to scale, with branch lengths in the same units as those of the
665 evolutionary distances used to infer the phylogenetic tree. The evolutionary distances were
666 computed using the Kimura 2-parameter method (Kimura, 1980) and are in the units of the number
667 of base substitutions per site. The rate variation among sites was modeled with a gamma
668 distribution (shape parameter = 1). The analysis involved 83 nucleotide sequences. All positions
669 with less than 95% site coverage were eliminated. That is, fewer than 5% alignment gaps, missing
670 data, and ambiguous bases were allowed at any position. There were a total of 524 positions in the
671 final dataset. Evolutionary analyses were conducted in MEGA6 (Tamura et al., 2013). Numbers in
672 parentheses are accession numbers of published sequences.

673

674




Fig. 1




Fig. 2




Fig. 3


Identification and characterization of lactic acid bacteria and yeasts of PDO Tuscan bread sourdough by culture dependent and independent methods

Michela Palla, Caterina Cristani, Manuela Giovannetti, Monica Agnolucci*

Department of Agriculture, Food and Environment, University of Pisa, Via del Borghetto 80, 56124 Pisa, Italy

*Corresponding author: M. Agnolucci, Department of Agriculture, Food and Environment, University of Pisa, Via del Borghetto 80, 56124 Pisa, Italy Phone: +39.0502216647, Fax: +39.0502210606, e-mail address: monica.agnolucci@unipi.it

Fig. S1. Electrophoresis of *AluI*, *HaeIII* and *HinfI* ARDRA patterns of LAB reference strains and isolates. 1: *L. sanfranciscensis* DSMZ 20451^T; 2: *L. brevis* DSMZ 20054^T; 3: *L. plantarum* IMA B23; 4: *L. curvatus* IMA LB51; 5: *L. fermentum* DSMZ 20052^T; 6: *L. panis* DSMZ 6035^T; 7-18: IMA 15LAB-IMA 26LAB; M: Marker 100bp (BioLabs).




Identification and characterization of lactic acid bacteria and yeasts of PDO Tuscan bread sourdough by culture dependent and independent methods

Michela Palla, Caterina Cristani, Manuela Giovannetti, Monica Agnolucci*

Department of Agriculture, Food and Environment, University of Pisa, Via del Borghetto 80, 56124 Pisa, Italy

*Corresponding author: M. Agnolucci, Department of Agriculture, Food and Environment, University of Pisa, Via del Borghetto 80, 56124 Pisa, Italy Phone: +39.0502216647, Fax: +39.0502210606, e-mail address: monica.agnolucci@unipi.it

Fig. S2. Electrophoresis of *Hae*III and *Hinf*I RFLP patterns of the ITS regions of yeast reference strains and isolates. 1: *S. cerevisiae* ATCC 32167; 2: *D. bruxellensis* IMA 1L; 3: *K. exigua* DBVPG 6956; 4: *C. milleri* DBVPG 6753^T; 5: *C. humilis* DBVPG 7219^T; 6: *C. humilis* DBVPG 6754; 7-20: IMA 19Y-IMA 35Y; M: Marker 100bp (BioLabs).




Identification and characterization of lactic acid bacteria and yeasts of PDO Tuscan bread sourdough by culture dependent and independent methods

Michela Palla, Caterina Cristani, Manuela Giovannetti, Monica Agnolucci*

Department of Agriculture, Food and Environment, University of Pisa, Via del Borghetto 80, 56124 Pisa, Italy

*Corresponding author: M. Agnolucci, Department of Agriculture, Food and Environment, University of Pisa, Via del Borghetto 80, 56124 Pisa, Italy Phone: +39.0502216647, Fax: +39.0502210606, e-mail address: monica.agnolucci@unipi.it

Fig. S3. DGGE profiles of the microbial community of PDO Tuscan bread sourdough. (a) LAB 16S rRNA gene V3-V4 region; 1, 2, 3: sourdough replicates; marker (M): ● *Lb. plantarum* IMA B23, ● *Lb. brevis* DSMZ 20054, ● *Lb. curvatus* IMA LB51, ● *Lb. fermentum* DSMZ 20052, ● *Lb. sanfranciscensis* DSMZ 20451, ● *Lb. panis* DSMZ 6035. (b) Yeast 26S rRNA gene D1/D2 region; 1, 2, 3: sourdough replicates; marker (M): ● *S. cerevisiae* ATCC 32167, ● *K. exigua* DBVPG 6956, ● *C. milleri* DBVPG 6753/*C. humilis* DBVPG 7219, ● *D. bruxellensis* IMA 1L. The numbers indicate sequenced DNA fragments and the colored circles their species affiliation.



Identification and characterization of lactic acid bacteria and yeasts of PDO Tuscan bread sourdough by culture dependent and independent methods

Michela Palla, Caterina Cristani, Manuela Giovannetti, Monica Agnolucci*

Department of Agriculture, Food and Environment, University of Pisa, Via del Borghetto 80, 56124 Pisa, Italy

*Corresponding author: M. Agnolucci, Department of Agriculture, Food and Environment, University of Pisa, Via del Borghetto 80, 56124 Pisa, Italy Phone: +39.0502216647, Fax: +39.0502210606, e-mail address: monica.agnolucci@unipi.it

Fig. S4. Dendrogram showing multiple sequence alignment of ITS1-5.8S-ITS2 region of *C. milleri* and *C. humilis* collected from GeneBank. The evolutionary history was inferred using the Neighbor-Joining method. Evolutionary analyses were conducted in MEGA6. For each sequence is reported its source of isolation and the relative number of *Hae*III restriction sites.

