

The canonical ring of a 3-connected curve *

Marco Franciosi, Elisa Tenni

Abstract

Let C be a Gorenstein curve which is either reduced or contained in a smooth algebraic surface.

We show that the canonical ring $R(C, \omega_C) = \bigoplus_{k \geq 0} H^0(C, \omega_C^{\otimes k})$ is generated in degree 1 if C is 3-connected and not (honestly) hyperelliptic; we show moreover that $R(C, L) = \bigoplus_{k \geq 0} H^0(C, L^{\otimes k})$ is generated in degree 1 if C is reduced and L is an invertible sheaf such that $\deg L|_B \geq 2p_a(B) + 1$ for every $B \subseteq C$.

keyword: algebraic curve, Noether's theorem, canonical ring

Mathematics Subject Classification (2010) 14H20, 14C20, 14H51

1 Introduction

Let C be a Gorenstein curve which is either reduced or contained in a smooth algebraic surface, ω_C be its dualizing sheaf of C and L be an invertible sheaf on C .

The main result of this paper is Theorem 3.3 stating that the canonical ring

$$R(C, \omega_C) = \bigoplus_{k \geq 0} H^0(C, \omega_C^{\otimes k})$$

is generated in degree 1 if C is a 3-connected and not honestly hyperelliptic curve (see Def. 2.1 and Def. 2.2). This is a generalization to singular curves of the classical Theorem of Noether for smooth curves (see [1, §III.2]) and can be regarded as a first step in a more general analysis of the Koszul groups $\mathcal{K}_{p,q}(C, \omega_C)$ of 3-connected curves (see [10] for the definition and the statement of the so called “*Green's conjecture*”). A detailed explanation of the role that the Koszul groups of smooth and singular curves play in the geometry of various moduli spaces can be found in [2].

*This research was partially supported by Italian MIUR through PRIN 2008 project “Geometria delle varietà algebriche e dei loro spazi di moduli”.

Additional motivation for the present work comes from the theory of surface fibrations, as shown by Catanese and Ciliberto in [3] and Reid in [15]. Indeed, given a surface fibration $f: S \rightarrow B$ over a smooth curve B , the *relative canonical algebra* $R(f) = \bigoplus_{n \geq 0} f_*(\omega_{S/B}^{\otimes n})$ gives important information on the geometry of the surface. It is clear that the behaviour of $R(f)$ depends on the canonical ring of every fibre.

The main result on the canonical ring for singular curves in the literature is the 1-2-3 conjecture, stated by Reid in [15] and proved in [7] and [13], which says that the canonical ring $R(C, \omega_C)$ of a connected Gorenstein curve of arithmetic genus $p_a(C) \geq 3$ is generated in degree 1, 2, 3, with the exception of a small number of cases. More recently in [8] the first author proved that the canonical ring is generated in degree 1 under the strong assumption that C is *even* (i.e., $\deg_B K_C$ is even on every subcurve $B \subseteq C$).

We remark that our result implies in particular that the canonical ring of a regular surface of general type is generated in degree ≤ 3 if there exists a curve $C \in |K_S|$ 3-connected and not honestly hyperelliptic (see [8, Thm. 1.2]). Moreover one can apply the same argument of Konno (see [13, thm. III]) and see that the relative canonical algebra of a relatively minimal surface fibration is generated in degree 1 if every fibre is 3-connected and not honestly hyperelliptic.

Our second result is Theorem 4.2 stating that the ring

$$R(C, L) = \bigoplus_{k \geq 0} H^0(C, L^{\otimes k})$$

considered as an algebra over $H^0(C, \mathcal{O}_C)$, is generated in degree 1 if C is reduced and L is an invertible sheaf such that $\deg L|_B \geq 2p_a(B) + 1$ for every $B \subseteq C$. This is a generalization of a Theorem of Castelnuovo (see [14]) on the projective normality of smooth projective curves. This result can be useful when dealing with properties (for instance Brill-Noether properties) of a family of smooth curves which degenerates to C .

In [7], [8], [13] the analysis of the canonical ring is based on the study of the Koszul groups $\mathcal{K}_{p,q}(C, \omega_C)$ (with p, q small) and their vanishing properties, together with some vanishing results for invertible sheaves of low degree.

In this paper we use a completely different approach. Our method is inspired by the arguments developed in a series of papers by Green and Lazarsfeld which appeared in the late '80s (see [10], [11]) and it is based on the generalization to singular curves of Clifford's Theorem given by the authors in [9].

Given an invertible sheaf L such that the map $H^0(C, L) \otimes H^0(C, L) \rightarrow H^0(C, L^{\otimes 2})$ fails to be surjective, we exhibit a 0-dimensional scheme S such that the map

$H^0(C, L) \otimes H^0(C, L) \rightarrow H^0(S, \mathcal{O}_S)$ induced by the restriction also fails to be surjective. Thus its dual map

$$\varphi: \text{Ext}^1(\mathcal{O}_S, \omega_C \otimes L^{-1}) \rightarrow \text{Hom}(H^0(C, L), H^1(C, \omega_C \otimes L^{-1}))$$

is not injective. From the analysis of an extension in the Kernel of φ we conclude that the cohomology of $\mathcal{I}_S \cdot L$ must satisfy some numerical conditions. This in turn contradicts Clifford's Theorem when $L = K_C$, or $\deg L|_B \geq 2p_a(B) + 1$ on every $B \subseteq C$.

Finally, we make an important comment on the role of numerical connection in generalizing Noether's theorem. By the results of [5] (see [5, §2, §3] or Theorem 2.3) and our main result Theorem 3.3 we have the following implications for a connected curve C

$$C \text{ 3-connected, not honestly hyperelliptic} \implies R(C, \omega_C) \text{ is generated in degree 1 and } \omega_C \text{ ample} \implies \omega_C \text{ is very ample}$$

If C is reduced it is known that the three properties are equivalent (see [4]). However this is false when C is not reduced. To see that the converse of the first implication fails one can take $C = 2F$, where F is a non hyperelliptic fibre of a surface fibration. In Ex. 3.4 below we will construct a curve with very ample canonical sheaf which fails Noether's Theorem, thus proving that the converse of the second implication is false. This examples support our belief that 3-connected curves are the most natural generalization of smooth curves when dealing with the properties of the canonical embedding.

Acknowledgments. The second author wishes to thank the Department of Mathematic of the University of Pisa, especially Rita Pardini, for providing an excellent research environment.

2 Notation and preliminary results

We work over an algebraically closed field \mathbb{K} of characteristic ≥ 0 .

Throughout this paper C will be a Gorenstein curve, either reduced or contained in a smooth algebraic surface X , in which case we allow C to be reducible and non reduced.

ω_C denotes the dualizing sheaf of C (see [12], Chap. III, §7), and $p_a(C)$ the arithmetic genus of C , $p_a(C) = 1 - \chi(\mathcal{O}_C)$. K_C denotes the canonical divisor.

Definition 2.1 *A curve C is honestly hyperelliptic if there exists a finite morphism $\psi: C \rightarrow \mathbb{P}^1$ of degree 2. (see [5, §3] for a detailed treatment).*

If A, B are subcurves of C such that $A + B = C$, then their product $A \cdot B$ is

$$A \cdot B = \deg_B(K_C) - (2p_a(B) - 2) = \deg_A(K_C) - (2p_a(A) - 2).$$

If C is contained in a smooth algebraic surface X this corresponds to the intersection product of curves as divisors on X .

Definition 2.2 C is m -connected if for every decomposition $C = A + B$ one has $A \cdot B \geq m$. C is numerically connected if it is 1-connected.

First we recall some useful results proved in [4] and [5].

Theorem 2.3 ([5] §2, §3) *Let C be a Gorenstein curve. Then*

- (i) *If C is 1-connected then $H^1(C, K_C) \cong \mathbb{K}$.*
- (ii) *If C is 2-connected and $C \not\cong \mathbb{P}^1$ then $|K_C|$ is base point free.*
- (iii) *If C is 3-connected and C is not honestly hyperelliptic (i.e., there does not exist a finite morphism $\psi: C \rightarrow \mathbb{P}^1$ of degree 2) then K_C is very ample.*

Proposition 2.4 ([5], Lemma 2.4) *Let C be a projective scheme of pure dimension 1, let \mathcal{F} be a coherent sheaf on C , and $\varphi: \mathcal{F} \rightarrow \omega_C$ a nonvanishing map of \mathcal{O}_C -modules. Set $\mathcal{J} = \text{Ann } \varphi \subset \mathcal{O}_C$, and write $B \subset C$ for the subscheme defined by \mathcal{J} . Then B is Cohen–Macaulay and φ has a canonical factorization of the form*

$$\mathcal{F} \rightarrow \mathcal{F}|_B \hookrightarrow \omega_B = \mathcal{H}om_{\mathcal{O}_C}(\mathcal{O}_B, \omega_C) \subset \omega_C,$$

where $\mathcal{F}|_B \hookrightarrow \omega_B$ is generically onto.

Proposition 2.5 ([4]) *Let C be a pure 1-dimensional projective scheme, let \mathcal{F} be a rank 1 torsion free sheaf on C .*

- (i) *If $\deg(\mathcal{F})|_B \geq 2p_a(B) - 1$ for every subcurve $B \subseteq C$ then $H^1(C, \mathcal{F}) = 0$.*
- (ii) *If \mathcal{F} is invertible and $\deg(\mathcal{F})|_B \geq 2p_a(B)$ for every subcurve $B \subseteq C$ then $|\mathcal{F}|$ is base point free.*
- (iii) *If \mathcal{F} is invertible and $\deg(\mathcal{F})|_B \geq 2p_a(B) + 1$ for every subcurve $B \subseteq C$ then \mathcal{F} is very ample on C .*

As we mentioned in the Introduction, our approach to the analysis of the ring $R(C, L) = \bigoplus_{k \geq 0} H^0(C, L^{\otimes k})$ for a line bundle L builds on the generalization of Clifford's Theorem proved by the authors in [9]. In the rest of this section we recall the main results we need from [9], namely, the notion of subcanonical cluster and Clifford's Theorem, and we prove some technical lemmas on the cohomology of rank one torsion free sheaves.

Definition 2.6 A cluster S of degree r is a 0-dimensional subscheme of C with length $\mathcal{O}_S = \dim_k \mathcal{O}_S = r$. A cluster $S \subset C$ is subcanonical if the space $H^0(C, \mathcal{I}_S \omega_C)$ contains a generically invertible section, i.e., a section s_0 which does not vanish on any subcurve of C .

Theorem 2.7 ([9], Theorem A) Let C be a Gorenstein 2-connected curve which is either reduced or contained in a smooth algebraic surface, and let $S \subset C$ be a subcanonical cluster.

Assume that S is a Cartier divisor or alternatively that there exists a generically invertible section $H \in H^0(C, \mathcal{I}_S K_C)$ such that $\text{div}(H) \cap \text{Sing}(C_{\text{red}}) = \emptyset$.

Then

$$h^0(C, \mathcal{I}_S K_C) \leq p_a(C) - \frac{1}{2} \deg(S).$$

Moreover if equality holds then the pair (S, C) satisfies one of the following assumptions:

- (i) $S = 0, K_C$;
- (ii) C is honestly hyperelliptic and S is a multiple of the honest g_2^1 ;
- (iii) C is 3-disconnected (i.e., there is a decomposition $C = A + B$ with $A \cdot B = 2$).

Remark 2.8 Let C and S be as in Theorem 2.7. Then Riemann-Roch implies that

$$h^0(C, \mathcal{I}_S K_C) + h^1(C, \mathcal{I}_S K_C) \leq p_a(C) + 1$$

and equality holds if one of the three cases listed in Theorem 2.7 is satisfied.

Remark 2.9 If $Z_0 \subset Z$ are clusters, then the natural restriction map $\mathcal{O}_Z \rightarrow \mathcal{O}_{Z_0}$ induces an inclusion $\text{Ext}^1(\mathcal{O}_{Z_0}, \mathcal{O}_C) \hookrightarrow \text{Ext}^1(\mathcal{O}_Z, \mathcal{O}_C)$.

Lemma 2.10 Let C be a Gorenstein curve and Z a cluster. Assume there exists an extension $\xi \in \text{Ext}^1(\mathcal{O}_Z, \mathcal{O}_C)$ such that $\xi \notin \text{Ext}^1(\mathcal{O}_{Z_0}, \mathcal{O}_C)$ for every proper subcluster $Z_0 \subsetneq Z$. Then the corresponding extension can be written as

$$0 \rightarrow \mathcal{O}_C \rightarrow \mathcal{H}om(\mathcal{I}_Z, \mathcal{O}_C) \rightarrow \mathcal{O}_Z \rightarrow 0.$$

Proof. Consider an extension corresponding to ξ

$$0 \rightarrow \mathcal{O}_C \rightarrow E_\xi \rightarrow \mathcal{O}_Z \rightarrow 0. \quad (1)$$

We prove first that E_ξ is torsion free. Indeed, if $\text{Tor}(E_\xi) \neq 0$ then there exists a subcluster $Z_0 \subset Z$ and a sheaf $E_0 \cong E_\xi / \text{Tor}(E_\xi)$ which fits in the following commutative diagram:

$$\begin{array}{ccccccc}
& & & \text{Tor}(E_\xi) & \xrightarrow{\cong} & \text{Tor}(E_\xi) & \\
& & & \downarrow & & \downarrow & \\
0 & \longrightarrow & \mathcal{O}_C & \longrightarrow & E_\xi & \longrightarrow & \mathcal{O}_Z \longrightarrow 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
0 & \longrightarrow & \mathcal{O}_C & \longrightarrow & E_0 & \longrightarrow & \mathcal{O}_{Z_0} \longrightarrow 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
& & 0 & & 0 & & 0
\end{array}$$

In particular there exists a proper subcluster Z_0 such that the extension corresponding to E_0 in $\text{Ext}^1(\mathcal{O}_{Z_0}, \mathcal{O}_C)$ corresponds to ξ , which is impossible.

Since E_ξ is a rank 1 torsion free sheaf it is reflexive, i.e., there is a natural isomorphism $\mathcal{H}om(\mathcal{H}om(E_\xi, \mathcal{O}_C), \mathcal{O}_C) \cong E_\xi$. Dualizing sequence (1), since $\mathcal{E}xt^1(\mathcal{O}_Z, \mathcal{O}_C) \cong \mathcal{O}_Z$ and $\mathcal{E}xt^1(E_\xi, \mathcal{O}_C) = 0$, we see that $\mathcal{H}om(E_\xi, \mathcal{O}_C) \cong \mathcal{I}_Z$, hence $E_\xi \cong \mathcal{H}om(\mathcal{I}_Z, \mathcal{O}_C)$. ■

Remark 2.11 Throughout the paper we repeatedly use the natural isomorphisms $\mathcal{O}_S \cong \mathcal{O}_S \cdot L$ and $H^0(S, \mathcal{O}_S)^* \cong \text{Ext}^1(\mathcal{O}_S, \omega_C \otimes L^{-1}) \cong \text{Ext}^1(\mathcal{O}_S \cdot L, \omega_C \otimes L^{-1})$ for every cluster S on C and for every invertible sheaf L .

A useful tool in the analysis of the multiplication map $H^0(C, L)^{\otimes 2} \rightarrow H^0(C, L^{\otimes 2})$ is the restriction to a suitable cluster S . Indeed the composition of the multiplication map $H^0(C, L)^{\otimes 2} \rightarrow H^0(C, L^{\otimes 2})$ with the evaluation map $H^0(C, L^{\otimes 2}) \rightarrow H^0(S, \mathcal{O}_S)$ yields a natural map $H^0(C, L) \otimes H^0(C, L) \rightarrow H^0(S, \mathcal{O}_S)$.

Lemma 2.12 *Let C be a Gorenstein curve and L an effective line bundle on C . Let S be a cluster such that the restriction map*

$$H^0(C, L) \otimes H^0(C, L) \rightarrow H^0(S, \mathcal{O}_S)$$

is not surjective. Then there exists a nonempty subcluster $S_0 \subseteq S$ such that

$$h^0(C, L) + h^1(C, L) \leq h^0(C, \mathcal{I}_{S_0}L) + h^1(C, \mathcal{I}_{S_0}L).$$

Proof. Let S be a cluster such that the restriction map

$$H^0(C, L) \otimes H^0(C, L) \rightarrow H^0(S, \mathcal{O}_S)$$

is not surjective. By Serre duality the dual map

$$\varphi: \text{Ext}^1(\mathcal{O}_S, \omega_C \otimes L^{-1}) \rightarrow \text{Hom}(H^0(C, L), H^1(C, \omega_C \otimes L^{-1}))$$

is not injective. The dual map φ is given as follows: consider an element $\xi \in \text{Ext}^1(\mathcal{O}_S, \omega_C \otimes L^{-1})$ and its corresponding extension

$$0 \rightarrow \omega_C \otimes L^{-1} \rightarrow E_\xi \rightarrow \mathcal{O}_S \rightarrow 0.$$

Let $c_\xi: H^0(S, \mathcal{O}_S) \rightarrow H^1(C, \omega_C \otimes L^{-1})$ be the connecting homomorphism induced by the extension. Then the restriction map $r: H^0(C, L) \rightarrow H^0(S, \mathcal{O}_S)$ induces a map $\varphi_\xi = c_\xi \circ r: H^0(C, L) \rightarrow H^1(C, \omega_C \otimes L^{-1})$ given as follows

$$\begin{array}{ccccccc} & & & & H^0(C, L) & & \\ & & & & \downarrow r & \searrow \varphi_\xi & \\ 0 & \longrightarrow & H^0(C, \omega_C \otimes L^{-1}) & \longrightarrow & H^0(C, E_\xi) & \xrightarrow{f_\xi} & H^0(S, \mathcal{O}_S) & \xrightarrow{c_\xi} & H^1(C, \omega_C \otimes L^{-1}) \end{array}$$

The map φ_ξ is precisely $\varphi(\xi) \in \text{Hom}(H^0(C, L), H^1(C, \omega_C \otimes L^{-1}))$. By definition $\varphi(\xi) = 0$ if and only if $\text{Im}(r) \subset \text{Im}(f_\xi)$. In particular if $\varphi(\xi) = 0$ then we have $\dim \text{Im}(r) \leq \dim \text{Im}(f_\xi)$ which implies that

$$h^0(C, L) + h^1(C, L) \leq h^0(C, E_\xi) + h^0(C, \mathcal{I}_S L). \quad (2)$$

In order to prove the Lemma let S_0 be minimal (with respect to the inclusion) among the subclusters of S for which the restriction $\text{Sym}^2 H^0(C, L) \rightarrow H^0(S_0, \mathcal{O}_{S_0})$ fails to be surjective. This implies that if $Z \subsetneq S_0$ is any subcluster then the map

$$\varphi_0: \text{Ext}^1(\mathcal{O}_Z, \omega_C \otimes L^{-1}) \rightarrow \text{Hom}(H^0(C, L), H^1(C, \omega_C \otimes L^{-1}))$$

is injective. Note that φ_0 factors through φ :

$$\begin{array}{ccc} \text{Ext}^1(\mathcal{O}_Z, \omega_C \otimes L^{-1}) & & \\ \downarrow & \searrow \varphi_0 & \\ \text{Ext}^1(\mathcal{O}_{S_0}, \omega_C \otimes L^{-1}) & \xrightarrow{\varphi} & \text{Hom}(H^0(C, L), H^1(C, \omega_C \otimes L^{-1})) \end{array}$$

By the minimality of S_0 if $\xi \in \text{Ext}^1(\mathcal{O}_{S_0}, \omega_C \otimes L^{-1})$ is in the kernel of φ , it must not belong to the image of $\text{Ext}^1(\mathcal{O}_Z, \omega_C \otimes L^{-1})$ for every $Z \subsetneq S_0$. The corresponding extension E_ξ is isomorphic to $\mathcal{H}om(\mathcal{I}_{S_0}, \mathcal{O}_C) \otimes \omega_C \otimes L^{-1} \cong \mathcal{H}om(\mathcal{I}_{S_0}L, \omega_C)$ thanks to Lemma 2.10. Thus $h^0(C, E_\xi) = h^1(C, \mathcal{I}_{S_0}L)$ by Serre duality. Inequality (2) becomes

$$h^0(C, L) + h^1(C, L) \leq h^0(C, \mathcal{I}_{S_0}L) + h^1(C, \mathcal{I}_{S_0}L).$$

■

3 Noether's Theorem for singular curves

The aim of this section is to prove Theorem 3.3, i.e., Noether's Theorem for singular curves. For the proof we use two main ingredients: a generalization of the *free pencil trick*, given in Lemma 3.2, and the surjectivity of the restriction map $H^0(C, \omega_C)^{\otimes 2} \rightarrow H^0(S, \mathcal{O}_S)$ for a suitable cluster S .

Lemma 3.1 *Let C be a Gorenstein curve which is either reduced or contained in a smooth algebraic surface. Assume that C is 2-connected and $p_a(C) \geq 2$. Let $H \in H^0(C, \omega_C)$ be a generic section.*

Then there exists a cluster S contained in $\text{div} H$ such that the following hold:

1. $h^0(C, \mathcal{I}_S \omega_C) = 2$
2. *the evaluation map $H^0(C, \mathcal{I}_S K_C) \otimes \mathcal{O}_C \rightarrow \mathcal{I}_S \omega_C$ is surjective.*

Proof. Since $|K_C|$ is base point free thanks to Theorem 2.3 we may assume that H is generically invertible and $\text{div} H$ is a length $2p_a(C) - 2$ cluster. Thus for every integer $v \in \{1, \dots, p_a(C)\}$ there exists at least one cluster $S_v \subseteq \text{div} H$ such that $h^0(C, \mathcal{I}_{S_v} \omega_C) = v$. In particular we may take a cluster S such that $h^0(C, \mathcal{I}_S \omega_C) = 2$ and S is maximal up to inclusion among the clusters contained in $\text{div} H$ with this property. S is the desired cluster. Indeed, if it were $S_0 \supsetneq S$ such that the image of the evaluation map $H^0(C, \mathcal{I}_{S_0} K_C) \otimes \mathcal{O}_C \rightarrow \mathcal{I}_{S_0} \omega_C$ was $\mathcal{I}_{S_0} \omega_C \subsetneq \mathcal{I}_S \omega_C$ then we would have $h^0(C, \mathcal{I}_{S_0} \omega_C) = 2$, contradicting the maximality of S . ■

Even though the sheaf $\mathcal{I}_S \omega_C$ defined in the above Lemma is not usually a line bundle with abuse of notation we will call it a *free pencil*.

Lemma 3.2 *Let the pair (C, S) be as in the previous lemma. Then the map*

$$H^0(C, \mathcal{I}_S \omega_C) \otimes H^0(C, \omega_C) \xrightarrow{m} H^0(C, \mathcal{I}_S \omega_C^{\otimes 2}) \quad (3)$$

is surjective.

Proof. Consider the evaluation map $H^0(C, \mathcal{I}_S \omega_C) \otimes \omega_C \xrightarrow{\text{ev}} \mathcal{I}_S \omega_C^{\otimes 2}$ and its kernel \mathcal{K} :

$$0 \rightarrow \mathcal{K} \rightarrow H^0(C, \mathcal{I}_S \omega_C) \otimes \omega_C \rightarrow \mathcal{I}_S \omega_C^{\otimes 2} \rightarrow 0. \quad (4)$$

The map (3) is surjective if and only if $h^1(C, \mathcal{K}) = 2$ since $h^1(C, \mathcal{I}_S \omega_C^{\otimes 2}) = 0$ by Proposition 2.5. In the rest of the proof we establish $h^1(C, \mathcal{K}) = 2$.

We have

$$\mathcal{K} \cong \mathcal{H}om(\mathcal{I}_S \omega_C, \omega_C).$$

Indeed consider a basis $\{x_0, x_1\}$ for $H^0(C, \mathcal{I}_S \omega_C)$ and define the map

$$\begin{aligned} \iota: \mathcal{H}om(\mathcal{I}_S \omega_C, \omega_C) &\rightarrow H^0(C, \mathcal{I}_S \omega_C) \otimes \omega_C \\ \varphi &\mapsto x_0 \otimes \varphi(x_1) - x_1 \otimes \varphi(x_0). \end{aligned}$$

Our aim is to check that ι is injective and $\iota(\mathcal{H}om(\mathcal{I}_S \omega_C, \omega_C))$ is precisely \mathcal{K} . It is clear that $\text{Im}(\iota) \subset \mathcal{K}$. Moreover ι is injective since the sheaf $\mathcal{I}_S \omega_C$ is generated by its sections x_0 and x_1 . It is straightforward to check that over the points $P \in C$ not belonging to S (where both the sheaves $\mathcal{H}om(\mathcal{I}_S \omega_C, \omega_C)$ and \mathcal{K} are line bundles), ι induces an isomorphism. Moreover computing the Euler characteristic we have

$$\chi(\mathcal{H}om(\mathcal{I}_S \omega_C, \omega_C)) = \chi(\mathcal{K}) = \deg S - (p_a(C) - 1)$$

hence the map ι induces an isomorphism between $\mathcal{H}om(\mathcal{I}_S \omega_C, \omega_C)$ and \mathcal{K} .

We know that

$$H^1(C, \mathcal{K})^* = \text{Hom}(\mathcal{K}, \omega_C) = H^0(C, \mathcal{H}om(\mathcal{K}, \omega_C))$$

It is easy to check that $\mathcal{I}_S \omega_C$ is reflexive, i.e.,

$$\mathcal{H}om(\mathcal{H}om(\mathcal{I}_S \omega_C, \omega_C), \omega_C) \cong \mathcal{I}_S \omega_C$$

thus $h^1(C, \mathcal{K}) = h^0(C, \mathcal{I}_S \omega_C) = 2$. ■

We may now prove our main theorem.

Theorem 3.3 *Let C be a Gorenstein curve which is either reduced or contained in a smooth algebraic surface. Assume that C is 3-connected, not honestly hyperelliptic and $p_a(C) \geq 3$. Then the map*

$$\text{Sym}^n H^0(C, \omega_C) \rightarrow H^0(C, \omega_C^{\otimes n})$$

is surjective for every $n \geq 0$.

Proof. It is already known that the canonical ring $R(C, \omega_C) = \bigoplus_{n \geq 0} H^0(C, \omega_C^{\otimes n})$ is generated in degree at most 2: see Konno [13, Prop. 1.3.3] and Franciosi [7, Th. C]. Notice that, even though both papers deal with the case of divisors on smooth surfaces, their proofs go through without changes to reduced Gorenstein curves. Thus to prove the theorem it is sufficient to show that the map in degree 2 is surjective:

$$H^0(C, \omega_C) \otimes H^0(C, \omega_C) \rightarrow H^0(C, \omega_C^{\otimes 2}).$$

We consider a free pencil $\mathcal{I}_S \omega_C$ (as in Lemma 3.1) and study the following commutative diagram:

$$\begin{array}{ccccc} H^0(C, \mathcal{I}_S \omega_C) \otimes H^0(C, \omega_C) & \hookrightarrow & H^0(C, \omega_C)^{\otimes 2} & \longrightarrow & H^0(S, \mathcal{O}_S) \otimes H^0(C, \omega_C) \\ \downarrow m & & \downarrow r & \searrow p & \downarrow \\ H^0(C, \mathcal{I}_S \omega_C^{\otimes 2}) & \hookrightarrow & H^0(C, \omega_C^{\otimes 2}) & \longrightarrow & H^0(S, \mathcal{O}_S) \end{array}$$

A simple diagram chase shows that if both the maps m and p are surjective, then the product map r is surjective too, proving Noether's theorem. Lemma 3.2 states precisely that the map m is surjective.

The map p must be surjective too: if not, we could apply Lemma 2.12 and conclude that there exists a nonempty subcanonical cluster $S_0 \subseteq S$, contained in a generic section in $H^0(C, \omega_C)$, such that

$$h^0(C, \mathcal{I}_{S_0} \omega_C) + h^1(C, \mathcal{I}_{S_0} \omega_C) \geq p_a(C) + 1.$$

By Theorem 2.7 and Remark 2.8 we know that this can not happen if C is 3-connected and not honestly hyperelliptic and $S_0 \neq \emptyset, K_C$. \blacksquare

Theorem 3.3 does not hold for every curve with very ample canonical sheaf, but only for the 3-connected ones. Indeed, Noether's Theorem may be false for canonical, non reduced and 3-disconnected curves, as shown in the following example.

Example 3.4 Let B be a smooth genus b curve with $b \geq 4$ and let D be a general effective divisor on B of degree $b + 3$. The linear system $|D|$ is very ample and induces an embedding of C in \mathbb{P}^3 (see [1, Ex. V.B.1]).

Define the ruled surface $X = \mathbb{P}_B(\mathcal{O}_B \oplus \mathcal{O}_B(D - K_B))$: the map $f: X \rightarrow B$ has a section Γ with selfintersection $(-b + 5)$ (see [12, §V.2] for the main numerical

properties). Consider the curve $C = 2\Gamma$: we have that $p_a(C) = b + 4$ and C is 2-disconnected (numerically disconnected if $b \geq 5$). By adjunction we have

$$K_C = (K_X + C)|_C = f^*(D)$$

and it is easy to check that it is very ample on C by analyzing the standard decomposition

$$0 \rightarrow \omega_\Gamma \rightarrow \omega_C \rightarrow \omega_{C|\Gamma} \rightarrow 0.$$

Since Γ is a section of $f: X \rightarrow B$, F induces an isomorphism $(B, \mathcal{O}_B(D)) \cong (\Gamma, \omega_{C|\Gamma})$. Therefore it is immediately seen that $|\omega_C|$ separates length 2 clusters.

The map

$$\mathrm{Sym}^2 H^0(C, \omega_C) \xrightarrow{q_0} H^0(C, \omega_C^{\otimes 2})$$

is not surjective, as one could see from the following diagram:

$$\begin{array}{ccccc} H^0(\Gamma, \omega_\Gamma) \otimes H^0(C, \omega_C) & \longrightarrow & H^0(C, \omega_C)^{\otimes 2} & \longrightarrow & H^0(\Gamma, \omega_{C|\Gamma}) \otimes H^0(\Gamma, \omega_{C|\Gamma}) \\ \downarrow & & \downarrow q & & \downarrow p \\ H^0(\Gamma, \omega_\Gamma \otimes \omega_C) & \longrightarrow & H^0(C, \omega_C^{\otimes 2}) & \longrightarrow & H^0(\Gamma, \omega_{C|\Gamma}^{\otimes 2}) \end{array}$$

Indeed if the map q_0 was surjective (hence q), then the map p would be surjective as well. Since $\omega_{C|\Gamma} \cong \mathcal{O}_B(D)$ the image of the map p is the same as the image of the map

$$p_0: \mathrm{Sym}^2 H^0(B, \mathcal{O}_B(D)) \rightarrow H^0(B, \mathcal{O}_B(2D))$$

which is not surjective, as one can easily check by computing the dimension of the two spaces.

4 Castelnuovo's Theorem for reduced curves

In this section we prove Theorem 4.2, a generalization of Castelnuovo's Theorem for reduced curves.

In the proof we will apply Lemma 2.12 and the following Proposition, which is a Clifford-type result for line bundles of high degree.

Proposition 4.1 *Let C be a Gorenstein reduced curve and L a line bundle on C such that*

$$\deg L|_B \geq 2p_a(B) + 1 \quad \text{for every } B \subset C.$$

If S is a cluster contained in a generic section $H \in H^0(C, L)$ then

$$h^0(C, \mathcal{I}_S L) + h^1(C, \mathcal{I}_S L) < h^0(C, L).$$

Proof. Notice at first that $H^1(C, L) = 0$ and $|L|$ is very ample by Proposition 2.5. Therefore a generic hyperplane section consists of $\deg L$ smooth points. Moreover by Riemann Roch Theorem we have

$$\begin{aligned} h^0(C, \mathcal{I}_S L) + h^1(C, \mathcal{I}_S L) < h^0(C, L) &\iff h^0(C, \mathcal{I}_S L) < h^0(C, L) - \frac{1}{2} \deg S \\ &\iff h^1(C, \mathcal{I}_S L) < \frac{1}{2} \deg S \end{aligned}$$

We argue by induction on the number of irreducible components of C . Suppose that C is irreducible or that the statement holds for every reduced curve with fewer components of C . If C is disconnected the statement is trivial, hence we may assume that C is connected. If $h^0(C, \mathcal{I}_S L) = 0$ or $h^1(C, \mathcal{I}_S L) = 0$ the result is trivial too, thus we may assume $h^0(C, \mathcal{I}_S L) > 0$ and $h^1(C, \mathcal{I}_S L) > 0$.

Suppose at first that there exists a proper subcurve $B \subset C$ such that

$$H^1(C, \mathcal{I}_S L) \cong H^1(B, \mathcal{I}_S L|_B).$$

By induction we have that $h^1(C, \mathcal{I}_S L) < \frac{1}{2} \deg S|_B \leq \frac{1}{2} \deg S$ and we may conclude.

If there is no subcurve $B \subset C$ as above (e.g. when C is irreducible) we can easily deduce from Proposition 2.4 that there exists a generically surjective map $\mathcal{I}_S L \hookrightarrow \omega_C$, hence we may assume that there exists a subcanonical cluster Z such that

$$\mathcal{I}_S L \cong \mathcal{I}_Z \omega_C.$$

Since S is contained in a generic section of $H^0(C, L)$ it is a Cartier divisor, hence Z is a Cartier divisor too.

If C is 2-connected we apply Theorem 2.7 and we conclude since

$$\begin{aligned} h^0(C, \mathcal{I}_C L) &= h^0(C, \mathcal{I}_Z \omega_C) \leq p_a(C) - \frac{1}{2} \deg Z \\ &= \frac{1}{2} \deg L + 1 - \frac{1}{2} \deg S < h^0(C, L) - \frac{1}{2} \deg S. \end{aligned}$$

If C is 2-disconnected, we can find a decomposition $C = C_1 + C_2$, such that $C_1 \cdot C_2 = 1$ and C_1 is 2-connected (see [6, Lemma A.4]). Thus we consider the following exact sequence:

$$0 \rightarrow \mathcal{I}_Z|_{C_1} \omega_{C_1} \rightarrow \mathcal{I}_Z \omega_C \rightarrow (\mathcal{I}_Z \omega_C)|_{C_2} \rightarrow 0.$$

We know by induction that

$$\begin{aligned} h^0(C_2, (\mathcal{I}_Z \omega_C)|_{C_2}) &= h^0(C, (\mathcal{I}_S L)|_{C_2}) < h^0(C_2, L|_{C_2}) - \frac{1}{2} \deg S|_{C_2} \\ &= -p_a(C_2) + 1 + \deg L|_{C_2} - \frac{1}{2} \deg S|_{C_2}. \end{aligned}$$

We apply Theorem 2.7 to $\mathcal{I}_{Z|C_1}\omega_{C_1}$ since the single point $C_1 \cap C_2$ is a base point for $|\omega_C|$, hence the space $H^0(C_1, \mathcal{I}_{Z|C_1}\omega_{C_1}) \cong H^0(C_1, \mathcal{I}_{Z|C_1}\omega_{C_1|C_1})$ contains an invertible section, that is $Z|_{C_1}$ is a subcanonical cluster. Thus

$$h^0(C_1, \mathcal{I}_{Z|C_1}\omega_{C_1}) \leq p_a(C_1) - \frac{1}{2} \deg Z|_{C_1} = \frac{1}{2} \deg L|_{C_1} - \frac{1}{2} \deg S|_{C_1} + \frac{1}{2}.$$

and we conclude since

$$\begin{aligned} h^0(C, \mathcal{I}_S L) &= h^0(C, \mathcal{I}_Z \omega_C) \leq h^0(C_2, (\mathcal{I}_Z \omega_C)|_{C_2}) + h^0(C_1, \mathcal{I}_{Z|C_1} \omega_{C_1}) \\ &< -p_a(C_2) + 1 + \deg L|_{C_2} - \frac{1}{2} \deg S|_{C_2} + \frac{1}{2} \deg L|_{C_1} - \frac{1}{2} \deg S|_{C_1} + \frac{1}{2} \\ &\leq h^0(C, L) - \frac{1}{2} \deg S. \end{aligned}$$

■

Theorem 4.2 *Let C be a Gorenstein reduced curve and let L be a line bundle on C such that*

$$\deg L|_B \geq 2p_a(B) + 1 \quad \text{for every } B \subset C.$$

Then the product map

$$\text{Sym}^n H^0(C, L) \rightarrow H^0(C, L^{\otimes n})$$

is surjective for every $n \geq 1$.

Proof. Notice at first that $H^1(C, L) = 0$ and L is very ample by Proposition 2.5.

If $n \geq 2$ the map

$$H^0(C, L^{\otimes n}) \otimes H^0(C, L) \rightarrow H^0(C, L^{\otimes(n+1)})$$

is surjective by [7, Prop. 1.5] since $H^1(C, L^{\otimes n} \otimes L^{-1}) = 0$. In order to prove the theorem we check that the map in degree 2 is surjective:

$$\text{Sym}^2 H^0(C, L) \rightarrow H^0(C, L^{\otimes 2}).$$

To this aim we consider a generic hyperplane section $S = \text{div} L$ and the following commutative diagram

$$\begin{array}{ccccc} H^0(C, \mathcal{I}_S L) \otimes H^0(C, L) & \hookrightarrow & H^0(C, L) \otimes H^0(C, L) & \longrightarrow & H^0(S, \mathcal{O}_S) \otimes H^0(C, L) \\ \downarrow & & \downarrow r & \searrow p & \downarrow \\ H^0(C, \mathcal{I}_S L^{\otimes 2}) & \hookrightarrow & H^0(C, L^{\otimes 2}) & \longrightarrow & H^0(S, \mathcal{O}_S) \end{array}$$

Notice that the first column is surjective since $\mathcal{I}_S L \cong \mathcal{O}_C$ while the second row is exact since $H^1(C, \mathcal{I}_S L^{\otimes 2}) \cong H^1(C, L) = 0$. A simple diagram chase shows that the map r is surjective if and only if the map p is surjective.

It is $h^0(C, \mathcal{I}_{S_0} L) + h^1(C, \mathcal{I}_{S_0} L) < h^0(C, L)$ for every subcluster $S_0 \subseteq S$ by Proposition 4.1, hence the map p must be surjective by Lemma 2.12. ■

Remark 4.3 If C is numerically connected our result implies that the embedded curve $\varphi_L(C) \subset \mathbb{P}H^0(C, L)^*$ is arithmetically Cohen-Macaulay.

References

- [1] E. Arbarello, M. Cornalba, P. A. Griffiths, J. Harris, *Geometry of Algebraic Curves, vol.1*, Springer, New York - Berlin - Heidelberg - Tokyo, 1985, 386 pp.
- [2] M. Aprodu, G. Farkas, Koszul Cohomology and Application to Moduli, in *Grassmannians, Moduli Spaces and Vector Bundles*, Clay Mathematics Proceedings, D. Ellwood and E. Previato editors, Vol. 14, 2011, 25–50.
- [3] F. Catanese, C. Ciliberto, *Surfaces with $p_g = q = 1$* , Problems in the theory of surfaces and their classification (Cortona, 1988), Sympos. Math. **XXXII**, Academic Press, London, 1991, 49–79.
- [4] F. Catanese, M. Franciosi, *Divisors of small genus on algebraic surfaces and projective embeddings*, Proc. of the 1993 Hirzebruch 65 Conference on Alg. Geom., Israel Math. Conf. Proc. **9** (1996), *Contemp. Math.*, AMS, 109–140.
- [5] F. Catanese, M. Franciosi, K. Hulek, M. Reid, *Embedding of curves and surfaces*, Nagoya Math. J. **154** (1999), pp. 185–220.
- [6] C. Ciliberto, P. Francia, M. Mendes Lopes, Remarks on the bicanonical map for surfaces of general type, *Math. Zeit.* **224** (1997), 137–166.
- [7] M. Franciosi, *Adjoint divisors on algebraic curves*, Adv. in Math. **186** (2004), 317–333.
- [8] M. Franciosi, *On the canonical ring of curves and surfaces*, arXiv: 1107.0579 [math.AG] (2011).
- [9] M. Franciosi, E. Tenni, *On Clifford's theorem for singular curves*, arXiv:1105.2253v1 [math.AG] (2011).
- [10] M. Green, *Koszul cohomology and the geometry of projective varieties*, J. Diff. Geom. **19** (1984), 125–171.

- [11] M. Green, L. Lazarsfeld, *On the projective normality of complete linear series on an algebraic curve*, Invent. Math. **83** (1986), 73–90.
- [12] R. Hartshorne, *Algebraic geometry*. Graduate Texts in Mathematics, No. 52. Springer-Verlag, New York-Heidelberg, 1977, xvi+496 pp.
- [13] K. Konno, *1-2-3 theorem for curves on algebraic surfaces*, J. reine angew. Math. **533** (2001), 171–205.
- [14] D. Mumford, *Varieties defined by quadratic equations*, in 'Questions on algebraic varieties', C.I.M.E., III Ciclo, Varenna, 1969, Ed. Cremonese, Rome, (1970), 30–100.
- [15] M. Reid, *Problems on pencils of low genus*, manuscript (1990), available at <http://www.warwick.ac.uk/masda/surf/more/atoms.pdf>

Marco Franciosi
Dipartimento di Matematica Applicata “U. Dini”, Università di Pisa
Via Buonarroti 1, I-56127 Pisa (Italy)
`franciosi@dma.unipi.it`

Elisa Tenni
Dipartimento di Matematica “L. Tonelli”, Università di Pisa
Largo Bruno Pontecorvo 5, I-56127 Pisa (Italy)
`tenni@mail.dm.unipi.it`