

# Electrostatic potentials at the nucleus for isoelectronic series of light atomic ions using the QMC method in relation to DFT

C. Amovilli<sup>a\*</sup> and N. H. March<sup>b,c,d</sup>

<sup>a</sup>*Dipartimento di Chimica e Chimica Industriale,  
Università di Pisa, Via G. Moruzzi 3, 56124 Pisa, Italy*

<sup>b</sup>*Abdus Salam ICTP, Trieste, Italy*

<sup>c</sup>*Department of Physics, University of Antwerp,  
Groenenborgerlaan 171, B-2020 Antwerp, Belgium*

<sup>d</sup>*Oxford University, Oxford, England*

February 4, 2015

## Abstract

The electrostatic potential at the nucleus is here calculated using the quantum Monte Carlo method. Both variational and diffusion-type data are presented for four different isoelectronic series of atomic ions, namely He, Li, Be and B. These results are then utilized to evaluate the ground-state energies of such atomic ions.

Keywords: Quantum Monte Carlo; Electrostatic potential at nuclei; He, Li, Be and B ionic series.

---

\*Corresponding author. e-mail:claudio.amovilli@unipi.it

# 1 Introduction

Early work of Foldy [1] was important in emphasizing an intimate correlation between the ground-state energy of neutral atoms of nuclear charge  $Ze$ , namely  $E(Z)$ , and the electrostatic potential created by the electrons, denoted as  $V_0$  below, evaluated at the assumed point nucleus. Before Foldy's study, such a correlation could be extracted analytically from the Thomas-Fermi (TF) semi-classical theory, beginning with the ground-state energy  $E(Z)$  of neutral atoms having nuclear charge  $Ze$  derived by Milne [2]. His result was explicitly

$$E(Z) = -0.7687Z^{7/3}\frac{e^2}{a_0} \quad : a_0 = \frac{\hbar^2}{me^2} . \quad (1)$$

The electron-nuclear potential energy  $U_{en}$  in the TF model is also known (see, for example, [3]) to be given by

$$U_{en}(Z) = \frac{7}{3}E(Z) . \quad (2)$$

But on physical grounds,  $U_{en}(Z)$  is simply the interaction energy of the charge  $Ze$  sitting in a potential  $V_0$ , ie

$$U_{en}(Z) = ZeV_0 . \quad (3)$$

Hence, using Eq.(3) in Eq.(2), the desired correlation is expressed by

$$V_0(Z) = \frac{7}{3}\frac{E(Z)}{Ze} = -1.79Z^{4/3}\frac{e}{a_0} \quad (4)$$

which is the precise semi-classical (non-relativistic) prediction for neutral atoms. Unfortunately, the self-consistent TF ground-state density  $n_{TF}(r)$  has two truly major difficulties: (i) it is infinite at the assumed point nucleus, being singular as  $r^{-3/2}$ , and (ii) the decay at large  $r$  is not exponential but has the much longer ranged  $r^{-6}$  form.

Politzer and Parr [4] proposed a generalization of Eq.(4) to read (see also Politzer chapter in Ref. [3] and Politzer et al [5])

$$E(Z, N) = \int_0^Z V_0(Z') dZ' . \quad (5)$$

However, it is now known that as  $Z'$  is varied through an isoelectronic series of atomic ions with  $N$  electrons, there is a critical (non-integral) atomic number, say  $Z_c(N)$ , at which, on further infinitesimal reduction, one electron ionizes, sometimes referred to, somewhat loosely, as a 'phase transition'. Hence, we shall rewrite Eq.(5) in our present study as

$$E(Z, N) = E(Z_c, N) + \int_{Z_c}^Z V_0(Z') dZ' . \quad (6)$$

Obviously  $V_0$  to be inserted in Eq.(6) is the central tool of the present work and this is naturally to be determined from the ground-state electron density  $n(Z, N; \mathbf{r})$  through the isoelectronic series of atomic ions with  $N$  electrons.

It is relevant at this point to emphasize that, at least in principle,  $E(Z, N)$  is determined solely by the ground-state electron density  $n(Z, N; \mathbf{r})$ , through the Hohenberg-Kohn theorem [6]. Unfortunately, the functional  $E[n(\mathbf{r})]$  remains unknown, and we bypass this fact here by invoking quantum Monte Carlo (QMC) calculations for the appropriate range of  $Z'$  to very accurately evaluate numerically the integral appearing in Eq.(6).

With this as background, we turn immediately to present the QMC results to insert into Eq.(6).

## 2 Computational details

As the variational Monte Carlo (VMC) wave function for the atomic ions, we employed a spin-free Slater-Jastrow form of the type

$$\Psi(\mathbf{r}_1, \mathbf{r}_2, \dots) = \Phi(\mathbf{r}_1, \mathbf{r}_2, \dots) J(r_1, r_2, \dots, r_{12}, \dots) . \quad (7)$$

The determinantal component is given by

$$\Phi = \sum_k D_k^\uparrow D_k^\downarrow d_k \quad (8)$$

where  $D_k^\uparrow$  and  $D_k^\downarrow$  are the Slater determinants written in terms of occupied orbitals of spin-up and spin-down electrons, respectively, and  $d_k$  are the mixing coefficients. The Jastrow correlation factor is the exponential of the sum

of three contributions which are functions of electron-nuclear (e-n), electron-electron (e-e), and pure three-body mixed e-e and e-n distances[7]. In all cases, we used a many determinant wave function with a flexible Slater-type basis set. We optimized all parameters at the VMC level in energy minimization and we performed the diffusion Monte Carlo (DMC) calculation with a time step of 0.05 a.u.

### 3 Discussion of results

We record in Tables 1-4 results for the ground-state energy  $E$  and for  $V_0$  from QMC calculations. Both VMC and DMC are here utilized, for  $N = 2$  to 5 (He- to B-like isoelectronic series of atomic ions).  $V_0$  values for the neutral atoms from VMC and DMC are in pretty good agreement. In particular for the B neutral atom  $V_0$  values are  $-11.41(1)$  and  $-11.39(1)$  Hartree for VMC and DMC respectively. The electron-nuclear potential energies  $U_{en}$  corresponding to these values follow from Eq.(3) for the neutral B atom as  $-57.07$  and  $-56.95$  Hartree.  $U_{en}/E = 2.316$  and  $2.310$ , these ratios being somewhat smaller than the TF prediction of  $7/3$  in Eq.(2). In the customary language of DFT, the ground-state energy  $E$  and the universal density functional  $F$ , on the minimum, are related by

$$E = F + U_{en} . \tag{9}$$

Hence for VMC, for neutral B,  $F = 32.43$  Hartree compared with  $F = 32.30$  Hartree for DMC. We anticipate that the DMC value for  $F$  will be somewhat more accurate than that from the variational approach. If  $T$  denotes the total kinetic energy including correlation kinetic energy, then the virial theorem would give as a useful approximation  $T = -E$ , in each case.

As to the ground-state density at the nucleus,  $n_0$  say, this is not an easy quantity to calculate accurately from QMC, though it is, in essence, an observable via the Mössbauer effect. However, Figure 1 shows our rough estimates for  $n_0$  from QMC. Apart from statistical error, our values are in agreement with literature data for neutral atoms [8, 9, 10, 11].

In Table 5 we show, for comparison, the separate magnitudes of the two terms on the right-hand-side (RHS) of Eq.(6) together with the total energy  $E(Z, Z)$  for neutral atoms. It is important here to make contact with the fairly recent work of Politzer et al [5]. These authors, however, did not integrate  $V_0(Z', N)$  from the critical value  $Z_c(N)$  to  $Z$ , but from a chosen value of  $Z'$ , namely  $N - 1$ . It was the existence of  $Z_c$ , at which  $V_0$  is continuous, but non-analytic, which prompted us to separate into the form of Eq.(6). Politzer et al analyzed a larger interval of nuclear charges. For the four neutral atoms of the present study, their results show an energy which is about 0.2 Hartree lower than the exact one. In our case, as shown in Table 5, the discrepancy is much lower, ranging from 0.0002 Hartree for He to 0.01 Hartree for B. It should be noted that they used the B3PW91 functional to compute  $V_0$  while here we use QMC.

It is worthy of note then that one could view Eq.(6) as a formally exact summation of the  $1/Z$  expansion[12] to all orders for the difference energy  $E(Z, N) - E(Z_c, N)$ . Unfortunately, so far no known exact analytic formula exists for  $Z_c(N)$ , though Cordero et al [13] give an empirical fit over a restricted range of  $N$ . Also the inequality (see, for example, [14])

$$N - 1 \geq Z_c(N) \geq N - 2 \tag{10}$$

is well established.

In summary, the main achievement of the present work is to exhibit extensive QMC calculations for four isoelectronic series of light elements. These are finally utilized to determine the ground-state energies  $E(Z, N)$  of such ions from Eq.(6).

### Acknowledgments

CA wishes to acknowledge partial support from MIUR (PRIN 2009). NHM began his contribution to the present study during a visit to the Chemistry Department of the University of Pisa. Thanks are due to Prof. C. Amovilli for generous hospitality. The work of NHM on the final form was brought to fruition during a subsequent visit to Abdus Salam ICTP, Trieste. Thanks are due to Prof. V. E. Kravtsov, for much stimulation and encouragement.

Finally, NHM is grateful to Profs. D. Lamoen and C. Van Alsenoy for making his continuing affiliation with the University of Antwerp possible.

## References

- [1] L. L. Foldy, *Phys. Rev.* **83**, 397 (1951).
- [2] E. A. Milne, *Proc. Camb. Phil. Soc.* **23**, 794 (1927).
- [3] N. H. March and B. M. Deb Editors, *The Single Particle Density in Physics and Chemistry*, (Academic Press, New York, 1987).
- [4] P. Politzer and R. G. Parr, *J. Chem. Phys.* **61**, 4258 (1974).
- [5] P. Politzer, Y. Ma, A. F. Jalbot and J. S. Murray, *Mol. Phys.* **103**, 2105 (2005).
- [6] P. Hohenberg and W. Kohn, *Phys. Rev.* **136**, 864 (1964).
- [7] C. Filippi and C. Umrigar, *J. Chem. Phys.* **105**, 213 (1996).
- [8] M. Hoffmann-Ostenhof, T. Hoffmann-Ostenhof and W. Thirring, *J. Phys. B: Atom. Molec. Opt. Phys.* **11**, L571 (1979).
- [9] J. Cioslowski, *Phys. Rev. A* **39**, 378 (1989).
- [10] C. Chen, *Eur. Phys. J. D* **56**, 303 (2010).
- [11] C. Nazé, S. Verdebout, P. Rynkun, G. Gaigalas, M. Godefroid and P. Jönsson, *arXiv*, 1406.1720v1 (2014).
- [12] D. Layzer, *Ann. Phys. NY* **8**, 271 (1959).
- [13] N. A. Cordero, N. H. March and J. A. Alonso, *Phys. Lett. A* **377**, 2955 (2013).
- [14] H. Hogreve, *J. Phys. B: Atom. Molec. Opt. Phys.* **31**, L439 (1998).

$Z$	$E(VMC)$	err	$V_0(VMC)$	err	$E(DMC)$	err	$ZV_0(DMC)$	err
0.88	-0.38517	0.00001	-0.8925	0.001	-0.38548	0.00001	-0.7867	0.002
0.89	-0.39398	0.00001	-0.9056	0.001	-0.39452	0.00001	-0.7991	0.002
0.90	-0.40306	0.00001	-0.9146	0.001	-0.40347	0.00001	-0.8213	0.002
0.905	-0.40784	0.00001	-0.9178	0.001	-0.40800	0.00001	-0.8270	0.002
0.907	-0.41029	0.00001	-1.1545	0.002	-0.41042	0.00002	-1.0189	0.002
0.91	-0.41378	0.00001	-1.1563	0.002	-0.41380	0.00002	-1.0349	0.002
0.911	-0.41495	0.00001	-1.1579	0.002	-0.41494	0.00002	-1.0427	0.002
0.9113	-0.41528	0.00001	-1.1589	0.001	-0.41524	0.00002	-1.0533	0.002
0.9115	-0.41559	0.00001	-1.1553	0.001	-0.41550	0.00002	-1.0452	0.002
0.912	-0.41598	0.00001	-1.1609	0.001	-0.41611	0.00002	-1.0614	0.002
0.915	-0.41957	0.00001	-1.1667	0.001	-0.41963	0.00002	-1.0756	0.002
0.92	-0.42547	0.00001	-1.1853	0.001	-0.42550	0.00002	-1.0845	0.002
0.93	-0.43743	0.00001	-1.2097	0.001	-0.43744	0.00002	-1.1244	0.002
0.96	-0.47479	0.00001	-1.2827	0.001	-0.47478	0.00002	-1.2307	0.002
1.00	-0.52773	0.00001	-1.3602	0.001	-0.52771	0.00002	-1.3655	0.002
1.10	-0.67475	0.00001	-1.5713	0.001	-0.67476	0.00002	-1.7379	0.002
1.30	-1.02980	0.00001	-1.9737	0.001	-1.02989	0.00002	-2.5669	0.003
1.50	-1.46519	0.00001	-2.3765	0.002	-1.46534	0.00002	-3.5699	0.004
1.70	-1.98061	0.00001	-2.7767	0.002	-1.98074	0.00002	-4.7168	0.005
1.95	-2.73736	0.00001	-3.2749	0.003	-2.73742	0.00002	-6.4033	0.006
1.98	-2.83657	0.00001	-3.3334	0.003	-2.83657	0.00002	-6.6097	0.007
2.00	-2.90364	0.00001	-3.3776	0.003	-2.90373	0.00004	-6.7484	0.007
2.01	-2.93757	0.00001	-3.4024	0.003	-2.93757	0.00003	-6.8330	0.007
2.03	-3.00590	0.00001	-3.4328	0.003	-3.00588	0.00003	-6.9968	0.007
2.05	-3.07503	0.00001	-3.4745	0.003	-3.07509	0.00003	-7.1379	0.007
2.10	-3.25133	0.00001	-3.5759	0.003	-3.25133	0.00004	-7.5318	0.008
2.30	-4.00664	0.00002	-3.9701	0.003	-4.00674	0.00003	-9.1471	0.008
2.65	-5.52082	0.00002	-4.6765	0.003	-5.52087	0.00003	-12.3791	0.009
3.00	-7.27989	0.00001	-5.3803	0.004	-7.27990	0.00003	-16.1282	0.0010
3.50	-10.21775	0.00001	-6.3776	0.005	-10.21780	0.00004	-22.3467	0.0017
4.00	-13.65555	0.00002	-7.3775	0.006	-13.65562	0.00005	-29.5164	0.0020
5.00	-22.03087	0.00003	-9.3803	0.007	-22.03134	0.00010	-46.8642	0.0027

Table 1: Quantum Monte Carlo results for the electrostatic potential at the nucleus for the He-like series of atomic ions for some thirty values (mostly non-integral) of nuclear charge  $Ze$ .

$Z$	$E(VMC)$	err	$V_0(VMC)$	err	$E(DMC)$	err	$ZV_0(DMC)$	err
1.95	-2.73407	0.00001	-3.3204	0.002	-2.73597	0.00001	-6.4446	0.002
2.01	-2.93414	0.00004	-3.4207	0.002	-2.93757	0.00001	-6.8702	0.002
2.05	-3.07421	0.00003	-3.5127	0.002	-3.07612	0.00001	-7.2125	0.002
2.10	-3.25455	0.00002	-3.6534	0.002	-3.25508	0.00001	-7.6669	0.002
2.30	-4.03046	0.00004	-4.1283	0.003	-4.03236	0.00006	-9.5009	0.003
2.50	-4.90202	0.00003	-4.5836	0.003	-4.90269	0.00003	-11.4711	0.002
2.70	-5.86398	0.00003	-5.0385	0.003	-5.86469	0.00002	-13.6113	0.002
2.81	-6.43108	0.00004	-5.2894	0.004	-6.43253	0.00002	-14.8618	0.002
2.84	-6.59192	0.00002	-5.3555	0.004	-6.59216	0.00002	-15.2186	0.002
2.845	-6.61868	0.00002	-5.3647	0.004	-6.61900	0.00002	-15.2732	0.002
2.85	-6.64559	0.00002	-5.3799	0.004	-6.64584	0.00002	-15.3327	0.002
2.855	-6.67250	0.00002	-5.3917	0.004	-6.67276	0.00002	-15.3879	0.002
2.86	-6.69947	0.00002	-5.3918	0.004	-6.69974	0.00002	-15.4451	0.002
3.00	-7.47779	0.00002	-5.7200	0.004	-7.47810	0.00002	-17.1661	0.003
3.50	-10.61908	0.00002	-6.8491	0.005	-10.61971	0.00003	-23.9474	0.004
4.00	-14.32433	0.00004	-7.9774	0.005	-14.32527	0.00010	-31.8723	0.0015
5.00	-23.42402	0.00006	-10.2136	0.007	-23.42581	0.00016	-51.1051	0.0020

Table 2: QMC results for Li-like isoelectronic series of atomic ions for about 20 values of fractional nuclear charge  $Ze$ .



$Z$	$E(VMC)$	err	$V_0(VMC)$	err	$E(DMC)$	err	$ZV_0(DMC)$	err
2.81	-6.42351	0.00003	-5.3520	0.004	-6.42963	0.00007	-14.9583	0.005
2.84	-6.58742	0.00002	-5.4496	0.004	-6.59033	0.00006	-15.5090	0.005
2.845	-6.61478	0.00003	-5.4832	0.006	-6.61785	0.00006	-15.5917	0.005
2.85	-6.64236	0.00004	-5.5108	0.004	-6.64480	0.00005	-15.7154	0.005
2.855	-6.67001	0.00003	-5.5267	0.004	-6.67227	0.00005	-15.7704	0.005
2.86	-6.69905	0.00003	-5.5644	0.004	-6.70082	0.00006	-15.8568	0.005
2.90	-6.91779	0.00005	-5.6429	0.004	-6.92404	0.00008	-16.3710	0.005
3.00	-7.49685	0.00004	-5.9034	0.004	-7.50053	0.00007	-17.6939	0.006
3.50	-10.76575	0.00005	-7.1637	0.005	-10.76827	0.00008	-25.0570	0.007
3.95	-14.24689	0.00006	-8.3000	0.006	-14.25006	0.00011	-32.7569	0.009
4.00	-14.66461	0.00007	-8.4101	0.006	-14.66791	0.00011	-33.6570	0.009
4.05	-15.08962	0.00007	-8.5547	0.006	-15.09286	0.00011	-34.5678	0.009
4.10	-15.52029	0.00007	-8.6760	0.006	-15.52382	0.00012	-35.5470	0.010
4.30	-17.30648	0.00008	-9.1839	0.006	-17.30963	0.00012	-39.4211	0.010
4.50	-19.19290	0.00008	-9.6728	0.006	-19.19638	0.00012	-43.5149	0.011
4.70	-21.17892	0.00009	-10.1850	0.007	-21.18294	0.00013	-47.8008	0.012
5.00	-24.34664	0.00007	-10.9277	0.007	-24.35036	0.00013	-54.6127	0.014

Table 3: QMC results for Be-like isoelectronic series of atomic ions for some fifteen values of fractional nuclear charge  $Ze$ .

$Z$	$E(VMC)$	err	$V_0(VMC)$	err	$E(DMC)$	err	$ZV_0(DMC)$	err
3.90	-13.83248	0.00006	-8.1922	0.005	-13.83736	0.00009	-31.8800	0.007
3.95	-14.24454	0.00006	-8.3012	0.005	-14.24887	0.00008	-32.7797	0.007
4.05	-15.08983	0.00008	-8.5864	0.006	-15.09302	0.00008	-34.7415	0.010
4.10	-15.52244	0.00006	-8.6979	0.006	-15.52767	0.00017	-36.1269	0.010
4.30	-17.34745	0.00007	-9.4313	0.006	-17.35668	0.00015	-40.5444	0.010
4.70	-21.34813	0.00008	-10.566	0.007	-21.35827	0.00018	-49.5728	0.012
5.00	-24.64189	0.00008	-11.414	0.007	-24.65164	0.00018	-56.9498	0.013

Table 4: QMC results for B-like isoelectronic series of atomic ions for seven values of fractional nuclear charge  $Ze$ .

$Z = N$	$Z_c$	$E_c(Z_c, N)$	$\int V_0$	$E(Z, Z)$	$E(DMC)$	ratio
2	0.911	-0.41494(2)	-2.4890	-2.9039	-2.90372	0.1667
3	2.000	-2.90372(4)	-4.5801	-7.4838	-7.47810	0.6340
4	2.856	-6.68112(5)	-7.9819	-14.6630	-14.66791	0.8370
5	4.000	-14.6679(10)	-9.9727	-24.6406	-24.65164	1.4708

Table 5: Displays magnitudes of the two terms on the RHS of Eq.(6).




Figure 1: Plot of the electron density at the nucleus  $n_0[Z, N]$  as a function of the fractional nuclear charge  $Z$  for the four series of atomic ions considered in this work. Neutral atoms correspond to the case  $N = Z$ . These are the only observables.