

REMARKS ABOUT BESICOVITCH COVERING PROPERTY IN CARNOT GROUPS OF STEP 3 AND HIGHER

ENRICO LE DONNE AND SÉVERINE RIGOT

ABSTRACT. We prove that the Besicovitch Covering Property (BCP) does not hold for some classes of homogeneous quasi-distances on Carnot groups of step 3 and higher. As a special case we get that, in Carnot groups of step 3 and higher, BCP is not satisfied for those homogeneous distances whose unit ball centered at the origin coincides with a Euclidean ball centered at the origin. This result comes in contrast with the case of the Heisenberg groups where such distances satisfy BCP.

1. INTRODUCTION

Covering theorems, among which is the Besicovitch Covering Property (BCP), see Definition 1.1 below, are known to be some of the fundamental tools of measure theory. More generally they turn out to be classical tools that play a crucial role in many problems in analysis and geometry. We refer for example to [4] and [7] for a more detailed introduction about covering theorems.

In contrast to the Euclidean case, the Heisenberg groups equipped with the commonly used (Cygan-)Korányi and Carnot-Carathéodory distances are known not to satisfy BCP ([5], [8], [9]). However, it has been recently proved that BCP holds in the Heisenberg groups equipped with those homogeneous distances whose unit ball centered at the origin coincides with a Euclidean ball centered at the origin ([6], see also Theorem 1.8 below).

For more general Carnot groups, BCP does not hold for Carnot-Carathéodory distances ([8]). Motivated by the question of whether one can find homogeneous (quasi-)distances on a given Carnot group for which BCP holds, we prove in the present paper that BCP does not hold for some classes of homogeneous quasi-distances on Carnot groups of step 3 and higher, see Theorem 1.6. As a particular case, we get that the main result in [6] do not extend to Carnot groups of step 3 and higher, that is, BCP is not satisfied when these groups are equipped with a homogeneous distance whose unit ball centered at the origin coincides with a Euclidean ball centered at the origin, see Corollary 1.7.

To state our results, we first recall the Besicovitch Covering Property in the general quasi-metric setting. Given a nonempty set X , we say that $d : X \times X \rightarrow [0, +\infty[$ is a quasi-distance on X if it is symmetric, $d(p, q) = 0$ if and only if $p = q$, and there exists a constant $C \geq 1$ such that $d(p, q) \leq C(d(p, p') + d(p', q))$ for all $p, p', q \in X$ (quasi-triangle inequality

Date: May 18, 2015.

2010 Mathematics Subject Classification. 28C15, 49Q15, 43A80.

Key words and phrases. Covering theorems, Carnot groups, Homogeneous quasi-distances.

The work of S.R. is supported by the ANR-12-BS01-0014-01 Geometrya.

with multiplicative constant C). We call (X, d) a quasi-metric space. When speaking of a ball B in (X, d) , it will be understood that B is a set of the form $B = B_d(p, r)$ for some $p \in X$ and some $r > 0$ where $B_d(p, r) := \{q \in X; d(q, p) \leq r\}$. Note that when d satisfies the quasi-triangle inequality with a multiplicative constant $C = 1$, then d is nothing but a distance on X .

Definition 1.1 (Besicovitch Covering Property). Let (X, d) be a quasi-metric space. We say that (X, d) satisfies the Besicovitch Covering Property (BCP) if there exists a constant $N \in \mathbb{N}$ such that the following holds. Let A be a bounded subset of X and \mathcal{B} be a family of balls such that each point of A is the center of some ball of \mathcal{B} , then there is a finite or countable subfamily $\mathcal{F} \subset \mathcal{B}$ such that the balls in \mathcal{F} cover A and every point in X belongs to at most N balls in \mathcal{F} , that is,

$$\mathbb{1}_A \leq \sum_{B \in \mathcal{F}} \mathbb{1}_B \leq N$$

where $\mathbb{1}_A$ denotes the characteristic function of the set A .

The Besicovitch Covering Property originates from the work of Besicovitch ([1], [2]). It is satisfied in the Euclidean space and more generally in any finite dimensional normed vector space.

Next, we recall the definition of Carnot groups and state the conventions and notations we shall use throughout this paper. A Carnot group G of step $s \geq 1$ is a connected and simply connected Lie group whose Lie algebra \mathfrak{g} is endowed with a stratification, $\mathfrak{g} = V_1 \oplus \cdots \oplus V_s$ where $[V_1, V_j] = V_{j+1}$ for $1 \leq j \leq s-1$ and $[V_1, V_s] = \{0\}$.

We set $n := \dim \mathfrak{g}$ and consider (X_1, \dots, X_n) a basis of \mathfrak{g} adapted to the stratification, i.e., $(X_{m_{j-1}+1}, \dots, X_{m_j})$ is a basis of V_j for $1 \leq j \leq s$ where $m_0 := 0$ and $m_j - m_{j-1} := \dim V_j$.

We identify G with \mathbb{R}^n via a choice of exponential coordinates of the first kind. Namely, for Carnot groups, the exponential map $\exp : \mathfrak{g} \rightarrow G$ is a diffeomorphism from \mathfrak{g} to G . We then identify $p = \exp(x_1 X_1 + \cdots + x_n X_n) \in G$ with (x_1, \dots, x_n) . The group law is then given by

$$(x_1, \dots, x_n) \cdot (x'_1, \dots, x'_n) := (x''_1, \dots, x''_n)$$

where $\exp(x''_1 X_1 + \cdots + x''_n X_n) = \exp(x_1 X_1 + \cdots + x_n X_n) \cdot \exp(x'_1 X_1 + \cdots + x'_n X_n)$. The coordinates (x''_1, \dots, x''_n) can be calculated via the Baker-Campbell-Hausdorff Formula from (x_1, \dots, x_n) , (x'_1, \dots, x'_n) , and the structural constants of the chosen basis.

Dilations $(\delta_\lambda)_{\lambda > 0}$ on G are given by

$$\delta_\lambda(x_1, \dots, x_n) := (\lambda^{\alpha_1} x_1, \dots, \lambda^{\alpha_n} x_n)$$

where $\alpha_i = j$ for $m_{j-1} + 1 \leq i \leq m_j$. These dilations form a one parameter group of group automorphisms.

Definition 1.2. We say that a quasi-distance d on G is *homogeneous* if d is left-invariant, i.e., $d(p \cdot q, p \cdot q') = d(q, q')$ for all $p, q, q' \in G$, and one-homogeneous with respect to the dilations, i.e., $d(\delta_\lambda(p), \delta_\lambda(q)) = \lambda d(p, q)$ for all $p, q \in G$ and all $\lambda > 0$.

Every homogeneous quasi-distance on a Carnot group G induces the topology of the group. Note also that any two homogeneous quasi-distances on a Carnot group G are

bi-Lipschitz equivalent. In particular, every homogeneous quasi-distance is bi-Lipschitz equivalent to every homogeneous distance.

One can characterize homogeneous quasi-distances by means of their unit ball centered at the origin. Namely, if d is a homogeneous quasi-distance on G and $K := B_d(0, 1)$, then 0 is in the interior of K , K is relatively compact, K is symmetric, i.e., $p \in K$ implies $p^{-1} \in K$, and for all $p \in G$ the set $\{\lambda > 0; \delta_{1/\lambda}(p) \in K\}$ is a closed sub-interval of $]0, +\infty[$. Conversely, if a subset K of G satisfies these assumptions, then

$$(1.3) \quad d(p, q) := \inf\{\lambda > 0; \delta_{1/\lambda}(p^{-1} \cdot q) \in K\}$$

defines a homogeneous quasi-distance on G . It is the homogeneous quasi-distance whose unit ball centered at the origin is K .¹

In particular, any set K of one of the following forms

$$(1.4) \quad K := \{(x_1, \dots, x_n) \in G; c_1|x_1|^{\gamma_1} + \dots + c_n|x_n|^{\gamma_n} \leq 1\}$$

or

$$(1.5) \quad K := \{x \in G; c_1\|\bar{x}_1\|_{d_1}^{\gamma_1} + \dots + c_s\|\bar{x}_s\|_{d_s}^{\gamma_s} \leq 1\}$$

for some $\gamma_i > 0$, $c_i > 0$, induces a homogeneous quasi-distance via (1.3). In (1.5), for $x = (x_1, \dots, x_n)$, we have set $\bar{x}_j := (x_{m_{j-1}+1}, \dots, x_{m_j})$, $d_j := \dim V_j$ and $\|\cdot\|_{d_j}$ denotes the Euclidean norm in \mathbb{R}^{d_j} .

Our main result is the following:

Theorem 1.6. *Let G be a Carnot group of step 3 or higher. Let K be a subset of G given by (1.4) or (1.5) and let d be the homogeneous quasi-distance induced by K via (1.3). Then BCP does not hold in (G, d) .*

Examples of homogeneous distances, i.e., satisfying the quasi-triangle inequality with a multiplicative constant $C = 1$, that satisfy the assumption of Theorem 1.6 have been given by Hebisch and Sikora. They proved in [3] that for any Carnot group G , there exists some $\alpha^* > 0$ such that, for all $0 < \alpha < \alpha^*$, the Euclidean ball $\{(x_1, \dots, x_n) \in G; |x_1|^2 + \dots + |x_n|^2 \leq \alpha^2\}$ with radius α induces a homogeneous distance on G via (1.3). For these distances, we have the following corollary.

Corollary 1.7. *Let G be a Carnot group of step 3 or higher and let d be a homogeneous distance on G whose unit ball centered at the origin is a Euclidean ball centered at the origin. Then BCP does not hold in (G, d) .*

As already mentioned, such homogeneous distances were our initial motivation and this corollary comes in contrast with the case of the Heisenberg groups, which are Carnot groups of step 2, due to the following result.

¹It may happen that a homogeneous quasi-distance on a Carnot group is not a continuous function on $G \times G$ with respect to the topological structure of the group. The fact that it induces the topology of the group only means that the unit ball centered at the origin contains the origin in its interior and that it is relatively compact. One can show that the quasi-distance is continuous on $G \times G$ if and only if its unit sphere at the origin is closed.

Theorem 1.8 ([6]). *Let \mathbb{H}^n be the n -th Heisenberg group and let d be a homogeneous distance on \mathbb{H}^n whose unit ball centered at the origin is a Euclidean ball centered at the origin. Then BCP holds in (G, d) .*

To our knowledge, the case of the Heisenberg groups are the only known examples of non-Abelian Carnot groups for which one can find homogeneous distances satisfying BCP, and the only known such distances are those considered in Theorem 1.8. This makes the Heisenberg groups peculiar cases as far as the validity of BCP for some homogeneous distance on a Carnot group is concerned. Theorem 1.6 indeed shows that natural analogues of these distances do not satisfy BCP as soon as the step of the group is 3 or higher. There are moreover some hints towards the fact that these results and this dichotomy between step 2 and step 3 or higher Carnot groups may generalize (and even in the more general setting of graded groups). One could indeed expect that homogeneous distances whose unit ball centered at the origin is a Euclidean ball centered at the origin satisfy BCP on Carnot groups of step 2 whereas there is no homogeneous distance for which BCP holds as soon as the step of the group is 3 or higher. This will be studied in a forthcoming paper.

The proof of Theorem 1.6 (see Section 3) is based on two main ingredients. First, we show that for a Carnot group equipped with a homogeneous quasi-distance whose unit ball centered at the origin is given by (1.4) (respectively (1.5)), the validity of BCP implies that $\gamma_1, \dots, \gamma_{m_1}$ (respectively γ_1) are bounded below by the step of the group, see Lemma 3.1. Hence, for Carnot groups of step 3 and higher, we get $\gamma_1, \dots, \gamma_{m_1} \geq 3$ (respectively $\gamma_1 \geq 3$) whenever BCP holds. Next, a reduction argument on the step of the group by taking a quotient allows us to reduce the problem to the case of the first Heisenberg group equipped with a homogeneous quasi-distance inherited from the original one. The fact that the quotient map is a submetry plays a key role here. Submetries are indeed particularly well adapted tools in this context. See Section 2 where we prove some of their properties related to the Besicovitch Covering Property. When $\gamma_1, \dots, \gamma_{m_1} \geq 3$ (respectively $\gamma_1 \geq 3$), the unit sphere of the homogeneous quasi-distance induced on the first Heisenberg group by the original one turns out to have vanishing Euclidean curvature at the poles. On the other hand, we know by [6] that, in the Heisenberg groups, BCP does not hold for homogeneous quasi-distances whose unit sphere has vanishing Euclidean curvature at the poles. In particular, BCP cannot hold for the inherited homogeneous quasi-distance. This implies in turn that BCP is not satisfied by the original distance.

2. WEAK BESICOVITCH COVERING PROPERTY AND SUBMETRIES

First, we introduce what we call here the Weak Besicovitch Covering Property (the terminology might not be standard).

Definition 2.1 (Family of Besicovitch balls). Let (X, d) be a quasi-metric space. We say that a family \mathcal{B} of balls in (X, d) is a *family of Besicovitch balls* if $\mathcal{B} = \{B = B_d(x_B, r_B)\}$ is a finite family of balls such that $x_B \notin B'$ for all $B, B' \in \mathcal{B}$, $B \neq B'$, and for which $\bigcap_{B \in \mathcal{B}} B \neq \emptyset$.

Definition 2.2 (Weak Besicovitch Covering Property). We say that a quasi-metric space (X, d) satisfies the Weak Besicovitch Covering Property (WBCP) if there exists a constant $Q \in \mathbb{N}$ such that $\text{Card } \mathcal{B} \leq Q$ for every family \mathcal{B} of Besicovitch balls in (X, d) .

If (X, d) satisfies BCP, then (X, d) satisfies WBCP. One can indeed take $Q = N$ where N is given by Definition 1.1.² We will prove in Section 3 that WBCP, and hence BCP, does not hold in Carnot groups of step 3 and higher equipped with homogeneous quasi-distances as in Theorem 1.6.

Submetries will play a key role in our arguments. In the rest of this section, we recall the definition of submetries and prove some of their properties to be used in the proof of Theorem 1.6, see Proposition 2.7 and Corollary 2.11.

Definition 2.3 (Submetry). Let (X, d_X) and (Y, d_Y) be quasi-metric spaces. We say that $\pi : X \rightarrow Y$ is a submetry if π is a surjective map such that

$$(2.4) \quad \pi(B_{d_X}(p, r)) = B_{d_Y}(\pi(p), r)$$

for all $p \in X$ and all $r > 0$.

Remark 2.5. Any submetry $\pi : (X, d_X) \rightarrow (Y, d_Y)$ is 1-Lipschitz. Indeed, given $p, q \in X$, set $r := d_X(p, q)$. We have $q \in B_{d_X}(p, r)$ hence $\pi(q) \in \pi(B_{d_X}(p, r)) = B_{d_Y}(\pi(p), r)$. Hence $d_Y(\pi(p), \pi(q)) \leq r = d_X(p, q)$.

The following characterization of submetries will be technically convenient. For subsets $A, B \subset X$, we consider here the distance $d_X(A, B)$ defined by $d_X(A, B) := \inf\{d_X(p, q); p \in A, q \in B\}$.

Proposition 2.6. *Let (X, d_X) and (Y, d_Y) be quasi-metric spaces. Let $\pi : (X, d_X) \rightarrow (Y, d_Y)$ be a surjective map. Then the following are equivalent:*

- (i) π is a submetry,
- (ii) for all $\hat{p} \in Y$, all $\hat{q} \in Y$ and all $p \in \pi^{-1}(\hat{p})$, there exists $q \in \pi^{-1}(\hat{q})$ such that

$$d_X(p, q) = d_Y(\hat{p}, \hat{q}) = d_X(\pi^{-1}(\hat{p}), \pi^{-1}(\hat{q})) = d_X(p, \pi^{-1}(\hat{q})) .$$

Proof. Assume that π is a submetry. Let $\hat{p} \in Y$, $\hat{q} \in Y$ and $p \in \pi^{-1}(\hat{p})$. Since π is 1-Lipschitz (see Remark 2.5), we have $d_Y(\hat{p}, \hat{q}) \leq d_X(p', q')$ for all $p' \in \pi^{-1}(\hat{p})$ and all $q' \in \pi^{-1}(\hat{q})$. It follows that

$$d_Y(\hat{p}, \hat{q}) \leq d_X(\pi^{-1}(\hat{p}), \pi^{-1}(\hat{q})) \leq d_X(p, \pi^{-1}(\hat{q})).$$

Set $r := d_Y(\hat{p}, \hat{q})$. We have $\hat{q} \in B_{d_Y}(\hat{p}, r) = \pi(B_{d_X}(p, r))$ hence one can find $q \in \pi^{-1}(\hat{q}) \cap B_{d_X}(p, r)$. Then we have $d_X(p, q) \leq r = d_Y(\hat{p}, \hat{q})$. All together, we get that

$$d_Y(\hat{p}, \hat{q}) \leq d_X(\pi^{-1}(\hat{p}), \pi^{-1}(\hat{q})) \leq d_X(p, \pi^{-1}(\hat{q})) \leq d_X(p, q) \leq d_Y(\hat{p}, \hat{q})$$

from which (ii) follows.

Conversely, assume that (ii) holds. Since π is assumed to be surjective, we only need to prove that (2.4) holds. Let $p \in X$ and $r > 0$. Let us first prove that $\pi(B_{d_X}(p, r)) \subset B_{d_Y}(\pi(p), r)$. Let $\hat{q} \in \pi(B_{d_X}(p, r))$. Then $\hat{q} = \pi(q)$ for some $q \in B_{d_X}(p, r)$ and it follows from (ii) that

$$d_Y(\pi(p), \hat{q}) = d_X(p, \pi^{-1}(\hat{q})) \leq d_X(p, q) \leq r ,$$

²WBCP is in general strictly weaker than BCP. However, BCP and WBCP are equivalent in doubling metric spaces.

i.e., $\hat{q} \in B_{d_Y}(\pi(p), r)$. Conversely, let $\hat{q} \in B_{d_Y}(\pi(p), r)$. It follows from (ii) that one can find $q \in \pi^{-1}(\hat{q})$ such that

$$d_X(p, q) = d_Y(\pi(p), \hat{q}) \leq r,$$

i.e., $q \in B_{d_X}(p, r)$. Hence $\hat{q} = \pi(q) \in \pi(B_{d_X}(p, r))$ and this concludes the proof. \square

The next proposition shows that submetries preserve the validity of WBCP.

Proposition 2.7. *Let (X, d_X) and (Y, d_Y) be quasi-metric spaces. Assume that there exists a submetry from (X, d_X) onto (Y, d_Y) . If (X, d_X) satisfies WBCP then (Y, d_Y) satisfies WBCP.*

Proof. Let $\pi : (X, d_X) \rightarrow (Y, d_Y)$ be a submetry. Let $\hat{\mathcal{B}} = \{B = B_{d_Y}(y_B, r_B)\}$ be a family of Besicovitch balls in (Y, d_Y) (see Definition 2.1). Let $\hat{p} \in \bigcap_{B \in \hat{\mathcal{B}}} B$ and fix some $p \in \pi^{-1}(\hat{p})$. Using Proposition 2.6, for each $B_{d_Y}(y_B, r_B) \in \hat{\mathcal{B}}$, one can find $x_B \in \pi^{-1}(y_B)$ such that $d_X(p, x_B) = d_Y(\hat{p}, y_B)$. It follows that $d_X(p, x_B) \leq r_B$ and hence $p \in \bigcap_{B \in \hat{\mathcal{B}}} B_{d_X}(x_B, r_B)$. On the other hand, since π is 1-Lipschitz (see Remark 2.5), we have $d_X(x_B, x_{B'}) \geq d_Y(y_B, y_{B'}) > \max(r_B, r_{B'})$ for all $B, B' \in \hat{\mathcal{B}}, B \neq B'$. It follows that $\{B_{d_X}(x_B, r_B); B \in \hat{\mathcal{B}}\}$ is a family of Besicovitch balls in (X, d_X) . Since (X, d_X) satisfies WBCP, we have $\text{Card } \hat{\mathcal{B}} \leq Q$ for some $Q \in \mathbb{N}$ (see Definition 2.2). Hence (Y, d_Y) satisfies WBCP as well. \square

In the next proposition we give a sufficient condition on the fibers of a surjective map that allows us to construct on the target space a quasi-distance for which this map is a submetry.

Proposition 2.8. *Let (X, d_X) be a quasi-metric space and Y be a nonempty set. Let $\pi : X \rightarrow Y$ be a surjective map. Assume that the fibers of π are parallel, i.e., assume that for all $\hat{p} \in Y$, all $\hat{q} \in Y$ and all $p \in \pi^{-1}(\hat{p})$, one can find $q \in \pi^{-1}(\hat{q})$ such that*

$$(2.9) \quad d_X(\pi^{-1}(\hat{p}), \pi^{-1}(\hat{q})) = d_X(p, q).$$

Then

$$d_Y(\hat{p}, \hat{q}) := d_X(\pi^{-1}(\hat{p}), \pi^{-1}(\hat{q}))$$

defines a quasi-distance on Y and π is a submetry from (X, d_X) onto (Y, d_Y) .

Proof. First, let us check that d_Y defines a quasi-distance on Y . Assume that $d_Y(\hat{p}, \hat{q}) = 0$. Then, by definition of d_Y and using (2.9), for all $p \in \pi^{-1}(\hat{p})$, one can find $q \in \pi^{-1}(\hat{q})$ such that $d_X(p, q) = d_Y(\hat{p}, \hat{q}) = 0$. This implies that $p = q$ and hence $\hat{p} = \hat{q}$. The fact that $d_Y(\hat{p}, \hat{q}) = d_Y(\hat{q}, \hat{p})$ is obvious from the definition of d_Y . Next, we check that d_Y satisfies the quasi-triangle inequality with the same multiplicative constant C as d_X . Let \hat{p}, \hat{q} and $\hat{p}' \in Y$. Let p' be some point in $\pi^{-1}(\hat{p}')$. Using (2.9), one can find $p \in \pi^{-1}(\hat{p})$ such that $d_Y(\hat{p}', \hat{p}) = d_X(p', p)$. Similarly, one can find $q \in \pi^{-1}(\hat{q})$ such that $d_Y(\hat{p}', \hat{q}) = d_X(p', q)$. Then we get that

$$\begin{aligned} d_Y(\hat{p}, \hat{q}) &= d_X(\pi^{-1}(\hat{p}), \pi^{-1}(\hat{q})) \leq d_X(p, q) \\ &\leq C(d_X(p, p') + d_X(p', q)) = C(d_Y(\hat{p}, \hat{p}') + d_Y(\hat{p}', \hat{q})). \end{aligned}$$

Finally, the fact that π is a submetry from (X, d_X) onto (Y, d_Y) follows from Proposition 2.6 together with (2.9). \square

We show in the following proposition that Proposition 2.8 can be applied to quotient maps from a topological group modulo a boundedly compact normal subgroup.

Proposition 2.10. *Let G be a topological group equipped with a left-invariant quasi-distance d that induces the topology of the group. Let $N \triangleleft G$ be a normal subgroup of G . Assume that N is boundedly compact. Then the cosets, i.e., the fibers of the quotient map $\pi : G \rightarrow G/N$, are parallel.*

Proof. Let $\hat{p}, \hat{q} \in G/N$ and $p \in \pi^{-1}(\hat{p})$. Since the quasi-distance on G is left-invariant and N is boundedly compact, any coset is boundedly compact as well. It follows that one can find $q \in \pi^{-1}(\hat{q})$ such that $d(p, q) = d(p, \pi^{-1}(\hat{q}))$. For each $\varepsilon > 0$, take $p' \in \pi^{-1}(\hat{p})$ and $q' \in \pi^{-1}(\hat{q})$ such that $d(p', q') \leq d(\pi^{-1}(\hat{p}), \pi^{-1}(\hat{q})) + \varepsilon$. By left-invariance of d and noting that $p \cdot (p')^{-1} \cdot q' \in \pi^{-1}(\hat{q})$, we get that

$$\begin{aligned} \varepsilon + d(\pi^{-1}(\hat{p}), \pi^{-1}(\hat{q})) &\geq d(p', q') = d(p, p \cdot (p')^{-1} \cdot q') \\ &\geq d(p, \pi^{-1}(\hat{q})) = d(p, q) \geq d(\pi^{-1}(\hat{p}), \pi^{-1}(\hat{q})). \end{aligned}$$

Since ε is arbitrary, it follows that $d(\pi^{-1}(\hat{p}), \pi^{-1}(\hat{q})) = d(p, q)$ and hence the fibers of the quotient map π are parallel. \square

The next corollary is a straightforward consequence of Proposition 2.8 and Proposition 2.10.

Corollary 2.11. *Let G be a topological group equipped with a left-invariant quasi-distance d that induces the topology of the group. Let $N \triangleleft G$ be a normal subgroup of G . Assume that N is boundedly compact. Let $\pi : G \rightarrow G/N$ denote the quotient map. Then*

$$d_{G/N}(\hat{p}, \hat{q}) := d(\pi^{-1}(\hat{p}), \pi^{-1}(\hat{q}))$$

defines a quasi-distance on G/N and π is a submetry from (G, d) onto $(G/N, d_{G/N})$.

3. PROOF OF THEOREM 1.6

This section is devoted to the proof of Theorem 1.6. We consider a Carnot group G of step s equipped with a homogeneous quasi-distance d whose unit ball centered at the origin is given by (1.4), i.e., can be described as

$$B_d(0, 1) = \{(x_1, \dots, x_n) \in G; c_1|x_1|^{\gamma_1} + \dots + c_n|x_n|^{\gamma_n} \leq 1\}$$

for some $\gamma_i > 0$, $c_i > 0$. The case of a homogeneous quasi-distance whose unit ball centered at the origin is given by (1.5) is similar and can be proved with the same arguments.

First, we prove that the validity of BCP implies that $\gamma_1, \dots, \gamma_{m_1}$ are bounded below by the step of the group. Recall that m_1 denotes the dimension of the first layer V_1 of the stratification of the Lie algebra \mathfrak{g} of G .

Lemma 3.1. *Assume that BCP holds in (G, d) . Then $\min(\gamma_1, \dots, \gamma_{m_1}) \geq s$.*

Proof. Let $1 \leq i \leq m_1$ be fixed and set $N_i := \{(x_1, \dots, x_n) \in G; x_k = 0 \text{ for } k \neq i, n\}$. Since BCP, and hence WBCP (see Definition 2.2), holds in (G, d) , WBCP also holds in (N_i, d_{N_i}) where d_{N_i} denotes the quasi-distance d restricted to N_i .³

On the other hand, N_i is an abelian subgroup of G that can be identified with \mathbb{R}^2 equipped with the usual addition, denoted by $+$, as a group law and with the family of dilations $\tilde{\delta}_\lambda(x, y) := (\lambda x, \lambda^s y)$ for $\lambda > 0$. With this identification, the quasi-distance d_{N_i} is then a left-invariant and one-homogeneous quasi-distance on $(\mathbb{R}^2, +, (\tilde{\delta}_\lambda)_{\lambda > 0})$ whose unit ball centered at the origin can be described as $\{(x, y) \in \mathbb{R}^2; c_i |x|^{\gamma_i} + c_n |y|^{\gamma_n} \leq 1\}$. It then follows from Lemma 3.2 below that $\gamma_i \geq s$. \square

Lemma 3.2. *Let \mathbb{R}^2 be equipped with the usual addition, denoted by $+$, as a group law and the family of dilations $\tilde{\delta}_\lambda(x, y) := (\lambda x, \lambda^s y)$ for some $s > 0$. Let ρ be a left-invariant and one-homogeneous quasi-distance on $(\mathbb{R}^2, +, (\tilde{\delta}_\lambda)_{\lambda > 0})$. Assume that the unit ball centered at the origin can be described as*

$$B_\rho(0, 1) = \{(x, y) \in \mathbb{R}^2; \alpha |x|^a + \beta |y|^b \leq 1\}$$

for some $a > 0, b > 0, \alpha > 0, \beta > 0$. If WBCP holds in (\mathbb{R}^2, ρ) , then $a \geq s$.

Proof. First, we note that we only need to consider the case $\alpha = \beta = 1$. Indeed, considering the group automorphism

$$f(x, y) := (\alpha^{1/a} x, \beta^{1/b} y),$$

which commutes with the dilations $\tilde{\delta}_\lambda$, then

$$\rho'(p, q) := \rho(f^{-1}(p), f^{-1}(q))$$

defines a homogeneous quasi-distance on $(\mathbb{R}^2, +, (\tilde{\delta}_\lambda)_{\lambda > 0})$ and $f : (\mathbb{R}^2, \rho) \rightarrow (\mathbb{R}^2, \rho')$, $f^{-1} : (\mathbb{R}^2, \rho') \rightarrow (\mathbb{R}^2, \rho)$ are submetries. It then follows from Proposition 2.7 that WBCP holds in (\mathbb{R}^2, ρ) if and only if WBCP holds in (\mathbb{R}^2, ρ') .

Thus let us assume that

$$B_\rho(0, 1) = \{(x, y) \in \mathbb{R}^2; |x|^a + |y|^b \leq 1\}.$$

Arguing by contradiction, let us assume that $0 < a < s$. We will prove that one can find $r > 1$ and a positive sequence $(\varepsilon_n)_{n \geq 1}$ decreasing to 0 such that, setting $p_n := (x_n, y_n)$ where

$$(3.3) \quad x_n = r^{-n} \quad \text{and} \quad y_n = \varepsilon_n^{-s} (1 - \varepsilon_n^a r^{-na})^{1/b},$$

the following hold. First,

$$(3.4) \quad 0 \in \partial B_\rho(p_n, \varepsilon_n^{-1})$$

for all $n \geq 1$. Second,

$$(3.5) \quad \rho(p_n, p_k) > \varepsilon_n^{-1}$$

for all $n \geq 2$ and all $1 \leq k \leq n - 1$. Since the sequence $(\varepsilon_n^{-1})_{n \geq 1}$ is increasing, we get from (3.5) that

$$\rho(p_n, p_k) > \max(\varepsilon_n^{-1}, \varepsilon_k^{-1})$$

³More generally, if Y is a subset of a quasi-metric space (X, d_X) that satisfies WBCP, then (Y, d_Y) also satisfies WBCP where d_Y denote the quasi-distance d_X restricted to Y .

for all $n \neq k$ hence $p_k \notin B_\rho(p_n, \varepsilon_n^{-1})$ for all $n \neq k$. Combining this with (3.4), we get that $\{B_\rho(p_n, \varepsilon_n^{-1}); n \in J\}$ is a family of Besicovitch balls for any $J \subset \mathbb{N}$ finite which gives a contradiction to the validity of WBCP in (\mathbb{R}^2, ρ) .

First, it follows from (3.3) that

$$|\varepsilon_n x_n|^a + |\varepsilon_n^s y_n|^b = \varepsilon_n^a r^{-na} + (1 - \varepsilon_n^a r^{-na}) = 1$$

hence $\rho(0, \tilde{\delta}_{\varepsilon_n}(p_n)) = 1$. By homogeneity it follows that $\rho(0, p_n) = \varepsilon_n^{-1}$ hence (3.4) holds for any fixed $r > 1$ and any positive sequence $(\varepsilon_n)_{n \geq 1}$. Hence it remains to find some $r > 1$ and some positive sequence $(\varepsilon_n)_{n \geq 1}$ decreasing to 0 such that (3.5) holds to conclude the proof.

Let $r > 1$ to be fixed later and set $\varepsilon_1 = 1$. By induction, assume that $\varepsilon_1 > \dots > \varepsilon_n$ have already been chosen. We are looking for $\varepsilon_{n+1} \in (0, \varepsilon_n)$ such that $\rho(p_{n+1}, p_k) > \varepsilon_{n+1}^{-1}$, i.e.,

$$(3.6) \quad \rho(\tilde{\delta}_{\varepsilon_{n+1}}(p_{n+1}), \tilde{\delta}_{\varepsilon_{n+1}}(p_k)) > 1$$

for all $1 \leq k \leq n$. We have

$$\begin{aligned} & |\varepsilon_{n+1}(x_k - x_{n+1})|^a + |\varepsilon_{n+1}^s(y_{n+1} - y_k)|^b \\ &= \varepsilon_{n+1}^a (r^{-k} - r^{-(n+1)a})^a + |(1 - \varepsilon_{n+1}^a r^{-(n+1)a})^{1/b} - \varepsilon_{n+1}^s \varepsilon_k^{-s} (1 - \varepsilon_k^a r^{-ka})^{1/b}|^b. \end{aligned}$$

Since $s > a > 0$, we have, for all $k \in \{1, \dots, n\}$ fixed,

$$(1 - \varepsilon_{n+1}^a r^{-(n+1)a})^{1/b} - \varepsilon_{n+1}^s \varepsilon_k^{-s} (1 - \varepsilon_k^a r^{-ka})^{1/b} = 1 - b^{-1} \varepsilon_{n+1}^a r^{-(n+1)a} + o(\varepsilon_{n+1}^a).$$

It follows that

$$\begin{aligned} & |\varepsilon_{n+1}(x_k - x_{n+1})|^a + |\varepsilon_{n+1}^s(y_{n+1} - y_k)|^b \\ &= 1 + \varepsilon_{n+1}^a ((r^{-k} - r^{-(n+1)a})^a - r^{-(n+1)a}) + o(\varepsilon_{n+1}^a) \\ &= 1 + \varepsilon_{n+1}^a r^{-ak} ((1 - r^{-(n+1)+k})^a - r^{-(n+1)+k}) + o(\varepsilon_{n+1}^a) \\ &\geq 1 + \varepsilon_{n+1}^a r^{-ak} ((1 - r^{-1})^a - r^{-a}) + o(\varepsilon_{n+1}^a). \end{aligned}$$

Hence, choosing $r > 1$ so that $(1 - r^{-1})^a - r^{-a} > 0$, we get that one can choose ε_{n+1} small enough so that

$$|\varepsilon_{n+1}(x_k - x_{n+1})|^a + |\varepsilon_{n+1}^s(y_{n+1} - y_k)|^b > 1$$

for all $1 \leq k \leq n$ which proves (3.6). \square

From now on, we assume that G is a Carnot group of step 3 or higher and we argue by contradiction, assuming that BCP holds in (G, d) . Hence, we have from Lemma 3.1 that

$$(3.7) \quad \min(\gamma_1, \dots, \gamma_{m_1}) \geq 3.$$

Let us consider $N := \exp(V_3 \oplus \dots \oplus V_s)$. Then N is a normal subgroup of G . The quotient group $\hat{G} := G/N$ can be identified with \mathbb{R}^{m_2} equipped with the group law

$$\hat{p} * \hat{p}' := \hat{\pi}([\hat{p}, 0] \cdot [\hat{p}', 0])$$

where $\hat{\pi}(x_1, \dots, x_n) := (x_1, \dots, x_{m_2})$ and where, for $p = (x_1, \dots, x_n) \in G$, we set $p := [\hat{p}, \tilde{p}]$ with $\hat{p} := \hat{\pi}(p)$ and $\tilde{p} := (x_{m_2+1}, \dots, x_n)$.

The group \hat{G} inherits from G a natural structure of Carnot group of step 2 with dilations given by

$$\hat{\delta}_\lambda(x_1, \dots, x_{m_2}) := (\lambda x_1, \dots, \lambda x_{m_1}, \lambda^2 x_{m_1+1}, \dots, \lambda^2 x_{m_2}) .$$

Since the exponential map is here a global diffeomorphism, N is boundedly compact in (G, d) and it follows from Corollary 2.11 that

$$d_{\hat{G}}(\hat{p}, \hat{q}) := d(\hat{\pi}^{-1}(\hat{p}), \hat{\pi}^{-1}(\hat{q}))$$

defines a quasi-distance on \hat{G} and $\hat{\pi} : (G, d) \rightarrow (\hat{G}, d_{\hat{G}})$ is a submetry. Hence, by Proposition 2.7, $(\hat{G}, d_{\hat{G}})$ satisfies WBCP.

Let us now check that $d_{\hat{G}}$ is the homogeneous quasi-distance on \hat{G} whose unit ball centered at the origin is given by

$$(3.8) \quad B_{d_{\hat{G}}}(0, 1) = \{(x_1, \dots, x_{m_2}) \in \hat{G}; c_1|x_1|^{\gamma_1} + \dots + c_{m_2}|x_{m_2}|^{\gamma_{m_2}} \leq 1\} .$$

The fact that $d_{\hat{G}}$ is left-invariant can be easily checked using the left invariance of d and the fact that $\hat{\pi}$ is a group homomorphism. The homogeneity of $d_{\hat{G}}$ with respect to dilations $\hat{\delta}_\lambda$ can be easily checked as well noting that $\hat{\pi}^{-1}(\hat{\delta}_\lambda(\hat{p})) = \delta_\lambda(\hat{\pi}^{-1}(\hat{p}))$ for all $\hat{p} \in \hat{G}$ and $\lambda > 0$ and using the homogeneity of d .

Let us now check that (3.8) holds. For $p = (x_1, \dots, x_n) \in G$, we have

$$\begin{aligned} d(0, p) &= \inf\{r > 0; \delta_{1/r}(p) \in B_d(0, 1)\} \\ &= \inf\{r > 0; f(r^{-\alpha_1}|x_1|, \dots, r^{-\alpha_n}|x_n|) \leq 1\} \end{aligned}$$

where $f : (\mathbb{R}^+)^n \rightarrow \mathbb{R}^+$ is given by $f(t_1, \dots, t_n) := c_1 t_1^{\gamma_1} + \dots + c_n t_n^{\gamma_n}$. This function being increasing with respect to the $(n - m_2)$ last coordinates, we have

$$d(0, [\hat{\pi}(p), 0]) \leq d(0, p)$$

for all $p \in G$. Together with Proposition 2.6, this implies that

$$d_{\hat{G}}(0, \hat{p}) = d(0, \hat{\pi}^{-1}(\hat{p})) = d(0, [\hat{p}, 0])$$

for all $\hat{p} \in \hat{G}$. Hence,

$$B_{d_{\hat{G}}}(0, 1) = \{(x_1, \dots, x_{m_2}) \in \hat{G}; f(|x_1|, \dots, |x_{m_2}|, 0, \dots, 0) \leq 1\}$$

which proves (3.8).

Let (Y_1, \dots, Y_{m_2}) be the basis of the Lie algebra $\hat{\mathfrak{g}}$ of \hat{G} inherited from the chosen basis (X_1, \dots, X_n) adapted to the stratification of \mathfrak{g} . Let us fix $i, j \in \{1, \dots, m_1\}$ such that $[Y_i, Y_j] \neq 0$. Set $\mathfrak{h} := \text{span}(Y_i, Y_j, [Y_i, Y_j])$ and $H := \exp \mathfrak{h}$. Then H is a subgroup of \hat{G} that can be identified with the first Heisenberg group. Recall that the first Heisenberg group is the Carnot group of step 2 whose stratification of the Lie algebra is given by $W_1 \oplus W_2$ where $\dim W_1 = 2$ and $\dim W_2 = 1$. Hence we can identify H with \mathbb{R}^3 equipped with the Heisenberg group structure given by

$$(x, y, z) \cdot (x', y', z') := (x + x', y + y', z + z' + \frac{1}{2}(xy' - x'y)) ,$$

where we identify $\exp(xY_i + yY_j + z[Y_i, Y_j])$ with (x, y, z) , and equipped with the family of dilations $((x, y, z) \mapsto (\lambda x, \lambda y, \lambda^2 z))_{\lambda > 0}$.

The quasi-distance d_H induced by the restriction of $d_{\hat{G}}$ on H is then a homogeneous quasi-distance whose unit ball centered at the origin is given by

$$B_{d_H}(0, 1) = \{(x, y, z) \in H; c_i|x|^{\gamma_i} + c_j|y|^{\gamma_j} + c_{m_1+1}|\xi_{m_1+1} z|^{\gamma_{m_1+1}} + \dots + c_{m_2}|\xi_{m_2} z|^{\gamma_{m_2}} \leq 1\}$$

where $[Y_i, Y_j] = \xi_{m_1+1}Y_{m_1+1} + \dots + \xi_{m_2}Y_{m_2}$.

Since WBCP holds in $(\hat{G}, d_{\hat{G}})$, WBCP also hold in (H, d_H) (see Footnote 3). On the other hand, we have $\gamma_i, \gamma_j \geq 3$ by (3.7). Near the north pole, i.e., the intersection of $\partial B_{d_H}(0, 1)$ with the positive z -axis, $B_{d_H}(0, 1)$ can thus be described as the subgraph $\{(x, y, z) \in H; z \leq \varphi(x, y)\}$ of a C^2 function φ whose first and second order partial derivatives vanish at the origin. Then it follows from [6, Theorem 6.1] that WBCP can not hold in (H, d_H) . Note that Theorem 6.1 in [6] holds not only for homogeneous distances but more generally for homogeneous quasi-distances (with the same proof). This gives a contradiction and concludes the proof of Theorem 1.6.

Acknowledgement. The second-named author is very grateful to the Department of Mathematics and Statistics of the University of Jyväskylä, where part of this work was done, for its hospitality.

REFERENCES

- [1] A.S. Besicovitch, *A general form of the covering principle and relative differentiation of additive functions*, Proc. Cambridge Philos. Soc. 41 (1945) 103–110.
- [2] A.S. Besicovitch, *A general form of the covering principle and relative differentiation of additive functions. II*, Proc. Cambridge Philos. Soc. 42 (1946) 1–10.
- [3] W. Hebisch, A. Sikora, *A smooth subadditive homogeneous norm on a homogeneous group*, Studia Math. 96 (1990), no. 3, 231–236.
- [4] J. Heinonen, *Lectures on analysis on metric spaces*, Universitext. Springer-Verlag, New York, 2001. x+140 pp.
- [5] A. Korányi, H.M. Reimann, *Foundations for the theory of quasiconformal mappings on the Heisenberg group*, Adv. Math. 111 (1995), no. 1, 1–87.
- [6] E. Le Donne, S. Rigot, *Besicovitch Covering Property for homogeneous distances in the Heisenberg groups*, preprint, <http://arxiv.org/abs/1406.1484>
- [7] P. Mattila, *Geometry of sets and measures in Euclidean spaces. Fractals and rectifiability*, Cambridge Studies in Advanced Mathematics, 44. Cambridge University Press, Cambridge, 1995. xii+343 pp.
- [8] S. Rigot, *Counter example to the Besicovitch covering property for some Carnot groups equipped with their Carnot-Carathéodory metric*, Math. Z. 248 (2004), no. 4, 827–848.
- [9] R.L. Sawyer, R.L. Wheeden, *Weighted inequalities for fractional integrals on Euclidean and homogeneous spaces*, Amer. J. Math. 114 (1992), no. 4, 813–874.

(Le Donne) DEPARTMENT OF MATHEMATICS AND STATISTICS, P.O. BOX 35, FI-40014, UNIVERSITY OF JYVÄSKYLÄ, FINLAND

E-mail address: ledonne@msri.org

(Rigot) LABORATOIRE DE MATHÉMATIQUES J.A. DIEUDONNÉ UMR CNRS 7351, UNIVERSITÉ NICE SOPHIA ANTIPOLIS, 06108 NICE CEDEX 02, FRANCE

E-mail address: rigot@unice.fr