

ASYMPTOTIC BEHAVIOR OF THE RIEMANNIAN HEISENBERG GROUP AND ITS HOROBOUNDARY

ENRICO LE DONNE, SEBASTIANO NICOLUSSI GOLO,
AND ANDREA SAMBUSETTI

ABSTRACT. The paper is devoted to the large scale geometry of the Heisenberg group \mathbb{H} equipped with left-invariant Riemannian metrics. We prove that two such metrics have bounded difference if and only if they are asymptotic, i.e., their ratio goes to one, at infinity. Moreover, we show that for every left-invariant Riemannian metric d on \mathbb{H} there is a unique subRiemannian metric d' for which $d - d'$ goes to zero at infinity, and we estimate the rate of convergence. As a first immediate consequence we get that the Riemannian Heisenberg group is at bounded distance from its asymptotic cone. The second consequence, which was our aim, is the explicit description of the horoboundary of the Riemannian Heisenberg group.

CONTENTS

1. Introduction	2
2. Preliminaries	6
3. Comparison between strictly subRiemannian metrics	9
4. Comparison between Riemannian and strictly subRiemannian metrics	13
5. The horoboundary	18
Appendix A. Length-minimizing curves for d_{CC} and d_R	22
References	24

Date: September 1, 2015.

2010 Mathematics Subject Classification. 20F69, 53C23, 53C17.

Key words and phrases. Heisenberg group, horoboundary, asymptotic cone, Riemannian geometry, sub-Riemannian geometry.

E.L.D. has been supported by the Academy of Finland project no. 288501. S.N.G. has been supported by the People Programme (Marie Curie Actions) of the European Union's Seventh Framework Programme FP7/2007-2013/ under REA grant agreement n. 607643.

1. INTRODUCTION

In large-scale geometry, various notions of space at infinity have received special interest for differently capturing the asymptotic geometric behavior. Two main examples of spaces at infinity are the asymptotic cone and the horoboundary. The description of the *asymptotic cone* for finitely generated groups is a crucial step in the algebraic characterization of groups of polynomial growth, [Mil68; Tit72; Gro81b; Wol68; Pan83; BLD13]. The notion of *horoboundary* has been formulated by Gromov [Gro81a], inspired by the seminal work of Busemann on the theory of parallels on geodesic spaces [Bus55]. The horoboundary has a fully satisfying visual description in the framework of $CAT(0)$ -spaces and of Gromov-hyperbolic spaces, [Gro87; Bal95; BH99]. It plays a major role in the study of dynamics and rigidity of negatively curved spaces, [Had98; Mos73; Gro87; Sul79; Pat76; Rob03]. The visual-boundary description breaks down for non-simply connected manifolds [DPS12] and when the curvature has variable sign, as we will make evident for the Riemannian Heisenberg group.

This paper contributes to the study of the asymptotic geometry of the simplest non-Abelian nilpotent group: the Heisenberg group. The asymptotic cone of the Heisenberg group equipped with a left-invariant Riemannian metric d_R is the Heisenberg group equipped with a Carnot-Carathéodory metric d_{CC} , see [Pan83] and also [BLD13]. Our contribution is a finer analysis of the asymptotic comparison of d_R and d_{CC} . This leads to the explicit knowledge of the (Riemannian) horoboundary. We remark that the Heisenberg group is not hyperbolic, hence one does not consider its visual boundary.

We recall the definition of horoboundary. Let (X, d) be a metric space. We consider the space of continuous real functions $\mathcal{C}(X)$ endowed with the topology of uniform convergence on compact sets. We denote by $\mathcal{C}(X)/\mathbb{R}$ the quotient with respect to the subspace of constant functions. The map $x \mapsto d(x, \cdot)$ induces an embedding $X \hookrightarrow \mathcal{C}(X)/\mathbb{R}$. The horoboundary of X is defined as $\partial_h X := \bar{X} \setminus X \subset \mathcal{C}(X)/\mathbb{R}$. See Section 5 for a detailed exposition.

The horoboundary has been investigated for finite-dimensional normed vector spaces, [Wal07], for Hilbert geometries, [Wal08], and for infinite graphs [WW06]. For non-simply connected, negatively curved manifolds it has been studied in [DPS12]. Nicas and Klein computed the horoboundary of the Heisenberg group when endowed with the Korany metric in [KN09], and with the metric d_{CC} in [KN10].

We will show that the horoboundary of the Heisenberg group endowed with a left-invariant Riemannian metric d_R coincides with the second case studied by Nicas and Klein, see Corollary 1.4. This will be an immediate consequence of our main result Theorem 1.3, which implies that the difference $d_R - d_{CC}$ converges to zero when evaluated

on points (p, q_n) with q_n being a sequence that leaves every compact set.

Remark 1.1. Another term for Carnot-Carathéodory metric is subRiemannian metric. In this paper we should discuss subRiemannian metrics that may actually be Riemannian. Therefore we follow the convention that subRiemannian geometry includes as a particular case Riemannian geometry. This is in agreement with several established references in the field, see [ABB15; Rif14; Jea14]. In the presence of a subRiemannian metric that is not Riemannian we shall use the term *strict subRiemannian*.

1.1. Detailed results. The Heisenberg group \mathbb{H} is the simply connected Lie group whose Lie algebra \mathfrak{h} is generated by three vectors X, Y, Z with only non-zero relation $[X, Y] = Z$. A left-invariant Riemannian metric d on \mathbb{H} is determined by a scalar product g on \mathfrak{h} ; a left-invariant *strictly* subRiemannian metric d is induced by a bracket generating plane $V \subset \mathfrak{h}$ and a scalar product g on V (see Section 2 for detailed exposition). In both cases we say that d is subRiemannian with *horizontal space* (V, g) , where $\dim V$ is either 2 or 3.

We are interested in the asymptotic comparison between these metrics. Given two left-invariant subRiemannian metrics d and d' on \mathbb{H} , we deal with three asymptotic behaviors, in ascending order of strength, each of which defines an equivalence relation among subRiemannian metrics:

- (i) $\lim_{d(p,q) \rightarrow \infty} \frac{d(p,q)}{d'(p,q)} = 1$;
- (ii) There exists $c > 0$ such that $|d(p, q) - d'(p, q)| < c$, for all p, q ;
- (iii) $\lim_{d(p,q) \rightarrow \infty} |d(p, q) - d'(p, q)| = 0$.

A first example of the implication (i) \Rightarrow (ii) was proved by Burago in [Bur94] for \mathbb{Z}^n -invariant metrics d on \mathbb{R}^n , by showing that d and the associated stable norm stay at bounded distance from each other. This result has been extended quantitatively for \mathbb{Z}^n -invariant metrics on geodesic metric spaces in [CS14]. Gromov and Burago asked for other interesting cases where the same implication holds. Another well-known case where (i) is equivalent to (ii) is that of hyperbolic groups. Beyond Abelian and hyperbolic groups, Krat proved the equivalence for word metrics on the discrete Heisenberg group $\mathbb{H}(\mathbb{Z})$, [Kra99]. For general subFinsler metrics on Carnot groups it has been proven in [BLD13], following [Sto98], that (i) is equivalent to the fact that the projections onto $\mathbb{H}/[\mathbb{H}, \mathbb{H}]$ of the corresponding unit balls coincide, see (c) below. Our first result shows that this last condition is equivalent to each one of (i) and (ii) in the case of the Heisenberg group endowed with subRiemannian metrics.

Theorem 1.2. *Let d and d' be two left-invariant subRiemannian metrics on \mathbb{H} whose horizontal spaces are (V, g) and (V', g') respectively.*

Let $\pi : \mathfrak{h} \rightarrow \mathfrak{h}/[\mathfrak{h}, \mathfrak{h}]$ be the quotient projection and $\hat{\pi} : \mathbb{H} \rightarrow \mathfrak{h}/[\mathfrak{h}, \mathfrak{h}]$ the corresponding group morphism.

Then the following assertions are equivalent:

- (a) there exists $c > 0$ such that $|d(p, q) - d'(p, q)| < c$, for all p, q ;
- (b) $\frac{d(p, q)}{d'(p, q)} \rightarrow 1$ when $d(p, q) \rightarrow \infty$;
- (c) $\hat{\pi}(\{p \in \mathbb{H} : d(0, p) \leq R\}) = \hat{\pi}(\{p \in \mathbb{H} : d'(0, p) \leq R\})$, for all $R > 0$, here 0 denotes the neutral element of \mathbb{H} ;
- (d) $\pi(\{v \in V : g(v, v) \leq 1\}) = \pi(\{v' \in V' : g'(v', v') \leq 1\})$;
- (e) there exists a scalar product \bar{g} on $\mathfrak{h}/[\mathfrak{h}, \mathfrak{h}]$ such that both $\pi|_V : (V, g) \rightarrow (\mathfrak{h}/[\mathfrak{h}, \mathfrak{h}], \bar{g})$ and $\pi|_{V'} : (V', g') \rightarrow (\mathfrak{h}/[\mathfrak{h}, \mathfrak{h}], \bar{g})$ are submetrics, i.e., they map balls to balls.

Next, we prove that in every class of the equivalence relation (iii) there is exactly one strictly subRiemannian metric. To every left-invariant subRiemannian metric d we define the *associated asymptotic metric* d' as follows. If d is Riemannian defined by a scalar product g on \mathfrak{h} , then d' is the strictly subRiemannian metric for which the horizontal space V is g -orthogonal to $[\mathfrak{h}, \mathfrak{h}]$ and the scalar product is $g|_V$. If d is strictly subRiemannian, then $d' = d$.

Theorem 1.3. *Let d and d' be two left-invariant subRiemannian metrics on \mathbb{H} . Their associated asymptotic metrics are the same if and only if*

$$(1) \quad \lim_{d(p, q) \rightarrow \infty} |d(p, q) - d'(p, q)| = 0.$$

Moreover, if (1) holds, then there is $C > 0$ such that

$$(2) \quad |d(p, q) - d'(p, q)| \leq \frac{C}{d(p, q)}, \quad \forall p, q \in \mathbb{H}.$$

We remark that the estimate (2) in Theorem 1.3 is sharp, as we will show in Remark 4.2.

The above result can be interpreted in terms of asymptotic cones. Namely, if d is a left-invariant Riemannian metric on \mathbb{H} and d' is the associated asymptotic metric, then (\mathbb{H}, d') is the asymptotic cone of (\mathbb{H}, d) . For the analogous result in arbitrary nilpotent groups see [Pan83]. By Theorem 1.3, more is true: (\mathbb{H}, d) is at bounded distance from (\mathbb{H}, d') . Notice that this consequence cannot be deduced by the similar results for discrete subgroups of the Heisenberg group in [Kra99] and [DS14], because the word metric is only quasi-isometric to the Riemannian one. Moreover, we remark that there are examples of nilpotent groups of step two that are not at bounded distance from their asymptotic cone, see [BLD13].

We now focus on the horoboundary. As a consequence of Theorem 1.3 and of the results of Klein-Nikas [KN10], we get:

Corollary 1.4. *If d_R is a left-invariant Riemannian metric on \mathbb{H} with associated asymptotic metric d_{CC} , then the horoboundary of (\mathbb{H}, d_R) coincides with the horoboundary of (\mathbb{H}, d_{CC}) ; hence, it is homeomorphic to a 2-dimensional closed disk \bar{D}^2 .*

More precisely, let g be the scalar product of d_R on \mathfrak{h} and $W \subset \mathfrak{h}$ the orthogonal plane to $[\mathfrak{h}, \mathfrak{h}]$. Define the norm $\|w\| := \sqrt{g(w, w)}$ on W . Fix a orthonormal basis (X, Y) for W and set $Z := [X, Y] \in [\mathfrak{h}, \mathfrak{h}]$, so that (X, Y, Z) is a basis of \mathfrak{h} . We identify $\mathfrak{h} \simeq \mathbb{H}$ via the exponential map, which is a global diffeomorphism. So, we write $p = w + zZ$ with $w \in W$ and $z \in \mathbb{R}$ for any point $p \in \mathbb{H}$. We say that a sequence of points $\{p_n\}_{n \in \mathbb{N}} \subset \mathbb{H}$ diverges if it leaves every compact set. Moreover, we shall use the following terminology for a diverging sequence of the form $p_n = w_n + z_n Z$:

- (1) *vertical divergence*, if there exists $M < \infty$ such that $\|w_n\| < M$ for all n ;
- (2) *non-vertical divergence with quadratic rate* $\nu \in [-\infty, +\infty]$, if w_n diverges and¹ $\lim_{n \rightarrow \infty} \frac{z_n}{4\|w_n\|^2} = -\nu$.

Then, according to [KN10] (see Corollary 5.6, 5.9 and 5.13 therein), we deduce the following description of the Riemannian horofunctions:

(v): a vertically diverging sequence $p_n = w_n + z_n Z$ converges to a horofunction h if and only if $w_n \rightarrow w_\infty$, and in this case

$$h(w + zZ) = \|w_\infty\| - \|w_\infty - w\|;$$

(nv): a non-vertically diverging sequence $p_n = w_n + z_n Z$ with quadratic rate ν converges to a horofunction h if and only if $\frac{w_n}{\|w_n\|} \rightarrow \hat{w}$, and then

$$h(w + zZ) = g(R_\vartheta(-\hat{w}), w)$$

where R_ϑ is the anti-clockwise rotation in W of angle $\vartheta = \mu^{-1}(\nu)$, and $\mu : [-\pi, \pi] \rightarrow \overline{\mathbb{R}}$ is the extended Gaveau function

$$\mu(\vartheta) := \frac{\vartheta - \sin \vartheta \cos \vartheta}{\sin^2(\vartheta)}.$$

Moreover, all the horofunctions of (\mathbb{H}, d_R) are of type **(v)** or **(nv)**, by Theorem 5.16 in [KN10]; it is also clear that neither is of both types.

In section 5 we will also determine the Busemann points of $\partial_h(\mathbb{H}, d_R)$, that is those horofunctions obtained by points diverging along quasi-geodesics (see Definition 5.1). We obtain, as in the subRiemannian case:

Corollary 1.5. *The Busemann points of (\mathbb{H}, d_R) are the horofunctions of type **(nv)** and can be identified to the boundary of the disk \bar{D}^2 .*

¹ From the paper [KN10], there is an extra 4 and a change of sign due to our different choice of coordinates.

The paper is organized as follows. In Section 2 we introduce the main objects and their basic properties. In Section 3 we estimate the difference between any two strictly subRiemannian left-invariant metrics on \mathbb{H} . In Section 4 we compare any Riemannian left-invariant metric on \mathbb{H} and its associated asymptotic metric. At the end of the section we shall prove Theorems 1.2 and 1.3. In Section 5 we concentrate on the horofunctions and we prove Corollaries 1.4 and 1.5. Appendix A is devoted to the explicit description of subRiemannian geodesics.

Acknowledgments. The initial discussions for this work were done at the ‘2013 Workshop on Analytic and Geometric Group Theory’ in Ventotene. We express our gratitude to the organizers: A. Iozzi, G. Kuhn and M. Sageev. We also thank the anonymous referee for several improving suggestions.

2. PRELIMINARIES

2.1. Definitions. The *first Heisenberg group* \mathbb{H} is the connected, simply connected Lie group associated to the Heisenberg Lie algebra \mathfrak{h} . The *Heisenberg Lie algebra* \mathfrak{h} is the only three dimensional nilpotent Lie algebra that is not commutative. It can be proven that, for any two linearly independent vectors $X, Y \in \mathfrak{h} \setminus [\mathfrak{h}, \mathfrak{h}]$, the triple $(X, Y, [X, Y])$ is a basis of \mathfrak{h} and $[X, [X, Y]] = [Y, [X, Y]] = 0$.

We denote by $\omega_{\mathbb{H}} : T\mathbb{H} \rightarrow \mathfrak{h}$ the left-invariant Maurer-Cartan form. Namely, denoting by 0 the neutral element of \mathbb{H} and identifying \mathfrak{h} with $T_0\mathbb{H}$, we have $\omega_{\mathbb{H}}(v) := dL_p^{-1}v$ for $v \in T_p\mathbb{H}$, where L_p is the left translation by p .

Let $\pi : \mathfrak{h} \rightarrow \mathfrak{h}/[\mathfrak{h}, \mathfrak{h}]$ be the quotient projection. Notice that $\mathfrak{h}/[\mathfrak{h}, \mathfrak{h}]$ is a commutative 2-dimensional Lie algebra. So the map π induces a Lie group epimorphism $\hat{\pi} : \mathbb{H} \rightarrow \mathfrak{h}/[\mathfrak{h}, \mathfrak{h}] \simeq \mathbb{H}/[\mathbb{H}, \mathbb{H}]$.

2.2. SubRiemannian metrics in \mathbb{H} . Let $V \subset \mathfrak{h}$ be a bracket generating subspace. We have only two cases: either $V = \mathfrak{h}$ or V is a plane and $\mathfrak{h} = V \oplus [\mathfrak{h}, \mathfrak{h}]$. In both cases the restriction of the projection $\pi|_V : V \rightarrow \mathfrak{h}/[\mathfrak{h}, \mathfrak{h}]$ is surjective. Let g be a scalar product on V and set the corresponding norm $\|v\| := \sqrt{g(v, v)}$ for $v \in V$.

An absolutely continuous curve $\gamma : [0, 1] \rightarrow \mathbb{H}$ is said *horizontal* if $\omega_{\mathbb{H}}(\gamma'(t)) \in V$ for almost every t . For a horizontal curve we have the *length*

$$\ell(\gamma) := \int_0^1 \|\omega_{\mathbb{H}}(\gamma'(t))\| dt.$$

A *subRiemannian metric* d is hence defined as

$$d(p, q) := \inf \{ \ell(\gamma) : \gamma \text{ horizontal curve from } p \text{ to } q \}.$$

SubRiemannian metrics on \mathbb{H} are complete, geodesic, and left-invariant. They are either Riemannian, when $V = \mathfrak{h}$, or *strictly subRiemannian*, when $\dim V = 2$. The pair (V, g) is called the *horizontal space of d* .

Since $\pi|_V : V \rightarrow \mathfrak{h}/[\mathfrak{h}, \mathfrak{h}]$ is surjective, it induces a norm $\|\cdot\|$ on $\mathfrak{h}/[\mathfrak{h}, \mathfrak{h}]$ such that $\pi : (V, \|\cdot\|) \rightarrow (\mathfrak{h}/[\mathfrak{h}, \mathfrak{h}], \|\cdot\|)$ is a submetry, i.e., for all $w \in \mathfrak{h}/[\mathfrak{h}, \mathfrak{h}]$ it holds $\|w\| = \inf\{\|v\| : \pi(v) = w\}$. Here we use the same notation for norms on V and on $\mathfrak{h}/[\mathfrak{h}, \mathfrak{h}]$, because there will be no possibility of confusion. The norm on $\mathfrak{h}/[\mathfrak{h}, \mathfrak{h}]$ is characterized by

$$(3) \quad \pi(\{v \in V : \|v\| \leq R\}) = \{w \in \mathfrak{h}/[\mathfrak{h}, \mathfrak{h}] : \|w\| \leq R\},$$

for all $R > 0$.

Proposition 2.1. *Let d be subRiemannian metric on \mathbb{H} with horizontal space (V, g) . Then for all $R > 0$*

$$\pi(\{v \in V : \|v\| \leq R\}) = \hat{\pi}(\{p \in \mathbb{H} : d(0, p) \leq R\}).$$

In particular, $\hat{\pi} : (\mathbb{H}, d) \rightarrow (\mathfrak{h}/[\mathfrak{h}, \mathfrak{h}], \|\cdot\|)$ is a submetry, i.e., for all $v, w \in \mathfrak{h}/[\mathfrak{h}, \mathfrak{h}]$

$$\|v - w\| = \inf\{d(p, q) : \hat{\pi}(p) = v, \hat{\pi}(q) = w\}.$$

Proof. \square Let $v \in V$ with $\|v\| \leq R$. Set $\gamma(t) := \exp(tv)$. Then $\gamma : [0, 1] \rightarrow \mathbb{H}$ is a horizontal curve with $d(0, \exp(v)) \leq \ell(\gamma) = \|v\| \leq R$. Since $\hat{\pi}(\exp(v)) = \pi(v)$, then we have proven this inclusion.

\square Let $p \in \mathbb{H}$ with $d(0, p) \leq R$ and let $\gamma : [0, T] \rightarrow \mathbb{H}$ be a d -length-minimizing curve from 0 to p parametrized by arc-length, so $T = d(0, p)$. Then $\hat{\pi} \circ \gamma : [0, T] \rightarrow \mathfrak{h}/[\mathfrak{h}, \mathfrak{h}]$ is a curve from 0 to $\pi(p)$ and

$$\begin{aligned} \|\hat{\pi}(p)\| &\leq \int_0^T \|(\hat{\pi} \circ \gamma)'(t)\| dt \\ &= \int_0^T \|\pi \circ \omega_{\mathbb{H}}(\gamma'(t))\| dt \\ &\leq \int_0^T \|\omega_{\mathbb{H}}(\gamma'(t))\| dt \\ &= \ell(\gamma) = d(0, p). \end{aligned}$$

In the first equality we used the fact that $\hat{\pi}$ is a morphism of Lie groups and its differential is π , i.e., $\omega_{\mathbb{H}/[\mathbb{H}, \mathbb{H}]} \circ d\hat{\pi} = \pi \circ \omega_{\mathbb{H}}$, where $\omega_{\mathbb{H}/[\mathbb{H}, \mathbb{H}]}$ is the Maurer-Cartan form of $\mathbb{H}/[\mathbb{H}, \mathbb{H}]$. \square

Proposition 2.2. *Let d, d' be two subRiemannian metrics on \mathbb{H} such that*

$$\lim_{p \rightarrow \infty} \frac{d(0, p)}{d'(0, p)} = 1.$$

Then

$$(4) \quad \hat{\pi}(\{p \in \mathbb{H} : d(0, p) \leq R\}) = \hat{\pi}(\{p \in \mathbb{H} : d'(0, p) \leq R\}).$$

Proof. Let $\|\cdot\|$ and $\|\cdot\|'$ be the norms on $\mathfrak{h}/[\mathfrak{h}, \mathfrak{h}]$ induced by d and d' , respectively. We will show that

$$(5) \quad \lim_{v \rightarrow \infty} \frac{\|v\|}{\|v\|'} = 1,$$

which easily implies $\|\cdot\| = \|\cdot\|'$, because for any fixed $v \in \mathfrak{h}/[\mathfrak{h}, \mathfrak{h}]$ one has $1 = \lim_{t \rightarrow \infty} \frac{\|tv\|}{\|tv\|'} = \frac{\|v\|}{\|v\|'}$. Moreover, the equality (4) follows from Proposition 2.1 combined with (3) and (5).

Since both maps $\hat{\pi} : (\mathbb{H}, d) \rightarrow (\mathfrak{h}/[\mathfrak{h}, \mathfrak{h}], \|\cdot\|)$ and $\hat{\pi} : (\mathbb{H}, d') \rightarrow (\mathfrak{h}/[\mathfrak{h}, \mathfrak{h}], \|\cdot\|')$ are submetries, for every $v \in \mathfrak{h}/[\mathfrak{h}, \mathfrak{h}]$ there are $p_v, p'_v \in \mathbb{H}$ such that $\hat{\pi}(p_v) = \hat{\pi}(p'_v) = v$, $\|v\| = d(0, p_v)$ and $\|v\|' = d'(0, p'_v)$.

Moreover it holds $\|v\|' \leq d'(0, p_v)$ and $\|v\| \leq d(0, p'_v)$, again because $\hat{\pi}$ is a submetry in both cases. Therefore

$$\frac{d(0, p_v)}{d'(0, p_v)} \leq \frac{\|v\|}{\|v\|'} \leq \frac{d(0, p'_v)}{d'(0, p'_v)}$$

Finally, if $v \rightarrow \infty$, then both $d(0, p_v)$ and $d(0, p'_v)$ go to infinity as well. The relation (5) is thus proven. \square

2.3. Balayage area and lifting of curves. Let $V \subset \mathfrak{h}$ be a two-dimensional subspace with $V \cap [\mathfrak{h}, \mathfrak{h}] = \{0\}$. Then $[\mathfrak{h}, \mathfrak{h}] = [V, V]$, i.e., V is bracket generating. Moreover, $\pi|_V : V \rightarrow \mathfrak{h}/[\mathfrak{h}, \mathfrak{h}]$ is an isomorphism.

If $\rho : [0, T] \rightarrow \mathfrak{h}/[\mathfrak{h}, \mathfrak{h}]$ is a curve with $\rho(0) = 0$, then there is a unique $\tilde{\rho} : [0, T] \rightarrow \mathbb{H}$ such that

$$\begin{cases} \tilde{\rho}(0) = 0, \\ \omega_{\mathbb{H}}(\tilde{\rho}'(t)) = \pi|_V^{-1}(\rho(t)'). \end{cases}$$

Since $(\pi \circ \tilde{\rho})' = \rho'$, then $\pi \circ \tilde{\rho} = \rho$. So, $\tilde{\rho}$ is called the *lift* of ρ .

The previous ODE system that defines $\tilde{\rho}$ can be easily integrated. Let $X, Y \in V$ be a basis, set $Z := [X, Y]$, so that (X, Y, Z) is a basis of \mathfrak{h} . Let $(x, y, z) = \exp(xX + yY + zZ)$ be the exponential coordinates on \mathbb{H} defined by (X, Y, Z) . Using the Baker-Campbell-Hausdorff formula, one shows that X, Y, Z induce the following left-invariant vector fields on \mathbb{H} :

$$\hat{X} = \partial_x - \frac{y}{2}\partial_z, \quad \hat{Y} = \partial_y + \frac{x}{2}\partial_z, \quad \hat{Z} = \partial_z.$$

Thanks to these vector fields, we can describe the Maurer-Cartan form as

$$\omega_{\mathbb{H}}(a\hat{X} + b\hat{Y} + c\hat{Z}) = aX + bY + cZ.$$

The lift of ρ is hence given by the ODE

$$\begin{cases} \tilde{\rho}'_1 = \rho'_1, \\ \tilde{\rho}'_2 = \rho'_2, \\ \tilde{\rho}'_3 = \frac{1}{2}(\rho_1\rho'_2 - \rho_2\rho'_1). \end{cases}$$

Take the coordinates (x, y) on $\mathfrak{h}/[\mathfrak{h}, \mathfrak{h}]$ given by the basis $(\pi(X), \pi(Y))$ and define the *balayage area* of a curve $\rho : [0, T] \rightarrow \mathfrak{h}/[\mathfrak{h}, \mathfrak{h}]$ as

$$(6) \quad \mathcal{A}(\rho) = \frac{1}{2} \int_{\rho} (x \, dy - y \, dx).$$

If $\rho(0) = 0$, then the balayage area of ρ corresponds to the signed area enclosed between the curve ρ and the line passing through 0 and $\rho(T)$.

It follows that

$$\tilde{\rho}(t) = (\rho_1(t), \rho_2(t), \mathcal{A}(\rho|_0^t)).$$

In an implicit form we can write

$$(7) \quad \tilde{\rho}(t) = \exp((\pi|_V)^{-1}(\rho(t)) + \mathcal{A}(\rho|_0^t)Z).$$

Notice that the lift $\tilde{\rho}$ of a curve ρ depends on the choice of V . Moreover, both the area and the Balayage area in $\mathfrak{h}/[\mathfrak{h}, \mathfrak{h}]$ depend on the choice of the basis (X, Y) . Nevertheless, once a plane $V \subset \mathfrak{h}$ is fixed, the lift $\tilde{\rho}$ does not depend on the choice of the basis X, Y .

If g is a scalar product on V and d is the corresponding strictly subRiemannian metric, the balayage area gives a characterization of d -length-minimizing curves. Let \bar{g} be the scalar product on $\mathfrak{h}/[\mathfrak{h}, \mathfrak{h}]$ induced by g . Then the d -length of a curve $\tilde{\rho} : [0, T] \rightarrow \mathbb{H}$ equals the length of $\rho = \pi \circ \tilde{\rho}$.

Therefore, given $p = (x, y, z) \in \mathbb{H}$, we have

$$d(0, p) = \inf \{ \ell(\rho) : \rho : [0, 1] \rightarrow \mathfrak{h}/[\mathfrak{h}, \mathfrak{h}], \rho(0) = 0, \rho(1) = \hat{\pi}(p), \mathcal{A}(\rho) = z \}.$$

This express the so-called *Dido's problem* in the plane, and the solutions are arc of circles. It degenerates into a line if $z = 0$. We can summarize the last discussion in the following result.

Lemma 2.3. *A curve $\tilde{\rho} : [0, 1] \rightarrow \mathbb{H}$ is d -length-minimizing from 0 to $p = (x, y, z)$ if and only if $\rho := \hat{\pi} \circ \tilde{\rho}$ is an arc of a circle from 0 to $\hat{\pi}(p)$ with $\mathcal{A}(\rho) = z$.*

3. COMPARISON BETWEEN STRICTLY SUBRIEMANNIAN METRICS

The present section is devoted to comparing strictly subRiemannian metrics. For such metrics, Proposition 3.1 gives the only non-trivial implication in Theorem 1.2. The general case will follow from Proposition 4.1.

Proposition 3.1. *Let d and d' be two strictly subRiemannian metrics on \mathbb{H} with horizontal spaces (V, g) and (V', g') , respectively. Suppose that there exists a scalar product \bar{g} on $\mathfrak{h}/[\mathfrak{h}, \mathfrak{h}]$ such that both*

$$\pi|_V : (V, g) \rightarrow (\mathfrak{h}/[\mathfrak{h}, \mathfrak{h}], \bar{g}) \quad \text{and} \quad \pi|_{V'} : (V', g') \rightarrow (\mathfrak{h}/[\mathfrak{h}, \mathfrak{h}], \bar{g})$$

are submetries.

Then

$$(8) \quad \sup_{p \in \mathbb{H}} |d(0, p) - d'(0, p)| < \infty.$$

Moreover, if $d \neq d'$, then

$$(9) \quad \limsup_{p \rightarrow \infty} |d(0, p) - d'(0, p)| > 0.$$

In the proof we will give the exact value of the supremum in (8). Indeed, by (11) and (12), we get $\sup_{p \in \mathbb{H}} |d(0, p) - d'(0, p)| = 2|h|$, where h is defined below. For (8), we will first prove that two of such subRiemannian metrics are isometric via a conjugation $x \mapsto gxg^{-1}$ for some $g \in \mathbb{H}$ and then we apply Lemma 3.2. For (9), we will give a sequence $p_n \rightarrow \infty$ and a constant $c > 0$ such that $|d(0, p_n) - d'(0, p_n)| > c$ for all $n \in \mathbb{N}$.

3.1. Proof of (8). Since $\dim V = \dim V' = 2$, then $\pi|_V$ and $\pi|_{V'}$ are isomorphisms. Therefore by the assumption they are isometries onto $(\mathfrak{h}/[\mathfrak{h}, \mathfrak{h}], \bar{g})$.

Let $X \in V \cap V'$ be with $g(X, X) = 1$. Then $g'(X, X) = 1$ as well.

Let $Y \in V$ be orthogonal to X with $g(Y, Y) = 1$. Then $Z := [X, Y] \neq 0$ and (X, Y, Z) is a basis of \mathfrak{h} .

Let $Y' := \pi|_{V'}^{-1}(\pi(Y)) \in V'$. Then $g'(Y', Y') = 1$ and $g'(X, Y') = 0$. Moreover, there is $h \in \mathbb{R}$ such that $Y' = Y + hZ$. In particular, $[X, Y'] = Z$.

Using the formula $\text{Ad}_{\exp(hX)}(v) = e^{\text{ad}_{hX}}v = v + h[X, v]$, we notice that

$$(10) \quad \begin{cases} \text{Ad}_{\exp(hX)}(X) = X, \\ \text{Ad}_{\exp(hX)}(Y) = Y', \\ \text{Ad}_{\exp(hX)}(Z) = Z. \end{cases}$$

In particular $\text{Ad}_{\exp(hX)}|_V : (V, g) \rightarrow (V', g')$ is an isometry.

Therefore, the conjugation

$$C_{\exp(hX)}(p) := \exp(hX) \cdot p \cdot \exp(hX)^{-1}$$

is an isometry $C_{\exp(hX)} : (\mathbb{H}, d) \rightarrow (\mathbb{H}, d')$.

We can now use the following Lemma 3.2 and get

$$(11) \quad \sup_{p \in \mathbb{H}} |d(0, p) - d'(0, p)| \leq 2|h|.$$

Lemma 3.2. *Let G be a group with neutral element e and let d, d' be two left-invariant distances on G . If there is $g \in G$ such that for all $p \in G$*

$$d'(e, p) = d(e, gpg^{-1}),$$

then for all $p \in G$

$$|d(e, p) - d'(e, p)| \leq 2 \min\{d(e, g), d'(e, g)\}.$$

Proof. Note that, since d is left invariant, then for all $a, b \in G$ we have $d(e, ab) \leq d(e, a) + d(e, b)$ and $d(e, a) = d(e, a^{-1})$. On the one side, we have $d'(e, p) = d(e, gpg^{-1}) \leq d(e, p) + 2d(e, g)$. On the other side, we have $d(e, p) = d(e, g^{-1}gpg^{-1}g) \leq d(e, g^{-1}) + d(e, gpg^{-1}) + d(e, g) = 2d(e, g) + d'(e, p)$. Hence $|d(e, p) - d'(e, p)| \leq 2d(e, g)$. By symmetry, we have also $|d(e, p) - d'(e, p)| \leq 2d'(e, g)$. \square

3.2. Proof of (9). We keep the same notation of the previous subsection. Up to switching V with V' , we can assume $h > 0$.

Let (x, y, z) be the exponential coordinates on \mathbb{H} induced by the basis (X, Y, Z) of \mathfrak{h} , i.e., $(x, y, z) = \exp(xX + yY + zZ) \in \mathbb{H}$. Similarly, on $\mathfrak{h}/[\mathfrak{h}, \mathfrak{h}]$ we have coordinates $(x, y) = x\pi(X) + y\pi(Y)$.

For $R > 0$, define

$$p_R := \left(0, 2R, \frac{\pi R^2}{2} + 2hR \right).$$

We will show that

$$(12) \quad \lim_{R \rightarrow \infty} d(0, p_R) - d'(0, p_R) = 2h.$$

Fix $R > 0$. Let $\gamma : [0, T] \rightarrow \mathbb{H}$ be a d' -minimizing curve from 0 to p_R . Then $\hat{\pi} \circ \gamma : [0, T] \rightarrow \mathfrak{h}/[\mathfrak{h}, \mathfrak{h}]$ is half circle of center $(0, R)$ and radius R . The balayage area of $\hat{\pi} \circ \gamma$ is

$$\mathcal{A}(\hat{\pi} \circ \gamma) = \frac{\pi R^2}{2}.$$

Let $\eta : [0, T] \rightarrow \mathbb{H}$ be the d -length-minimizing curve from 0 to p_R . Then $\hat{\pi} \circ \eta : [0, T] \rightarrow \mathfrak{h}/[\mathfrak{h}, \mathfrak{h}]$ is an arc of a circle of radius S_R whose




FIGURE 1. Curves in $\mathfrak{h}/[\mathfrak{h}, \mathfrak{h}]$ for the proof of (9).

balayage area is

$$(13) \quad \mathcal{A}(\hat{\pi} \circ \eta) = \frac{\pi R^2}{2} + 2Rh = \mathcal{A}(\hat{\pi} \circ \gamma) + 2Rh.$$

It is clear that $S_R > R$ and that the circle of $\hat{\pi} \circ \eta$ has center (μ_R, R) for some $\mu_R > 0$. So we have

$$S_R^2 = R^2 + \mu_R^2.$$

It is also clear from the picture that

$$(14) \quad \frac{\pi S_R^2}{2} + 2R\mu_R \leq \mathcal{A}(\hat{\pi} \circ \eta) \leq \frac{\pi S_R^2}{2} + 2S_R\mu_R.$$

Now, let's look at the lengths. First of all, notice that $\ell_{d'}(\gamma) = \ell(\hat{\pi} \circ \gamma)$ and $\ell_d(\eta) = \ell(\hat{\pi} \circ \eta)$. For one curve we have

$$\ell(\hat{\pi} \circ \gamma) = \pi R,$$

for the other we have the estimate

$$\pi S_R + 2\mu_R \leq \ell(\hat{\pi} \circ \eta),$$

which is clear from the picture. Hence

$$\begin{aligned} \liminf_{R \rightarrow \infty} d(0, p) - d'(0, p) &= \liminf_{R \rightarrow \infty} \ell(\hat{\pi} \circ \eta) - \ell(\hat{\pi} \circ \gamma) \\ &\geq \lim_{R \rightarrow \infty} \pi S_R + 2\mu_R - \pi R \\ &= \lim_{R \rightarrow \infty} \pi(S_R - R) + 2\mu_R. \end{aligned}$$

We claim that

$$(15) \quad \lim_{R \rightarrow \infty} \pi(S_R - R) + 2\mu_R = 2h.$$

Let us start by checking that,

$$(16) \quad \mu_R < h.$$

Indeed, from the first inequality of (14) together with (13) it follows

$$\frac{\pi S_R^2}{2} + 2R\mu_R \leq \frac{\pi R^2}{2} + 2hR.$$

Since $S_R > R$, then

$$0 < \frac{\pi S_R^2}{2} - \frac{\pi R^2}{2} \leq 2R(h - \mu_R),$$

i.e., the inequality (16).

From the second inequality of (14) together with (13) we get

$$\frac{\pi R^2}{2} + 2Rh \leq \frac{\pi S_R^2}{2} + 2S_R\mu_R.$$

Using the facts $\mu_R \leq h$ and $S_R \leq R + \mu_R \leq R + h$, from the above inequality one gets

$$(17) \quad \begin{aligned} 0 \leq 2(h - \mu_R) &\leq (S_R - R) \frac{1}{R} \left(\frac{\pi}{2} (S_R + R) + 2\mu_R \right) \\ &\leq (S_R - R) \left(\pi + \frac{h}{R} \left(\frac{\pi}{2} + 2 \right) \right) \end{aligned}$$

Moreover, since $h^2 \geq \mu_R^2 = S_R^2 - R^2 = (S_R - R)(S_R + R)$, we also have

$$(18) \quad \lim_{R \rightarrow \infty} (S_R - R) = 0.$$

Finally, from (18) and (17) we obtain (15), as claimed. This completes the proof of (12) and of Proposition 3.1.

4. COMPARISON BETWEEN RIEMANNIAN AND STRICTLY SUBRIEMANNIAN METRICS

Let d_R be a Riemannian metric on \mathbb{H} with horizontal space (\mathfrak{h}, g) .

Let $V \subset \mathfrak{h}$ be the plane orthogonal to $[\mathfrak{h}, \mathfrak{h}]$ and let d_{CC} be the strictly subRiemannian metric on \mathbb{H} with horizontal space $(V, g|_V)$.

Fix a basis (X, Y, Z) for \mathfrak{h} such that (X, Y) is an orthonormal basis of $(V, g|_V)$ and $Z = [X, Y]$. The matrix representation of g with respect to (X, Y, Z) is

$$g = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & \zeta^2 \end{pmatrix}$$

where $\zeta > 0$.

Let d_{CC} be the strictly subRiemannian metric on \mathbb{H} with horizontal space $(V, g|_V)$.

Our aim in this section is to prove the following proposition.

Proposition 4.1. *If $d_{CC}(0, p)$ is large enough, then:*

$$(19) \quad 0 \leq d_{CC}(0, p) - d_R(0, p) \leq \frac{4\pi^2}{\zeta^2} \frac{1}{d_{CC}(0, p) - \frac{2^{3/2}\pi}{\zeta}}.$$

In particular it holds

$$(20) \quad \lim_{p \rightarrow \infty} |d_{CC}(0, p) - d_R(0, p)| = 0.$$

For the proof of this statement, we need to know length-minimizing curves for d_R and d_{CC} , and a few properties of those, see the exposition in the Appendix A.

Proof. Let (x, y, z) be the exponential coordinates on \mathbb{H} induced by the basis (X, Y, Z) of \mathfrak{h} , i.e., $(x, y, z) = \exp(xX + yY + zZ) \in \mathbb{H}$. Fix $p = (p_1, p_2, p_3) \in \mathbb{H}$.

Notice that both d_R and d_{CC} are generated as length metrics using the same length measure ℓ , with the difference that d_R minimizes the

length among all the curves, while d_{CC} takes into account only the curves tangent to V . This implies that

$$\forall p, q \in \mathbb{H} \quad d_{CC}(p, q) \geq d_R(p, q),$$

therefore we get the first inequality in (19). We need to prove the second inequality of (19).

If $p \in \{z = 0\}$, then $d_{CC}(0, p) = d_R(0, p)$ by Corollary A.7, and the claim is true.

Suppose $p \notin \{z = 0\}$ and let $\gamma : [0, T] \rightarrow \mathbb{H}$ be a d_R -length minimizing curve from $0 = \gamma(0)$ to $p = \gamma(T)$. Since $p \notin \{z = 0\}$ and since we supposed that $d_{CC}(0, p)$ is large enough, then by Corollary A.5 we can parametrize γ in such a way that γ is exactly in the form expressed in TYPE II in Proposition A.2 for some $k > 0$ and $\theta \in \mathbb{R}$.

By Corollary A.4 it holds

$$(21) \quad kT \leq 2\pi.$$

Moreover, by Corollary A.8

$$(22) \quad d_R(0, p) = \|\omega_{\mathbb{H}}(\gamma')\| \cdot T = \sqrt{1 + \frac{k^2}{\zeta^2}} \cdot T.$$

Let $\eta : [0, T] \rightarrow \mathbb{H}$ be the d_{CC} -length-minimizing curve corresponding to γ as shown in Corollary A.8. Then we know that $d_{CC}(0, \eta(T)) = \ell(\eta) = T$, and

$$(23) \quad p = \gamma(T) = \eta(T) + (0, 0, \frac{kT}{\zeta^2}).$$

Hence by Corollary A.8 and (21)

$$d_{CC}(0, p) \leq d_{CC}(0, \eta(T)) + d_{CC}(\eta(T), \gamma(T)) \leq T + \frac{2^{3/2}\pi}{\zeta},$$

i.e.,

$$(24) \quad \frac{1}{T} \leq \frac{1}{d_{CC}(0, p) - \frac{2^{3/2}\pi}{\zeta}}.$$

Since η is a d_{CC} -rectifiable curve, then $\eta(T)_3 = \mathcal{A}(\hat{\pi} \circ \eta)$, where $\eta(T)_3$ is the third coordinate of the point in the exponential coordinates. Since $\hat{\pi} \circ \gamma = \hat{\pi} \circ \eta$, then we have by (23)

$$(25) \quad p_3 = \mathcal{A}(\hat{\pi} \circ \gamma) + \frac{kT}{\zeta^2}.$$

Notice that $\hat{\pi} \circ \gamma$ is an arc of a circle in $\mathfrak{h}/[\mathfrak{h}, \mathfrak{h}]$ of radius $\frac{1}{k}$, see Proposition A.2.

Now we want to define a horizontal curve $\tilde{\rho} : [-\epsilon, T + \epsilon] \rightarrow \mathbb{H}$, where $\epsilon > 0$ has to be chosen, such that $\tilde{\rho}(-\epsilon) = 0$ and $\tilde{\rho}(T + \epsilon) = p$. We first define a curve $\rho : [-\epsilon, T + \epsilon] \rightarrow \mathfrak{h}/[\mathfrak{h}, \mathfrak{h}]$ and then take its lift to \mathbb{H} .




FIGURE 2. Curves for Case 1 and Case 2.

For the definition of ρ we follow two different strategies for two different cases:

CASE 1. Suppose that $\hat{\pi} \circ \gamma$ doesn't cover the half of the circle, i.e., $T \leq \frac{\pi}{k}$. Set $\lambda = \hat{\pi}(p) \in \mathfrak{h}/[\mathfrak{h}, \mathfrak{h}]$. Then T is smaller than the circle of diameter $\|\lambda\|$, i.e.,

$$(26) \quad \|\lambda\| \geq \frac{T}{\pi}.$$

Let $\lambda^\perp \in \mathfrak{h}/[\mathfrak{h}, \mathfrak{h}]$ be the unit vector perpendicular to λ and forming an angle smaller than $\pi/2$ with the arc $\hat{\pi} \circ \gamma$. Let $\epsilon > 0$ such that

$$(27) \quad \epsilon \cdot \|\lambda\| = \frac{kT}{\zeta^2}.$$

Now, define $\rho : [-\epsilon, T + \epsilon] \rightarrow \mathfrak{h}/[\mathfrak{h}, \mathfrak{h}]$ as

$$\rho(t) = \begin{cases} (t + \epsilon)\lambda^\perp & \text{for } -\epsilon \leq t \leq 0 \\ \epsilon\lambda^\perp + \hat{\pi} \circ \gamma(t) & \text{for } 0 \leq t \leq T \\ \epsilon\lambda^\perp + \hat{\pi} \circ \gamma(T) - (t - T)\lambda^\perp & \text{for } T \leq t \leq T + \epsilon \end{cases}$$

Notice that

$$\mathcal{A}(\rho) = \mathcal{A}(\hat{\pi} \circ \gamma) + \epsilon \cdot \|\lambda\| = \mathcal{A}(\hat{\pi} \circ \gamma) + \frac{kT}{\zeta^2} \stackrel{(25)}{=} p_3$$

and that $\rho(T + \epsilon) = \hat{\pi} \circ \gamma(T) = \hat{\pi}(p)$. Then the horizontal lift $\tilde{\rho} : [-\epsilon, T + \epsilon] \rightarrow \mathbb{H}$ of ρ is a d_{CC} -rectifiable curve from 0 to p .

CASE 2. Suppose that $\hat{\pi} \circ \gamma$ covers more than half of the circle. Let $\lambda \in \mathfrak{h}/[\mathfrak{h}, \mathfrak{h}]$ be the diameter of the circle that contains 0. Since T is

shorter than the whole circle, then

$$(28) \quad \|\lambda\| \geq \frac{T}{\pi}.$$

Let λ^\perp be the unit vector perpendicular to λ and forming an angle smaller than $\pi/2$ with the arc $\hat{\pi} \circ \gamma$. Let $\epsilon > 0$ be such that

$$(29) \quad \epsilon \cdot \|\lambda\| = \frac{kT}{\zeta^2}.$$

Now, define $\rho : [-\epsilon, T + \epsilon] \rightarrow \mathfrak{h}/[\mathfrak{h}, \mathfrak{h}]$ as

$$\rho(t) = \begin{cases} (t + \epsilon)\lambda^\perp & \text{for } -\epsilon \leq t \leq 0 \\ \epsilon\lambda^\perp + \hat{\pi} \circ \gamma(t) & \text{for } 0 \leq t \leq \frac{\pi\|\lambda\|}{2} \\ \epsilon\lambda^\perp + \lambda - (t - \frac{\pi\|\lambda\|}{2})\lambda^\perp & \text{for } \frac{\pi\|\lambda\|}{2} \leq t \leq \frac{\pi\|\lambda\|}{2} + \epsilon \\ \hat{\pi} \circ \gamma(t - \epsilon) & \text{for } \frac{\pi\|\lambda\|}{2} + \epsilon \leq t \leq T + \epsilon \end{cases}$$

where we used the fact $\lambda = \hat{\pi} \circ \gamma(\frac{\pi\|\lambda\|}{2})$. Notice that

$$\mathcal{A}(\rho) = \mathcal{A}(\hat{\pi} \circ \gamma) + \epsilon \cdot \|\lambda\| = \mathcal{A}(\hat{\pi} \circ \gamma) + \frac{kT}{\zeta^2} = p_3.$$

Then the horizontal lift $\tilde{\rho} : [-\epsilon, T + \epsilon] \rightarrow \mathbb{H}$ of ρ is a d_{CC} -rectifiable curve from 0 to p .

In both cases $\tilde{\rho}$ is a horizontal curve from 0 to p of length

$$(30) \quad \ell(\tilde{\rho}) = T + 2\epsilon$$

Moreover, from (26) and (27) (respectively (28) and (29)) we get

$$(31) \quad \epsilon = \frac{kT}{\zeta^2\|\lambda\|} \leq \frac{kT}{\zeta^2} \frac{\pi}{T} \stackrel{(21)}{\leq} \frac{2\pi}{\zeta^2} \frac{\pi}{T} = \frac{2\pi^2}{\zeta^2 T}$$

Finally using in order (22), (30), (31), (24)

$$\begin{aligned} d_{CC}(0, p) - d_R(0, p) &\leq \ell(\tilde{\rho}) - \sqrt{1 + \frac{k^2}{\zeta^2}} \cdot T \leq \\ &\leq T + 2\epsilon - \sqrt{1 + \frac{k^2}{\zeta^2}} \cdot T \leq 2\epsilon \leq 2 \frac{2\pi^2}{\zeta^2 T} \leq \frac{4\pi^2}{\zeta^2} \frac{1}{d_{CC}(0, p) - \frac{2^{3/2}\pi}{\zeta}}. \end{aligned}$$

□

Remark 4.2. The inequality (2) is sharp. Indeed, for $z \rightarrow \infty$, we have the asymptotic equivalence

$$(32) \quad d_{CC}(0, (0, 0, z)) - d_R(0, (0, 0, z)) \sim \frac{4\pi^2}{\zeta^2} \frac{1}{d_{CC}(0, (0, 0, z))}.$$

Proof of (32). We claim that, for $z > 0$ large enough,

$$(33) \quad d_R(0, (0, 0, z)) = 2\sqrt{\pi}\sqrt{z - \frac{\pi}{\zeta^2}}.$$

Let $\gamma : [0, T] \rightarrow \mathbb{H}$ be a d_R -length-minimizing curve from 0 to $(0, 0, z)$. Since z is large, we assume that γ is of (TYPE II), see Proposition A.2, for some $k > 0$ and $\theta = 0$. Since the end point is on the Z axis, we have

$$(34) \quad kT = 2\pi$$

and $z = \frac{T}{2k} + \frac{kT}{\zeta^2}$, from which follows

$$(35) \quad T^2 = 4\pi \left(z - \frac{2\pi}{\zeta^2} \right).$$

We know also the length of γ (see Corollary A.8) and so we get

$$\begin{aligned} d_R(0, (0, 0, z)) = \ell(\gamma) &= T \|\omega_{\mathbb{H}}(\gamma')\| = T \sqrt{1 + \frac{k^2}{\zeta^2}} = \sqrt{T^2 + \frac{4\pi^2}{\zeta^2}} \\ &= \sqrt{4\pi \left(z - \frac{2\pi}{\zeta^2} \right) + \frac{4\pi^2}{\zeta^2}} = 2\sqrt{\pi}\sqrt{z - \frac{\pi}{\zeta^2}}. \end{aligned}$$

Claim (33) is proved. From Corollary A.6 we get $d_{CC}(0, (0, 0, z)) = 2\sqrt{\pi}\sqrt{z}$ and

$$\begin{aligned} d_{CC}(0, (0, 0, z)) - d_R(0, (0, 0, z)) &= 2\sqrt{\pi} \left(\sqrt{z} - \sqrt{z - \frac{\pi}{\zeta^2}} \right) \\ &= \frac{2\sqrt{\pi}}{\sqrt{z}} \frac{\frac{\pi}{\zeta^2}}{1 + \sqrt{1 - \frac{\pi}{\zeta^2 z}}} = \frac{1}{2\sqrt{\pi}\sqrt{z}} \frac{4\pi^2}{\zeta^2} \frac{1}{1 + \sqrt{1 - \frac{\pi}{\zeta^2 z}}}. \end{aligned}$$

□

We are now ready to give the proof of the main theorems:

Proof of Theorem 1.2. The implication (a) \Rightarrow (b) is trivial. The implication (b) \Rightarrow (c) is proven in Proposition 2.2. The equivalence (c) \Leftrightarrow (d) follows from Proposition 2.1. The assertion (e) is a restatement of (d). For (d) \Rightarrow (a) one uses Proposition 4.1 in order to reduce to the case when both d and d' are strictly subRiemannian and then one applies Proposition 3.1. □

Proof of Theorem 1.3. This is a consequence of Propositions 4.1 and of the sharpness result (9) of Proposition 3.1. □

5. THE HOROBOUNDARY

Let (X, d) be a geodesic space and $\mathcal{C}(X)$ the space of continuous functions $X \rightarrow \mathbb{R}$ endowed with the topology of the uniform convergence on compact sets. The map $\iota : X \hookrightarrow \mathcal{C}(X)$, $(\iota(x))(y) := d(x, y)$, is an embedding, i.e., a homeomorphism onto its image.

Let $\mathcal{C}(X)/\mathbb{R}$ be the topological quotient of $\mathcal{C}(X)$ with kernel the constant functions, i.e., for every $f, g \in \mathcal{C}(X)$ we set the equivalence relation $f \sim g \Leftrightarrow f - g$ is constant.

Then the map $\hat{\iota} : X \hookrightarrow \mathcal{C}(X)/\mathbb{R}$ is still an embedding. Indeed, since the map $\mathcal{C}(X) \rightarrow \mathcal{C}(X)/\mathbb{R}$ is continuous and open, we only need to show that $\hat{\iota}$ is injective: if $x, x' \in X$ are such that $\iota(x) - \iota(x')$ is constant, then one takes $z \in X$ such that $d(x, z) = d(x', z)$, which exists because (X, d) is a geodesic space, and checks that

$$d(x, x') = \iota(x)(x') - \iota(x')(x') = \iota(x)(z) - \iota(x')(z) = 0.$$

Define the *horoboundary* of (X, d) as

$$\partial_h X := cl(\hat{\iota}(X)) \setminus \hat{\iota}(X) \subset \mathcal{C}(X)/\mathbb{R},$$

where $cl(\hat{\iota}(X))$ is the topological closure.

Another description of the horoboundary is possible. Fix $o \in X$ and set

$$\mathcal{C}(X)_o := \{f \in \mathcal{C}(X) : f(o) = 0\}.$$

Then the restriction of the quotient projection $\mathcal{C}(X)_o \rightarrow \mathcal{C}(X)/\mathbb{R}$ is an isomorphism of topological vector spaces. Indeed, one easily checks that it is both injective and surjective, and that its inverse map is $[f] \mapsto f - f(o)$, where $[f] \in \mathcal{C}(X)/\mathbb{R}$ is the class of equivalence of $f \in \mathcal{C}(X)$.

Hence, we can identify $\partial_h X$ with a subset of $\mathcal{C}(X)_o$. More explicitly: $f \in \mathcal{C}(X)_o$ belongs to $\partial_h X$ if and only if there is a sequence $p_n \in X$ such that $p_n \rightarrow \infty$ (i.e., for every compact $K \subset X$ there is $N \in \mathbb{N}$ such that $p_n \notin K$ for all $n > N$) and the sequence of functions $f_n \in \mathcal{C}(X)_o$,

$$(36) \quad f_n(x) := d(p_n, x) - d(p_n, o),$$

converge uniformly on compact sets to f .

Proof of Corollary 1.4. Let us first remark that if d, d' are two geodesic distances on X and

$$(37) \quad \lim_{d(p,q)+d'(p,q) \rightarrow \infty} |d'(p, q) - d(p, q)| = 0.$$

then

$$\partial_h(X, d') = \partial_h(X, d).$$

Indeed, first of all the space $\mathcal{C}(X)_o$ depends only on the topology of X . Moreover, if $f \in \partial_h(X, d)$, let $p_n \in X$ be a sequence as in (36) and set $f'_n(x) := d'(p_n, x) - d'(p_n, o)$. Then

$$|f'_n(x) - f_n(x)| \leq |d'(p_n, x) - d(p_n, x)| + |d'(p_n, o) - d(p_n, o)|,$$

and as a consequence of (37) we get $f'_n \rightarrow f$ uniformly on compact sets. This shows $\partial_h(X, d) \subset \partial_h(X, d')$. The other inclusion follows by the symmetry of (37) in d and d' .

Now, if d_R and d_{CC} are metrics on \mathbb{H} like in Corollary 1.4, then (37) is easily satisfied thanks to Theorem 1.3, and therefore $\partial_h(\mathbb{H}, d_R) = \partial_h(\mathbb{H}, d_{CC})$ if the Riemannian metric d_R and the subRiemannian metric d_{CC} are compatible. The conclusion follows from [KN10]. \square

The Busemann points in the boundary $\partial_h(X, d)$ are usually defined as the horofunctions associated to sequences of points (p_n) diverging to infinity along rays or “almost geodesic rays”. However, in literature there are different definitions of almost geodesic rays, according to the generality of the metric space (X, d) under consideration ([Haa96], [Rie02], [DPS12]). A map $\gamma : I = [0, +\infty) \rightarrow (X, d)$ into a complete length space is called

- a *quasi-ray*, if the length excess

$$\Delta_N(\gamma) = \sup_{t, s \in [N, +\infty)} \ell(\gamma; t, s) - d(\gamma(t), \gamma(s))$$

tends to zero for $N \rightarrow +\infty$;

- an *almost geodesic ray*, if

$$\Theta_N(\gamma) = \sup_{t, s \in [N, +\infty)} d(\gamma(t), \gamma(s)) + d(\gamma(s), \gamma(0)) - t$$

tends to zero for $N \rightarrow +\infty$.

(Notice that the second definition depends on the parametrization, while the first one is intrinsic). We will use here a notion of Busemann points which is more general than both of them:

Definition 5.1. A sequence of points (p_n) diverging to infinity in (X, d) is said to diverge *almost straightly* if for all $\epsilon > 0$ there exists L such that for every $n \geq m \geq L$ we have

$$(38) \quad d(p_L, p_m) + d(p_m, p_n) - d(p_L, p_n) < \epsilon$$

It is easy to verify that points diverging along a quasi-ray or along an almost-geodesic ray diverge almost straightly. We then define a *Busemann point* as a horofunction f which is the limit of a sequence $f_n(x) = d(p_n, x) - d(p_n, o)$, for points (p_n) diverging to infinity almost straightly.

To prove Corollary 1.5, we need the following lemma. We remind that a metric space is *boundedly compact* if closed balls are compact.

Lemma 5.2. *Let (X, d) be a boundedly compact geodesic space, $o \in X$ and $\{p_n\}_{n \in \mathbb{N}} \subset X$ a sequence of points diverging almost straightly. Then:*

- (i) *the sequence $f_n(x) = d(p_n, x) - d(p_n, o)$ converges uniformly on compacts to a horofunction f ;*

$$(ii) \lim_{n \rightarrow \infty} f(p_n) + d(o, p_n) = 0.$$

Proof. Since the 1-Lipschitz functions f_n are uniformly bounded on compact sets and (X, d) is boundedly compact, then the family $\{f_n\}_{n \in \mathbb{N}}$ is pre-compact with respect to the uniform convergence on compact sets. Hence, if we prove that there is a unique accumulation point, then we obtain that the whole sequence $\{f_n\}_{n \in \mathbb{N}}$ converges.

So, let $g, g' \in \mathcal{C}(X)$ and let $\{f_{n_k}\}_{k \in \mathbb{N}}$ and $\{f_{n'_k}\}_{k \in \mathbb{N}}$ be two subsequences of $\{f_n\}_{n \in \mathbb{N}}$ such that $f_{n_k} \rightarrow g$ and $f_{n'_k} \rightarrow g'$ uniformly on compact sets. We claim

$$(39) \quad \forall \epsilon > 0 \exists R_\epsilon \in \mathbb{R} \forall x \in X \quad |g'(x) + R_\epsilon - g(x)| \leq \epsilon.$$

Let $\epsilon > 0$. Let $L \in \mathbb{N}$ be such that (38) holds for every $n \geq m \geq L$. Define for $x \in X$

$$\begin{aligned} g_L(x) &:= \lim_{k \rightarrow \infty} d(p_{n_k}, x) - d(p_{n_k}, p_L) = g(x) - g(p_L) \\ g'_L(x) &:= \lim_{k \rightarrow \infty} d(p_{n'_k}, x) - d(p_{n'_k}, p_L) = g'(x) - g'(p_L). \end{aligned}$$

Then for $n_i \geq n'_j \geq L$, we get for all $x \in X$

$$\begin{aligned} d(p_{n_i}, x) - d(p_{n_i}, p_L) - d(p_{n'_j}, x) + d(p_{n'_j}, p_L) \\ \leq d(p_{n_i}, p_{n'_j}) - d(p_{n_i}, p_L) + d(p_{n'_j}, p_L) \leq \epsilon. \end{aligned}$$

By taking the limit $i \rightarrow \infty$ and $j \rightarrow \infty$, we obtain for all $x \in X$

$$g_L(x) - g'_L(x) \leq \epsilon.$$

By the symmetry of the argument, also $g'_L(x) - g_L(x) \leq \epsilon$ holds. Therefore for all $x \in X$

$$\epsilon \geq |g'_L(x) - g_L(x)| = |g'(x) - g(x) + g(p_L) - g'(p_L)|.$$

Setting $R_\epsilon = g(p_L) - g'(p_L)$, we conclude the proof of claim (39).

It is now easy to conclude (i) from (39). Indeed, taking $x = o$, we have $|R_\epsilon| \leq \epsilon$, therefore for all $\epsilon > 0$ and for all $x \in X$ $|g(x) - g'(x)| \leq 2\epsilon$, i.e., $g = g'$. This completes the proof of (i).

To prove assertion (ii), fix $\epsilon > 0$ and let $L \in \mathbb{N}$ be as above. Then we have for all $n \geq m \geq L$

$$\begin{aligned} 0 &\leq d(p_m, p_n) - d(p_n, o) + d(p_m, o) \\ &= d(p_m, p_n) + d(p_L, p_m) - d(p_L, p_n) + \\ &\quad + d(p_L, p_n) - d(p_n, o) - d(p_L, p_m) + d(p_m, o) \\ &\leq \epsilon + d(p_L, p_n) - d(p_n, o) - d(p_L, p_m) + d(p_m, o). \end{aligned}$$

Taking first the limit $n \rightarrow \infty$ and then $m \rightarrow \infty$ in the above lines, we obtain the estimate

$$\begin{aligned} 0 &\leq \liminf_{m \rightarrow \infty} f(p_m) + d(p_m, o) \\ &\leq \limsup_{m \rightarrow \infty} f(p_m) + d(p_m, o) \leq \epsilon + f(p_L) - f(p_L) = \epsilon. \end{aligned}$$

Since $\epsilon > 0$ is arbitrary, then (ii) holds true. \square

Then, the proof of Corollary 1.5 runs similarly to Theorem 6.5 of [KN10].

Proof of Corollary 1.5. The horofunctions of type **(nv)** clearly are Busemann points, as they are limits, in particular, of the Riemannian geodesic rays which are the horizontal half-lines issued from the origin and which are always minimizing, see Proposition A.2 and Corollary A.3 in the Appendix. On the other hand, consider a horofunction of type **(v)**, $h_u = (v, z) = |u| - |u - v|$, for $u \in W$. Assume that there exists an almost straightly diverging sequence of points $p_n = v_n + z_n Z$ converging to h_u . By Lemma 5.2 (ii), we deduce that

$$\lim_{n \rightarrow \infty} f_u(p_n) + d_R(o, p_n) = \lim_{n \rightarrow \infty} |u| - |u - v_n| + d_R(o, p_n) = 0,$$

hence $\{v_n\}_{n \in \mathbb{N}}$ is necessarily an unbounded sequence. By Corollary 1.4 and the following description of horofunctions, it follows that h_u should be of type **(nv)**, a contradiction. \square

5.0.1. *Concluding remarks.* The Riemannian Heisenberg group shows a number of counterintuitive features which is worth to stress:

(i) in view of Corollary 1.4, all Riemannian metrics on \mathbb{H} with the same associated asymptotic metric have the same Busemann functions, though they are not necessarily isometric (in contrast, notice that all strictly subRiemannian metrics on \mathbb{H} are isometric). However, this is not surprising, because all left-invariant Riemannian metrics on \mathbb{H} are homothetic.

(ii) there exist diverging sequences of points $\{p_n\}_{n \in \mathbb{N}}$ that *visually converge* to a limit direction v (that is, the minimizing geodesics γ_n from o to p_n tend to a limit, minimizing geodesic γ_v with initial direction v), but whose associated limit point $h_{\{p_n\}}$ is not given by the limit point $\gamma_v(+\infty)$ of γ_v . This happens for all vertically divergent sequences $\{p_n\}_{n \in \mathbb{N}}$, as the limit geodesic γ_v is horizontal in this case (see Proposition A.2 and Corollary A.4 in the Appendix).

(iii) there exist diverging trajectories $\{p_n\}_{n \in \mathbb{N}}$, $\{q_n\}_{n \in \mathbb{N}}$ staying at bounded distance from each other, but defining different limit points (e.g., vertically diverging sequences of points with different limit horofunctions).

(iv) it is not true that, for a cocompact group of isometries G of (\mathbb{H}, d_R) , the limit set of G (which is the set of accumulation points of an orbit Gx_0 in $\partial(\mathbb{H}, d_R)$) equals the whole Gromov boundary; for instance, the discrete Heisenberg group $G = \mathbb{H}(\mathbb{Z})$, has a limit set equal to the set of all Busemann points, plus a discrete subset of the interior of the disk boundary \bar{D}^2 . Also, the limit set may depend on the choice of the base point $x_0 \in \mathbb{H}$.

(v) The functions appearing in **(nv)** coincide with the Busemann functions of a Euclidean plane in the direction $R_{\vartheta}(\hat{v}_{\infty})$; that is, the horofunction $h(v, z)$ associated to a diverging sequence $P_n = (v_n, z_n)$ of (\mathbb{H}, d_R) is obtained just by dropping the vertical component z of the argument, and then applying to v the usual Euclidean Busemann function in the direction which is opposite to the limit direction of the v_n 's, rotated by an angle ϑ depending on the quadratic rate of divergence of the sequence (ϑ is zero for points diverging sub-quadratically, and $\vartheta = \pm\pi$ when the divergence is sup-quadratical).

These properties mark a remarkable difference to the theory of non-positively curved, simply connected spaces.

APPENDIX A. LENGTH-MINIMIZING CURVES FOR d_{CC} AND d_R

In the Heisenberg group, locally length-minimizing curves are smooth solutions of an Hamiltonian system both in the Riemannian and in the subRiemannian case. Locally length-minimizing curves are also called *geodesics*.

Let d_R be a Riemannian metric on \mathbb{H} with horizontal space (\mathfrak{h}, g) . Let $V \subset \mathfrak{h}$ be the plane orthogonal to $[\mathfrak{h}, \mathfrak{h}]$ and let d_{CC} be the strictly subRiemannian metric on \mathbb{H} with horizontal space $(V, g|_V)$.

Fix a basis (X, Y, Z) for \mathfrak{h} such that (X, Y) is an orthonormal basis of $(V, g|_V)$ and $Z = [X, Y]$. Set $\zeta = \sqrt{g(Z, Z)}$. The basis (X, Y, Z) induces the exponential coordinates (x, y, z) on \mathbb{H} , i.e., $(x, y, z) = \exp(xX + yY + zZ)$. We will work in this coordinate system.

The Riemannian and subRiemannian length-minimizing curves are known and we recall their parametrization in the following two propositions. SubRiemannian geodesics can be found with different notation in [Cap+07]. Riemannian geodesics are found in [Mar97], in different coordinates and with parametrization by arc-length.

Proposition A.1 (subRiemannian geodesics). *All the non-constant locally length-minimizing curves of d_{CC} starting from 0 and parametrized by arc-length are the following: given $k \in \mathbb{R} \setminus \{0\}$ and $\theta \in \mathbb{R}$*

- (TYPE I) *The horizontal lines $t \mapsto (t \cos \theta, t \sin \theta, 0)$;*
 (TYPE II) *The curves $t \mapsto (x(t), y(t), z(t))$ given by*

$$\begin{cases} x(t) = \frac{1}{k} (\cos \theta (\cos(kt) - 1) - \sin \theta \sin(kt)) \\ y(t) = \frac{1}{k} (\sin \theta (\cos(kt) - 1) + \cos \theta \sin(kt)) \\ z(t) = \frac{1}{2k} t - \frac{1}{2k^2} \sin(kt). \end{cases}$$

Here the derivative at $t = 0$ is $(-\sin \theta, \cos \theta, 0)$.

Proposition A.2 (Riemannian geodesics). *All non-constant locally length-minimizing curves of d_R parametrized by a multiple of arc-length and starting from 0 are the following: given $k \in \mathbb{R} \setminus \{0\}$ and $\theta \in \mathbb{R}$*

- (TYPE 0) *The vertical line $t \mapsto (0, 0, t)$;*
 (TYPE I) *The horizontal lines $t \mapsto (t \cos \theta, t \sin \theta, 0)$;*
 (TYPE II) *The curves $t \mapsto (x(t), y(t), z(t))$ given by*

$$\begin{cases} x(t) = \frac{1}{k} (\cos \theta (\cos(kt) - 1) - \sin \theta \sin(kt)) \\ y(t) = \frac{1}{k} (\sin \theta (\cos(kt) - 1) + \cos \theta \sin(kt)) \\ z(t) = \frac{1}{2k} t - \frac{1}{2k^2} \sin(kt) + \frac{k}{\zeta^2} t. \end{cases}$$

Here the derivative at $t = 0$ is $(-\sin \theta, \cos \theta, \frac{k}{\zeta^2})$, which has Riemannian length $\sqrt{1 + \frac{k^2}{\zeta^2}}$.

The expression of geodesics helps us to prove the following facts.

Corollary A.3. *The horizontal lines of TYPE I are globally d_R - and d_{CC} -length-minimizing curves.*

Corollary A.4. *Both d_R - and locally d_{CC} -length-minimizing curves γ of TYPE II are not minimizing from 0 to $\gamma(t)$ if $|t| > \frac{2\pi}{k}$.*

Proof. This statement depends on the fact that in both cases, if we fix $k \in \mathbb{R} \setminus \{0\}$, then for all θ the corresponding length-minimizing curves $\gamma_{k,\theta}$ of TYPE II meet each other at the point $\gamma_{k,\theta}(2\pi/k)$. \square

Corollary A.5. *The locally d_R -length-minimizing curve γ of TYPE 0, $t \mapsto (0, 0, t)$, is not minimizing from 0 to $\gamma(t)$ for $|t| > \frac{2\pi}{\zeta^2}$.*

Proof. For $k > 0$ let γ_k be the d_R -length-minimizing curve of TYPE II with this k and $\theta = 0$. Then $(\gamma_k)_3(\frac{2\pi}{k}) = \frac{\pi}{k^2} + \frac{2\pi}{\zeta^2}$. Letting $k \rightarrow \infty$ we obtain $\hat{z} := \frac{2\pi}{\zeta^2}$. This means that for every $\epsilon > 0$ there is $z \leq \hat{z} + \epsilon$ and $k > 0$ such that $\gamma_k(\frac{2\pi}{k}) = (0, 0, z)$. Therefore $t \mapsto (0, 0, t)$ cannot be minimizing after z , and therefore after \hat{z} . \square

Corollary A.6. *If $p = (x, y, p_3)$ and $q = (x, y, q_3)$, then*

$$d_{CC}(p, q) = 2\sqrt{\pi} \cdot \sqrt{|p_3 - q_3|}.$$

Proof. First suppose $p = 0$: we have to prove that $d_{CC}(0, (0, 0, z)) = 2\sqrt{\pi}\sqrt{|z|}$. This is done by looking at the length-minimizing curves: they come from complete circle of perimeter $2\pi R = d$ and area $\pi R^2 = |z|$, so that $d_{CC}(0, (0, 0, z)) = 2\pi\sqrt{\frac{|z|}{\pi}} = 2\sqrt{\pi}\sqrt{|z|}$. The general case follows from the left-invariance of d_{CC} :

$$\begin{aligned} d_{CC}((x, y, p_3), (x, y, q_3)) &= d_{CC}(0, (x, y, p_3)^{-1}(x, y, q_3)) \\ &= d_{CC}(0, (0, 0, q_3 - p_3)). \end{aligned}$$

□

Corollary A.7. *If $p \in \{z = 0\}$, then*

$$d_{CC}(0, p) = d_R(0, p)$$

Corollary A.8. *d_R - and d_{CC} -length-minimizing curves of TYPE II are in bijection via the following rule: If $\eta : [0, T] \rightarrow \mathbb{H}$ is a d_{CC} -length-minimizing curve of TYPE II, then*

$$\gamma(t) = \eta(t) + \left(0, 0, \frac{kt}{\zeta^2}\right)$$

is d_R -length-minimizing of TYPE II, where $k \in \mathbb{R}$ is given by η . Moreover, it holds

$$\|\omega_{\mathbb{H}}(\gamma')\|^2 = 1 + \frac{k^2}{\zeta^4}$$

and

$$d_{CC}(\gamma(t), \eta(t)) = 2\sqrt{\pi} \sqrt{\frac{kt}{\zeta^2}}$$

Proof. All the statements come directly from the expression of the geodesics. Notice that a d_{CC} -length-minimizing curve η of TYPE II is parametrized by arc-length, i.e., $\|\omega_{\mathbb{H}}(\eta')\| \equiv 1$. On the other hand, the corresponding d_R -length-minimizing curve γ has derivative $\omega_{\mathbb{H}}(\gamma') = \omega_{\mathbb{H}}(\eta') + \frac{k}{\zeta^2}Z$, where $\omega_{\mathbb{H}}(\eta')$ is orthogonal to Z . Hence $\|\omega_{\mathbb{H}}(\gamma')\|^2 = 1 + \frac{k^2}{\zeta^4}$ □

REFERENCES

- [ABB15] A. A. Agrachëv, D. Barilari, and U. Boscain. *Introduction to Riemannian and sub-Riemannian geometry*. 2015.
- [Bal95] W. Ballmann. *Lectures on spaces of nonpositive curvature*. Vol. 25. DMV Seminar. With an appendix by Misha Brin. Birkhäuser Verlag, Basel, 1995, pp. viii+112. ISBN: 3-7643-5242-6.
- [BH99] M. R. Bridson and A. Haefliger. *Metric spaces of non-positive curvature*. Vol. 319. Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]. Springer-Verlag, Berlin, 1999, pp. xxii+643. ISBN: 3-540-64324-9.
- [BLD13] E. Breuillard and E. Le Donne. “On the rate of convergence to the asymptotic cone for nilpotent groups and subFinsler geometry”. In: *Proc. Natl. Acad. Sci. USA* 110.48 (2013), pp. 19220–19226. ISSN: 1091-6490.
- [Bur94] D. Burago. “Periodic metrics”. In: *Seminar on Dynamical Systems (St. Petersburg, 1991)*. Vol. 12. Progr. Non-linear Differential Equations Appl. Birkhäuser, Basel, 1994, pp. 90–95.

- [Bus55] H. Busemann. *The geometry of geodesics*. Academic Press Inc., New York, N. Y., 1955, pp. x+422.
- [Cap+07] L. Capogna et al. *An introduction to the Heisenberg group and the sub-Riemannian isoperimetric problem*. Vol. 259. Progress in Mathematics. Birkhäuser Verlag, Basel, 2007, pp. xvi+223. ISBN: 978-3-7643-8132-5; 3-7643-8132-9.
- [CS14] F. Cerocchi and A. Sambusetti. “Quantitative Bounded Distance Theorem and Margulis’ Lemma for Z^n actions with applications to homology”. In: *ArXiv e-prints* (Dec. 2014). arXiv: [1412.6516](#).
- [DPS12] F. Dal’bo, M. Peigné, and A. Sambusetti. “On the horoboundary and the geometry of rays of negatively curved manifolds”. In: *Pacific J. Math.* 259.1 (2012), pp. 55–100. ISSN: 0030-8730.
- [DS14] M. Duchin and M. Shapiro. “Rational growth in the Heisenberg group”. In: *ArXiv e-prints* (Nov. 2014). arXiv: [1411.4201](#).
- [Gro81a] M. Gromov. “Hyperbolic manifolds, groups and actions”. In: *Riemann surfaces and related topics: Proceedings of the 1978 Stony Brook Conference (State Univ. New York, Stony Brook, N. Y., 1978)*. Vol. 97. Ann. of Math. Stud. Princeton Univ. Press, Princeton, N.J., 1981, pp. 183–213.
- [Gro81b] M. Gromov. “Groups of polynomial growth and expanding maps”. In: *Inst. Hautes Études Sci. Publ. Math.* 53 (1981), pp. 53–73. ISSN: 0073-8301.
- [Gro87] M. Gromov. “Hyperbolic groups”. In: *Essays in group theory*. Vol. 8. Math. Sci. Res. Inst. Publ. Springer, New York, 1987, pp. 75–263.
- [Haa96] A. Haas. “Dirichlet points, Garnett points, and infinite ends of hyperbolic surfaces. I”. In: *Ann. Acad. Sci. Fenn. Math.* 21.1 (1996), pp. 3–29. ISSN: 0066-1953.
- [Had98] J. Hadamard. “Les surfaces à courbures opposées et leurs lignes géodésiques”. In: *Jorn. de Math.* 4.5 (1898), pp. 27–74.
- [Jea14] F. Jean. *Control of nonholonomic systems: from sub-Riemannian geometry to motion planning*. Springer Briefs in Mathematics. Springer, Cham, 2014, pp. x+104. ISBN: 978-3-319-08689-7; 978-3-319-08690-3.
- [KN09] T. Klein and A. Nicas. “The horofunction boundary of the Heisenberg group”. In: *Pacific J. Math.* 242.2 (2009), pp. 299–310. ISSN: 0030-8730.
- [KN10] T. Klein and A. Nicas. “The horofunction boundary of the Heisenberg group: the Carnot-Carathéodory metric”. In: *Conform. Geom. Dyn.* 14 (2010), pp. 269–295. ISSN: 1088-4173.

- [Kra99] S. A. Krat. “Asymptotic properties of the Heisenberg group”. In: *Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI)* 261.G geom. i Topol. 4 (1999), pp. 125–154, 268. ISSN: 0373-2703.
- [Mar97] V. Marenich. “Geodesics in Heisenberg groups”. In: *Geom. Dedicata* 66.2 (1997), pp. 175–185. ISSN: 0046-5755.
- [Mil68] J. Milnor. “A note on curvature and fundamental group”. In: *J. Differential Geometry* 2 (1968), pp. 1–7. ISSN: 0022-040X.
- [Mos73] G. D. Mostow. *Strong rigidity of locally symmetric spaces*. Annals of Mathematics Studies, No. 78. Princeton University Press, Princeton, N.J.; University of Tokyo Press, Tokyo, 1973, pp. v+195.
- [Pan83] P. Pansu. “Croissance des boules et des géodésiques fermées dans les nilvariétés”. In: *Ergodic Theory Dynam. Systems* 3.3 (1983), pp. 415–445. ISSN: 0143-3857.
- [Pat76] S. J. Patterson. “The limit set of a Fuchsian group”. In: *Acta Math.* 136.3-4 (1976), pp. 241–273. ISSN: 0001-5962.
- [Rie02] M. A. Rieffel. “Group C^* -algebras as compact quantum metric spaces”. In: *Doc. Math.* 7 (2002), 605–651 (electronic). ISSN: 1431-0635.
- [Rif14] L. Rifford. *Sub-Riemannian geometry and optimal transport*. Springer Briefs in Mathematics. Springer, Cham, 2014, pp. viii+140. ISBN: 978-3-319-04803-1; 978-3-319-04804-8.
- [Rob03] T. Roblin. “Ergodicité et équidistribution en courbure négative”. In: *Mém. Soc. Math. Fr. (N.S.)* 95 (2003), pp. vi+96. ISSN: 0249-633X.
- [Sto98] M. Stoll. “On the asymptotics of the growth of 2-step nilpotent groups”. In: *J. London Math. Soc. (2)* 58.1 (1998), pp. 38–48. ISSN: 0024-6107.
- [Sul79] D. Sullivan. “The density at infinity of a discrete group of hyperbolic motions”. In: *Inst. Hautes Études Sci. Publ. Math.* 50 (1979), pp. 171–202. ISSN: 0073-8301.
- [Tit72] J. Tits. “Free subgroups in linear groups”. In: *J. Algebra* 20 (1972), pp. 250–270. ISSN: 0021-8693.
- [Wal07] C. Walsh. “The horofunction boundary of finite-dimensional normed spaces”. In: *Math. Proc. Cambridge Philos. Soc.* 142.3 (2007), pp. 497–507. ISSN: 0305-0041.
- [Wal08] C. Walsh. “The horofunction boundary of the Hilbert geometry”. In: *Adv. Geom.* 8.4 (2008), pp. 503–529. ISSN: 1615-715X.
- [Wol68] J. A. Wolf. “Growth of finitely generated solvable groups and curvature of Riemannian manifolds”. In: *J. Differential Geometry* 2 (1968), pp. 421–446. ISSN: 0022-040X.

- [WW06] C. Webster and A. Winchester. “Busemann points of infinite graphs”. In: *Trans. Amer. Math. Soc.* 358.9 (2006), 4209–4224 (electronic). ISSN: 0002-9947.

(Le Donne, Nicolussi Golo) UNIVERSITY OF JYVASKYLA, FINLAND

(Sambusetti) SAPIENZA UNIVERSITÀ DI ROMA, ITALY