

EPIPHYTIC LICHENS AS INDICATORS OF ENVIRONMENTAL QUALITY AROUND A MUNICIPAL SOLID WASTE LANDFILL (C ITALY)

5 Luca Paoli^{1,*}, Alice Grassi¹, Andrea Vannini¹, Ivana Maslaňáková², Ivana Biľová², Martin Bačkor²,
Adelmo Corsini³, Stefano Loppi¹

¹ Department of Life Sciences, University of Siena, via P.A. Mattioli 4, I-53100, Siena, Italy;

² Department of Botany, Institute of Biology and Ecology, P.J. Šafárik University in Košice, Mánesova 23, SK-04001
Košice, Slovakia;

10 ³ Biologist, Pistoia, Italy

*corresponding author: Luca Paoli

Tel. (+39) 0577 235408 – Fax (+39) 0577 232896 – email: paoli4@unisi.it

15 **Abstract**

Epiphytic lichens have been used as indicators of environmental quality around a municipal
solid waste landfill in C Italy. An integrated approach, using the diversity of epiphytic lichens, as
well as element bioaccumulation and physiological parameters in the lichen *Flavoparmelia*
caperata (L.) Hale was applied along a transect from the facility. The results highlighted the
20 biological effects of air pollution around the landfill. The Index of Lichen Diversity (ILD) increased
and the content of heavy metals (Cr, Cd, Cu, Fe, Ni and Zn) decreased with distance from the
landfill. Clear stress signals were observed in lichens growing in front of the facility, i.e.
discoloration, necrosis, membrane lipid peroxidation, lower ergosterol content, higher
dehydrogenase activity. Decreased photosynthetic efficiency, altered chlorophyll integrity and
25 production of secondary metabolites were also found. The results suggested that lichens can be
profitably used as bioindicators of environmental quality around landfills.

Keywords: air pollution, biomonitoring, chlorophyll, heavy metals, *Flavoparmelia caperata*,
Lichen Diversity Values, secondary metabolites

30

1. Introduction

There is a high public concern for the environmental and health impact related to waste
management. Source reduction, reuse, recycling, composting, incineration and landfilling are part of
35 the integrated system for waste management. The European Commission considers landfilling as
the least preferable option, which should be limited to the necessary minimum; however, it is still
the main waste disposal method throughout Europe.

Recent legislation prescribes strict rules for waste disposal in landfills (European Union, Directive
2008/98/EC of the European Parliament and of the European Council of 19 November 2008).
40 Nevertheless, municipal solid waste (MSW) landfills may potentially contaminate fresh- and
especially groundwater, soil, air and the biota. MSW landfills release a variety of air pollutants and
among them landfill gas and particulate matter are particularly relevant. Emission of landfill gas
poses environmental concern owing to the content of methane, carbon dioxide, nitrogen oxides and
a mixture of contaminants, including volatile organic compounds (Chiriac et al., 2011; Sun et al.,
45 2013). Dumping activities are also a source of particulate matter, whose composition reflects the
nature of the waste and the waste layer (terrain) used to stabilize the surface. Around landfill sites,
enhanced concentrations of particulate-bound metals were found due to wind erosion, resuspension
from the surface of the site, road surface dust and resuspension of deposited particulate matter on a
road surface, refuse truck emissions e.g., exhausts, tire wear dust, brake wear dust (Chalvatzaki et
50 al., 2010; 2014; Han et al., 2011).

Recent environmental legislation prescribes objectives and rules for air quality to protect human health and the environment “paying particular attention to sensitive populations, and the environment as a whole, to improve the monitoring and assessment of air quality including the deposition of pollutants” (Directive 2008/50/EC of the European Parliament and of the Council of 21 May 2008 on ambient air quality and cleaner air for Europe). Under the Air Quality Directive, biomonitoring of air pollution can contribute to the implementation of environmental policy on air quality and atmospheric pollution control, providing consistent data for environmental management (Pirintsos and Loppi, 2008). In this context, the use of lichens as bioindicators of environmental quality provides a special viewpoint on the atmospheric environment (Nimis et al., 2002).

Lichens are symbiotic organisms composed by green algae (or cyanobacteria) and fungi. Lichen metabolism depends on the mineral uptake from the atmosphere; therefore these organisms are effective in trapping trace elements from the surrounding environment. They grow very slowly, do not have stomata or cuticle regulating air exchanges and accumulate contaminants over the whole surface. Changes in the composition of lichen communities, trace element content and physiological status are useful bioindicators, providing spatial and temporal evidence for trends in ambient pollution burdens (Loppi, 2014).

The diversity of epiphytic lichens proved to be a useful and robust indicator of air quality (Loppi et al., 2002). The concentrations of heavy metals in lichen thalli resulted correlated with their environmental levels (Bari et al., 2001; Sloof, 1995). In addition, several parameters (e.g., chlorophyll *a* fluorescence emission, chlorophyll integrity, membrane lipid peroxidation, alteration of secondary metabolites, ergosterol content, dehydrogenase activity) have been introduced into experimental lichenology for the qualitative assessment of environmental stress (Bačkor, 2011). Only few lichen based studies reported on the biological effects of air pollution determined by different waste management strategies: waste incineration (Loppi et al., 1995, 2000; Paoli et al., 2015; Tretiach et al., 2011); landfilling (Paoli et al., 2012), industrial composting (Paoli et al., 2014).

A comprehensive biomonitoring programme should integrate several methods distributed along the biomonitoring chain, allowing to detect exposure, threads and impacts also when waste management is concerned. Therefore, among these methods, changes in the diversity of epiphytic lichens provide a useful method for monitoring the overall status of an ecosystem and the impacts of emissions; measuring the accumulation of airborne pollutants in lichen thalli (e.g., of heavy metals) can be used to assess spatial and temporal exposure; measuring physiological responses allows detecting early effects of ongoing processes and threads, before their consequences are apparent at higher levels (Bealey et al., 2008; Paoli et al., 2015).

In the present study, epiphytic lichens have been used as indicators of environmental quality around a MSW landfill, addressing specific questions: does the landfill influences i) the diversity of epiphytic lichens, (ii) the bioaccumulation of selected heavy metals, and (iii) the physiological status of lichens?

2. Material and Methods

2.1 Study area

The investigated landfill (43°52'52" N, 10°53'21" E, ca. 60 m asl) is located in Tuscany, C Italy. Operating since 1996, it extends over an area of 160,000 m² and actually it is foreseen to last until 2027, when the overall volume will be 3,010,000 m³. The authorized daily capacity is 420 tons. The disposed material consists of municipal solid, hazardous and non hazardous wastes and the material for landfill cover. Wastes may include scraps of paper, plastics and metals, packing, spent tires, textile products, building materials, ashes from municipal solid waste incinerators, polluted terrain from environment reclamation, etc.

The landfill site is located over an impermeable natural clay layer; bottom and side boundaries may

vary according to the period of cultivation. They generally include protective layers, such as a compact clay layer (100 cm), geotextile membranes, gravel (50 cm), geomembranes, non-woven fabric (1200 g/m²), pulper products (50 cm). Landfill covers consist of a waste layer (terrain) to stabilize the surface, drainage systems, compact clay (20 cm), soil bentonite and a vegetative soil layer (up to 100 cm). A grassy mantle and/or reafforestation with local vegetation complete the recovery of the site after closing of each parcel. Systems for leachate treatment, and for gas recovery, collection and treatment, are in operation and landfill gas is used to support energy needs. Atmospheric concentrations of several pollutants of environmental concern in the vicinity of the landfill were available through instrumental monitoring (source Pistoambiente). Referring to the period corresponding to our study (spring 2012 – spring 2013), values were within the following ranges (hourly concentrations, µg/Nm³): As (0.14–0.73), Cd (0.14–0.16), Cr (12–29), Cu (0.6–8.4), Ni (6.4–18.4), Pb (3.3–4.8), V (0.18–0.25) and HF (0.10 – 0.35), HCl (0–6.3), NO_x (25–82), SO_x (0.1–6.3).

The landfill is surrounded to the N, W and S by a vegetation belt dominated by *Quercus cerris* and *Q. pubescens*. Vineyards, olive plantations and woodlands characterize the hilly surroundings, while to the E lowland, several plant nurseries and inhabited areas prevail. The climate of the area is intermediate between sub-oceanic and sub-Mediterranean, with mean annual rainfall over 1300 mm and mean annual temperature of 14.1°C.

2.2 Experimental design

To assess possible biological effects caused by the presence of the landfill, sampling sites were classified into three groups according to the distance from the landfill: 1) sites directly facing the landfill; 2) sites located at about 200 m from the landfill; 3) sites located at about 1500 m from the landfill.

The above distances were selected based on previous studies (Paoli et al., 2012): in particular, sites of group 2) correspond to the outer margin of the vegetation belt surrounding the landfill, which roughly ranges up to 200 m. The distance of 1500 m corresponds to sites with negligible impact. For each group, three sampling sites were investigated in the period May 2012 – May 2013 for lichen diversity as well as element bioaccumulation and physiological status in native lichen thalli of the species *Flavoparmelia caperata* (L.) Hale. Sampling sites are represented by circular plots of 60 m diameter. Since the sampling sites at about 1500 m from the landfill belongs to the regular monitoring network of 500 m × 500 m installed in the surrounding territory, for the homogeneity of data distribution, the plot of 60 m diameter has been selected within this level (see next paragraph). The diversity of epiphytic lichens and the accumulation of selected elements were investigated as part of prescriptions implemented to improve the environmental performance of the facility and enhance health protection (Eco-Management and Audit Scheme, EMAS).

2.3 Assessment of the lichen diversity

The diversity of epiphytic lichens was measured according to the standardized protocol of Asta et al. (2002). An Index of Lichen Diversity (ILD, also named LDV "Lichen Diversity Value" see e.g. CEN standard EN 16413 "Ambient air - Biomonitoring with lichens - Assessing epiphytic lichen diversity) was computed using a sampling grid consisting of four 50×10 cm² ladders, each divided into five 10×10 cm² units. The grid was positioned systematically on the N, E, S and W cardinal sides of the bole of each tree, at a height of 1 m above ground. The ILD of the tree corresponded to the sum of frequencies of epiphytic lichens in the grid and the ILD of each monitoring site was the arithmetic mean of the ILD measured for each sampled tree. Trees for epiphytic lichen sampling (*Q. cerris* and *Q. pubescens*) were deemed suitable if well lit, with girth >60 cm, trunk near straight, not damaged and without parts with >25% cover of bryophytes. Three trees were sampled in each sampling site. In case of identification problems during field sampling, specimens were collected and identified later in the laboratory. Nomenclature follows the online database *Italic* (Nimis and

Martellos, 2008).

2.4 Bioaccumulation of trace elements

155 The bioaccumulation of selected elements in native lichens was investigated according to Bargagli
and Nimis (2002). Thalli of the foliose lichen *F. caperata* were collected for element analysis. The
species *F. caperata* was chosen because of its wide distribution in the study area and because it had
been already used in bioaccumulation studies (Loppi et al., 2004; Paoli et al., 2012). At each
160 sampling site, 10-30 thalli growing on the bole of 3-10 *Q. cerris* or *Q. pubescens* trees were
harvested from all cardinal exposures, at 1-2 m from ground. Tree boles were deemed suitable for
lichen harvesting if almost straight, not damaged and without parts with >25% cover of bryophytes.
In the laboratory, samples were carefully cleaned under a stereoscopic microscope to remove
extraneous material deposited onto the surface, such as moss samples, bark pieces and soil particles.
Only the peripheral part of the thalli (up to 5 mm from lobe tips) was selected for the analysis; in *F.*
165 *caperata* this part roughly corresponds to the last year of growth and can be easily separated from
the bark, being distinguishable by a paler colour and absence of rhizinae. Concentrations of selected
elements of toxicological concern (As, Cd, Cr, Ni, Pb, V, Zn) and Fe (being associated to soil
contamination of the samples) were determined by ICP-MS (Perkin Elmer – Sciex, Elan 6100) as
reported in Paoli et al. (2012) and expressed on a dry weight basis ($\mu\text{g/g dw}$). Analytical quality was
checked by analysing the Standard Reference Material IAEA-336 ‘lichen’. Precision of analysis
170 was estimated by the coefficient of variation of 4 replicates and was found to be within 10% for all
elements. Three replicates were measured at each site.

2.5 Physiological parameters

175 The following parameters were used for the assessment of the physiological status of the lichens:
chlorophyll integrity, photosynthetic efficiency, membrane lipid peroxidation, viability, ergosterol
content and secondary metabolites.

2.5.1 Chlorophyll integrity

180 Photosynthetic pigments were measured as described in Pisani et al. (2007). Two extraction cycles,
45 min each, were run in a warm bath (65°C), using 5 mL of DMSO. Absorbance of the extracts
was measured using a UV-visible spectrophotometer (Agilent 8453). Chlorophyll integrity was
expressed by the ratio between the absorbance at 435 and 415 nm ($\text{OD}_{435}/\text{OD}_{415}$). A decrease of this
ratio reflects chlorophyll *a* degradation to phaeophytin *a* (Garty et al., 2000). Five replicates were
measured at each site.

185

2.5.2 Photosynthetic efficiency

The photosynthetic efficiency of the samples was assessed in terms of chlorophyll (Chl) *a*
fluorescence emission, that was analysed by the classical physiological indicator F_v/F_m ,
representing the potential quantum yield of primary photochemistry, where $F_v = (F_m - F_0)$ is the
190 variable fluorescence and F_0 and F_m are minimum and maximum Chl *a* fluorescence. Samples were
lightened 1 sec with a saturating 3000 $\mu\text{mol/m}^2\text{s}$ light pulse and fluorescence emission was
recorded for 1 sec. Samples were measured with a Plant Efficiency Analyzer (Handy PEA,
Hansatech Ltd, Norfolk, UK). Ten replicates were measured at each site.

2.5.3 Membrane lipid peroxidation

195 Membrane lipid peroxidation was estimated using the thiobarbituric acid reactive substances
(TBARS) assay, as reported in Pisani et al. (2011). TBARS are a decomposition product of
polyunsaturated fatty acids which are produced during peroxidation of membrane lipids (Mittler,
2002). Their content increases upon exposure of lichen thalli to high concentrations of toxic
200 elements (Bačkor et al., 2010; Pisani et al., 2011). Five replicates were measured at each site, results

were expressed as $\mu\text{mol/g}$ (dw).

2.5.4 Sample viability

205 Triphenyltetrazolium chloride (TTC) reduction to triphenylformazan (TPF) is a good indicator of dehydrogenase activity (dark respiration) and was used to assess sample viability according to Bačkor and Fahselt (2005). Results were expressed as absorbance units/g (dw). Five replicates were measured at each site.

2.5.5 Ergosterol content

210 Ergosterol content in lichens is sensitive to the exposure to heavy metals, which likely reduces the integrity of cell membranes of the mycobiont. Samples (100 mg) were measured as indicated in Dahlman et al. (2002). Ergosterol absorption at 280 nm was measured with a UV detector (Ecom LCD 2084). As ergosterol is sensitive to light, all steps were conducted almost in the dark. Three replicates were measured at each site.

215

2.5.6 Secondary metabolites

Secondary metabolites (caperatic and usnic acid) were measured as indicated by Bačkor et al. (2011). Three replicates were measured at each site.

220 2.6 Data interpretation and statistical analysis

Significance of differences ($P < 0.05$) concerning total ILD, element accumulation and physiological parameters was checked by ANOVA, using the HSD Tukey test for post-hoc comparisons. The ILD values were interpreted in terms of air pollution according to the following scale (Paoli and Loppi, 2008): 0 = very high (lichen desert), 1-40 = high, 41-80 = moderate, 81-120 = low, >120 = negligible. Metal concentrations were interpreted in terms of air pollution based on a scale (Table 1) desumed from Bargagli and Nimis (2002). The Spearman correlation test was used to investigate correlations between lichen diversity, physiological parameters and heavy metals. Prior to the analysis, data were transformed on a 0 (min) - 1 (max) scale.

230 3. Results

3.1 Lichen diversity (ILD)

The lichen diversity values and the list of the species recorded along with their average frequency, are summarized in Table 2. The Index of Lichen Diversity (ILD) significantly increased moving
235 from the landfill to the group of sites at 1500 m from the landfill ($F=23.89$ $P=0.000$). ILD values indicated conditions of high to moderate air pollution in front of the landfill, moderate to low air pollution at 200 m, and low to negligible air pollution at 1500 m. Thirty-seven epiphytic lichens were recorded. The average number of species per sampled tree was 9 at the landfill, 11 at 200 m, 16 at 1500 m. The highest contribution to lichen frequencies were determined by crustose, tolerant species (e.g., *Candelariella xanthostigma*, *Lepraria* sp.). The lichen vegetation along the transect was dominated by mesophilous and acidophilous or subneutrophilous species, while only few species were nitrophilous (Table 2). The frequency of the widespread foliose lichen *F. caperata* clearly decreased approaching the landfill, while species typically occurring in relatively undisturbed areas (*Catillaria nigroclavata*, *Collema subflaccidum*, *Phlyctis argena*, *Physcia clementei*, *Physconia perisidiosa*, *P. servitii*) were present only at 200 and 1500 m.

3.2 Heavy metals

A notable statistically significant decrease in the accumulation of trace elements in the lichen *F. caperata* was observed along with distance from the landfill (Table 3). Values measured in samples
250 collected in front of the landfill indicated high air pollution for Cr and Ni, moderate air pollution for

Cd, Cu, Fe and Zn, low air pollution for As and Pb. Values measured in samples collected at 200 m indicated moderate air pollution for Ni, low air pollution for Cd, Cr, Cu, Fe and Zn, very low air pollution for As and Pb. Values for all elements but Ni measured in samples collected at 1500 m indicated a condition of very low air pollution; moderate air pollution emerged for Ni.

255

3.3 Physiological parameters

The investigated physiological parameters allowed detecting stress symptoms in the lichens collected in front of the landfill (Table 4). Lower chlorophyll integrity ($OD_{435/415}$) and higher dehydrogenase activity were found at the sites directly facing the landfill. Peroxidated membrane lipids (TBARS) and lowered ergosterol content were found up to 200 m from the landfill. Trends were not so clear for secondary metabolites, but overall caperatic acid decreased and usnic acid increased approaching the landfill. Photosynthetic efficiency (F_v/F_m) was not affected by the landfill, although lower values were occasionally measured in lichen thalli facing the landfill. Observations carried out on *F. caperata* revealed signs of discoloration and necrosis in samples collected in front of the cultivated area. No sign of damage was observed at other sites. In addition, a notable increase of the number of dying thalli on the bark of *Quercus* trees in front of a new parcel under cultivation was observed. These thalli are also widely parasitized by lichenicolous fungi.

260

265

3.4 Correlations between ILD, physiological parameters and heavy metals

Spearman rank correlation coefficients between the investigated parameters are shown in Table 5. In general, the lichen diversity (ILD) was negatively correlated with the content of heavy metals and TBARS and positively with chlorophyll integrity and the content of secondary metabolites. The content of heavy metals (but Ni) was positively correlated with that of TBARS and negatively with chlorophyll integrity. A negative correlation emerged between all investigated elements and usnic acid and between the content of Fe, Cd, Zn, As, Pb and caperatic acid. In addition, Fe, Cd, Zn, As, Pb were positively correlated with dehydrogenase activity. Concerning physiological parameters, $OD_{435/415}$ was positively correlated with the content of ergosterol and secondary metabolites, caperatic acid negatively with TBARS and usnic acid negatively with dehydrogenase activity. Chlorophyll *a* fluorescence emission was not significantly correlated with any of the other parameters.

275

280

4. Discussion

The fact that the lichen communities (diversity, number of species, presence of sensitive lichens) steadily indicated an improvement of environmental conditions, from high-moderate pollution to low-negligible effects according to the distance from the source, suggested that the dumping activities were the cause of the worsening of conditions compared with the surrounding area. This evidence agrees with a condition of environmental alteration limited to the sites directly facing the landfill, as reported by Paoli et al. (2012). In addition, the lichen diversity was negatively correlated with the content of heavy metals, whose depositions decreased with distance from the landfill.

285

Recent studies including a biodiversity assessment reported the use of higher plants and lichens as bioindicators around waste management facilities (Kotovicová et al., 2011; Paoli et al., 2014; Vaverková et al., 2012). For instance, vascular plants (and few lichen species) were used as biological indicators in Czech Republic to assess the potential impact of landfill sites on the surrounding soil and the atmospheric environment, based on presence of species, biomass production and the and occurrence of stress symptoms (Kotovicová et al., 2011; Vaverková et al., 2012). Data suggested a limited impact around landfill sites operating in compliance with the most modern and strictest requirements and standards. An assessment of lichen diversity around a composting plant of organic waste in C Italy revealed a local eutrophication due to atmospheric NH_3 depositions, which enhanced the diffusion of nitrophilous species (Paoli et al., 2014).

295

Airborne particulate matter arising from landfilling operations may be highly enriched in heavy

300

metals depending on the nature of the materials and the resuspension from the surface of the site (Chalvatzaki et al., 2014). A study carried out around a MSW landfill near Mexico City revealed that the resuspended surface dust from the landfill heavily influenced the composition of airborne particulate in the surrounding area (Vega et al., 2001). A study carried out around a MSW composting site in Crete (Greece) revealed dust emissions by erosion of the piles and subsequent depositions with maximum concentrations at a distance of 25-75 m downwind of the piles in the direction of prevailing wind. Concentrations above the 24 h limit for the protection of human health were occasionally measured at distances up to 275 m from the source (Chalvatzaki et al., 2012). Compared with the surrounding environment, the elemental content of our lichens around the landfill was enriched in Cd, Cr, Cu, Fe, Ni and Zn. Consistently, the analysis of airborne emissions from a landfill in the UK, indicated Cr, Fe, Ni, Pb and Zn as the main components of the particulate matter around the site (Koshy et al., 2009). Similarly, a study carried out with the lichen *Pseudevernia furfuracea* near Rome around the largest European MSW landfill showed a significant bioaccumulation of As, Cd, Cr, Cu, Ni, Pb and Zn (Protano et al., 2014). In our study, the heavy metal content decreased by 37-71% moving from sites directly facing the landfill to sites 200 m away. In this process it is noteworthy the buffering role of the vegetation belt dominated by oak trees surrounding the landfill in limiting the spreading of airborne particulate matter. Green belts are known sinks for air pollutants, including gases and particulate matter, and trees are able to act as filters through leaf adsorption, deposition of aerosols and particulate on leaf surfaces and fallout on leeward side of vegetation belts (Smith, 1990). Heavy metals originating from landfill operations likely affected the physiological status of *F. caperata*, reducing chlorophyll integrity and increasing membrane lipid peroxidation and dark respiration. In addition, higher chlorophyll degradation and membrane lipid peroxidation were correlated with sites facing the landfill characterized by lower lichen diversity. The fact that in lichens chlorophyll degradation is correlated with the accumulation of heavy metals is well documented (Garty et al., 2000). Moreover, the toxic effect of Cd, Cr, Cu and Ni on chlorophyll integrity and TBARS content has been specifically tested in lichens (Bačkor et al., 2009; 2010; Unal et al., 2010). Ergosterol, a sterol component of fungal cell membranes, can be affected by exposure to heavy metals around a waste landfill, as suggested by our results. Ergosterol is considered a good indicator of mycobiont viability, as its content correlates with the amount of metabolically active fungal cells (Ekblad et al., 1998) and laboratory treatments of lichen thalli with heavy metals (i.e. Cd, Cu and Ni) significantly reduced ergosterol content according to the dose (Bačkor et al., 2009; 2010). This is in agreement with a higher content of TBARS found in the thalli facing the landfill. Our samples increased the rate of respiration close to the landfill, as suggested by the level of dehydrogenase respiratory activity. This parameter is often taken as indicator of viability upon exposure to several stressors, including heavy metals (Bačkor and Fahsel, 2005), so that a decreased activity of respiratory dehydrogenases is negatively correlated with high doses of elements (Bačkor, 2011; Pisani et al., 2011). However, within a certain degree, their increase could be stress-induced and linked to the activity of other enzymes with antioxidant activity involved in the detoxification of heavy metals. The negative correlations between the content of heavy metals and the content of usnic and caperatic acid suggest that heavy metals affected the production of secondary lichen metabolites. It is noteworthy that usnic acid, a yellow cortical pigment exhibiting antibiotic effects, protects the lichen against bacteria and fungi and caperatic acid, a medullary compound, may have a similar role. Therefore, it is highly likely that *F. caperata* samples close to the landfill are parasitized by lichenicolous fungi since at these sites they have decreased protections against these parasites. Interestingly, the ratio between cortical and medullary secondary metabolites was found to increase in lichen samples transplanted in proximity of a local soil contamination source by waste dumping in Slovakia (Lackovičová et al., 2013). Similarly, due to the decrease of medullary caperatic acid, this ratio also increased in our samples, being 0.13 at a distance of 1500 and 200 m and 0.60 at the

landfill. All the above evidences suggest that an integrated use of lichen based methods, i.e. methods with a stronger link to source attribution (e.g., bioaccumulation), methods reflecting ongoing processes (e.g., physiological responses) and their environmental effects (e.g., biodiversity assessments), could contribute to the definition of environmental quality based on the biological effects of different waste management practices on sensitive components of the ecosystems (Paoli et al., 2015).

5. Conclusions

This study showed that lichens can profitably be used as bioindicators of the biological effects of air pollution around a municipal solid waste landfill and that they depict the status of environmental quality around such facilities. Lichens showed an impoverished flora and an increased accumulation of heavy metals at sites facing the landfill. In addition, also the physiological status was affected: lower chlorophyll integrity and higher membrane lipid peroxidation, as well as alteration of secondary metabolites were found in the investigated *F. caperata* samples. However, these results also highlighted that the atmospheric impact caused by the landfill is spatially limited, also due to the buffering action of a green oak belt surrounding the landfill, suggesting that successful and sustainable waste management may be associated with satisfactory environmental quality.

References

- Asta, J., Erhardt, W., Ferretti, M., Fornasier, F., Kirschbaum, U., Nimis, P.L., Purvis, O.W., Pirintsos, S., Scheidegger, C., Van Haluwyn, C., Wirth, V., 2002. Mapping lichen diversity as an indicator of environmental quality. In: Nimis, P.L., Scheidegger, C., Wolseley, P.A. (Eds.), *Monitoring with Lichens – Monitoring Lichens*. Kluwer Academic Publishers, pp. 273–279.
- Bačkor, M., 2011. Lichens and heavy metals: toxicity and tolerance. Pavol Jozef Šafárik University in Košice.
- Bačkor, M., Kováčik, J., Dzubaj, A., Bačkorová, M., 2009. Physiological comparison of copper toxicity in the lichens *Peltigera rufescens* (Weis) Humb. and *Cladina arbuscula* subsp. *mitis* (Sandst.) Ruoss. *Plant Growth Regul.* 58, 279–286.
- Bačkor, M., Kováčik, J., Piovar, J., Pisani, T., Loppi, S., 2010. Physiological aspects of cadmium and nickel toxicity in the lichens *Peltigera rufescens* and *Cladina arbuscula* subsp. *mitis*. *Water Air Soil Pollut.* 207, 253–262.
- Bačkor, M., Péli, E. R., Vantová, I., 2011. Copper tolerance in the macrolichens *Cladonia furcata* and *Cladina arbuscula* subsp. *mitis* is constitutive rather than inducible. *Chemosphere* 85, 106–113.
- Bačkor, M., Fahselt, D., 2005. Tetrazolium reduction as an indicator of environmental stress in lichens and isolated bionts. *Environ Exp Bot* 53, 125–133.
- Bargagli, R., Nimis, P.L., 2002. Guidelines for the use of epiphytic lichens as biomonitors of atmospheric deposition of trace elements. In: Nimis, P.L., Scheidegger, C., Wolseley, P.A. (Eds.), *Monitoring with Lichens – Monitoring Lichens*. Kluwer Academic Publishers, pp. 295–299.
- Bari, A., Rosso, A., Minciardi, M.R., Troiani, F., Piervittori, R., 2001. Analysis of heavy metals in atmospheric particulates in relation to their bioaccumulation in explanted *Pseudevernia furfuracea* thalli. *Environ Monit Assess* 69, 205–220.
- Bealey, W.J., Long, S., Spurgeon, D.J., Leith, I., Cape, J.N., 2008. Review and implementation study of biomonitoring for assessment of air quality outcomes. *Science Report SC030175/SR2*. Environment Agency, Bristol, pp. 1–170.
- Chalvatzaki, E., Aleksandropoulou, V., Lazaridis, M., 2014. A case study of landfill workers exposure and dose to particulate matter-bound metals. *Water Air Soil Pollut* 225, 1782.

- Chalvatzaki, E., Aleksandropoulou, V., Glytsos, T., Lazaridis, M., 2012. The effect of dust emissions from open storage piles to particle ambient concentration and human exposure. *Waste Manage*, 32, 2456–2468.
- 405 Chalvatzaki, E., Kopanakis, I., Kontaksakis, M., Glytsos, T., Kalogerakis, N., Lazaridis, M. 2010. Measurements of particulate matter concentrations at a landfill site (Crete, Greece). *Waste Manage*, 11, 2058–2064.
- Chiriac, R., De Araujos Morais, J., Carre, J., Bayard, R., Chovelon, J.M., Gourdon, R., 2011. Study of the VOC emissions from a municipal solid waste storage pilot-scale cell: Comparison with biogases from municipal waste landfill site. *Waste Manage* 31, 2294–2301.
- 410 Dahlman, L., Zetherström, M., Sundberg, B., Näsholm, T., Palmqvist, K., 2002. Measuring ergosterol and chitin in lichens. In: Kranner, I., Beckett, R., Varma, A. (Eds.). *Protocols in Lichenology: Culturing. Biochemistry. Ecophysiology and Use in Biomonitoring*. Springer-Verlag, pp. 348–362.
- 415 Ekblad, A., Wallander, H., Nasholm, T., 1998. Chitin and ergosterol combined to measure total and living biomass in ectomycorrhizae. *New Phytol* 138, 143–149.
- Garty, J., Weissman, L., Tamir, O., Beer, S., Cohen, Y., Karnieli, A., Orlovsky, L., 2000. Comparison of five physiological parameters to assess the vitality of the lichen *Ramalina lacera* exposed to air pollution. *Physiol Plantarum* 109, 410–418.
- 420 Han, S., Youn, J.S., Jung, Y.W., 2011. Characterization of PM₁₀ and PM_{2.5} source profiles for resuspended road dust collected using mobile sampling methodology. *Atmos Environ* 45, 3343–3351.
- Koshy, L., Jones, T., BéruBé, K., 2009. Characterization and bioreactivity of respirable airborne particles from a municipal landfill. *Biomarkers* 14, 49–53.
- 425 Kotovicová, J., Toman, F., Vaverková, M., Stejskal, B., 2011. Evaluation of waste landfills impact on the environment with the use of bioindicators. *Pol J Environ Stud*, 20, 371–377.
- Lackovičová, A., Guttová, A., Bačkor, M., Pišut P., Pišut, I., 2013. Response of *Evernia prunastri* to urban environmental conditions in Central Europe after the decrease of air pollution. *Lichenologist* 45, 89–100.
- 430 Loppi, S., 2014. Lichens as sentinels for air pollution at remote alpine areas (Italy). *Environ Sci Poll Res* 41, 2563–2571.
- Loppi, S., Frati, F., Paoli, L., Bigagli, V., Rosetti, C., Bruscoli, C., Corsini, A., 2004. Biodiversity of epiphytic lichens and heavy metal contents of *Flavoparmelia caperata* thalli as indicators of temporal variations of air pollution in the town of Montecatini Terme (central Italy). *Sci Tot Environ* 326, 113–122.
- 435 Loppi, S., Ivanov, D., Boccardi, R., 2002. Biodiversity of epiphytic lichens and air pollution in the town of Siena (central Italy). *Environ Pollut* 116, 123–128.
- Loppi, S., Putorti, E., De Dominicis, V., Barbaro, A., 1995. Lichens as bioindicators of air quality near a municipal solid waste incineration plant in central Italy. *Allionia* 33, 121–129.
- 440 Loppi, S., Putorti, E., Pirintsos, S.A., De Dominicis, V., 2000. Accumulation of heavy metals in epiphytic lichens near a municipal solid waste incinerator (central Italy). *Environ Monit Assess* 61, 361–371.
- Mittler, R., 2002. Oxidative stress, antioxidants and stress tolerance. *Trends Plant Sci* 7, 405–410.
- Nimis, P.L., Scheidegger, C., Wolseley, P.A. (Eds.), 2002. *Monitoring with Lichens – Monitoring Lichens*. NATO Science Series. Kluwer Academic Publishers, Dordrecht, Netherlands, pp. 408.
- 445 Nimis, P.L., Martellos, S., 2008. ITALIC – the Information System on Italian Lichens. Version 4.0. IN4.0/1. University of Trieste, Dept. of Biology. <http://dbiodbs.univ.trieste.it/>.
- Paoli, L., Benesperi, R., Proietti Pannunzi, D., Corsini, A., Loppi, S., 2014. Biological effects of ammonia released from a composting plant assessed with lichens. *Environ Sci Pollut Res* 21, 5861–5872.
- 450

- Paoli, L., Corsini, A., Bigagli, V., Vannini, J., Bruscoli, C., Loppi, S., 2012. Long-term biological monitoring of environmental quality around a solid waste landfill assessed with lichens. *Environ Pollut* 161, 70–75.
- 455 Paoli, L., Munzi, S., Guttová, A., Sardella, G., Loppi, S., 2015. Lichens as suitable indicators of the biological effects of atmospheric pollutants around a municipal solid waste incinerator (S Italy) *Ecol Indic* 52, 362–370.
- Paoli, L., Loppi, S., 2008. A biological method to monitor early effects of the air pollution caused by the industrial exploitation of geothermal energy. *Environ Pollut* 155, 383–388.
- 460 Pirintsos, S.A., Loppi, S., 2008. Biomonitoring atmospheric pollution: the challenge of times in environmental policy on air quality. *Environ Pollut* 151, 269–271.
- Pisani, T., Munzi, S., Paoli, L., Bačkor, M., Loppi, S., 2011. Physiological effects of arsenic in the lichen *Xanthoria parietina* (L.) Th. Fr. *Chemosphere* 82, 963–969.
- Pisani, T., Paoli, L., Gaggi, C., Pirintsos, S.A., Loppi, S., 2007. Effects of high temperature on epiphytic lichens: issues for consideration in a changing climate scenario. *Plant Biosyst* 141, 1–6.
- 465 Protano, C., Guidotti, M., Owczarek, M., Fantozzi, L., Blasi, G., Vitali, M., 2014. Polycyclic aromatic hydrocarbons and metals in transplanted lichen (*Pseudevernia furfuracea*) at sites adjacent to a solid waste landfill in Central Italy. *Arch Environ Contam Toxicol* 66, 471–481.
- 470 Sloof, J.E., 1995. Lichens as quantitative biomonitors for atmospheric trace-element deposition using transplant. *Atmos Environ* 29, 11–20.
- Smith, W.H., 1990. *Air Pollution and Forests: Interaction between air contaminants and forest ecosystems*. Second Edition, Springer-Verlag, New York.
- Sun, Y., Wang, Y.-N., Sun, X., Wu, H., Zhang, H., 2013. Production characteristics of N₂O during stabilization of municipal solid waste in an intermittent aerated semi-aerobic bioreactor landfill. *Waste Manage* 33, 2729–2736.
- 475 Tretiach, M., Candotto Carniel, F., Loppi, S., Carniel, A., Bortolussi, A., Mazzilis, D., Del Bianco, C., 2011. Lichen transplants as a suitable tool to identify mercury pollution from waste incinerator: a case study from NE Italy. *Environ Monit Assess* 175, 589–600.
- 480 Unal, D., Isik, N.O., Sukatar, A., 2010. Effects of chromium VI stress on photosynthesis, chlorophyll integrity, cell viability, and proline accumulation in lichen *Ramalina farinacea*. *Russ J Plant Physiol* 57, 664–669.
- Vaverková, M., Toman, F., Kotovicová, J., 2012. Research into the occurrence of some plant species as indicators of landfill impact on the environment. *Pol J Environ Stud* 21, 755–762.
- 485 Vega, E., Mugica, V., Reyes, E., Sanchez, G., Chow, J.C., Watson, J.G., 2001. Chemical composition of fugitive dust emitters in Mexico City. *Atmos Environ* 35, 4033–4039.

490

495

500

Table 1. Scale for the interpretation of trace element concentrations in lichens ($\mu\text{g/g dw}$) desumed after Bargagli and Nimis (2002).

air pollution	As	Cd	Cr	Cu	Fe	Ni	Pb	Zn
very low	<0.4	<0.3	<2	<9	<400	<1.5	<10	<35
low	1.2	0.6	4	15	800	3.0	25	65
moderate	1.9	1.2	6	25	1200	5.0	55	95
high	2.5	2.1	13	40	1600	7.0	95	135
very high	>2.5	>2.1	>13	>40	>1600	>7	>95	>135

505

510

515

520

525

530

535

540

545 Table 2. Index of Lichen Diversity (ILD) and list of lichen species. ILD was interpreted in terms of air pollution according to paragraph 2.6. Significant differences between sites ($P < 0.05$) are indicated by a different letter. Pr = percentage of relevés where each species has been recorded, Fr = frequency expressed as percentage. Nomenclature of the species follows Nimis and Martellos (2008); " nitrophilous lichen.

	landfill		200 m		1500 m	
Index of Lichen Diversity (impact of air pollution)	45 ± 12 ^c (high - moderate)		72 ± 13 ^b (moderate - low)		113 ± 28 ^a (low - negligible)	
Lichen species	Pr	Fr	Pr	Fr	Pr	Fr
<i>Lepraria</i> sp.	89	61	86	74	50	32
<i>Candelariella reflexa</i> "	78	53	58	39	67	53
<i>Flavoparmelia caperata</i>	42	16	69	51	75	50
<i>Parmelia sulcata</i>	28	6	50	21	17	3
<i>Cladonia coniocraea</i>	22	13	36	21	8	2
<i>Leprocaulon microscopicum</i>	22	14	19	9		
<i>Physcia adscendens</i> "	22	8	3	3	92	57

<i>Punctelia borreri</i>	19	8	39	17	75	62
<i>Lecanora strobilina</i>	17	8	56	27		
<i>Normandina pulchella</i>	17	8	22	9	50	45
<i>Parmotrema perlatum</i>	17	3	22	7	8	2
<i>Flavoparmelia soledians</i>	11	7	14	7	58	45
<i>Melanelixia fuliginosa</i>	8	3	56	42	8	5
<i>Hyperphyscia adglutinata</i> "	8	4	11	4	33	15
<i>Candelaria concolor</i> "	8	2	3	1	50	48
<i>Candelariella xanthostigma</i>	8	5				
<i>Pertusaria albescens</i>	3	1	14	6	25	5
<i>Evernia prunastri</i>	3	1	6	2	8	2
<i>Ramalina farinacea</i>	3	1	6	1		
<i>Xanthoria parietina</i> "	3	1			17	3
<i>Aplotomma turgida</i>	3	1				
<i>Physcia tenella</i> "			11	4		
<i>Phlyctis argena</i>			8	4	42	27
<i>Lecanora symmicta</i>			8	3		
<i>Lecidella elaeochroma</i> "			6	2	33	18
<i>Parmelina tiliacea</i>			6	2	33	17
<i>Amandinea punctata</i> "			6	4	8	5
<i>Catillaria nigroclavata</i>			3	2		
<i>Lecanora carpinea</i> "			3	1		
<i>Physconia servitii</i>					42	28
<i>Lecanora chlarotera</i> "					33	12
<i>Physconia grisea ssp.grisea</i> "					33	8
<i>Collema subflaccidum</i>					25	7
<i>Physconia perisidiosa</i>					17	3
<i>Phaeophyscia orbicularis</i> "					8	8
<i>Physcia clementei</i>					8	2

Table 3. Content of trace elements ($\mu\text{g/g}$) in the lichen *Flavoparmelia caperata* as a function of distance from the landfill. Heavy metals were interpreted in terms of air pollution according to the intervals given in Table 1.

Elements	landfill	200 m	1500 m	ANOVA
As	0.49 ± 0.10^a low	0.31 ± 0.05^b very low	0.18 ± 0.031^c very low	$F=29.12$ $P=0.000$
Cd	0.62 ± 0.01^a moderate	0.37 ± 0.01^b low	0.24 ± 0.03^c very low	$F=32.35$ $P=0.000$
Cr	7.5 ± 4.2^a high	2.2 ± 0.3^b low	1.3 ± 0.2^b very low	$F=13.32$ $P=0.003$
Cu	17.8 ± 6.8^a moderate	9.9 ± 1.0^b low	6.6 ± 0.1^b very low	$F=13.08$ $P=0.000$
Fe	1092 ± 413^a moderate	525 ± 20^b low	275 ± 7^b very low	$F= 18.67$ $P=0.000$
Ni	6.0 ± 1.6^a high	3.6 ± 0.3^b moderate	4.4 ± 0.2^{ab} moderate	$F=13.82$ $P=0.000$
Pb	22.9 ± 12.6^a low	6.9 ± 2.4^b very low	3.1 ± 0.1^b very low	$F=13.61$ $P=0.000$
Zn	79.5 ± 27.1^a moderate	47.5 ± 6.1^b low	33.6 ± 2.4^b very low	$F=13.02$ $P=0.000$

5

565

570

575

580

585 Table 4. Physiological parameters in the lichen *Flavoparmelia caperata* as a function of distance from the landfill: chlorophyll integrity ($OD_{435/415}$), potential quantum yield of PSII as indicator of photosynthetic efficiency (F_v/F_M), dehydrogenase activity as indicator of lichen viability (A_{492}/g dw), TBARS, thiobarbituric acid reactive substances ($\mu\text{mol}/g$ dw), ergosterol content (mg/g dw), caperatic and usnic acid ($\%$ dw). Significant differences between sites ($P < 0.05$) are indicated by a different letter.

590

Parameters	landfill	200 m	1500 m	ANOVA
$OD_{435/415}$	0.95 ± 0.02^a	1.02 ± 0.04^b	1.07 ± 0.03^b	$F=25.68$ $P=0.000$
F_v/F_M	0.68 ± 0.10	0.73 ± 0.04	0.76 ± 0.02	$F=2.026$ $P=0.161$

Dehydrogenase	7.9 ± 3.4^a	5.3 ± 1.3^b	4.1 ± 1.6^b	$F=3.79 P=0.042$
TBARS	19.8 ± 4.6^a	14.7 ± 3.5^a	7.3 ± 1.4^b	$F=12.41 P=0.000$
ergosterol	0.39 ± 0.09^a	0.43 ± 0.05^a	0.64 ± 0.11^b	$F=8.639 P=0.007$
caperatic acid (% dw)	4.7 ± 3.2^a	7.7 ± 0.9^{ab}	8.6 ± 0.8^b	$F=3.74 P=0.048$
usnic acid (% dw)	1.37 ± 0.14^a	1.07 ± 0.31^c	1.19 ± 0.31^{ab}	$F=10.08 P=0.003$

Table 5. Spearman correlation coefficients between the investigated parameters: Index of Lichen Diversity (ILD), trace elements (Cr, Fe, Ni, Cu, Zn, As, Cd, Pb), chlorophyll integrity (OD_{435/415}), thiobarbituric acid reactive substances (TBARS), dehydrogenase activity, caperatic and usnic acid, ergosterol content, potential quantum yield of PSII (F_v/F_M).

595

	ILD	Cr	Fe	Ni	Cu	Zn	As	Cd	Pb	OD _{435/415}	TBARS	Dehydrogenase	Caperatic acid	Usnic acid	Ergosterol
ILD	-														
Cr	-0.76***	-													
Fe	-0.78***	0.99***	-												
Ni	-0.53*	0.56*	0.66**	-											
Cu	-0.75***	0.99***	0.99***	0.53*	-										
Zn	-0.75***	0.92***	0.93***	n.s.	0.95***	-									
As	-0.75***	1.00***	0.99***	0.54*	1.00***	0.93***	-								
Cd	-0.74***	0.89***	0.90***	0.69**	0.87***	0.75***	0.88***	-							
Pb	-0.91***	0.87***	0.90***	0.61*	0.86***	0.90***	0.86***	0.81***	-						
OD_{435/415}	0.61*	-0.68**	-0.68**	n.s.	-0.70***	-0.78***	-0.68**	-0.55*	-0.78***	-					
TBARS	-0.51*	0.76***	0.73***	n.s.	0.76***	0.69**	0.76***	0.67**	0.58**	n.s.	-				
Dehydrogenase	n.s.	n.s.	0.52*	n.s.	n.s.	0.53*	0.50*	0.54*	0.54*	n.s.	n.s.	-			
Caperatic acid	0.54*	n.s.	-0.51*	n.s.	n.s.	-0.57*	-0.51*	-0.56*	-0.65*	0.74**	n.s.	-0.77**	-		
Usnic acid	0.63*	-0.82***	-0.81***	-0.73**	-0.81***	-0.72**	-0.81***	-0.79***	-0.64*	0.63*	-0.66**	n.s.	n.s.	-	
Ergosterol	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	0.63*	n.s.	n.s.	n.s.	n.s.	-
F_v/F_M	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.