

SHORT NOTE

Assessing the ecological status of Italian lagoons using a biomass-based index

Michele Mistri^a, Angel Borja^b, Floriana Aleffi^c, Claudio Lardicci^d, Davide Tagliapietra^e, Cristina Munari^a

^a*Department of Chemical and Pharmaceutical Sciences, University of Ferrara, Via Fossato di Mortara 17, 44121 Ferrara, Italy*

^b*AZTI-Tecnalia, Marine Research Division, Herrera Kaia, Portualdea s/n, 20110 Pasaia, Spain*

^c*ARPA FVG, Osservatorio Alto Adriatico, Via La Marmora 13, 34139 Trieste, Italy*

^d*Department of Biology, University of Pisa, Via Derna 1, 56126 Pisa, Italy*

^e*CNR-ISMAR, Arsenale di Venezia, Tesa 104, 30122 Venezia, Italy*

**Corresponding author: msm@unife.it (M. Mistri)*

Tel: +39 0532 455736; Fax: +39 0532 249761

Abstract

We compared the performance of abundance- and biomass-based M-AMBI in the 13 major Italian lagoons, using a benthic dataset constituted by 208 sampling sites. The relative importance of ecological groups changed when using abundance or biomass, sometimes leading to an improved ecological status classification. Being biomass more ecologically relevant than abundance, the adoption of a biomass-based index may better describe the ecological status of lagoons, where the community is naturally disturbed and dominated by tolerant and opportunist species.

Keywords:

Ecological status, M-AMBI, M-bMAMBI, Lagoons, Mediterranean Sea

Ecological indicators are an effective way to characterize marine ecosystem health, and their number is rapidly increasing. Among the biological quality elements highlighted by the European Directive 2000/60/EC, benthic macrofauna is known to be probably the best effective indicator of pollution stress, as it shows predictive responses to different levels of natural and anthropogenic impact. Based on the Pearson & Rosenberg (1978) paradigm several biotic indices have been proposed in recent years. The marine biotic index AMBI/M-AMBI (Borja et al., 2000; Muxika et al., 2007) is probably the most widely used benthic index all over the world. In Europe, for example, many Countries have officially adopted the index for the description of ecological quality of coastal waters (Bulgaria, France, Germany, Italy, Romania, Slovenia and Spain; Borja et al., 2009; Birk et al., 2012). AMBI relies on the calculation of the biotic coefficient, which is based in turn on the proportion of disturbance-sensitive taxa and is expressed on a continuous scale ranging from 0 (best status) to 6 (worst status). The AMBI approach follows a model (Grall and Glemarec, 1997) which categorizes benthic invertebrates into five ecological groups (from EG-I, sensitive, to EG-V, first order opportunists), depending on their dominance along a gradient of organic enrichment. Recently, Warwick et al. (2010) suggested to estimate AMBI using biomass (bAMBI) and production (pAMBI). This because in an assemblage the abundance of a species can be relatively a poor measure of its functional importance, particularly in stressed situations when the insensitive species tend to be small bodied opportunists (Warwick et al., 2010). Muxika et al. (2012) successfully assessed the proposed modification to AMBI along the Basque coast (northern Spain), showing that those AMBI modifications were highly correlated and thus useful to assess the benthic quality status, if boundaries between quality classes were re-determined. More recently, Mistri and Munari (2015) tested the performance of biomass-based AMBI (bAMBI and pAMBI) in transitional ecosystems, finding good agreement between the response of all biomass-based indices and disturbance expressed by the severity of pressures. The use of a biomass-based index for the assessment of the ecological quality status in transitional systems is not trivial, since several studies (Magni et al., 2009; Munari and Mistri, 2010; Sigovini et al., 2013; Prato et al., 2014) suggested

that the use of indices based on species tolerance/sensitivity need to be adapted where the community is naturally disturbed and dominated by small-sized opportunists. The use of biomass in calculating M-AMBI (M-bAMBI) was tested by Cai et al. (2014, 2015) in the assessment of the benthic quality status of Bohai Bay (north of China), a shallow water basin receiving industrial and municipal wastewater from coastal cities (Cai et al., 2014). Those authors found that M-bAMBI seemed more effective than M-AMBI in indicating human pressures of the Bay (Cai et al., 2015).

Since lagoons are often characterized by high benthic biomass (McLusky, 1989), and biomass is a measure of ecosystem functions (Warwick et al., 2010), in this note we explore the performance of biomass-based M-AMBI (M-bAMBI) in the most important Italian lagoons, at which the disturbance status was known. We assembled a data set of macrofaunal counts from 13 large Italian lagoons (Fig. 1) occurring along a cline of 7° of latitude (between 45°44'N and 39°56'N). Along Italian coasts there are almost 170 lagoons, but 140 of them have a surface area <10 km². With the exclusion of Orbetello Lagoon and Stagno di Tortoli (Tyrrhenian Sea), all the largest lagoons (e.g. Grado-Marano, Venice, Po Delta, Comacchio, Lesina) are located along the Western Adriatic coasts. Our data set comprises all main Adriatic and Tyrrhenian Italian lagoons (Caleri, Marinetta, Barbamarco, Canarin, Scardovari and Goro are those in the present Po Delta). In Table 1, major geographical and ecological features of the 13 lagoons are shown. A total of 103 sites, representative of the different habitats found within each lagoon, were sampled repeatedly over time, for a total of 208 sampling points. In Table 1 the main benthic community parameters at each of the 13 lagoons (mean number of species, S, and mean diversity, H') are shown. Over 400 macrobenthic taxa were gathered at the 208 sampling points, with annelids displaying the highest number of taxa, followed by crustaceans and molluscs. Most species found in the 13 lagoons are cosmopolitan (i.e. with a wide geographical distribution, such as *Capitella capitata*, *Polydora ciliata*, *Streblospio shrubsolii*, and *Hediste diversicolor*). Several endemic species were also found (such as *Corophium orientale*, *Microdeutopus algicola*, *Pectinaria koreni* and *Ampihthoe riedli*), together with many non-indigenous species (such as *Anadara inaequalis*, *A. demiri*, *Arcuatula*

senhousia, *Ruditapes philippinarum*, *Rhithropanopeus harrisi*, *Dyspanopeus sayii* and *Grandidierella japonica*), which were most common in the Po Delta lagoons.

Pressures (Supplementary Material, Table S1) were quantified (1: low, 2: medium and 3: high) for each location and sampling station, as partial pressure, total pressure and as a pressure index (PI), following an approach close to that proposed by Aubry and Elliott (2006), based upon best professional judgment. According to Borja et al. (2011) the total pressure was the sum of partial pressures, and the pressure index was calculated as an average value of the pressures. Abundance (M-AMBI) and biomass (M-bAMBI) based indices were calculated using AMBI 5.0 software (freely available at <http://ambi.azti.es>). Reference conditions were those reported by the Italian Act 260/10, which considers three lagoon typologies, with different reference conditions, as a function of tidal range and salinity (Table 2). Regression between AMBI-based indices and Pressure Index (PI) was performed to analyze the agreement in the pollution classification, and significance was assessed through regression ANOVA. In Fig. 2 the relationship between PI and abundance-based indices (AMBI and M-AMBI) and biomass-based indices (bAMBI and M-bAMBI) at the 103 lagoonal stations is shown (M-AMBI: $F=53.6$; $P<0.001$; AMBI: $F=10.4$; $P<0.01$; M-bAMBI: $F=39.4$; $P<0.001$; bAMBI: $F=19.7$; $P<0.001$). The relationship between M-AMBI and M-bAMBI was fitted using a trend line (Fig. 3; $F=116.9$; $P<0.001$). The formula for the trend line was then used to calculate values of M-bAMBI corresponding to M-AMBI values separating status categories. Agreement between classification obtained through M-AMBI and M-bAMBI was determined by considering only two ecological status: “Undisturbed” and “Disturbed”. The undisturbed status was determined when the derived status was High/Good, and scored as “1”. Disturbed status corresponded to Moderate/Poor/Bad, and scored as “0”. The non-parametric Wilcoxon pairs test was used to assess agreement or disagreement between M-AMBI and M-bAMBI classification on the undisturbed and disturbed status of sites on a statistical basis, and showed a mismatch of 22.6% of cases in which one index classified the status of stations as “Undisturbed” and the other “Disturbed” ($Z=4.76$; $P<0.001$). Compared to M-AMBI, there were

more sites assigned to an undisturbed status in M-bAMBI. In Fig. 4, the ecological status (ES) gathered through M-AMBI and M-bAMBI is shown. At certain lagoons, differences in ES were minimal, e.g. at Comacchio, where the benthic community is in such a state of deep suffering (Munari and Mistri, 2014) that the ES was unsatisfactory, either by using abundance or biomass. At other lagoons (e.g. Grado Marano, Venice and Po Delta), instead, a slight improvement in the ES can be noticed, with some sites changing their ES from Moderate to Good.

The AZTI index (Borja et al., 2000) bases its functioning on dividing benthic species into previously defined ecological groups (EG-I to EG-V), and then determining the respective proportion of the different groups in the benthic community. Its calculation rely on the relative decrease of sensitive species (EG-I) confronted with increasing disturbance in the sediment or, conversely, the increase of species that are resistant or indifferent to disturbance (EG-II and EG-III), or that are even encouraged by such conditions like the opportunist species (EG-IV and EG-V) that proliferate when the sediment is rich in organic matter. Different level of disturbance was present at our 103 sampling station, and this was reflected by the relative abundance of ecological groups with, e.g., EG-I dominant at some stations at Grado Marano, Baiona, Lesina and Tortoli lagoons, and EG-V dominant, e.g., at Comacchio, and some sites in the Po Delta and Orbetello (Fig. 5). The different distribution of EG-I to EG-V taxa in our 208 sites was obviously reflected by M-AMBI scores. We are however aware that AMBI/M-AMBI, considering the abundances of stress-tolerant species to detect anthropogenic impacts, does not take into account the fact that tolerant species may also be tolerant of natural stressors (the "Estuarine Quality Paradox", Dauvin and Ruellet, 2009). Warwick et al. (2010) suggested to use production data instead of abundance because production is the most common measure of ecosystem function. However, given the difficulty of production assessment, biomass is often used as a proxy measure (terHorst and Munguia, 2008). Since biomass (and thus production) is more ecologically relevant than abundance, its use to derive AMBI-based indices is intriguing, especially in lagoons where the community is naturally disturbed and dominated by tolerant and opportunist species.

Assessing the ES of Bohai Bay (Yellow Sea, China), Cai et al. (2014) found M-AMBI and M-bAMBI to produce quite similar results, with slight lower M-bAMBI values. Those authors justified lower M-bAMBI values with the lower diversity values calculated with biomass compared with those calculated using density. We also found H' calculated with biomass to be generally lower than H' calculated with abundance (Fig. 6), however, differently from Cai et al. (2014) we found slight lower M-AMBI values. In their study on the Basque Coast, Muxika et al. (2012) found that the distributions of ecological groups' dominances were very similar when biomass was used instead of abundance. Conversely, in our lagoonal data set, the proportion of ecological groups into the community varied greatly if we considered abundance or biomass-based data (Fig. 5). In certain cases the use of biomass instead of abundance resulted in a better ES classification, despite the "Moderate/Good" boundary for M-bAMBI was higher than that for M-AMBI (0.739 vs 0.710). For example, the site 4.09 in the Caleri lagoon (Po Delta) scored the ES "Moderate" (EQR=0.694) by M-AMBI, but "Good" (EQR=0.926) by M-bAMBI. At this site, considering abundance, the benthic community was numerically dominated by EG-III (60.5%), followed by and EG-V (20.8), and EG-IV (14.2%). Numerically dominant species were *Streblospio shrubsolii* (48.7% of the whole community abundance), *Capitella capitata* (15.9%), *Polydora ciliata* (14.2%), and *Hydroides dianthus* (7.9%). These figures changed when we considered biomass, since *S. shrubsolii* constituted 11.3% of the total biomass, *C. capitata* (3.7%), *Polydora ciliata* (3.2%), and *Hydroides dianthus* (0.7%). Conversely, *Palaemon serratus* constituted 19.8% of community biomass (but only 0.3% of community abundance), and *Neanthes succinea* constituted 14.5% of total biomass (1.3% of total abundance). Lagoonal macrobenthos is mainly composed by species tolerant to the naturally disturbed conditions, which include highly variable temperature, salinity and oxygen concentration (Elliott and Quintino, 2007). Because of the dominance in terms of abundance of tolerant species, the use of abundance-based indices does not always guarantee a correct assessment of ES, as it occurs in the marine environment.

Italian lagoons exhibit different and peculiar characteristics depending on their geographical, hydrodynamic and ecological features: these variations generate composite gradients that involve salinity, marine water renewal (e.g. residence time), nutrients, turbidity and sediment structure. (Tagliapietra et al., 2009). Superimposed to this huge variability there is a century-old human activity. Coastal lagoons represent important and fragile ecosystems in the coastal landscape, providing key ecosystems services such as water quality improvement, fisheries resources, habitat and food for migratory and resident animals. For the evaluation of their ecological status, the measure adopted (abundance or biomass) can lead to changes in results. Because of lagoonal inherent variability that leads to the coexistence of many species spanning all ecological groups, the relative contribution of the different macrobenthic taxa can vary greatly depending on the adopted unit. The concept of ecological status must take into account the structure and function of the lagoon ecosystem, and results from this study suggest that a biomass-based index may better describe the ecological status of Italian lagoons.

References

- Aubry, A., Elliott, M., 2006. The use of environmental integrative indicators to assess seabed disturbance in estuaries and coasts: application to the Humber Estuary, UK. *Mar. Pollut. Bull.* 53, 175-185.
- Birk, S., Bonne, W., Borja, A., Brucet, S., Courrat, A., Poikane, S., Solimini, A., van de Bund, W., Zampoukas, N., Hering, D., 2012. Three hundred ways to assess Europe's surface waters: An almost complete overview of biological methods to implement the Water Framework Directive. *Ecol. Ind.* 18, 31-41.
- Borja, A., Franco, J., Pérez, V., 2000. A marine biotic index to establish the ecological quality of soft-bottom benthos within European estuarine and coastal environments. *Mar. Pollut. Bull.* 40, 1100-1114.
- Borja, A., Miles, A., Occhipinti-Ambrogi, A., Berg, T., 2009. Current status of macroinvertebrate methods used for assessing the quality of European marine waters: implementing the Water Framework Directive. *Hydrobiologia* 633, 181-196.
- Borja, A., Barbone, E., Basset, A., Borgersen, G., Brkljacic, M., Elliott, M., Garmendia, J.M., Marques, J.C., Mazik, K., Muxika, I., Neto, J.M., Norling, K., Rodríguez, J.G., Rosati, I., Rygg, B., Teixeira, H., Trayanova, A., 2011. Response of single benthic metrics and multi-metric methods to anthropogenic pressure gradients, in five distinct European coastal and transitional ecosystems. *Mar. Pollut. Bull.* 62, 499-513.
- Cai, W., Borja, A., Liu, L., Meng, W., Muxika, I., Rodríguez, J.G., 2014. Assessing benthic health under multiple human pressures in Bohai Bay (China), using density and biomass in calculating AMBI and M-AMBI. *Mar. Ecol.* 35, 180-192.
- Cai, W., Borja, A., Liu, K., Zhu, Y., Zhou, J., Liu, L., 2015. Assessing the benthic quality status of the Bohai Bay (China) with proposed modifications of M-AMBI. *Acta Oceanol. Sin.* 34, 111-121.

- Dauvin, J.C., Ruellet, T., 2009. The estuarine quality paradox. Is it possible to define an ecological quality status for specific modified and naturally stressed estuarine ecosystems? *Mar. Pollut. Bull.* 59, 38-47.
- Elliott, M., Quintino, V., 2007. The estuarine quality paradox, environmental homeostasis and the difficulty of detecting anthropogenic stress in naturally stressed areas. *Mar. Pollut. Bull.* 54, 640-645.
- European Community, 2000. Directive 2000/60/EC of the European Parliament and of the Council of 23 October 2000 establishing a framework for community action in the field of water policy. *Official Journal of the European Communities* 43 (L327), Bruxelles.
- Grall, J., Glemarec, M., 1997. Using biotic indices to estimate macrobenthic community perturbations in the Bay of Brest. *Estuar. Coast. Shelf Sci.* 44, 43-53.
- Magni, P., Tagliapietra, D., Lardicci, C., Balthis, L., Castelli, A., Como, S., Frangipane, G., Giordani, G., Hyland, J., Maltagliati, F., Pessa, G., Rismondo, A., Tataranni, M., Tomassetti, P., Viaroli, P., 2009. Animal-sediment relationships: evaluating the Pearson–Rosenberg paradigm in Mediterranean coastal lagoons. *Mar. Pollut. Bull.* 58, 478-486.
- Mistri, M., Munari, C., 2015. The performance of biomass-based AMBI in lagoonal ecosystems. *Mar. Pollut. Bull.* 99, 126-137.
- Munari C., Mistri, M., 2014. Traditional management of lagoons for fishery can be inconsistent with restoration purposes: the Valli di Comacchio study case. *Chem. Ecol.* 30, 653-665.
- Muxika, I., Borja, A., Bald, J., 2007. Using historical data, expert judgement and multivariate analysis in assessing reference conditions and benthic ecological status, according to the European Water Framework Directive. *Mar. Pollut. Bull.* 55,16-29.
- Muxika, I., Somerfield, P.J., Borja, A., Warwick, R.M., 2012. Assessing proposed modifications to the AZTI marine biotic index (AMBI), using biomass and production. *Ecol. Ind.* 12, 96-104.
- Pearson, T.H., Rosenberg, R., 1978. Macrobenthic succession in relation to organic enrichment and pollution of the marine environment. *Oceanogr. Mar. Biol. Annu. Rev.* 16, 229-311.

- Prato S., La Valle P., De Luca E., Lattanzi L., Migliore G., Morgana J.G., Munari C., Nicoletti L., Izzo G., Mistri M., 2014. The "one-out, all-out" principle entails the risk of imposing unnecessary restoration costs: a study case in two Mediterranean coastal lakes. *Mar. Pollut. Bull.* 80, 30-40.
- Sigovini, M., Keppel, E., Tagliapietra, D., 2013. M-AMBI revisited: looking inside a widely-used benthic index. *Hydrobiologia* 717, 41-50.
- Tagliapietra, D., Sigovini, M., Volpi Ghirardini, A., 2009. A review of terms and definitions to categorise estuaries, lagoons and associated environments. *Mar. Freshw. Res.* 60, 497-509.
- terHorst, C.P., Munguia, P., 2008. Measuring ecosystem function: consequences arising from variation in biomass-productivity relationship. *Community Ecology* 9, 39-44.
- Warwick, R.M., Clarke, K.R., Somerfield, P.J., 2010. Exploring the marine biotic index (AMBI): variations on a theme by Ángel Borja. *Mar. Pollut. Bull.* 60, 554-559.

Figure Legends

Fig. 1. Location of the considered lagoons.

Fig. 2. Relationship between Pressure Index (PI) and abundance-based (AMBI and M-AMBI) and biomass-based (bAMBI and M-bAMBI) indices.

Fig. 3. Relationship between M-AMBI and M-bAMBI.

Fig. 4. Ecological status at the various lagoons through M-AMBI (dark grey) and M-bAMBI (light grey).

Fig. 5. Concentration of EG-I to V at each sampling site considering abundance or biomass (GrMar: Grado Marano; V-C: Venice Central; V-Chi: Venice Chioggia; V-PdR: Venice Palude della Rosa; Po Delta: Caleri, Marinetta, Barbamarco, Canarin, Scardovari, Goro; Com: Comacchio; Bai: Baiona; Les: Lesina; Orb: Orbetello; Tor: Tortoli).

Fig. 6. Relationship between abundance-based and biomass-based diversity (H').



Fig. 1




Fig. 2




Fig. 3




Fig. 4.




Fig. 5




Fig. 6

Table 1.

Major geographical and ecological features of the 13 studied lagoons

Lagoon	Latitude	Longitude	Area km ²	Mean depth m	Salinity	Typology	Sampling sites	Tot n samples	S (± SD)	H' (± SD)
Grado										
Marano	45°42'N	13°20'E	160	1.5	poly/euhaline	mt	21	21	25.6 ± 14.3	2.9 ± 1.0
Venice	45°24'N	12°19'E	500	2.5	poly/euhaline	mt	20	43	18.4 ± 12.4	2.1 ± 0.6
Caleri	45°05'N	12°18'E	11	2.0	meso/polyhaline	mt	4	8	14.0 ± 6.8	1.8 ± 0.6
Marinetta	45°03'N	12°21'E	10	0.8	meso/polyhaline	mt	6	12	14.6 ± 4.1	1.3 ± 0.8
Barbamarco	45°00'N	12°27'E	8	0.8	meso/polyhaline	mt	2	4	15.5 ± 8.7	1.6 ± 0.5
Canarin	44°55'N	12°29'E	10	0.8	meso/polyhaline	mt	3	6	12.7 ± 2.3	2.0 ± 0.9
Scardovari	44°51'N	12°24'E	32	1.5	meso/polyhaline	mt	5	10	22.3 ± 4.4	2.3 ± 0.5
Goro	44°49'N	12°18'E	37	2.0	meso/polyhaline	mt	16	16	26.2 ± 6.3	1.9 ± 0.9
Comacchio	44°36'N	12°10'E	117	0.8	euhaline	nt	4	22	9.6 ± 4.7	1.7 ± 0.7
Baiona	44°30'N	12°14'E	12	1.0	polyhaline	mt	3	6	32.5 ± 3.5	3.0 ± 0.2
Lesina	41°52'N	15°26'E	51	0.8	meso/polyhaline	nt	4	12	14.4 ± 4.3	2.3 ± 0.5
Orbetello	42°26'N	11°11'E	27	1.5	polyhaline	nt	9	36	13.1 ± 6.6	2.0 ± 0.7
Tortoli	39°56'N	09°40'E	3	1.0	poly/euhaline	nt	6	12	25.3 ± 8.5	3.1 ± 0.8

Table 2.

Reference conditions for various lagoon typologies.

Tidal range	Salinity	AMBI	H'	S
Not tidal	-	1.85	3.3	25
Microtidal	Oligo-meso-poly	2.14	3.4	28
Microtidal	Eu-iper	0.63	4.23	46