Breaking of the gauge symmetry in lattice gauge theories

Claudio Bonati,¹ Andrea Pelissetto,² and Ettore Vicari¹

¹Dipartimento di Fisica dell'Università di Pisa and INFN Largo Pontecorvo 3, I-56127 Pisa, Italy

²Dipartimento di Fisica dell'Università di Roma Sapienza and INFN Sezione di Roma I, I-00185 Roma, Italy

(Dated: July 30, 2021)

We study perturbations that break gauge symmetries in lattice gauge theories. As a paradigmatic model, we consider the three-dimensional Abelian-Higgs (AH) model with an N-component scalar field and a noncompact gauge field, which is invariant under U(1) gauge and SU(N) transformations. We consider gauge-symmetry breaking perturbations that are quadratic in the gauge field, such as a photon mass term, and determine their effect on the critical behavior of the gauge-invariant model, focusing mainly on the continuous transitions associated with the charged fixed point of the AH field theory. We discuss their relevance and compute the (gauge-dependent) exponents that parametrize the departure from the critical behavior (continuum limit) of the gauge-invariant model. We also address the critical behavior of lattice AH models with broken gauge symmetry, showing an effective enlargement of the global symmetry, from U(N) to O(2N), which reflects a peculiar cyclic renormalization-group flow in the space of the lattice AH parameters and of the photon mass.

Introduction. Gauge symmetries play a key role in the construction of the theoretical models of fundamental interactions [1, 2] and in the description of emergent phenomena in condensed-matter and statistical physics [3–8]. They may be exact, as in the Standard Model of fundamental interactions, or effectively emerge at low energies, as in some many-body systems. Effectively emergent gauge symmetries have also been discussed in the context of fundamental interactions, see, e.g., Refs. [3, 9–11]. In this case, they may arise from microscopic interactions of different nature, such as string models [12].

To correctly interpret experimental results in terms of models with an emergent gauge symmetry, a solid understanding of the effects of gauge-symmetry violations is essential. This issue is crucial in the context of analog quantum simulations, for example, when controllable atomic systems are engineered to effectively reproduce the dynamics of gauge-symmetric theoretical models, with the purpose of obtaining physical information from the experimental study of their quantum dynamics in laboratory. Several proposals of artificial gauge-symmetry realizations have been reported, see, e.g., Refs. [13, 14] and references therein (see also Refs. [15–20] for some experimental realizations), in which the gauge symmetry is expected to effectively emerge in the low-energy dynamics. A possible strategy is that of adding a *penalty* term to the Hamiltonian, that suppresses the interactions violating the gauge symmetry. This strategy assumes that gauge-symmetry breaking (GSB) terms become negligible at low energies, thereby making the dynamics effectively gauge invariant in this limit [13, 21, 22]. In spite of the relevance of these issues, there is at present little understanding of the effects of GSB perturbations on the continuum limit of quantum or statistical systems with gauge symmetries, or equivalently on the critical behavior close to continuous transitions, where long-range correlations develop, realizing the corresponding quantum field theory.

In this paper we address this problem by considering three-dimensional (3D) lattice gauge theories, obtained by discretizing the action of corresponding quantum field theories. We study the role of GSB perturbations at the critical transitions of gauge-invariant models, to understand whether and when they are relevant, i.e. they break gauge invariance in the low-energy or large-distance behavior (continuum limit). If this is the case, GSB terms may lead to different continuum limits, as we shall see.

The model. As a paradigmatic model, we consider the 3D scalar electrodynamics or Abelian-Higgs (AH) field theory, with an N-component complex scalar field $\Phi(\boldsymbol{x})$ coupled to the electromagnetic field $A_{\mu}(\boldsymbol{x})$. Its Lagrangian density reads [2]

$$\mathcal{L} = |D_{\mu}\Phi|^{2} + w \,\Phi^{*}\Phi + \frac{u}{4} \,(\Phi^{*}\Phi)^{2} + \frac{1}{4g^{2}} \,(\partial_{\mu}A_{\nu} - \partial_{\nu}A_{\mu})^{2} \,,$$
(1)

where $D_{\mu} \equiv \partial_{\mu} + iA_{\mu}$. The AH theory is invariant under U(1) gauge and SU(N) global transformations. Its 3D renormalization-group (RG) flow has a stable charged (with nonzero gauge coupling) fixed point (FP) for $N \geq N_c$ [23, 24], with $N_c = 7(2)$ [25, 26]. According to the RG theory [27–30], the charged FP is expected to describe the critical behavior, and therefore the continuum limit, of U(1) gauge models with SU(N) global symmetry.

Lattice representations of the continuum theory (1) differ for the topological nature of the lattice gauge field. One can either use the real field $A_{x,\mu}$ as in the continuum theory (noncompact model) or the link variables $\lambda_{x,\mu} \in U(1)$ (compact model, corresponding to $e^{iA_{x,\mu}}$). In this work we mostly consider the 3D noncompact AH (ncAH) model defined on cubic lattices of size L^3 , which has a continuous transition line for $N > N_c$, along which the continuum limit is described by the 3D AH field theory (1) [25, 31, 32]. The fundamental fields are unitlength N-component complex vectors \mathbf{z}_x ($\bar{\mathbf{z}}_x \cdot \mathbf{z}_x = 1$) defined on the lattice sites \mathbf{x} and real fields $A_{x,\mu}$ defined on the lattice links. The lattice action is

$$S_{\text{AH}}(\boldsymbol{z}, \boldsymbol{A}) = -J N \sum_{\boldsymbol{x}, \mu} 2 \operatorname{Re} \left(\bar{\boldsymbol{z}}_{\boldsymbol{x}} \cdot \lambda_{\boldsymbol{x}, \mu} \, \boldsymbol{z}_{\boldsymbol{x}+\hat{\mu}} \right) \qquad (2)$$
$$+ \frac{1}{4g_0^2} \sum_{\boldsymbol{x}, \mu\nu} (\Delta_{\mu} A_{\boldsymbol{x}, \nu} - \Delta_{\nu} A_{\boldsymbol{x}, \mu})^2 \,,$$

where $\lambda_{\boldsymbol{x},\mu} \equiv e^{iA_{\boldsymbol{x},\mu}}$, g_0 is the lattice gauge coupling, $\hat{\mu}$ are unit vectors along the lattice directions, and $\Delta_{\mu}A_{\boldsymbol{x},\nu} = A_{\boldsymbol{x}+\hat{\mu},\nu} - A_{\boldsymbol{x},\nu}$. The action S_{AH} has a global SU(N) symmetry, $\boldsymbol{z}_{\boldsymbol{x}} \to V\boldsymbol{z}_{\boldsymbol{x}}$ with $V \in \text{SU}(N)$, and a local U(1) gauge symmetry, $\boldsymbol{z}_{\boldsymbol{x}} \to e^{i\theta_{\boldsymbol{x}}}\boldsymbol{z}_{\boldsymbol{x}}$ and $A_{\boldsymbol{x},\mu} \to A_{\boldsymbol{x},\mu} + \theta_{\boldsymbol{x}} - \theta_{\boldsymbol{x}+\hat{\mu}}$. We consider C^* boundary conditions [25, 33, 34] (see also App. A), to remove the degeneracy under $A_{\boldsymbol{x},\mu} \to A_{\boldsymbol{x},\mu} + 2\pi n_{\mu}$ with $n_{\mu} \in \mathbb{Z}$, obtaining well defined expectation values for gauge-invariant operators $O(\boldsymbol{z}, \boldsymbol{A})$,

$$\langle O(\boldsymbol{z}, \boldsymbol{A}) \rangle = \frac{\sum_{\{\boldsymbol{z}, \boldsymbol{A}\}} O(\boldsymbol{z}, \boldsymbol{A}) \ e^{-S_{\text{AH}}(\boldsymbol{z}, \boldsymbol{A})}}{\sum_{\{\boldsymbol{z}, \boldsymbol{A}\}} e^{-S_{\text{AH}}(\boldsymbol{z}, \boldsymbol{A})}} \,. \tag{3}$$

The phase diagram of the ncAH model (2) with N > 2 is characterized by a Coulomb phase for small J (shortranged scalar and long-ranged gauge correlations), a Higgs phase for large J and small g_0 (condensed scalarfield and gapped gauge correlations), and a molecular phase for large J and g_0 (condensed scalar-field and long-ranged gauge correlations) [25]. They are separated by three transition lines, which are continuous or of first order depending on N. In particular, for $N > N_c = 7(2)$, the ncAH model undergoes continuous transitions between the Coulomb and Higgs (CH) phases, for $0 < g_0^2 \lesssim 4$. The corresponding critical behavior is described by the charged FP of the 3D AH field theory [25]. For $g_0 \to 0$, one has $A_{\boldsymbol{x},\mu} \to 1$ modulo gauge transformations, so that one recovers the O(2N) vector model. We consider the gauge-invariant bilinear operator

$$Q_{\boldsymbol{x}}^{ab} = \bar{z}_{\boldsymbol{x}}^{a} z_{\boldsymbol{x}}^{b} - \frac{1}{N} \delta^{ab} , \qquad (4)$$

which transforms as $Q_x \to V^{\dagger} Q_x V$ under global SU(N) transformations. It provides an effective order parameter for the spontaneous breaking of the global SU(N) symmetry.

GSB perturbations. We study how perturbations breaking the U(1) gauge symmetry affect the CH critical behavior. In this exploratory study we consider the quadratic perturbation

$$P_M = \frac{r}{2} \sum_{\boldsymbol{x},\mu} A_{\boldsymbol{x},\mu}^2 \,, \tag{5}$$

which can be interpreted as a photon mass term. Such a mass term is generally introduced as an infrared regulator in perturbative computations in quantum electrodynamics [2]. We also consider the local quadratic operators

$$P_L = \frac{a}{2} \sum_{\boldsymbol{x}} \left(\sum_{\mu} \Delta_{\mu} A_{\boldsymbol{x},\mu} \right)^2, \quad P_A = \frac{b}{2} \sum_{\boldsymbol{x}} \left(\sum_{\mu} n_{\mu} A_{\boldsymbol{x},\mu} \right)^2, \quad (6)$$

2

where n_{μ} is an arbitrary unit vector. When added to the ncAH action, i.e., if one considers $S = S_{AH} + P_{\#}$, all quadratic terms defined in Eqs. (5) and (6) break gauge invariance, leaving a global U(N) symmetry $z_x \to U z_x$, $U \in U(N)$. However, they affect the critical behavior quite differently. The mass term (5) is expected to be relevant at the CH transitions, since it drastically changes the long-distance properties of the gauge-field correlations. In particular, the Coulomb phase disappears in the presence of a photon mass. Therefore, as soon as the perturbation is turned on (r > 0), the system is expected to flow out of the charged AH FP. On the other hand, the quadratic terms P_L and P_A , cf. Eq. (6), may be interpreted as the result of the Fadeev-Popov procedure for a gauge fixing [2], being related to the Lorentz $(\partial_{\mu}A_{\mu}=0)$ and axial $(\boldsymbol{n}\cdot\boldsymbol{A}=0)$ gauge fixing [35], respectively. If they are the only GSB perturbations present in the model, they are expected to be irrelevant for gaugeinvariant correlations (more precisely, their presence does not change gauge-invariant expectation values). However, as we shall see below, they play a role, when they are added to the action together with the mass term (5), as they make the limit $r \to 0$ well defined.

Relevance of the GSB perturbations. To characterize the strength of the perturbation P_M , we compute the corresponding RG dimension $y_r > 0$. This exponent provides information on how to scale r to keep GSB effects small. Indeed, when the correlation length ξ increases, approaching the continuum limit, one should decrease r faster than ξ^{-y_r} to ensure that GSB effects are negligible. We estimate y_r by finite-size scaling (FSS) analyses of Monte Carlo (MC) data. We consider the correlation function $\langle \operatorname{Tr} Q_x Q_y \rangle$ of the operator Q_x defined in Eq. (4), and the corresponding second-moment correlation length ξ . We consider RG-invariant quantities R, such as $R_{\xi} = \xi/L$ and the Binder parameter $U = \langle \mu_2^2 \rangle / \langle \mu_2 \rangle^2$, where $\mu_2 = \sum_{x,y} \operatorname{Tr} Q_x Q_y$. At continuous transitions driven by the parameter J, they are expected to behave as [30]

$$R(L, J, g_0) \approx f_R(X) + O(L^{-\omega}), \ X = (J - J_c) L^{1/\nu}, \ (7)$$

where ν is the length-scale critical exponent, and $\omega > 0$ is the exponent controlling the leading scaling corrections. It is also useful to consider the FSS relation [36]

$$U = F_U(R_{\xi}) + O(L^{-\omega}), \qquad (8)$$

where F_U is a universal function independent of any normalization. To estimate y_r , we consider the behavior of the RG invariant quantities R in the presence of the GSB term (5). In the large-L limit, we expect [37]

$$R(L, J, g_0, r) \approx \mathcal{F}_R(X, Y), \qquad Y = r L^{y_r}, \qquad (9)$$

which holds provided [38] that $y_r > 1/\nu$, where ν is the thermal exponent of the gauge model (along the CH transition line we have $1/\nu = 1.387(6)$, 1.247(12) for N = 15, 25, respectively). Eq. (9) is the usual FSS relation for

a multicritical point in systems with a global symmetry. However, in the present case its validity is not obvious, given that the mass term P_M is not well defined in the (r = 0) gauge-invariant noncompact theory: averages of the mass term can only be computed in the presence of a maximal gauge fixing [35, 39], such as the axial (using C^* conditions) or Lorentz ones. For these reasons, we consider three different actions with GSB terms:

M1:
$$S_1 = S_{AH} + P_M$$
, (10)

M2:
$$S_2 = S_{AH} + P_L + P_M$$
, (11)

M3:
$$S_3 = S_{AH} + P_M$$
 with $A_{x,3} = 0$, (12)

where M2 can be associated with the Lorentz gauge, and M3 is defined imposing the axial gauge. We expect Eq. (9) to be well defined in models M2 and M3, while its validity in model M1 is instead not clear.

Numerical estimates of the RG dimensions. We performed MC simulations for N = 15 and N = 25 along the CH transition line (estimates of the critical points and exponents can be found in Ref. [25]), for the three models M#, see App. A for details. The results confirm that P_M is relevant. Indeed, for fixed r, there is a clear departure from the gauge-invariant (r = 0) critical behavior. In Fig. 1 we show results for N = 25 at the critical point. The exponent y_r is estimated by fitting the data at J_c to Eq. (9), setting X = 0. We obtain $y_r = 1.4(1)$ for M2 [for both a = 1 and a = 10, cf. Eq. (6)] and $y_r = 2.55(5)$ for M3. We also mention that if we apply Eq. (9) to U computed in M1 without gauge fixing, we obtain the effective estimate $y_r \approx 1.4$, see top Fig. 1, confirming the relevance of P_M along the CH transition line. Analogous results are obtained for N = 15, in particular $y_r = 2.55(10)$ for M3. The exponent y_r turns out to depend on the gauge fixing, indicating that the gauge fixing influences the RG properties of the mass perturbation. Apparently, gauge-dependent modes, that are controlled by the gauge fixing term, are crucial in determining the effects of the photon mass term. Note that y_r is quite large, therefore the corresponding GSB perturbation must decrease rapidly with L —faster than L^{-y_r} —to keep GSB effects under control.

Critical behaviors in the presence of finite GSB terms. We now address the behavior of the ncAH model in the presence of a finite GSB term such as the photon mass. Also for finite r we expect a transition at a finite value $J_c(r)$, with $J_c(r = 0) = J_c$, where J_c is the CH transition point in the gauge-invariant model. Since the charged fixed point is unstable with respect to P_M , we expect the transition to belong to a different universality class, which should only depend on the global symmetry of the model. Although the global symmetry group for r > 0 is U(N), we will now argue that continuous transitions at $J_c(r)$ are characterized by a larger O(2N)invariance group. We note that, since gauge fields are not expected to be relevant for $r \neq 0$, one can use the standard Landau-Ginzburg-Wilson (LGW) approach [27–30] to predict the critical behavior. Since the gauge sym-




FIG. 1: Data of U at the critical point $J_c \approx 0.295515$ of the ncAH model for N = 25 and $g_0^2 = 2.5$, as a function of $Y = rL^{y_r}$. Results for models M1 (without gauge fixing, top) and M3 (axial gauge, bottom).

metry is broken, $z_{\boldsymbol{x}}$ represents the microscopic orderparameter field. Therefore, the LGW basic field is an *N*-component complex vector $\Psi(\boldsymbol{x})$. The Lagrangian is the sum of the kinetic term $|\partial_{\mu}\Psi|^2$ and of the most general U(*N*)-invariant quartic potential:

$$\mathcal{L}_{\text{LGW}} = \partial_{\mu} \Psi^* \cdot \partial_{\mu} \Psi + w \ \Psi^* \cdot \Psi + \frac{u}{4} \left(\Psi^* \cdot \Psi \right)^2.$$
(13)

It is easy to check that \mathcal{L}_{LGW} is actually O(2N) invariant. Indeed, there are no dimension-2 and 4 U(N) invariant operators that break the O(2N) symmetry. The lowest-dimension operators that are not O(2N) symmetric have dimension six close to four dimensions—for instance, $(\text{Im } \Phi^* \cdot \partial_\mu \Phi)^2$ —and thus they are expected to be irrelevant at the 3D O(2N) FP. Therefore, the critical behavior of generic vector systems with global U(N) invariance (without gauge symmetries) is expected to belong to the O(2N) universality class, implying an effective enlargement of the global symmetry of the critical modes (restricted only to the critical region).

The above analysis can be extended to lattice AH models with compact gauge fields (cAH), using the link variables $\lambda_{\boldsymbol{x},\mu} \in \mathrm{U}(1)$ and the pure gauge action $S_{\lambda} =$ $-g_0^{-2} \sum_{\boldsymbol{x},\mu \neq \nu} \operatorname{Re} \lambda_{\boldsymbol{x},\mu} \lambda_{\boldsymbol{x}+\hat{\mu},\nu} \bar{\lambda}_{\boldsymbol{x}+\hat{\nu},\mu} \bar{\lambda}_{\boldsymbol{x},\nu}$ in Eq. (2). Unlike ncAH models, cAH models with $N \geq 2$ present only two phases, separated by a disorder-order transition line where gauge correlations are not critical [31]. Since the scalar fields turns out to be the only critical degrees of freedom, the effective description of the transitions is provided by the SU(N)-invariant LGW Φ^4 theory with a matrix gauge-invariant order parameter, corresponding to Q_x^{ab} in Eq. (4) [31, 40]. For N = 2 this LGW theory has a stable O(3) vector FP, thus predicting O(3) continuous transitions [30] for any gauge coupling $g_0 > 0$, including $g_0 \to \infty$ [for $g_0 \to 0$, instead, the model becomes equivalent to the O(4) vector model]. This has been also confirmed numerically [31]. Gauge invariance can be broken by adding $P_M = -r \sum_{\boldsymbol{x},\mu} \operatorname{Re} \lambda_{\boldsymbol{x},\mu}$, which plays the role of a photon mass for $\lambda_{\boldsymbol{x},\mu}$ close to 1. When




FIG. 2: Estimates of U versus R_{ξ} for the model M1 with N = 25, $g_0^2 = 2.5$, r = 1. We also report results for the gauge-invariant model (r = 0) and the O(50) vector model (full line, obtained by large-L extrapolations of MC data for the appropriate spin-2 correlations, see Ref. [31] and App. A). The results for r = 1 appear to converge toward the O(50) universal curve, consistently with $O(L^{-\omega})$ corrections with $\omega \approx 1$, supporting the RG prediction reported in the text.

the gauge symmetry is effectively broken (as discussed in Ref. [41], this requires r to be sufficiently large), the critical behavior should be described by the LGW theory (13), which predicts that continuous transitions belong to the O(4) vector universality class.

The RG predictions at fixed r are confirmed by numerical results for both ncAH and cAH models. In Fig. 2 we plot U versus R_{ξ} for the ncAH model with N = 25 and r = 1. The data around the critical point $J_c(r)$ are expected to converge to a universal curve, cf. Eq. (8), which can be compared with the analogous curves of models that belong to known universality classes. The data approach the O(2N) vector universal curve (obtained using an appropriate operator that corresponds to Q_x^{ab} in the O(2N) model [31]), confirming the LGW RG argument. For the cAH model with N = 2, we observe an asymptotic O(4) vector behavior for r = 1 and r = 2.25, in agreement with the general arguments in App. A.

Various classes of GSB perturbations. On the basis of the results presented in this paper, we may distinguish three classes of GSB perturbations. (i) First, there are GSB perturbations that are relevant at the stable FP of the lattice gauge-invariant theory. They drive the system out of criticality and may give rise to a different critical behavior. The photon mass term (5) plays this role along the CH line in the ncAH model. (ii) A second class corresponds to gauge fixings and GSB perturbations like those appearing in Eq. (6). If they are the only GSB terms present in the model, they are irrelevant: gaugeinvariant observables are unchanged. However, if they are present together with some relevant GSB perturbation, they play a role: the RG flow close to the charged FP depends both on the gauge-fixing and on the relevant perturbation. This may be due to the fact that a gauge fixing is needed to make non-gauge-invariant correlations




FIG. 3: Sketch of the cyclic RG flow of the ncAH model in the space of the parameters J, g_0 and r, showing an unusual loop between the O(2N) and the charged FP C(N).

well defined in the gauge-invariant theory or to the role of gauge-dependent modes that are sensitive to gauge fixings. (iii) GSB perturbations associated with RG operators with negative RG dimensions, whose effects are suppressed in the critical (continuum) limit.

When the added GSB perturbations are relevant, the lattice system may develop a different critical behavior or continuum limit. This is the case of the ncAH model with a photon mass term, which has a global U(N) invariance. Quite interestingly, the transitions in this model belong to the O(2N) vector universality class, with an effective enlargement of the global symmetry at the transition. This symmetry enlargement is expected in any model in which the GSB perturbation is relevant and it preserves the global U(N) symmetry.

Cyclic RG flow. It is worth noting that the above results lead to a peculiar RG flow, see Fig. 3 for a sketch in the coupling space (J, g_0, r) . For $g_0^2 \to 0$, the gauge fields are frozen, and the model is equivalent to the O(2N) vector model, whose critical behavior is controlled by the corresponding O(2N) FP. If the gauge interactions are turned on, i.e., one sets $g_0 > 0$ keeping r = 0, the systems flows towards the charged FP of the AH field theory, which is stable for any $0 < g_0^2 \lesssim 4$. Finally, if a photon mass is added, i.e., one sets r > 0, since the charged FP is unstable under this perturbation, the RG flow goes back to the O(2N) FP, which is now stable, independently of q_0 and r. This RG behavior can be hardly reconciled with an irreversibility of the RG flow, analogous to that generally associated with the monotonic properties implied by the *c*-theorem of 2D critical systems [42, 43], see also Refs. [44–49] for similar proposals in 3D systems [50].

Conclusions. In conclusion, we have studied the effect of GSB perturbations on the critical behavior—or, equivalently, the continuum limit—of gauge-invariant theories. The behavior at charged FP turns out to be more complicated than that observed when global symmetries are broken. In particular, we observe apparent violations of universality. For instance, the RG dimension of the same GSB perturbation appears to depend on local gaugefixing conditions, a result that, we believe, should be further investigated. Moreover, GSB perturbations give rise to unexpected phenomena, like the cyclic RG flow sketched in Fig. 3.

Several extensions are called for, to achieve a satisfactory understanding of the problem and to identify its universal features, such as the study of other lattice gauge theories—in particular, it would be interesting to extend the analysis to the nonabelian gauge groups— and of other classes of GSB perturbations, for example preserving residual discrete gauge subgroups (such approximations may be useful for analog simulations). It would also be important to rephrase and extend the present results to quantum Hamiltonian systems [51] (see Refs. [52, 53] for recent works addressing issues related to GSB effects and the approach to the continuum limit).

Acknowledgments

Numerical simulations have been performed on the CSN4 cluster of the Scientific Computing Center at INFN-PISA.

Appendix A: The numerical analyses

In this appendix we provide some details on the numerical computations reported in the paper.

1. The models

In most of the simulations we have considered the noncompact Abelian-Higgs (ncAH) model. The fundamental fields are unit-length N-component complex vectors $\boldsymbol{z}_{\boldsymbol{x}}$ $(\bar{\boldsymbol{z}}_{\boldsymbol{x}} \cdot \boldsymbol{z}_{\boldsymbol{x}} = 1)$ defined on the lattice sites and real fields $A_{\boldsymbol{x},\mu}$ defined on the lattice links. The lattice action is

$$S_{\rm ncAH}(\boldsymbol{z}, \boldsymbol{A}) = S_{\boldsymbol{z}}(\boldsymbol{z}, \boldsymbol{A}) + S_{nc}(\boldsymbol{A}), \qquad (A1)$$

where

$$S_{z}(\boldsymbol{z}, \boldsymbol{A}) = -J N \sum_{\boldsymbol{x}, \mu} 2 \operatorname{Re} \left(\bar{\boldsymbol{z}}_{\boldsymbol{x}} \cdot \lambda_{\boldsymbol{x}, \mu} \, \boldsymbol{z}_{\boldsymbol{x}+\hat{\mu}} \right), \quad (A2)$$
$$S_{nc}(\boldsymbol{A}) = \frac{1}{4g_{0}^{2}} \sum_{\boldsymbol{x}, \mu\nu} (\Delta_{\mu} A_{\boldsymbol{x}, \nu} - \Delta_{\nu} A_{\boldsymbol{x}, \mu})^{2},$$

 $\lambda_{\boldsymbol{x},\mu} \equiv e^{iA_{\boldsymbol{x},\mu}}, g_0$ is the lattice gauge coupling, $\hat{\mu}$ are unit vectors along the lattice directions, and $\Delta_{\mu}A_{\boldsymbol{x},\nu} = A_{\boldsymbol{x}+\hat{\mu},\nu} - A_{\boldsymbol{x},\nu}$.

We have also considered the lattice compact AH model (cAH), with action

$$S_{cAH}(\boldsymbol{z}, \boldsymbol{\lambda}) = S_{\boldsymbol{z}}(\boldsymbol{z}, \boldsymbol{\lambda}) + S_{c}(\boldsymbol{\lambda}),$$
 (A3)

where $S_z(\boldsymbol{z}, \boldsymbol{\lambda})$ is given by Eq. (A2), while

$$S_{c}(\boldsymbol{\lambda}) = -g_{0}^{-2} \sum_{\boldsymbol{x},\mu \neq \nu} \operatorname{Re} \lambda_{\boldsymbol{x},\mu} \lambda_{\boldsymbol{x}+\hat{\mu},\nu} \,\bar{\lambda}_{\boldsymbol{x}+\hat{\nu},\mu} \,\bar{\lambda}_{\boldsymbol{x},\nu} \,. \quad (A4)$$

To simulate the ncAH model, it is not possible to use periodic boundary conditions, since all gauge-invariant observables associated to loops that wrap around the lattice are not bounded and their average values are ill-defined. As in our previous work [25], we consider C^* boundary conditions. They are used here for the cAH model too, although periodic boundary conditions would be appropriate, as well. We consider cubic lattices of size L, so that C^* boundary conditions amount to the identifications (see Ref. [25] for a thorough discussion)

$$A_{\boldsymbol{x}+L\hat{\nu},\mu} = -A_{\boldsymbol{x},\mu} , \qquad \boldsymbol{z}_{\boldsymbol{x}+L\hat{\nu}} = \bar{\boldsymbol{z}}_{\boldsymbol{x}} . \qquad (A5)$$

To be consistent with Eq. (A5), local gauge transformations are defined by $A_{\boldsymbol{x},\mu} \to A_{\boldsymbol{x},\mu} + \alpha(\boldsymbol{x} + \hat{\mu}) - \alpha(\boldsymbol{x})$, with an antiperiodic function $\alpha(\boldsymbol{x})$: $\alpha(\boldsymbol{x} + L\hat{\nu}) = -\alpha(\boldsymbol{x})$. As a consequence, observables that involve a nontrivial wrapping around the lattice are not gauge invariant.

 C^* boundary conditions are very convenient when implementing axial gauges. Indeed, it is possible to fix $A_{x,3} = 0$ —or, more generally, $\sum_{\mu} n_{\mu} A_{x,\mu} = 0$ — on all lattice sites, at variance with the case of periodic boundary conditions. From the explicit construction discussed in Ref. [25], it follows that no residual gauge freedom is left once $A_{x,3} = 0$ is enforced on all lattice sites. It is important to note that also the Lorentz gauge is a maximal gauge, with no residual gauge freedom. Indeed, suppose the opposite, i.e., that there are two different gauge configurations $A_{x,\mu}^{(1)}$ and $A_{x,\mu}^{(2)}$ that are related by a gauge transformation, $A_{x,\mu}^{(2)} = A_{x,\mu}^{(1)} + \Delta_{\mu}\alpha(x)$, and that both satisfy the condition $\sum_{\mu} \Delta_{\mu} A_{x,\mu}^{(i)} = 0$. The function $\alpha(x)$ must satisfy

$$\sum_{\mu} \Delta_{\mu} [\Delta_{\mu} \alpha(\boldsymbol{x})] = 0, \qquad (A6)$$

which implies that α is a zero eigenmode of the lattice Laplacian. For C^* boundary conditions there is no zero mode, as α is antiperiodic, proving that $A_{\boldsymbol{x},\mu}^{(1)} = A_{\boldsymbol{x},\mu}^{(2)}$. For periodic boundary conditions, there is one zero mode, $\alpha(\boldsymbol{x}) = C$, where C is space independent, so that also in this case $A_{\boldsymbol{x},\mu}^{(1)} = A_{\boldsymbol{x},\mu}^{(2)}$. We have considered the ncAH model for N = 15 and

We have considered the ncAH model for N = 15 and 25, fixing in both cases $g_0^2 = 2.5$. For this value of g_0 the ncAH model undergoes a continuous transition for $J = J_c$. As discussed in Ref. [25], such transition is controlled by the charged fixed point (FP) of the Abelian-Higgs field theory. We have performed simulations for $J = J_c$ on lattices of size $L \leq 96$ (N = 25) and $L \leq 64$ (N = 15). For N = 15 we used the estimate J_c reported in Ref. [25], see Table I. For N = 25 we used an improved estimate. We performed additional simulations for $J \approx J_c$ on larger lattices (while in Ref. [25] we limited ourselves to lattices with $L \leq 64$, here we consider values of L up to L = 96) and reanalyzed the data. The result is $J_c = 0.295515(4)$, which is consistent with the estimate $J_c = 0.295511(4)$ reported in Ref. [25]. We have also considered the cAH model for N = 2, the only case where

N	g_0^2		J_c
	25	2.5	0.295515(4)
	15	2.5	0.309798(6)
	2	∞	0.7102(1)

TABLE I: Critical values J_c of the coupling J for the values of g_0^2 used in the present simulations. The value of J_c for N = 25 is an improvement of the estimate of Ref. [25]. Results for N = 15 and N = 2 are taken from Ref. [25] (note that $1/g_0^2$ was named κ), and Ref. [40], respectively.




FIG. 4: Estimates of R_{ξ} (top) and U (bottom) at the critical point $J_c \approx 0.309798$ of the ncAH model for N = 15 and $g_0^2 = 2.5$, as a function of rL^{y_r} , for model M3 (axial gauge).

a continuous transition is present. The parameter g_0^2 does not play any role [31] and we have therefore set $1/g_0^2 =$ 0 (no gauge action); the estimate of the corresponding critical value J_c is also reported in Table I.

Beside simulations of the gauge model, we have also performed simulations of the O(2N) spin model with action S_z and $\lambda_{x,\mu} = 1$, measuring the same quantities we compute in the gauge model (see Appendix B of Ref. [31] for a discussion of the relation between correlation functions of the CP^{N-1} order parameter Q^{ab} and spin-two correlations in the vector O(2N) spin model). We considered N = 2 and N = 25, determining U and R_{ξ} , and in particular the universal curve $U = F(R_{\xi})$.

2. Technical details: simulations and data analysis

In the Monte Carlo simulations we use an overrelaxation algorithm, obtained by combining Metropolis updates of the scalar and of the gauge fields and microcanonical updates of the scalar field. The latter moves are obtained by generalizing the usual reflection moves used in O(N) models. We perform a Metropolis update of the z and gauge fields every 5 (on the largest lattices, every 10) microcanonical updates of the scalar field. Trial states for the Metropolis updates are generated by adding a random number to $A_{x,\mu}$ and by multiplying z_x by a random 2×2 unitary matrix close to the identity; in both cases, we tune the update to have an acceptance rate of approximately 30%.

The typical statistics (discarding thermalization) of each data point varies in the range $4 \times 10^5 \cdot 2 \times 10^6$. Errors are estimated by using a combination of jackknife and blocking procedures. In all cases, the autocorrelation times were at most of the order of 10^2 updates.

To compute the estimates of y_r , we have assumed that U and R_{ξ} satisfy the scaling relation

$$R(r,L) = f(rL^{y_r}) + L^{-\omega}g(rL^{y_r})$$
(A7)

at the critical point $J = J_c$. In the fits we have approximated the scaling functions with polynomials and we have analyzed U and R_{ξ} together, looking for a value of y_r the minimizes the sum of the residuals for the two observables. The value of ω is unknown. In the absence of GSB terms, we expect ω to be close to 1 (for $N = \infty$ we have $\omega = 1$), but, once GSB terms are added, new irrelevant operators may appear and ω may be smaller. For this reason we have determined y_r for values of ω in the interval 0.2-1. Except for model M1, the ω dependence is at most of the same order of the statistical errors.

3. Results for N = 15

To investigate the N dependence of the exponent y_r , beside considering the ncAH model with N = 25 we have also considered the same model for N = 15. We have only studied model M3 (axial gauge), obtaining

$$y_r = 2.55(10),$$
 (A8)

which should be compared with the result for N = 25, $y_r = 2.55(5)$. The N dependence is apparently small, much smaller than the statistical errors. In Fig. 4 we show R_{ξ} and U against rL^{y_r} (the analogous plot of U for N = 25 is shown in the main text). The ratio R_{ξ} shows a very nice scaling behavior with small scaling corrections, which increase as rL^{y_r} increases. The Binder parameter shows larger corrections, that have the opposite behavior: they decrease as rL^{y_r} increases.

4. Results for the compact model with N = 2

As discussed in the main text, we also performed an exploratory investigation of the effect of explicit gauge breaking terms using the compact discretization. We studied the N = 2 cAH model, which is the only one in which a continuous transition is present [31]. In this case we defined the photon mass operator as


$$P_M = -r \sum_{\boldsymbol{x},\mu} \operatorname{Re} \lambda_{\boldsymbol{x},\mu} , \qquad (A9)$$

and performed simulations using the axial gauge. We set $\lambda_{x,3} = 1$ (this the analogue of the condition $A_{x,3} = 0$ used in the noncompact case) on all sites. Note that this is possible as we use C^* boundary conditions also for the compact model.

As for N = 25 we studied the behavior of the model with action $S_{cAH} + P_M$ for two finite values of r, r = 1and r = 2.25. In Fig. 5 we report the Binder parameter versus R_{ξ} . Data for r = 2.25 are perfectly consistent with an O(4) behavior, confirming the expected symmetry enlargement. The results for r = 1 instead are still quite far from the O(4) curve, although they show the correct trend. Except for values of R_{ξ} where U has a maximum, as L increases from 16 to 64, the data move towards the O(4) universal curve. Close to the maximum, the behavior is nonmonotonic, but the data start moving towards the O(4) curve for $L \geq 48$. A thorough investigation of the compact model will be reported in a forthcoming publication [41].

Phys. B **915**, 135 (2017).

- [10] D. Foerster, H. B. Nielsen, N. Ninomiya, Dynamical stability of local gauge symmetry, Phys. Lett. B 94, 135 (1980).
- [11] J. Iliopoulos, D. V. Nanopoulos, and T. N. Tomaras, Infrared stability of anti-grandunification, Phys. Lett. B 94, 141 (1980).
- [12] J. Polchinski, String theory, (Cambridge University Press, 1998).
- [13] E. Zohar, J. I. Cirac, and B. Reznik, Quantum simulations of lattice gauge theories using ultracold atoms in optical lattices, Rep. Prog. Phys. 79, 014401 (2015).
- [14] M. C. Bañuls, *et al*, Simulating lattice gauge theories with quantum technologies, Eur. Phys. J. D 74, 165 (2020).
- [15] E. A. Martinez, *et al*, Real-time dynamics of lattice gauge theories with a few-qubit quantum computer, Nature 534, 516 (2016).
- [16] H. Bernien, et al, Probing many-body dynamics on a 51atom quantum simulator, Nature 551, 579 (2017).
- [17] N. Klco, et al, Quantum-classical computation of Schwinger model dynamics using quantum computers Phys. Rev. A 98, 032331 (2018).
- [18] C. Schweizer, et al, Floquet approach to Z_2 lattice gauge



2.0

1.9

FIG. 5: Estimates of U versus R_{ξ} for the cAH model M1 with $N = 2, g_0^2 = \infty$: results for r = 1 (top panel), and r = 2.25 (bottom panel). The critical couplings are $J_c \approx 0.552$ and $J_c \approx 0.328$ for r = 1 and r = 2.25 respectively. We also report results for the gauge-invariant model (r = 0) and the O(4) vector model (full line, obtained by large-L extrapolations of MC data for the appropriate spin-2 correlations, see Ref. [31]).

- S. Weinberg, *The Quantum Theory of Fields*, (Cambridge University Press, 2005).
- [2] J. Zinn-Justin, Quantum Field Theory and Critical Phenomena, fourth edition (Clarendon Press, Oxford, 2002).
- [3] X.-G. Wen, Quantum field theory of many-body systems: from the origin of sound to an origin of light and electrons, (Oxford University Press, 2004)
- [4] P. W. Anderson, Superconductivity: Higgs, Anderson and all that, Nat. Phys. 11, 93 (2015).
- [5] S. Gazit, F. F. Assaad, S. Sachdev, A. Vishwanath, and C. Wang, Confinement transition of Z₂ gauge theories coupled to massless fermions: emergent QCD₃ and SO(5) symmetry, Proceedings of the National Academy of Sciences 115, E6987 (2018).
- [6] S. Sachdev, Topological order, emergent gauge fields, and Fermi surface reconstruction, Rep. Prog. Phys. 82, 014001 (2019).
- [7] S. Sachdev, H. D. Scammell, M. S. Scheurer, and G. Tarnopolsky, Gauge theory for the cuprates near optimal doping, Phys. Rev. B 99, 165126 (2019).
- [8] H. Goldman, R. Sohal, and E. Fradkin, Landau-Ginzburg Theories of Non-Abelian Quantum Hall States from Non-Abelian Bosonization, Phys. Rev. B 100, 115111 (2019).
- [9] C. Wetterich, Gauge symmetry from decoupling, Nucl.

theories with ultracold atoms in optical lattices, Nature Phys. 15, 1168 (2019).

- [19] F. Görg, K. Sandholzer, J. Minguzzi, R. Desbuquois, M. Messer, and T. Esslinger, Realization of densitydependent Peierls phases to engineer quantized gauge fields coupled to ultracold matter, Nature Phys. 15, 1161 (2019).
- [20] A. Mil, et al, A scalable realization of local U(1) gauge invariance in cold atomic mixtures, Science 367, 1128 (2020).
- [21] E. Zohar and E. Reznik, Confinement and Lattice Quantum-Electrodynamic Electric Flux Tubes Simulated with Ultracold Atoms, Phys. Rev. Lett. 107, 275301 (2011).
- [22] M. C. Bañuls and K. Cichy, Review on novel methods for lattice gauge theories, Rep. Prog. Phys. 83, 024401 (2020).
- [23] B. I. Halperin, T. C. Lubensky, and S. K. Ma, First-Order Phase Transitions in Superconductors and Smectic-A Liquid Crystals, Phys. Rev. Lett. **32**, 292 (1974).
- [24] M. Moshe and J. Zinn-Justin, Quantum field theory in the large N limit: A review, Phys. Rep. 385, 69 (2003).
- [25] C. Bonati, A. Pelissetto, and E. Vicari, Lattice Abelian-Higgs model with noncompact gauge fields, Phys. Rev. B 103, 085104 (2021).
- [26] B. Ihrig, N. Zerf, P. Marquard, I. F. Herbut, and M. M. Scherer, Abelian Higgs model at four loops, fixed-point collision and deconfined criticality, Phys. Rev. B 100, 134507 (2019).
- [27] K. G. Wilson and J. Kogut, The renormalization group and the ϵ expansion, Phys. Rep. **12**, 77 (1974).
- [28] M. E. Fisher, The renormalization group in the theory of critical behavior, Rev. Mod. Phys. 47, 543 (1975).
- [29] K. G. Wilson, The renormalization group and critical phenomena, Rev. Mod. Phys 55, 583 (1983).
- [30] A. Pelissetto and E. Vicari, Critical Phenomena and Renormalization Group Theory, Phys. Rep. 368, 549 (2002).
- [31] A. Pelissetto and E. Vicari, Multicomponent compact Abelian-Higgs lattice models, Phys. Rev. E 100, 042134 (2019).
- [32] If one considers the model with compact fields, a critical behavior associated with the charged AH FP is only observed with fields of higher charge $q \ge 2$ [54].
- [33] A. S. Kronfeld and U. J. Wiese, SU(N) gauge theories with C periodic boundary conditions. 1. Topological structure, Nucl. Phys. B 357, 521 (1991).
- [34] B. Lucini, A. Patella, A. Ramos and N. Tantalo, Charged hadrons in local finite-volume QED+QCD with C* boundary conditions, JHEP 02, 076 (2016).
- [35] We consider gauge fixings defined by the condition $G_x[A] = 0$ for all sites x. We assume that $G_x[A]$ is a local *linear* combination of the fields $A_{x,\mu}$ and that the gauge fixing is maximal (no gauge freedom is left after the introduction of the gauge fixing). Using the usual Fadeev-Popov procedure, we can replace the gauge-fixing with a term $\exp(-a\sum_x G_x[A]^2)$, without changing the expectation values of gauge-invariant operators.
- [36] C. Bonati, A. Pelissetto, and E. Vicari, Phase Diagram, Symmetry Breaking, and Critical Behavior of Three-Dimensional Lattice Multiflavor Scalar Chromodynamics, Phys. Rev. Lett. **123**, 232002 (2019); Threedimensional lattice multiflavor scalar chromodynamics: Interplay between global and gauge symmetries, Phys.

Rev. D 101. 034505 (2020).

- [37] M. E. Fisher, The renormalization group in the theory of critical behavior Rev. Mod. Phys. 46, 597 (1974); (erratum) 47, 543 (1975); Scaling Axes and the Spin-Flop Bicritical Phase Boundaries, Phys. Rev. Lett. 34, 1634 (1975).
- [38] If $y_r < 1/\nu$, Eq. (9) still holds but one has to replace X with the appropriate linear scaling field (see Ref. [37]), $X = (J - ar - J_c)L^{1/\nu}$. The constant *a* is fixed by the requirement that $J_c(r) = J_c + ar + O(r^2)$. Here $J_c(r)$ is the transition value for the model at fixed *r* and $J_c = J_c(0)$. If $y_r < 1/\nu$, the behavior for $J = J_c$ is controlled by $X = -arL^{1/\nu}$ and thus, fits of *R* to functions of rL^{y_r} would give $y_r = 1/\nu$: no information on the relevance or irrelevance of the perturbation would be obtained.
- [39] M. Creutz, Quarks, Gluons and Lattices (Cambridge University press, 1985).
- [40] A. Pelissetto and É. Vicari, Three-dimensional ferromagnetic CP^{N-1} models, Phys. Rev. E **100**, 022122 (2019).
- [41] C. Bonati, A. Pelissetto and E.Vicari, Lattice gauge theories in the presence of a linear gauge-symmetry breaking, Phys. Rev. E 104, 014140 (2021).
- [42] A. B. Zamolodchikov, Irreversibility of the Flux of the Renormalization Group in a 2D Field Theory, JETP Lett. 43, 730 (1986).
- [43] J. Cardy, Scaling and renormalization in statistical physics, (Cambridge University Press, 1996).
- [44] S. S. Pufu, The F-theorem and F-maximization, J. Phys. A: Math. Theor. 50, 443008 (2017).
- [45] T. Grover, Entanglement Monotonicity and the Stability of Gauge Theories in Three Spacetime Dimensions, Phys. Rev. Lett. **112**, 151601 (2014).
- [46] H. Casini and M. Huerta, Renormalization group running of the entanglement entropy of a circle, Phys. Rev. D 85, 125016 (2012).
- [47] I. R. Klebanov, S. S. Pufu, S. Sachdev, and B. R. Safdi, Entanglement entropy of 3-d conformal gauge theories with many flavors, J. High Energy Phys. 05 (2012) 036.
- [48] I. R. Klebanov, S. S. Pufu, and B. R. Safdi, F-theorem without supersymmetry, J. High Energy Phys. 10 (2011) 038.
- [49] R. C. Myers and A. Sinha, Holographic c-theorems in arbitrary dimensions, J. High Energy Phys. 01 (2011) 125.
- [50] It is worth mentioning that some mechanisms that allow cyclic RG flows even in the presence of local monotonicity have also been proposed; see, e.g., T. L. Curtright, X. Jin, and C. K. Zachos, Renormalization Group Flows, Cycles, and c-Theorem Folklore, Phys. Rev. Lett. **108**, 131601 (2012).
- [51] J. B. Kogut, An Introduction to Lattice Gauge Theory and Spin Systems, Rev. Mod. Phys. 51, 659 (1979).
- [52] M. Van Damme, J. C. Halimeh, and P. Hauke, Gaugesymmetry violation quantum phase transition in lattice gauge theories, arXiv:2010.07338.
- [53] T. V. Zache, M. Van Damme, J. C. Halimeh, P. Hauke, and D. Banerjee, Achieving the continuum limit of quantum link lattice gauge theories on quantum devices, arXiv:2104.00025.
- [54] C. Bonati, A. Pelissetto, and E. Vicari, Higher-charge three-dimensional compact lattice Abelian-Higgs models, Phys. Rev. E 102, 062151 (2020).