



Study of J/ψ meson production inside jets in pp collisions at $\sqrt{s} = 8$ TeV

The CMS Collaboration*

CERN, Switzerland



ARTICLE INFO

Article history:

Received 3 October 2019
 Received in revised form 31 March 2020
 Accepted 1 April 2020
 Available online 7 April 2020
 Editor: M. Doser

Keywords:

CMS
 J/ψ Mesons
 Jets

ABSTRACT

A study of the production of prompt J/ψ mesons contained in jets in proton-proton collisions at $\sqrt{s} = 8$ TeV is presented. The analysis is based on data corresponding to an integrated luminosity of 19.1 fb^{-1} collected with the CMS detector at the LHC. For events with at least one observed jet, the angular separation between the J/ψ meson and the jet is used to test whether the J/ψ meson is part of the jet. The analysis shows that most prompt J/ψ mesons having energy above 15 GeV and rapidity $|y| < 1$ are contained in jets with pseudorapidity $|\eta_{\text{jet}}| < 1$. The differential distributions of the probability to have a J/ψ meson contained in a jet as a function of jet energy for a fixed J/ψ energy fraction are compared to a theoretical model using the fragmenting jet function approach. The data agree best with fragmenting jet function calculations that use a long-distance matrix element parameter set in which prompt J/ψ mesons are predicted to be unpolarized. This technique demonstrates a new way to test predictions for prompt J/ψ production using nonrelativistic quantum chromodynamics.

© 2020 Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

1. Introduction

The mechanism for producing J/ψ mesons as bound states of charm quark pairs ($c\bar{c}$) in hadronic collisions has been under intensive experimental and theoretical study since the 1974 discovery of the J/ψ meson in proton-nucleon collisions [1] and in e^+e^- annihilations [2]. The early theoretical descriptions of the hadronic production process considering only color-singlet production [3,4] were at odds with the differential cross section measurements as a function of the J/ψ transverse momentum $p_T^{J/\psi}$ made by experimenters at the Fermilab Tevatron [5] for $p_T^{J/\psi} > 6$ GeV. A new theoretical approach, nonrelativistic quantum chromodynamics (NRQCD), was used to address the problem [6–8]. The NRQCD model includes both color-singlet and color-octet amplitudes for the $c\bar{c}$ system that ultimately produces the J/ψ meson. It proved to be capable of explaining the cross section data, using phenomenological parameters called long-distance matrix elements (LDMEs) that are adjusted to describe J/ψ meson production data. Within the NRQCD factorization assumption, the LDME parameters are process independent. However, each determination of an LDME set can choose a specific collection of J/ψ meson production data and J/ψ meson kinematic requirements. Furthermore, different LDME sets that describe the production data may have different predictions for the J/ψ meson polarization [9]. Experiments [10,11]

have shown that the prompt J/ψ meson polarization at large $p_T^{J/\psi}$ (> 12 GeV) is small. Recent NRQCD studies extend the range of experimental input to include low- p_T data and attempt to make global fits to the full set of charmonium information. A review of these studies can be found in Ref. [12].

A remaining theoretical problem is to determine the mechanism by which a $c\bar{c}$ system in an angular momentum state and quark color configuration $^{2S+1}L_J^n$ hadronizes into a J/ψ meson. Here, S , L , and J are the spin, orbital, and total angular momentum quantum numbers of the $c\bar{c}$ system. Its color state is labeled by n , with $n = 1$ or 8 referring to a color-singlet or color-octet configuration, respectively. The J/ψ meson has $J = S = 1$ and $n = 1$. The analysis described in this Letter combines the experimental measurement of J/ψ mesons contained in jets with a theoretical approach based on the fragmenting jet function (FJF) model [13]. The FJF model postulates that the $c\bar{c}$ pair is not produced directly in the hard scattering, but is a fragmentation product of a high- p_T jet. The model uses the methodology of NRQCD to compute the cross section contributions for all relevant $^{2S+1}L_J^n$ terms. Each cross section term has a characteristic relation between the jet energy E_{jet} and its fraction carried by the J/ψ meson: $z = E_{J/\psi}/E_{\text{jet}}$.

A study of J/ψ mesons contained in jets in the rapidity region $y_{J/\psi} > 2$, dominated by charm fragmentation for large z , has been reported by the LHCb Collaboration [14]. The LHCb analysis, which measured the z distribution integrated over jet energy, does not have the sensitivity to LDME parameter sets that characterizes this analysis.

* E-mail address: cms-publication-committee-chair@cern.ch.

The data for this analysis were collected by the CMS detector in proton-proton (pp) collisions from the CERN LHC, corresponding to an integrated luminosity of 19.1 fb^{-1} at $\sqrt{s} = 8 \text{ TeV}$. It is the first experimental study of prompt J/ψ mesons contained in jets produced in the gluon-dominated central rapidity region, where the FJF theory for gluonic jet fragmentation applies.

2. Theoretical framework

The hadronization process is nonperturbative. It is handled in the FJF approach by an NRQCD expansion of the fragmentation function for a jet initially produced in a hard scattering at high energy. The observables are E_{jet} and z . Following Ref. [13], the differential cross section for dijet production, with one jet fragmenting to a J/ψ meson, can be written symbolically as

$$\frac{d^2\sigma(E_{\text{jet}}; z)}{dE_{\text{jet}} dz} = \sum_{A, B, i, j} f_{A/p} f_{B/p} d\sigma_{ABij}(c\bar{c}X, n, \mathcal{J}_j) \otimes \mathcal{F}_S \otimes \mathcal{G}_i^{J/\psi}(E_{\text{jet}}, z|R, \mu). \quad (1)$$

In this expression, A and B are the partons in the colliding protons with fractional flavor content $f_{A/p}, f_{B/p}$, respectively, while i and j are the outgoing partons. The symbolic hard-scattering cross section $d\sigma_{ABij}(c\bar{c}X, n, \mathcal{J}_j)$ produces the fragmenting jet from outgoing parton i and the recoil jet \mathcal{J}_j from outgoing parton j . The fragmenting jet produces a $c\bar{c}$ system characterized by S, L, J , and n quantum numbers, plus an inclusive hadronic state X that represents the remainder of the jet. The function \mathcal{F}_S controls the evolution of the fragmenting system down to the energy scale μ equal to the mass of the $c\bar{c}$ system, to allow the development of jet structure from soft gluons. The nonperturbative fragmentation of the $c\bar{c}$ system into the observed J/ψ meson is described by the function $\mathcal{G}_i^{J/\psi}(E_{\text{jet}}, z|R, \mu)$, where E_{jet} is determined in a cone of angular radius R .

The type of parton i that produces the fragmenting jet, and ultimately the J/ψ meson, depends on the jet rapidity region. In the central rapidity region covered by this analysis, gluon fragmentation dominates [15]. The FJF expression for $\mathcal{G}_i^{J/\psi}$ sums over all contributing partons, but the light flavor contributions are suppressed and can be neglected. In Ref. [13], the small central charm quark fragmentation contribution was absorbed into the ${}^3S_1^1$ contribution to gluon fragmentation, so $\mathcal{G}^{J/\psi}$ in this Letter represents only gluon fragmentation.

In Ref. [16], the authors updated the work of Ref. [13] to make an explicit computation of the perturbative dijet double-differential cross section, followed by the fragmentation of one of the jets to a J/ψ meson. They integrated over the kinematic variables of the second jet to give an FJF expression for the absolute differential cross section to produce a jet of energy E_{jet} that fragments into a J/ψ meson carrying energy fraction z of the parent jet energy along with the remaining fragments. In the NRQCD decomposition of $\mathcal{G}^{J/\psi}$ for central J/ψ meson hadroproduction with $p_T > 10 \text{ GeV}$, four FJF terms are relevant: ${}^3S_1^1, {}^1S_0^8, {}^3S_1^8,$ and ${}^3P_J^8$. Only the ${}^1S_0^8$ term has all angular momenta equal to zero in the $c\bar{c}$ rest frame. If this NRQCD term were to dominate the jet fragmentation process, then the J/ψ meson would be produced unpolarized.

3. The CMS detector

The central feature of the CMS apparatus is a superconducting solenoid of 6 m internal diameter, providing a magnetic field of 3.8 T. Within the solenoid volume are a silicon pixel and strip tracker, a lead tungstate crystal electromagnetic calorimeter, and a

brass and scintillator hadron calorimeter, each composed of a barrel and two endcap sections. When combining information from the entire detector, the jet energy resolution amounts typically to 15% at 10 GeV and 8% at 100 GeV [17]. Muons are detected in gas-ionization chambers embedded in the steel flux-return yoke outside the solenoid, covering the pseudorapidity range $|\eta| < 2.4$. The silicon tracker measures charged particles within the pseudorapidity range $|\eta| < 2.5$. It consists of 1440 silicon pixel and 15 148 silicon strip detector modules. For nonisolated particles of $1 < p_T < 10 \text{ GeV}$ and $|\eta| < 1.4$, the track resolutions are typically 1.5% in p_T and 25–90 (45–150) μm in the transverse (longitudinal) impact parameter [18]. Matching muons to tracks measured in the silicon tracker results in a relative transverse momentum resolution, for muons with $20 < p_T < 100 \text{ GeV}$, of 1.3–2.0% in the barrel [19]. Events of interest are selected using a two-tiered trigger system [20]. The first level, composed of custom hardware processors, uses information from the calorimeters and muon detectors to select events at a rate of around 100 kHz within a fixed time interval of less than 4 μs . The second level, known as the high-level trigger (HLT), consists of a farm of processors running a version of the full event reconstruction software optimized for fast processing. This reduces the event rate to around 1 kHz before data storage. A more detailed description of the CMS detector, together with a definition of the coordinate system used and the relevant kinematic variables, can be found in Ref. [21].

4. Event selection and background subtraction

The experimental methods follow those used by previous CMS analyses of inclusive J/ψ and $\Upsilon(nS)$ production at $\sqrt{s} = 7 \text{ TeV}$ [22–26]. The event selection is based on a dimuon trigger involving the silicon tracker and muon systems. The trigger requires two oppositely charged muons with rapidity of the dimuon system $|y| < 1.25$ and its invariant mass range $2.7 < m_{\mu\mu} < 3.5 \text{ GeV}$. The three-dimensional fit to the dimuon vertex must have a χ^2 probability (the p -value of the χ^2 returned by the fit) $> 0.5\%$. Only dimuon pairs in which the muons bend away from each other in the magnetic field are used to allow a precise dimuon efficiency determination. The dimuon p_T trigger threshold varied from 5 to 9 GeV during the data-taking period. The primary event vertex is defined as the one with the largest summed p_T of its associated tracks.

The offline selection requires a dimuon pair with $p_T > 10 \text{ GeV}$, $|y| < 1$, energy $E > 15 \text{ GeV}$, and vertex fit χ^2 probability $> 1\%$. In order to guarantee agreement to within 3% between the single-muon efficiencies from control samples and from simulation, each muon must have $p_T^\mu > 6 \text{ GeV}$ and $|\eta_\mu| < 2.1$, or $p_T^\mu > 5 \text{ GeV}$ and $|\eta_\mu| < 0.8$. The muon candidate must satisfy the CMS “tight” muon quality requirements on the number of tracker hits, the muon track fit quality, and the distance along the beam line from the primary event vertex [19]. No muon isolation requirements are applied, because we look for $J/\psi + \text{jet}$ associations. The J/ψ signal invariant mass range is $2.95 < m_{\mu\mu} < 3.20 \text{ GeV}$. After the data selection, we observe at most one J/ψ candidate per event.

The trigger does not use any information about jets in the event. Jets are reconstructed from particle-flow objects [27], using an anti- k_T algorithm with a distance parameter of 0.5 [28], as implemented in the FASTJET package [29]. The jet response has been corrected to the particle level [17]. Although the J/ψ candidate is not a particle-flow object, its decay muons are. This does not exclude jets that consist only of a J/ψ meson. However, such jets constitute less than 10^{-4} of this sample and do not affect the results presented here. The jet properties include the energy E_{jet} , the transverse momentum magnitude p_T^{jet} , the number of constituents, and the number of included muons. Each bunch crossing in the data produces, on average, 14 reconstructed pp vertices,

corresponding to 21 interactions per bunch crossing. The extra interactions produce so-called pileup distortions, which are corrected using the procedure described in Ref. [17]. For this analysis, the jet selection requirements are $p_T^{\text{jet}} > 25$ GeV and $|\eta_{\text{jet}}| < 1$.

The J/ψ event candidates are classified as prompt, nonprompt, or combinatorial. Nonprompt events include those J/ψ mesons that come from decays of b hadrons. Combinatorial candidates are accidental pairings of an identified μ^+ and a μ^- such that the dimuon invariant mass falls within the signal mass interval. The nonprompt background is strongly reduced by applying a selection on the variable Σ_{TD} , which is the sum of the squares of the significance (in units of standard deviations) of the transverse distance of closest approach of each muon track to the primary vertex. The Σ_{TD} distribution has a sharp peak near zero from prompt events and a long tail at larger Σ_{TD} from nonprompt sources, which we fit with an exponential function. From a prompt J/ψ meson Monte Carlo (MC) sample, we find that $>99\%$ of the events have $\Sigma_{TD} < 10$. The simulated Σ_{TD} shape agrees with that in data for this region, so we require $\Sigma_{TD} < 10$ to define the prompt dimuon events. In the J/ψ data, the exponential function that describes the nonprompt background is extrapolated into the range $\Sigma_{TD} < 10$ to estimate the fraction of nonprompt events in the prompt signal mass range. This is $(5.7 \pm 0.1)\%$. The events in the prompt signal mass range also contain combinatorial background, which is determined by interpolating the $m_{\mu\mu}$ low (2.70–2.90 GeV) and high (3.25–3.50 GeV) sideband regions. We find that the combinatorial background fraction in the prompt signal mass range is $(1.4 \pm 0.2)\%$. The quoted uncertainties in the backgrounds are statistical only. All distributions shown in this Letter have the nonprompt and combinatorial backgrounds subtracted. After background subtraction, there are 1.63×10^6 prompt J/ψ meson candidates.

5. Association of jets and J/ψ mesons

The analysis makes no restriction on the number of jets that pass the jet selection requirements, which we term “observed jets”. For jet requirements $p_T^{\text{jet}} > 25$ GeV and $|\eta_{\text{jet}}| < 1$, the fractions of J/ψ meson events that have 0, 1, 2, or 3 observed jets are $(55.12 \pm 0.06)\%$, $(34.03 \pm 0.05)\%$, $(9.58 \pm 0.02)\%$, and $(1.27 \pm 0.08)\%$, respectively, where the uncertainties are statistical only. For events with at least one observed jet, the association of a J/ψ meson with a jet is made using the angular separation $\Delta R = \sqrt{(\eta_{\text{jet}} - \eta_{\mu\mu})^2 + (\phi_{\text{jet}} - \phi_{\mu\mu})^2}$. Here, η_{jet} ($\eta_{\mu\mu}$) and ϕ_{jet} ($\phi_{\mu\mu}$) are the pseudorapidity and azimuthal angle (modulo π), respectively, of the jet (dimuon) direction. The ΔR distribution for the best-matched jet is sharply peaked at zero, as seen for events with one observed jet in Fig. 1 (upper). The J/ψ meson and the jet are defined as associated if $\Delta R < 0.5$. Furthermore, if both decay muons from the J/ψ meson are among the objects that comprise the jet, we say that the J/ψ meson is a constituent of the jet.

When there are two observed jets in the event, further evidence that J/ψ meson production comes primarily from jets is shown in Fig. 1 (lower). This plot shows ΔR for the J/ψ meson with respect to each observed jet in two-jet events. The higher-energy (leading) jet has ΔR_l , the lower-energy (subleading) one ΔR_{sl} . Note that the energy labels here play no role in the analysis; the jets need only to pass the jet p_T and $|\eta_{\text{jet}}|$ requirements given above. The J/ψ meson is not required to come from either jet. The clusters of events in Fig. 1 (lower), near $(\Delta R_l, \Delta R_{sl}) = (0, \pi)$ and $(\pi, 0)$, show that $(94.1 \pm 0.1)\%$ of the time, the J/ψ meson is a constituent of one of the two jets in the event. In events with a J/ψ meson and two jets, the mean and RMS deviation of the distribution of the number of jet constituents, charged and neutral, for the fragmenting jet (25 ± 8) and the recoil jet (29 ± 8) are similar. The shapes of the jet energy spectra for the jet containing the J/ψ meson and the recoil




Fig. 1. The distributions of (upper) $\Delta R(\text{jet}, J/\psi)$ for one-jet events and (lower) $\Delta R_l(\text{leading jet}, J/\psi)$ vs. $\Delta R_{sl}(\text{subleading jet}, J/\psi)$ for two-jet events.

jet are indistinguishable. The difference in the probability for the J/ψ meson to be a jet constituent in the one- and two-jet cases, along with a discussion of the small excess for $2.4 < \Delta R < 3.5$ in Fig. 1 (upper), will be addressed in Section 12.

6. Experimental application of the FJF approach

The authors of Refs. [13,16] emphasize that experimental sensitivity to the FJF terms in jet fragmentation comes from measuring the jet energy dependence of the function \mathcal{G} in Eq. (1) at fixed z . In the FJF framework, the dependence of the fragmenting jet differential cross section on the J/ψ properties comes solely through the z variable. Integrating Eq. (1) over z gives the single-jet differential cross section for the production of J/ψ mesons contained in jets, as a function of E_{jet} . This is used as a normalization term in Ref. [16], where the differential cross section for a jet to fragment to a J/ψ meson with the energy fraction z is calculated for jets having $p_T^{\text{jet}} > 25$ GeV and pseudorapidity $|\eta_{\text{jet}}| < 1.2$. The resulting J/ψ meson is required to have energy above 15 GeV and rapidity $|y_{J/\psi}| < 1$. The jet fragmentation cross section is normalized by integrating over the z range 0.3–0.8. The authors showed that the jet energy dependence of the normalized FJF terms is insensitive to the exact z range used. At a fixed z value, called z_1 , the ratio of the fragmenting jet differential cross section due to a single FJF term i to the sum of the cross section integrals for $0.3 < z < 0.8$ for all FJF terms is termed $(d\tilde{\sigma}_i/dE_{\text{jet}} dz)|_{z_1}$ in Ref. [16]. The sum of this ratio over all four FJF terms is denoted as $(d\tilde{\sigma}/dE_{\text{jet}} dz)|_{z_1}$. For a given LDME parameter set, each of the four FJF terms is different. Also, changing the LDME parameter set changes the FJF predictions for the four terms.

The authors of Ref. [16] cite next-to-leading order (NLO) calculations [30–33] to argue that the $p_T^{J/\psi}$ range for the three z_1 values used in this analysis is large enough that the 3S_1

NRQCD term cannot contribute to the sum. Therefore, in computing $(d\tilde{\sigma}/dE_{\text{jet}} dz)|_{z_1}$ to compare to these data, only the three color-octet terms are included. However, in the low- z region included in the normalizing integral, the ${}^3S_1^1$ NRQCD term can play a role and is included in their calculation for $0.3 < z < 0.8$.

The experimental proxy for $(d\tilde{\sigma}/dE_{\text{jet}} dz)|_{z_1}$, evaluated for a jet energy bin centered at E_c , is called $\Xi(E_c; z_1)$:

$$\Xi(E_c; z_1) \equiv \frac{N(E_c; z_1)}{\int_{0.3}^{0.8} N(E_c; z) dz}, \quad (2)$$

where $N(E_c; z_1)$ is the number of events having a J/ψ meson contained in a jet for a z interval Δ_z centered on z_1 in that E_{jet} bin. The number of events is weighted to correct for the J/ψ meson efficiency and acceptance, as described in Section 7, as well as corrected for jet efficiency and jet energy resolution, as described in Section 8. We use a z interval $\Delta_z = \pm 0.025$ around z_1 , which is small enough to be insensitive to z variations in Ξ and large enough to provide a reasonable number of events in each E_{jet} bin.

7. Efficiency corrections for J/ψ mesons

Measuring the properties of events when a J/ψ meson is a jet constituent requires an event-by-event J/ψ meson efficiency correction. Each entry in the signal or background event distributions has an event weight, defined as $1/\epsilon_{J/\psi}$. The dimuon acceptance times efficiency $\epsilon_{J/\psi}$ is determined using a simulated sample of unpolarized J/ψ meson events, uniformly distributed in 1 GeV wide p_T bins and uniformly distributed over $|y_{J/\psi}| < 1.5$. Only the J/ψ meson is simulated; studies [25,26] show that using a complete PYTHIA [34] event simulation does not change the efficiency results. The $J/\psi \rightarrow \mu^+\mu^-$ decay is simulated using EVTGEN [35]; radiative effects are treated by PHOTOS [36]; and the detector response to the two muons is simulated using the GEANT4-based [37] CMS simulation program. The simulated J/ψ meson must pass the quality requirements listed in Section 4. The total efficiency $\epsilon_{J/\psi}$ varies with the rapidity and transverse momentum of the J/ψ meson because the muon reconstruction, dimuon vertex reconstruction, and dimuon trigger efficiencies depend on these variables. There is also an HLT trigger inefficiency if two muons in the event have a small angular separation. This is also taken from simulation and checked against data taken using a single-muon trigger.

8. Jet energy corrections and unfolding

A crucial part of the analysis is measuring the energy of the jet that contains the J/ψ meson. To test whether there might be an influence on the jet energy distribution due to the presence of the J/ψ meson, we study the two-jet events shown in Fig. 1 (lower). The energy distributions of the fragmenting jet and the recoil jet are compared for $0.3 < z < 0.8$ and for z ranges of 0.40–0.45, 0.50–0.55, and 0.60–0.65. The shapes of the measured energy distributions of the recoil and fragmenting jets for each sample are indistinguishable. There is no evidence that having a J/ψ meson as a constituent affects the jet energy distribution.

The jet energy distributions are compared to the FJF model predictions in bins of jet energy. Experimentally, the jet energy bin width ΔE_{jet} is constrained by the finite jet energy resolution of the CMS apparatus, which must be unfolded. We use $\Delta E_{\text{jet}} = 8$ GeV. The D'Agostini unfolding method [38] from the ROOUNFOLD package [39] is used to extract the unsmearred Ξ distribution. The procedure uses an input generator-level jet energy distribution (truth distribution) derived from PYTHIA light-quark or gluon jets. Simulation shows that for measured jet energy $E_{\text{jet}} > 44$ GeV, the jet reconstruction efficiency exceeds 98.5% and is consistent with being energy independent. Thus, 44 GeV is the lowest jet energy

considered in the unfolding procedure. The unfolding procedure uses the CMS jet energy resolution and jet finding efficiency [27] to determine the unfolded jet energy matrix and the MISS matrix. The latter is filled for events that fail the jet efficiency test or fall outside the unfolded jet energy window 44–140 GeV. The method was validated using several different simulated jet energy input truth distributions, including a fit to the PYTHIA shape using the sum of exponentials and the raw data itself in a bootstrap approach. There was no change in the unfolded distributions that exceeded $\sigma_{\text{stat}}/4$ for any jet energy bin. Based on unfolding studies in simulation that used three to six iterations, we found that four unfolding iterations gave stable matches to the simulation events and showed no sensitivity to different choices for the input truth distribution. Based on the simulation results, the unfolded jet energy range is $56 < E_{\text{jet}} < 120$ GeV. This range is stable when the input distribution is changed. Henceforth, E_{jet} will refer to the unfolded quantity, unless otherwise noted.

The unfolded jet energy distributions for the $\Xi(E; z)$ functions have bin-to-bin correlations that affect the statistical uncertainty in Ξ for each jet energy bin. The statistical uncertainties are evaluated by repeating the unfolding procedure 250 times, forming the covariance matrix, and determining the uncertainty for each jet energy bin. The statistical uncertainties computed by this procedure are 0.02 to 0.06%. The unfolding in z is dominated by the E_{jet} resolution. The changes in z from the unfolding procedure for the region of interest (0.40–0.65) are less than 0.01 in z . Therefore, the measured z values are used in the $\Xi(E_{\text{jet}}; z)$ determinations.

9. Systematic uncertainties

The systematic uncertainties arise from the determination of the event weight, based on the J/ψ meson and muon properties, and from a bias in the J/ψ -jet association, discussed below. The systematic uncertainty in the jet energy scale is small compared to the jet energy resolution used in the unfolding. Varying the jet energy by the jet energy scale systematic ($< 2.2\%$) uncertainty before the unfolding made no change in the Ξ results.

The CMS studies at $\sqrt{s} = 8$ TeV using a tag-and-probe method [22,23] show that, for the offline requirements used in this analysis, the ratio of the single-muon efficiency in data and MC simulation is consistent with unity within $< 3\%$, independent of p_T^μ [40]. The tracking efficiency in data and simulation agree to within 1% per track. The dimuon vertex and trigger simulation also have 1% systematic uncertainties. The dimuon HLT trigger inefficiency varies with $p_T^{J/\psi}$ in the range 4.5–7.5%. For the few dimuons with $p_T > 60$ GeV, it can go up to 15%. The difference between unity (no loss) and the simulated HLT trigger efficiency is assigned as the HLT systematic uncertainty for each event. All of the above systematic uncertainties are added in quadrature to determine the total systematic uncertainty in the weight for each event. To estimate the impact of the weight systematic uncertainty on the $\Xi(E_{\text{jet}}; z_1)$ function, two additional $\Xi(E_{\text{jet}}; z_1)$ functions are made for each z_1 . One uses distributions in which the weight for each event is raised by one standard deviation; in the other, the event weight is lowered by one standard deviation. The shifted $\Xi(E_{\text{jet}}; z_1)$ values are compared to the unshifted value in each energy bin. The relative systematic uncertainty in the event weight ranges from 0.2 to 0.9% of the standard-weight $\Xi(E_{\text{jet}}; z_1)$ values.

In addition, there is a selection bias in the J/ψ meson and jet association that disfavors the configuration when the difference $\eta_{\text{jet}} - \eta_{J/\psi}$ has the opposite sign to η_{jet} . The bias originates from events that are lost in the selection on $|\eta_{\text{jet}} - \eta_{J/\psi}|$ and is evaluated from data. The number of events per E_{jet} bin in the biased region is rescaled to match the yield in the unbiased region. Half of the difference between the measured and corrected number of events in each E_{jet} bin is assigned as its bias systematic uncertainty. The




Fig. 2. Comparison of $\Xi(E_{\text{jet}}; z_1)$ versus E_{jet} from data with FJF predictions of the total differential cross section, each normalized to unit area, for the BCKL (squares), BK (inverted triangles), and Chao (diamonds) LDME parameter sets. The three z_1 ranges are (upper left) $z_1 = 0.425$; (upper right) $z_1 = 0.525$; (lower) $z_1 = 0.625$. The curves show the detailed energy dependence of the predictions. The vertical bars on the data points are the quadrature sum of the statistical and systematic uncertainties.

weight and bias systematic uncertainties are added in quadrature to obtain the systematic uncertainty in $\Xi(E_{\text{jet}}; z_1)$, which ranges from 0.3 to 1.0%. These uncertainties are then added in quadrature with the uncertainty in the unfolding procedure discussed in the previous section.

10. The FJF predictions of the jet energy spectrum

In this analysis, we use three z_1 values: 0.425, 0.525, and 0.625. These are the centers of three nonoverlapping z subregions with $\Delta z = 0.05$ from the measurement region $0.3 < z < 0.8$. In these three z regions, the FJF terms have different jet energy distributions for a given LDME parameter set. The authors of Ref. [16] supplied tables of the normalized differential cross section terms $(d\tilde{\sigma}_i/dE_{\text{jet}} dz)|_{z_1}$, computed for $\sqrt{s} = 8$ TeV and jet radius $R = 0.5$. The cone algorithm used for the theoretical calculation does not introduce a systematic effect since there are no background or pileup sources in the theory. As described in Section 6, we compare the data to sum of the $^1S_0^8$, $^3S_1^8$, and $^3P_J^8$ FJF functions for the LDME parameter sets from Bodwin, Chung, Kim, and Lee (BCKL) [30], Butenschoen and Kniehl (BK) [41], and Chao, et al. (Chao) [42]. The LDME parameter sets for these three studies are derived from different selections of J/ψ meson production measurements, e.g., the BK set includes electroproduction data and uses a lower J/ψ meson p_T limit than is used in the hadroproduction-only selection of the BCKL and Chao sets. All groups report that their LDME sets yield J/ψ meson differential cross sections that agree with the J/ψ meson production data on which the extractions were based.

11. Comparison of data with FJF total cross section predictions

In this section we compare the data with the prediction for the FJF total differential cross section from each of the three LDME sets. Fig. 2 shows the normalized jet energy distributions for the data and the FJF total cross section predictions for each LDME set

Table 1

The χ^2 value and the associated p -value (in parentheses) for 7 degrees of freedom from the comparison of the data and the predictions for the total FJF cross section shape at $z_1 = 0.425$, 0.525, and 0.625, using the BCKL, the BK, and the Chao LDME parameter sets.

	0.425	0.525	0.625
BCKL	22.2 (0.23%)	11.0 (14%)	10.7 (15%)
BK	59.6 (<0.001%)	60.1 (<0.001%)	64.0 (<0.001%)
Chao	267 (<0.001%)	96 (<0.001%)	164 (<0.001%)

at each of the three z_1 values used in the analysis. The uncertainties in the data include the statistical and systematic components added in quadrature. For each z_1 , the bin-averaged FJF values are used to calculate the χ^2 for the comparison of the FJF total differential cross section prediction to the data. The LDME calculations from Refs. [30,41,42] have normalization uncertainties, as shown in Ref. [16]. The FJF calculations give the ratio of the cross section in a small- z region to the cross section integral over a wide z range. The uncertainty in the predicted FJF values due to the theory normalization uncertainty is almost completely correlated for the numerator and denominator of the ratio. The resulting theoretical uncertainty is negligible compared to the experimental uncertainty in the $\Xi(E_{\text{jet}}; z_1)$ values. We therefore ignore it in computing the χ^2 values to match data and theory. The χ^2 value and the associated p -value for comparison of data to each LDME set are given in Table 1. An a priori decision was made that a model prediction is an acceptable match to the data only if the χ^2 p -value is larger than 0.1% for seven degrees of freedom. Otherwise, we say that the model does not match the data. For all three z_1 ranges, the FJF predictions using the BCKL LDME parameters match data. Neither the BK nor the Chao LDME parameter sets describe these jet + constituent J/ψ data for any z_1 value.

The observation that these new data on J/ψ meson production as constituents of jets match the FJF predictions for the BCKL LDME parameter set and reject two others validates the FJF ap-

proach to treating jets as a major source of J/ψ production in the gluon-rich central region in pp interactions. It also demonstrates that the BCKL LDME parameter set can describe new features of J/ψ hadronic production at large $p_T^{J/\psi}$. The BCKL LDME parameters were developed from a completely different data set than these $J/\psi + \text{jet}$ data, so there is no a priori reason to expect them to have predicted these measurements. The BCKL parameters are known to predict small J/ψ polarization [30], in agreement with experiment [10,11] for the range of $p_T^{J/\psi}$ values selected in this analysis (10–40 GeV). Because this analysis studies only high- p_T J/ψ meson production and shows that the BCKL LDME parameters describe the process and other sets do not, it suggests a tension between high- p_T J/ψ results and global charmonium studies [12].

12. Total fraction of J/ψ mesons from jets

In this section, we determine whether jets are the major source of prompt energetic J/ψ mesons ($E_{J/\psi} > 15$ GeV) in the central region ($|\eta_{\text{jet}}| < 1$). Here, E_{jet} refers to the measured jet energy before unfolding. As shown in Fig. 1 (upper), for events with a J/ψ meson and only one observed jet, $(84.0 \pm 0.1)\%$ of the J/ψ candidates are within $\Delta R < 0.5$ of that jet. This is consistent with jets being the dominant source of J/ψ production in this kinematic range when there is at least one observed jet in the event. However, events with one or more observed jets having $p_T^{\text{jet}} > 25$ GeV account for only $(44.9 \pm 0.1)\%$ of the prompt J/ψ meson sample.

To understand the source of J/ψ meson events with no jets passing the $p_T^{\text{jet}} > 25$ GeV requirement, termed zero-jet events, we note that a jet that has a constituent J/ψ meson can fail the p_T^{jet} threshold even though the J/ψ meson is observed. For instance, when the p_T^{jet} threshold is raised from 25 to 30 GeV, the fraction of zero-jet events with an identified J/ψ meson increases from 55 to 65%. For one-jet events in data with p_T^{jet} thresholds of 30, 35, and 40 GeV, the observed jet is found within $\Delta R < 0.5$ of the J/ψ meson in the event $(84.0 \pm 0.2)\%$ of the time, i.e., the probability of a jet having a constituent J/ψ meson is independent of p_T^{jet} . Only jets with $E_{\text{jet}} > 44$ GeV pass the $p_T^{\text{jet}} > 25$ GeV requirement with 100% efficiency over the range $0 < |\eta_{\text{jet}}| < 1$. Jets having $E_{\text{jet}} < 44$ GeV can contain observed J/ψ mesons with $E_{J/\psi} > 15$ GeV, but some of these jets will not pass the $p_T^{\text{jet}} > 25$ GeV requirement.

In order to correct for this effect, we fit the E_{jet} distribution for jets containing a J/ψ meson to the sum of two exponential functions in the range $44 < E_{\text{jet}} < 150$ GeV. We use the fit to extrapolate the number of jets containing a J/ψ meson to lower E_{jet} values, in order to estimate the number of jets with a constituent J/ψ meson that would be present in the lower-energy region for full p_T acceptance. Jet reconstruction efficiency corrections are not applied at this stage. The FJF model is valid for $z < 0.8$ [13]. In the data, only $(1.3 \pm 0.1)\%$ of jets having a constituent J/ψ meson have $z > 0.8$; we truncate the model at $z = 0.8$, setting a limit of $E_{\text{jet}} > 19$ GeV for the extrapolation. Some jets in the $E_{\text{jet}} = 25$ –44 GeV range have sufficiently large polar angles to pass the $p_T^{\text{jet}} > 25$ GeV requirement. These are subtracted from the extrapolation to avoid double counting. The number of jets from extrapolation in each 1 GeV wide jet energy bin i is corrected for the jet reconstruction efficiency ϵ_i to predict the total number N_i of jets with energy E_i .

In order to contribute to the data sample, a jet with energy E_i must produce a J/ψ meson with energy E_j . The probability P_j for the J/ψ meson to have energy E_j is taken from the results of this analysis, normalized to unity for 55 bins covering the range $15 < E_{J/\psi} < 70$ GeV. The total number A_i of jets with energy E_i

that contain a J/ψ meson with energy fraction $z_{ij} = E_j/E_i$ in the range 0.3–0.8 is

$$A_i = N_i \sum_{j=1}^{55} P_j w(z_{ij}). \quad (3)$$

The function $w(z_{ij})$ is the probability that a jet of energy E_i will contain a J/ψ meson having energy E_j . To proceed, we need a specific model for the jet and J/ψ kinematics. We use the jet fragmentation model in Ref. [13] for $E_{\text{jet}} = 50$ GeV. The probability is zero for $z > 0.8$. The model predicts that (43 ± 3) (stat)% of the J/ψ mesons should be accompanied by zero observed jets, compared to 55% found in the data.

There are systematic uncertainties in this result. In a private communication, the authors of Ref. [13] also provided a z probability calculation for $E_{\text{jet}} = 20$ GeV. The model prediction for the number of zero-jet events using the 20 GeV z probability calculation differs by 3% from the 50 GeV result. This difference is taken as the systematic uncertainty in the z fragmentation probability. The uncertainty in the MC prediction of the low-energy jet efficiency is 13%. We also made a closure test by using the model to predict the number of observed jets lost when the jet p_T threshold was raised from 25 to 40 GeV. The model prediction agrees with the actual number of lost jets to within $(3.5 \pm 0.1)\%$. However, there is a jet energy dependence in the matching between the data and the prediction. Extrapolating the bin-by-bin jet energy dependence of that difference into the 19–44 GeV range, the closure study gives a 7% systematic uncertainty in the predicted number of zero-jet events having jet energies less than 44 GeV. Adding the systematic uncertainties in quadrature, the predicted fraction of zero-jet events with a J/ψ meson as a constituent of a jet with $p_T^{\text{jet}} < 25$ GeV is (43 ± 3) (stat) ± 7 (syst)%.

If we apply this reasoning to results from Section 5, the small peak in the range $2.5 < \Delta R < 3.4$ in Fig. 1 (upper) is actually the recoil jet in a dijet pair for which the parent jet of the J/ψ meson was not observed. This increases the fraction of J/ψ mesons that are constituents of a jet in the one-jet sample from $(84.0 \pm 0.1)\%$ to $(94.3 \pm 0.1)\%$. With this interpretation, and the results from Section 5, we find that the one- and two-jet fractions for a jet to have a constituent J/ψ meson are both essentially 94%. The overall fraction of J/ψ mesons that come from jets is, then, $0.94 \cdot 45\% = 42\%$ from events with one or more observed jets, plus 43% from the zero-jet sample. While the zero-jet model is simple, it passes an experimental closure test. Also, it follows the trend of the data as the jet p_T requirement is raised in steps from 25 to 40 GeV. Using it, we conclude that (85 ± 3) (stat) ± 7 (syst)% of the J/ψ mesons within our kinematic acceptance, $E_{J/\psi} > 15$ GeV and $|y_{J/\psi}| < 1$, are constituents of jets with $E_{\text{jet}} > 19$ GeV and $|\eta_{\text{jet}}| < 1$.

13. Summary

The first analysis has been presented comparing data for prompt J/ψ mesons produced as constituents of central gluonic jets with a theoretical analysis based on the fragmenting jet function (FJF) approach. The term prompt means that the J/ψ meson is consistent with originating from the primary vertex. In the FJF model, the jet fragments into a $c\bar{c}$ system in an angular momentum state and quark color configuration $^{2S+1}L_J^n$, plus other hadrons. Here, S , L , and J are the spin, orbital, and total angular momentum quantum numbers of the $c\bar{c}$ system and n indicates a color-singlet ($n = 1$) or color-octet ($n = 8$) configuration. The FJF analysis uses the nonrelativistic quantum chromodynamics (NRQCD) approach to compute the cross section for the formation of a J/ψ meson from the $c\bar{c}$ system for four specific S , J , L , and n configurations: 1S_0 , 3S_1 , $^3S_1^1$, and $^3P_J^8$.

The data were collected by the CMS Collaboration in proton-proton collisions at $\sqrt{s} = 8$ TeV, corresponding to an integrated luminosity of 19.1 fb^{-1} . The kinematic selections for the analysis are $E_{J/\psi} > 15$ GeV, $|y_{J/\psi}| < 1$, $p_T^{\text{jet}} > 25$ GeV, and $|\eta_{\text{jet}}| < 1$. In z ranges 0.40–0.45, 0.50–0.55, and 0.60–0.65, where z is the J/ψ meson fraction of the jet energy, the shape of the measured differential cross section as a function of E_{jet} for J/ψ meson production as a jet constituent is compared to the FJF prediction for this quantity, using three different long-distance matrix element (LDME) parameter sets. The FJF predictions using the Bodwin, Chung, Kim, and Lee (BCKL) [30] LDME parameters match the data for all three z ranges. In contrast, the FJF predictions for the LDME parameter sets from Butenschoen and Kniehl (BK) [41] and Chao, et al., (Chao) [42] disagree with the data for all three z ranges. This establishes the ability of the FJF analysis to describe J/ψ meson production from central gluonic jets and the ability of this kind of experiment to distinguish among different sets of LDME parameters, all of which describe inclusive J/ψ meson production for their choice of data. The BCKL LDME set, constructed using inclusive hadronic production data with $p_T^{J/\psi} > 10$ GeV, not only describes the production of high- p_T J/ψ mesons as constituents of jets but also predicts small J/ψ meson polarization.

When a jet is observed in an event, the fraction of J/ψ mesons that are jet constituents is $(94.2 \pm 0.1)\%$, averaged over one- and two-jet events. Using a simple model to estimate the fraction of J/ψ mesons that are constituents of jets that fail the analysis p_T^{jet} requirement, we find that jets are the source of $(85 \pm 3 \text{ (stat)} \pm 7 \text{ (syst)})\%$ of the J/ψ mesons produced in the kinematic region probed in this study. Interpreting the results in the framework of the FJF model, jet fragmentation accounts for almost all prompt J/ψ mesons produced at large $p_T^{J/\psi}$. The data are consistent with an NRQCD treatment of the FJF process using the BCKL parameter set.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

We thank Ira Rothstein, Matthew Baumgart, and Prashant Shrivastava for their help in creating the tables of the predictions from their fragmenting jet function model to compare with our data. We thank Nora Brambilla for editorial input and theoretical guidance.

We congratulate our colleagues in the CERN accelerator departments for the excellent performance of the LHC and thank the technical and administrative staffs at CERN and at other CMS institutes for their contributions to the success of the CMS effort. In addition, we gratefully acknowledge the computing centers and personnel of the Worldwide LHC Computing Grid for delivering so effectively the computing infrastructure essential to our analyses. Finally, we acknowledge the enduring support for the construction and operation of the LHC and the CMS detector provided by the following funding agencies: BMBWF and FWF (Austria); FNRS and FWO (Belgium); CNPq, CAPES, FAPERJ, FAPERGS, and FAPESP (Brazil); MES (Bulgaria); CERN; CAS, MOST, and NSFC (China); COLCIENCIAS (Colombia); MSES and CSF (Croatia); RPF (Cyprus); SENESCYT (Ecuador); MoER, ERC IUT, and ERDF (Estonia); Academy of Finland, MEC, and HIP (Finland); CEA and CNRS/IN2P3 (France); BMBF, DFG, and HGF (Germany); GSRT (Greece); NKFI (Hungary); DAE and DST (India); IPM (Iran); SFI (Ireland); INFN (Italy); MSIP and NRF (Republic of Korea); MES (Latvia); LAS (Lithuania); MOE and UM (Malaysia); BUAP, CINVESTAV, CONACYT, LNS, SEP,

and UASLP-FAI (Mexico); MOS (Montenegro); MBIE (New Zealand); PAEC (Pakistan); MSHE and NSC (Poland); FCT (Portugal); JINR (Dubna); MON, ROSATOM, RAS, RFBR, and NRC KI (Russia); MESTD (Serbia); SEIDI, CPAN, PCTI, and FEDER (Spain); MoSTR (Sri Lanka); Swiss Funding Agencies (Switzerland); MST (Taipei); ThEPCenter, IPST, STAR, and NSTDA (Thailand); TUBITAK and TAEK (Turkey); NASU and SFFR (Ukraine); STFC (United Kingdom); DOE and NSF (USA).

Individuals have received support from the Marie-Curie program and the European Research Council and Horizon 2020 Grant, contract No. 675440 (European Union); the Leventis Foundation; the Alfred P. Sloan Foundation; the Alexander von Humboldt Foundation; the Belgian Federal Science Policy Office; the Fonds pour la Formation à la Recherche dans l'Industrie et dans l'Agriculture (FRIA-Belgium); the Agentschap voor Innovatie door Wetenschap en Technologie (IWT-Belgium); the F.R.S. - FNRS and FWO (Belgium) under the "Excellence of Science – EOS" – be.h project n. 30820817; The Ministry of Education, Youth and Sports (MEYS) of the Czech Republic; the Lendület ("Momentum") Program and the János Bolyai Research Scholarship of the Hungarian Academy of Sciences, the New National Excellence Program ÚNKP, the NKFI research grants 123842, 123959, 124845, 124850, and 125105 (Hungary); the Council of Science and Industrial Research, India; the HOMING PLUS program of the Foundation for Polish Science, cofinanced from European Union, Regional Development Fund, the Mobility Plus program of the Ministry of Science and Higher Education, the National Science Center (Poland), contracts Harmonia 2014/14/M/ST2/00428, Opus 2014/13/B/ST2/02543, 2014/15/B/ST2/03998, and 2015/19/B/ST2/02861, Sonata-bis 2012/07/E/ST2/01406; the National Priorities Research Program by Qatar National Research Fund; the Programa Estatal de Fomento de la Investigación Científica y Técnica de Excelencia María de Maeztu, grant MDM-2015-0509 and the Programa Severo Ochoa del Principado de Asturias; the Thalís and Aristeia programs cofinanced by EU-ESF and the Greek NSRF; the Rachadapisek Sompot Fund for Postdoctoral Fellowship, Chulalongkorn University and the Chulalongkorn Academic into Its 2nd Century Project Advancement Project (Thailand); The Welch Foundation, contract C-1845; and the Weston Havens Foundation (USA).

References

- [1] J.J. Aubert, et al., E598, Experimental observation of a heavy particle J, Phys. Rev. Lett. 33 (1974) 1404, <https://doi.org/10.1103/PhysRevLett.33.1404>.
- [2] J.-E. Augustin, et al., Discovery of a narrow resonance in e^+e^- annihilation, Phys. Rev. Lett. 33 (1974) 1406, <https://doi.org/10.1103/PhysRevLett.33.1406>.
- [3] R. Baier, R. Ruckl, Hadronic production of J/ψ and $\Upsilon(1S)$: transverse momentum distributions, Phys. Lett. B 102 (1981) 364, [https://doi.org/10.1016/0370-2693\(81\)90636-5](https://doi.org/10.1016/0370-2693(81)90636-5).
- [4] J.-P. Lansberg, On the mechanisms of heavy-quarkonium hadroproduction, Eur. Phys. J. C 61 (2009) 693, <https://doi.org/10.1140/epjc/s10052-008-0826-9>, arXiv:0811.4005.
- [5] F. Abe, et al., CDF, J/ψ and $\psi(2S)$ production in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV, Phys. Rev. Lett. 79 (1997) 572, <https://doi.org/10.1103/PhysRevLett.79.572>.
- [6] G.T. Bodwin, E. Braaten, G.P. Lepage, Rigorous QCD analysis of inclusive annihilation and production of heavy quarkonium, Phys. Rev. D 51 (1995) 1125, <https://doi.org/10.1103/PhysRevD.51.1125>, arXiv:hep-ph/9407339, Erratum: <https://doi.org/10.1103/PhysRevD.55.5853>.
- [7] P. Cho, A.K. Leibovich, Color-octet quarkonia production, Phys. Rev. D 53 (1996) 150, <https://doi.org/10.1103/PhysRevD.53.150>.
- [8] P. Cho, A.K. Leibovich, Color-octet quarkonia production. II, Phys. Rev. D 53 (1996) 6203, <https://doi.org/10.1103/PhysRevD.53.6203>.
- [9] P. Faccioli, V. Knünz, C. Lourenco, J. Seixas, H.K. Wöhri, Quarkonium production in the LHC era: a polarized perspective, Phys. Lett. B 736 (2014) 98, <https://doi.org/10.1016/j.physletb.2014.07.006>, arXiv:1403.3970.
- [10] A. Abulencia, et al., CDF, Polarizations of J/ψ and $\psi(2S)$ mesons produced in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV, Phys. Rev. Lett. 99 (2007) 132001, <https://doi.org/10.1103/PhysRevLett.99.132001>, arXiv:0704.0638.
- [11] CMS Collaboration, Measurement of the prompt J/ψ and $\psi(2S)$ polarizations in pp collisions at $\sqrt{s} = 7$ TeV, Phys. Lett. B 727 (2013) 381, <https://doi.org/10.1016/j.physletb.2013.10.055>, arXiv:1307.6070.

- [12] J.-P. Lansberg, *New observables in inclusive production of quarkonium*, arXiv:1903.09185, 2019.
- [13] M. Baumgart, A.K. Leibovich, T. Mehen, I.Z. Rothstein, *Probing quarkonium production mechanisms with jet substructure*, J. High Energy Phys. 11 (2014) 003, [https://doi.org/10.1007/JHEP11\(2014\)003](https://doi.org/10.1007/JHEP11(2014)003), arXiv:1406.2295.
- [14] LHCb Collaboration, *Study of J/ψ production in jets*, Phys. Rev. Lett. 118 (2017) 192001, <https://doi.org/10.1103/PhysRevLett.118.192001>, arXiv:1701.05116.
- [15] J. Pumplin, D.R. Stump, J. Huston, H.L. Lai, P.M. Nadolsky, W.K. Tung, *New generation of parton distributions with uncertainties from global QCD analysis*, J. High Energy Phys. 07 (2002) 012, <https://doi.org/10.1088/1126-6708/2002/07/012>, arXiv:hep-ph/0201195.
- [16] L. Dai, P. Shrivastava, *Quarkonium polarization and the long distance matrix elements hierarchies using jet substructure*, Phys. Rev. D 96 (2017) 036020, <https://doi.org/10.1103/PhysRevD.96.036020>, arXiv:1707.08629.
- [17] CMS Collaboration, *Determination of jet energy calibration and transverse momentum resolution in CMS*, J. Instrum. 6 (2011) P11002, <https://doi.org/10.1088/1748-0221/6/11/P11002>, arXiv:1107.4277.
- [18] CMS Collaboration, *Description and performance of track and primary-vertex reconstruction with the CMS tracker*, J. Instrum. 9 (2014) P10009, <https://doi.org/10.1088/1748-0221/9/10/P10009>, arXiv:1405.6569.
- [19] CMS Collaboration, *Performance of CMS muon reconstruction in pp collision events at $\sqrt{s} = 7$ TeV*, J. Instrum. 7 (2012) P10002, <https://doi.org/10.1088/1748-0221/7/10/P10002>, arXiv:1206.4071.
- [20] CMS Collaboration, *The CMS trigger system*, J. Instrum. 12 (2017) P01020, <https://doi.org/10.1088/1748-0221/12/01/P01020>, arXiv:1609.02366.
- [21] CMS Collaboration, *The CMS experiment at the CERN LHC*, J. Instrum. 3 (2008) S08004, <https://doi.org/10.1088/1748-0221/3/08/S08004>.
- [22] CMS Collaboration, *Prompt and non-prompt J/ψ production in pp collisions at $\sqrt{s} = 7$ TeV*, Eur. Phys. J. C 71 (2011) 1575, <https://doi.org/10.1140/epjc/s10052-011-1575-8>, arXiv:1011.4193.
- [23] CMS Collaboration, *Upsilon production cross-section in pp collisions at $\sqrt{s} = 7$ TeV*, Phys. Rev. D 83 (2011) 112004, <https://doi.org/10.1103/PhysRevD.83.112004>, arXiv:1012.5545.
- [24] CMS Collaboration, *J/ψ and $\psi(2S)$ production in pp collisions at $\sqrt{s} = 7$ TeV*, J. High Energy Phys. 02 (2012) 011, [https://doi.org/10.1007/JHEP02\(2012\)011](https://doi.org/10.1007/JHEP02(2012)011), arXiv:1111.1557.
- [25] CMS Collaboration, *Measurement of J/ψ and $\psi(2S)$ prompt double-differential cross sections in pp collisions at $\sqrt{s} = 7$ TeV*, Phys. Rev. Lett. 114 (2015) 191802, <https://doi.org/10.1103/PhysRevLett.114.191802>, arXiv:1502.04155.
- [26] CMS Collaboration, *Measurements of the $Y(1S)$, $Y(2S)$, and $Y(3S)$ differential cross sections in pp collisions at $\sqrt{s} = 7$ TeV*, Phys. Lett. B 749 (2015) 14, <https://doi.org/10.1016/j.physletb.2015.07.037>, arXiv:1501.07750.
- [27] CMS Collaboration, *Particle-flow reconstruction and global event description with the CMS detector*, J. Instrum. 12 (2017) P10003, <https://doi.org/10.1088/1748-0221/12/10/P10003>, arXiv:1706.04965.
- [28] M. Cacciari, G.P. Salam, G. Soyez, *The anti- k_r jet clustering algorithm*, J. High Energy Phys. 04 (2008) 063, <https://doi.org/10.1088/1126-6708/2008/04/063>, arXiv:0802.1189.
- [29] M. Cacciari, G.P. Salam, G. Soyez, *FastJet user manual*, Eur. Phys. J. C 72 (2012) 1896, <https://doi.org/10.1140/epjc/s10052-012-1896-2>, arXiv:1111.6097.
- [30] G.T. Bodwin, H.S. Chung, U.-R. Kim, J. Lee, *Fragmentation contributions to J/ψ production at the Tevatron and the LHC*, Phys. Rev. Lett. 113 (2014) 022001, <https://doi.org/10.1103/PhysRevLett.113.022001>, arXiv:1403.3612.
- [31] N. Brambilla, et al., *Heavy quarkonium: progress, puzzles, and opportunities*, Eur. Phys. J. C 71 (2011) 1534, <https://doi.org/10.1140/epjc/s10052-010-1534-9>, arXiv:1010.5827.
- [32] J.P. Lansberg, *QCD corrections to J/ψ polarisation in pp collisions at RHIC*, Phys. Lett. B 695 (2011) 149, <https://doi.org/10.1016/j.physletb.2010.10.054>, arXiv:1003.4319.
- [33] M. Butenschoen, B.A. Kniehl, *Reconciling J/ψ production at HERA, RHIC, Tevatron, and LHC with NRQCD factorization at next-to-leading order*, Phys. Rev. Lett. 106 (2011) 022003, <https://doi.org/10.1103/PhysRevLett.106.022003>, arXiv:1009.5662.
- [34] T. Sjostrand, S. Mrenna, P.Z. Skands, *PYTHIA 6.4 physics and manual*, J. High Energy Phys. 05 (2006) 026, <https://doi.org/10.1088/1126-6708/2006/05/026>, arXiv:hep-ph/0603175.
- [35] D.J. Lange, *The EvtGen particle decay simulation package*, Nucl. Instrum. Methods A 462 (2001) 152, [https://doi.org/10.1016/S0168-9002\(01\)00089-4](https://doi.org/10.1016/S0168-9002(01)00089-4).
- [36] E. Barberio, B. van Eijk, Z. Was, *Photos: a universal Monte Carlo for QED radiative corrections in decays*, Comput. Phys. Commun. 66 (1991) 115, [https://doi.org/10.1016/0010-4655\(91\)90012-A](https://doi.org/10.1016/0010-4655(91)90012-A).
- [37] G. Collaboration, (GEANT4), *geant 4 - a simulation toolkit*, Nucl. Instrum. Methods A 506 (2003) 250, [https://doi.org/10.1016/S0168-9002\(03\)01368-8](https://doi.org/10.1016/S0168-9002(03)01368-8).
- [38] G. D'Agostini, *A multidimensional unfolding method based on Bayes' theorem*, Nucl. Instrum. Methods A 362 (1995) 487, [https://doi.org/10.1016/0168-9002\(95\)00274-X](https://doi.org/10.1016/0168-9002(95)00274-X).
- [39] T. Auye, *Unfolding algorithms and tests using RooUnfold*, in: *Proceedings, PHYSTAT 2011 Workshop on Statistical Issues Related to Discovery Claims in Search Experiments and Unfolding*, CERN, Geneva, Switzerland, 2011, p. 313, arXiv:1105.1160.
- [40] CMS Collaboration, *Muon ID performance: low- p_T muon efficiencies*, Technical Report CMS DP-2014/020, 2014, https://twiki.cern.ch/twiki/pub/CMSPublic/PhysicsResultsMUO/DP2014_020.pdf.
- [41] M. Butenschoen, B.A. Kniehl, *Next-to-leading-order tests of NRQCD factorization with J/ψ yield and polarization*, Mod. Phys. Lett. A 28 (2013) 1350027, <https://doi.org/10.1142/S0217732313500272>, arXiv:1212.2037.
- [42] K.-T. Chao, Y.-Q. Ma, H.-S. Shao, K. Wang, Y.-J. Zhang, *J/ψ polarization at hadron colliders in nonrelativistic QCD*, Phys. Rev. Lett. 108 (2012) 242004, <https://doi.org/10.1103/PhysRevLett.108.242004>, arXiv:1201.2675.

The CMS Collaboration

A.M. Sirunyan[†], A. Tumasyan

Yerevan Physics Institute, Yerevan, Armenia

W. Adam, F. Ambroggi, T. Bergauer, J. Brandstetter, M. Dragicevic, J. Erö, A. Escalante Del Valle, M. Flechl, R. Frühwirth¹, M. Jeitler¹, N. Krammer, I. Krätschmer, D. Liko, T. Madlener, I. Mikulec, N. Rad, J. Schieck¹, R. Schöfbeck, M. Spanring, D. Spitzbart, W. Waltenberger, C.-E. Wulz¹, M. Zarucki

Institut für Hochenergiephysik, Wien, Austria

V. Drugakov, V. Mossolov, J. Suarez Gonzalez

Institute for Nuclear Problems, Minsk, Belarus

M.R. Darwish, E.A. De Wolf, D. Di Croce, X. Janssen, A. Lelek, M. Pieters, H. Rejeb Sfar, H. Van Haevermaet, P. Van Mechelen, S. Van Putte, N. Van Remortel

Universiteit Antwerpen, Antwerpen, Belgium

F. Blekman, E.S. Bols, S.S. Chhibra, J. D'Hondt, J. De Clercq, D. Lontkovskiy, S. Lowette, I. Marchesini, S. Moortgat, Q. Python, K. Skovpen, S. Tavernier, W. Van Doninck, P. Van Mulders

Vrije Universiteit Brussel, Brussel, Belgium

D. Beghin, B. Bilin, H. Brun, B. Clerbaux, G. De Lentdecker, H. Delannoy, B. Dorney, L. Favart, A. Grebenyuk, A.K. Kalsi, A. Popov, N. Postiau, E. Starling, L. Thomas, C. Vander Velde, P. Vanlaer, D. Vannerom

Université Libre de Bruxelles, Bruxelles, Belgium

T. Cornelis, D. Dobur, I. Khvastunov², M. Niedziela, C. Roskas, D. Trocino, M. Tytgat, W. Verbeke, B. Vermassen, M. Vit, N. Zaganidis

Ghent University, Ghent, Belgium

O. Bondu, G. Bruno, C. Caputo, P. David, C. Delaere, M. Delcourt, A. Giammanco, V. Lemaitre, A. Magitteri, J. Prisciandaro, A. Saggio, M. Vidal Marono, P. Vischia, J. Zobec

Université Catholique de Louvain, Louvain-la-Neuve, Belgium

F.L. Alves, G.A. Alves, G. Correia Silva, C. Hensel, A. Moraes, P. Rebello Teles

Centro Brasileiro de Pesquisas Fisicas, Rio de Janeiro, Brazil

E. Belchior Batista Das Chagas, W. Carvalho, J. Chinellato³, E. Coelho, E.M. Da Costa, G.G. Da Silveira⁴, D. De Jesus Damiao, C. De Oliveira Martins, S. Fonseca De Souza, L.M. Huertas Guativa, H. Malbouisson, J. Martins⁵, D. Matos Figueiredo, M. Medina Jaime⁶, M. Melo De Almeida, C. Mora Herrera, L. Mundim, H. Nogima, W.L. Prado Da Silva, L.J. Sanchez Rosas, A. Santoro, A. Sznajder, M. Thiel, E.J. Tonelli Manganote³, F. Torres Da Silva De Araujo, A. Vilela Pereira

Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

C.A. Bernardes^a, L. Calligaris^a, T.R. Fernandez Perez Tomei^a, E.M. Gregores^b, D.S. Lemos, P.G. Mercadante^b, S.F. Novaes^a, Sandra S. Padula^a

^a *Universidade Estadual Paulista, São Paulo, Brazil*

^b *Universidade Federal do ABC, São Paulo, Brazil*

A. Aleksandrov, G. Antchev, R. Hadjiiska, P. Iaydjiev, M. Misheva, M. Rodozov, M. Shopova, G. Sultanov

Institute for Nuclear Research and Nuclear Energy, Bulgarian Academy of Sciences, Sofia, Bulgaria

M. Bonchev, A. Dimitrov, T. Ivanov, L. Litov, B. Pavlov, P. Petkov

University of Sofia, Sofia, Bulgaria

W. Fang⁷, X. Gao⁷, L. Yuan

Beihang University, Beijing, China

Z. Hu, Y. Wang

Department of Physics, Tsinghua University, Beijing, China

M. Ahmad, G.M. Chen, H.S. Chen, M. Chen, C.H. Jiang, D. Leggat, H. Liao, Z. Liu, S.M. Shaheen⁸, A. Spiezia, J. Tao, E. Yazgan, H. Zhang, S. Zhang⁸, J. Zhao

Institute of High Energy Physics, Beijing, China

A. Agapitos, Y. Ban, G. Chen, A. Levin, J. Li, L. Li, Q. Li, Y. Mao, S.J. Qian, D. Wang, Q. Wang

State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing, China

M. Xiao

Zhejiang University, Hangzhou, China

C. Avila, A. Cabrera, C. Florez, C.F. González Hernández, M.A. Segura Delgado

Universidad de Los Andes, Bogota, Colombia

J. Mejia Guisao, J.D. Ruiz Alvarez, C.A. Salazar González, N. Vanegas Arbelaez

Universidad de Antioquia, Medellin, Colombia

D. Giljanović, N. Godinovic, D. Lelas, I. Puljak, T. Sculac

University of Split, Faculty of Electrical Engineering, Mechanical Engineering and Naval Architecture, Split, Croatia

Z. Antunovic, M. Kovac

University of Split, Faculty of Science, Split, Croatia

V. Brigljevic, S. Ceci, D. Ferencek, K. Kadija, B. Mesic, M. Roguljic, A. Starodumov⁹, T. Susa

Institute Rudjer Boskovic, Zagreb, Croatia

M.W. Ather, A. Attikis, E. Erodotou, A. Ioannou, M. Kolosova, S. Konstantinou, G. Mavromanolakis, J. Mousa, C. Nicolaou, F. Ptochos, P.A. Razis, H. Rykaczewski, D. Tsiakkouri

University of Cyprus, Nicosia, Cyprus

M. Finger¹⁰, M. Finger Jr.¹⁰, A. Kveton, J. Tomsa

Charles University, Prague, Czech Republic

E. Ayala

Escuela Politecnica Nacional, Quito, Ecuador

E. Carrera Jarrin

Universidad San Francisco de Quito, Quito, Ecuador

Y. Assran^{11,12}, S. Elgammal¹²

Academy of Scientific Research and Technology of the Arab Republic of Egypt, Egyptian Network of High Energy Physics, Cairo, Egypt

S. Bhowmik, A. Carvalho Antunes De Oliveira, R.K. Dewanjee, K. Ehataht, M. Kadastik, M. Raidal, C. Veelken

National Institute of Chemical Physics and Biophysics, Tallinn, Estonia

P. Eerola, L. Forthomme, H. Kirschenmann, K. Osterberg, M. Voutilainen

Department of Physics, University of Helsinki, Helsinki, Finland

F. Garcia, J. Havukainen, J.K. Heikkilä, T. Järvinen, V. Karimäki, M.S. Kim, R. Kinnunen, T. Lampén, K. Lassila-Perini, S. Laurila, S. Lehti, T. Lindén, P. Luukka, T. Mäenpää, H. Siikonen, E. Tuominen, J. Tuominiemi

Helsinki Institute of Physics, Helsinki, Finland

T. Tuuva

Lappeenranta University of Technology, Lappeenranta, Finland

M. Besancon, F. Couderc, M. Dejardin, D. Denegri, B. Fabbro, J.L. Faure, F. Ferri, S. Ganjour, A. Givernaud, P. Gras, G. Hamel de Monchenault, P. Jarry, C. Leloup, E. Locci, J. Malcles, J. Rander, A. Rosowsky, M.Ö. Sahin, A. Savoy-Navarro¹³, M. Titov

IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette, France

S. Ahuja, C. Amendola, F. Beaudette, P. Busson, C. Charlot, B. Diab, G. Falmagne, R. Granier de Cassagnac, I. Kucher, A. Lobanov, C. Martin Perez, M. Nguyen, C. Ochando, P. Paganini, J. Rembser, R. Salerno, J.B. Sauvan, Y. Sirois, A. Zabi, A. Zghiche

Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, Institut Polytechnique de Paris, France

J.-L. Agram¹⁴, J. Andrea, D. Bloch, G. Bourgatte, J.-M. Brom, E.C. Chabert, C. Collard, E. Conte¹⁴, J.-C. Fontaine¹⁴, D. Gelé, U. Goerlach, M. Jansová, A.-C. Le Bihan, N. Tonon, P. Van Hove

Université de Strasbourg, CNRS, IPHC UMR 7178, Strasbourg, France

S. Gadrat

Centre de Calcul de l'Institut National de Physique Nucleaire et de Physique des Particules, CNRS/IN2P3, Villeurbanne, France

S. Beauceron, C. Bernet, G. Boudoul, C. Camen, A. Carle, N. Chanon, R. Chierici, D. Contardo, P. Depasse, H. El Mamouni, J. Fay, S. Gascon, M. Gouzevitch, B. Ille, Sa. Jain, F. Lagarde, I.B. Laktineh, H. Lattaud, A. Lesauvage, M. Lethuillier, L. Mirabito, S. Perries, V. Sordini, L. Torterotot, G. Touquet, M. Vander Donckt, S. Viret

Université de Lyon, Université Claude Bernard Lyon 1, CNRS-IN2P3, Institut de Physique Nucléaire de Lyon, Villeurbanne, France

T. Toriashvili¹⁵

Georgian Technical University, Tbilisi, Georgia

I. Bagaturia¹⁶

Tbilisi State University, Tbilisi, Georgia

C. Autermann, L. Feld, M.K. Kiesel, K. Klein, M. Lipinski, D. Meuser, A. Pauls, M. Preuten, M.P. Rauch, C. Schomakers, J. Schulz, M. Teroerde, B. Wittmer

RWTH Aachen University, I. Physikalisches Institut, Aachen, Germany

A. Albert, M. Erdmann, S. Erdweg, T. Esch, B. Fischer, S. Ghosh, T. Hebbeker, K. Hoepfner, H. Keller, L. Mastrolorenzo, M. Merschmeyer, A. Meyer, P. Millet, G. Mocellin, S. Mondal, S. Mukherjee, D. Noll, A. Novak, T. Pook, A. Pozdnyakov, T. Quast, M. Radziej, Y. Rath, H. Reithler, M. Rieger, J. Roemer, A. Schmidt, S.C. Schuler, A. Sharma, S. Wiedenbeck, S. Zaleski

RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany

G. Flügge, W. Haj Ahmad¹⁷, O. Hlushchenko, T. Kress, T. Müller, A. Nehr Korn, A. Nowack, C. Pistone, O. Pooth, D. Roy, H. Sert, A. Stahl¹⁸

RWTH Aachen University, III. Physikalisches Institut B, Aachen, Germany

M. Aldaya Martin, P. Asmuss, I. Babounikau, H. Bakhshiansohi, K. Beernaert, O. Behnke, U. Behrens, A. Bermúdez Martínez, D. Bertsche, A.A. Bin Anuar, K. Borras¹⁹, V. Botta, A. Campbell, A. Cardini, P. Connor, S. Consuegra Rodríguez, C. Contreras-Campana, V. Danilov, A. De Wit, M.M. Defranchis, C. Diez Pardos, D. Domínguez Damiani, G. Eckerlin, D. Eckstein, T. Eichhorn, A. Elwood, E. Eren, E. Gallo²⁰, A. Geiser, J.M. Grados Luyando, A. Grohsjean, M. Guthoff, M. Haranko, A. Harb, A. Jafari, N.Z. Jomhari, H. Jung, A. Kasem¹⁹, M. Kasemann, H. Kaveh, J. Keaveney, C. Kleinwort, J. Knolle, D. Krücker, W. Lange, T. Lenz, J. Leonard, J. Lidrych, K. Lipka, W. Lohmann²¹, R. Mankel, I.-A. Melzer-Pellmann, A.B. Meyer, M. Meyer, M. Missiroli, G. Mittag, J. Mnich, A. Mussgiller, V. Myronenko, D. Pérez Adán, S.K. Pflitsch, D. Pitzl, A. Raspereza, A. Saibel, M. Savitskyi, V. Scheurer, P. Schütze, C. Schwanenberger, R. Shevchenko, A. Singh, H. Tholen, O. Turkot, A. Vagnerini, M. Van De Klundert, G.P. Van Onsem, R. Walsh, Y. Wen, K. Wichmann, C. Wissing, O. Zenaiev, R. Zlebick

Deutsches Elektronen-Synchrotron, Hamburg, Germany

R. Aggleton, S. Bein, L. Benato, A. Benecke, V. Blobel, T. Dreyer, A. Ebrahimi, A. Fröhlich, C. Garbers, E. Garutti, D. Gonzalez, P. Gunnellini, J. Haller, A. Hinzmann, A. Karavdina, G. Kasieczka, R. Klanner, R. Kogler, N. Kovalchuk, S. Kurz, V. Kutzner, J. Lange, T. Lange, A. Malara, J. Multhaupt, C.E.N. Niemeyer, A. Perieanu, A. Reimers, O. Rieger, C. Scharf, P. Schleper, S. Schumann, J. Schwandt, J. Sonneveld, H. Stadie, G. Steinbrück, F.M. Stober, M. Stöver, B. Vormwald, I. Zoi

University of Hamburg, Hamburg, Germany

M. Akbiyik, C. Barth, M. Baselga, S. Baur, T. Berger, E. Butz, R. Caspart, T. Chwalek, W. De Boer, A. Dierlamm, K. El Morabit, N. Faltermann, M. Giffels, P. Goldenzweig, A. Gottmann, M.A. Harrendorf, F. Hartmann¹⁸, U. Husemann, S. Kudella, S. Mitra, M.U. Mozer, D. Müller, Th. Müller, M. Musich, A. Nürnberg, G. Quast, K. Rabbertz, M. Schröder, I. Shvetsov, H.J. Simonis, R. Ulrich, M. Wassmer, M. Weber, C. Wöhrmann, R. Wolf

Karlsruher Institut fuer Technologie, Karlsruhe, Germany

G. Anagnostou, P. Asenov, G. Daskalakis, T. Gerasis, A. Kyriakis, D. Loukas, G. Paspalaki

Institute of Nuclear and Particle Physics (INPP), NCSR Demokritos, Aghia Paraskevi, Greece

M. Diamantopoulou, G. Karathanasis, P. Kontaxakis, A. Manousakis-katsikakis, A. Panagiotou, I. Papavergou, N. Saoulidou, A. Stakia, K. Theofilatos, K. Vellidis, E. Vourliotis

National and Kapodistrian University of Athens, Athens, Greece

G. Bakas, K. Kousouris, I. Papakrivopoulos, G. Tsipolitis

National Technical University of Athens, Athens, Greece

I. Evangelou, C. Foudas, P. Gianneios, P. Katsoulis, P. Kokkas, S. Mallios, K. Manitaras, N. Manthos, I. Papadopoulos, J. Strologas, F.A. Triantis, D. Tsitsonis

University of Ioánnina, Ioánnina, Greece

M. Bartók²², R. Chudasama, M. Csanad, P. Major, K. Mandal, A. Mehta, M.I. Nagy, G. Pasztor, O. Surányi, G.I. Veres

MTA-ELTE Lendület CMS Particle and Nuclear Physics Group, Eötvös Loránd University, Budapest, Hungary

G. Bencze, C. Hajdu, D. Horvath²³, F. Sikler, T.Á. Vámi, V. Veszpremi, G. Vesztergombi[†]

Wigner Research Centre for Physics, Budapest, Hungary

N. Beni, S. Czellar, J. Karancsi²², A. Makovec, J. Molnar, Z. Szillasi

Institute of Nuclear Research ATOMKI, Debrecen, Hungary

P. Raics, D. Teyssier, Z.L. Trocsanyi, B. Ujvari

Institute of Physics, University of Debrecen, Debrecen, Hungary

T. Csorgo, W.J. Metzger, F. Nemes, T. Novak

Eszterhazy Karoly University, Karoly Robert Campus, Gyongyos, Hungary

S. Choudhury, J.R. Komaragiri, P.C. Tiwari

Indian Institute of Science (IISc), Bangalore, India

S. Bahinipati²⁴, C. Kar, G. Kole, P. Mal, V.K. Muraleedharan Nair Bindhu, A. Nayak²⁵, D.K. Sahoo²⁴, S.K. Swain

National Institute of Science Education and Research, HBNI, Bhubaneswar, India

S. Bansal, S.B. Beri, V. Bhatnagar, S. Chauhan, R. Chawla, N. Dhingra, R. Gupta, A. Kaur, M. Kaur, S. Kaur, P. Kumari, M. Lohan, M. Meena, K. Sandeep, S. Sharma, J.B. Singh, A.K. Viridi, G. Walia

Panjab University, Chandigarh, India

A. Bhardwaj, B.C. Choudhary, R.B. Garg, M. Gola, S. Keshri, Ashok Kumar, S. Malhotra, M. Naimuddin, P. Priyanka, K. Ranjan, Aashaq Shah, R. Sharma

University of Delhi, Delhi, India

R. Bhardwaj²⁶, M. Bharti²⁶, R. Bhattacharya, S. Bhattacharya, U. Bhawandeep²⁶, D. Bhowmik, S. Dutta, S. Ghosh, M. Maity²⁷, K. Mondal, S. Nandan, A. Purohit, P.K. Rout, G. Saha, S. Sarkar, T. Sarkar²⁷, M. Sharan, B. Singh²⁶, S. Thakur²⁶

Saha Institute of Nuclear Physics, HBNI, Kolkata, India

P.K. Behera, P. Kalbhor, A. Muhammad, P.R. Pujahari, A. Sharma, A.K. Sikdar

Indian Institute of Technology Madras, Madras, India

D. Dutta, V. Jha, V. Kumar, D.K. Mishra, P.K. Netrakanti, L.M. Pant, P. Shukla

Bhabha Atomic Research Centre, Mumbai, India

T. Aziz, M.A. Bhat, S. Dugad, G.B. Mohanty, N. Sur, R. Kumar Verma

Tata Institute of Fundamental Research-A, Mumbai, India

S. Banerjee, S. Bhattacharya, S. Chatterjee, P. Das, M. Guchait, S. Karmakar, S. Kumar, G. Majumder, K. Mazumdar, N. Sahoo, S. Sawant

Tata Institute of Fundamental Research-B, Mumbai, India

S. Chauhan, S. Dube, V. Hegde, B. Kansal, A. Kapoor, K. Kothekar, S. Pandey, A. Rane, A. Rastogi, S. Sharma

Indian Institute of Science Education and Research (IISER), Pune, India

S. Chenarani²⁸, E. Eskandari Tadavani, S.M. Etesami²⁸, M. Khakzad, M. Mohammadi Najafabadi, M. Naseri, F. Rezaei Hosseinabadi

Institute for Research in Fundamental Sciences (IPM), Tehran, Iran

M. Felcini, M. Grunewald

University College Dublin, Dublin, Ireland

M. Abbrescia^{a,b}, R. Aly^{a,b,29}, C. Calabria^{a,b}, A. Colaleo^a, D. Creanza^{a,c}, L. Cristella^{a,b}, N. De Filippis^{a,c}, M. De Palma^{a,b}, A. Di Florio^{a,b}, L. Fiore^a, A. Gelmi^{a,b}, G. Iaselli^{a,c}, M. Ince^{a,b}, S. Lezki^{a,b}, G. Maggi^{a,c}, M. Maggi^a, G. Miniello^{a,b}, S. My^{a,b}, S. Nuzzo^{a,b}, A. Pompili^{a,b}, G. Pugliese^{a,c}, R. Radogna^a, A. Ranieri^a, G. Selvaggi^{a,b}, L. Silvestris^a, R. Venditti^a, P. Verwilligen^a

^a INFN Sezione di Bari, Bari, Italy

^b Università di Bari, Bari, Italy

^c Politecnico di Bari, Bari, Italy

G. Abbiendi^a, C. Battilana^{a,b}, D. Bonacorsi^{a,b}, L. Borgonovi^{a,b}, S. Braibant-Giacomelli^{a,b}, R. Campanini^{a,b}, P. Capiluppi^{a,b}, A. Castro^{a,b}, F.R. Cavallo^a, C. Ciocca^a, G. Codispoti^{a,b}, M. Cuffiani^{a,b}, G.M. Dallavalle^a, F. Fabbri^a, A. Fanfani^{a,b}, E. Fontanesi^{a,b}, P. Giacomelli^a, C. Grandi^a, L. Guiducci^{a,b}, F. Iemmi^{a,b}, S. Lo Meo^{a,30}, S. Marcellini^a, G. Masetti^a, F.L. Navarria^{a,b}, A. Perrotta^a, F. Primavera^{a,b}, A.M. Rossi^{a,b}, T. Rovelli^{a,b}, G.P. Siroli^{a,b}, N. Tosi^a

^a INFN Sezione di Bologna, Bologna, Italy

^b Università di Bologna, Bologna, Italy

S. Albergo^{a,b,31}, S. Costa^{a,b}, A. Di Mattia^a, R. Potenza^{a,b}, A. Tricomi^{a,b,31}, C. Tuve^{a,b}

^a INFN Sezione di Catania, Catania, Italy

^b Università di Catania, Catania, Italy

G. Barbagli^a, A. Cassese, R. Ceccarelli, K. Chatterjee^{a,b}, V. Ciulli^{a,b}, C. Civinini^a, R. D'Alessandro^{a,b}, E. Focardi^{a,b}, G. Latino^{a,b}, P. Lenzi^{a,b}, M. Meschini^a, S. Paoletti^a, G. Sguazzoni^a, L. Viliani^a

^a INFN Sezione di Firenze, Firenze, Italy

^b Università di Firenze, Firenze, Italy

L. Benussi, S. Bianco, D. Piccolo

INFN Laboratori Nazionali di Frascati, Frascati, Italy

M. Bozzo^{a,b}, F. Ferro^a, R. Mulargia^{a,b}, E. Robutti^a, S. Tosi^{a,b}

^a *INFN Sezione di Genova, Genova, Italy*

^b *Università di Genova, Genova, Italy*

A. Benaglia^a, A. Beschi^{a,b}, F. Brivio^{a,b}, V. Ciriolo^{a,b,18}, S. Di Guida^{a,b,18}, M.E. Dinardo^{a,b}, P. Dini^a, S. Gennai^a, A. Ghezzi^{a,b}, P. Govoni^{a,b}, L. Guzzi^{a,b}, M. Malberti^a, S. Malvezzi^a, D. Menasce^a, F. Monti^{a,b}, L. Moroni^a, G. Ortona^{a,b}, M. Paganoni^{a,b}, D. Pedrini^a, S. Ragazzi^{a,b}, T. Tabarelli de Fatis^{a,b}, D. Zuolo^{a,b}

^a *INFN Sezione di Milano-Bicocca, Milano, Italy*

^b *Università di Milano-Bicocca, Milano, Italy*

S. Buontempo^a, N. Cavallo^{a,c}, A. De Iorio^{a,b}, A. Di Crescenzo^{a,b}, F. Fabozzi^{a,c}, F. Fienga^a, G. Galati^a, A.O.M. Iorio^{a,b}, L. Lista^{a,b}, S. Meola^{a,d,18}, P. Paolucci^{a,18}, B. Rossi^a, C. Sciacca^{a,b}, E. Voevodina^{a,b}

^a *INFN Sezione di Napoli, Napoli, Italy*

^b *Università di Napoli 'Federico II', Napoli, Italy*

^c *Università della Basilicata, Potenza, Italy*

^d *Università G. Marconi, Roma, Italy*

P. Azzi^a, N. Bacchetta^a, D. Bisello^{a,b}, A. Boletti^{a,b}, A. Bragagnolo^{a,b}, R. Carlin^{a,b}, P. Checchia^a, P. De Castro Manzano^a, T. Dorigo^a, U. Dosselli^a, F. Gasparini^{a,b}, U. Gasparini^{a,b}, A. Gozzelino^a, S.Y. Hoh^{a,b}, P. Lujan^a, M. Margoni^{a,b}, A.T. Meneguzzo^{a,b}, J. Pazzini^{a,b}, M. Presilla^b, P. Ronchese^{a,b}, R. Rossin^{a,b}, F. Simonetto^{a,b}, A. Tiko^a, M. Tosi^{a,b}, M. Zanetti^{a,b}, P. Zotto^{a,b}, G. Zumerle^{a,b}

^a *INFN Sezione di Padova, Padova, Italy*

^b *Università di Padova, Padova, Italy*

^c *Università di Trento, Trento, Italy*

A. Braghieri^a, D. Fiorina, P. Montagna^{a,b}, S.P. Ratti^{a,b}, V. Re^a, M. Ressegotti^{a,b}, C. Riccardi^{a,b}, P. Salvini^a, I. Vai^{a,b}, P. Vitulo^{a,b}

^a *INFN Sezione di Pavia, Pavia, Italy*

^b *Università di Pavia, Pavia, Italy*

M. Biasini^{a,b}, G.M. Bilei^a, D. Ciangottini^{a,b}, L. Fanò^{a,b}, P. Lariccia^{a,b}, R. Leonardi^{a,b}, G. Mantovani^{a,b}, V. Mariani^{a,b}, M. Menichelli^a, A. Rossi^{a,b}, A. Santocchia^{a,b}, D. Spiga^a

^a *INFN Sezione di Perugia, Perugia, Italy*

^b *Università di Perugia, Perugia, Italy*

K. Androsov^a, P. Azzurri^a, G. Bagliesi^a, V. Bertacchi^{a,c}, L. Bianchini^a, T. Boccali^a, R. Castaldi^a, M.A. Ciocci^{a,b}, R. Dell'Orso^a, G. Fedi^a, L. Giannini^{a,c}, A. Giassi^a, M.T. Grippo^a, F. Ligabue^{a,c}, E. Manca^{a,c}, G. Mandorli^{a,c}, A. Messineo^{a,b}, F. Palla^a, A. Rizzi^{a,b}, G. Rolandi³², S. Roy Chowdhury, A. Scribano^a, P. Spagnolo^a, R. Tenchini^a, G. Tonelli^{a,b}, N. Turini, A. Venturi^a, P.G. Verdini^a

^a *INFN Sezione di Pisa, Pisa, Italy*

^b *Università di Pisa, Pisa, Italy*

^c *Scuola Normale Superiore di Pisa, Pisa, Italy*

F. Cavallari^a, M. Cipriani^{a,b}, D. Del Re^{a,b}, E. Di Marco^{a,b}, M. Diemoz^a, E. Longo^{a,b}, B. Marzocchi^{a,b}, P. Meridiani^a, G. Organtini^{a,b}, F. Pandolfi^a, R. Paramatti^{a,b}, C. Quaranta^{a,b}, S. Rahatlou^{a,b}, C. Rovelli^a, F. Santanastasio^{a,b}, L. Soffi^{a,b}

^a *INFN Sezione di Roma, Rome, Italy*

^b *Sapienza Università di Roma, Rome, Italy*

N. Amapane^{a,b}, R. Arcidiacono^{a,c}, S. Argiro^{a,b}, M. Arneodo^{a,c}, N. Bartosik^a, R. Bellan^{a,b}, A. Bellora, C. Biino^a, A. Cappati^{a,b}, N. Cartiglia^a, S. Cometti^a, M. Costa^{a,b}, R. Covarelli^{a,b}, N. Demaria^a, B. Kiani^{a,b}, C. Mariotti^a, S. Maselli^a, E. Migliore^{a,b}, V. Monaco^{a,b}, E. Monteil^{a,b}, M. Monteno^a, M.M. Obertino^{a,b},

L. Pacher^{a,b}, N. Pastrone^a, M. Pelliccioni^a, G.L. Pinna Angioni^{a,b}, A. Romero^{a,b}, M. Ruspa^{a,c}, R. Sacchi^{a,b},
R. Salvatico^{a,b}, V. Sola^a, A. Solano^{a,b}, D. Soldi^{a,b}, A. Staiano^a

^a INFN Sezione di Torino, Torino, Italy

^b Università di Torino, Torino, Italy

^c Università del Piemonte Orientale, Novara, Italy

S. Belforte^a, V. Candelise^{a,b}, M. Casarsa^a, F. Cossutti^a, A. Da Rold^{a,b}, G. Della Ricca^{a,b}, F. Vazzoler^{a,b},
A. Zanetti^a

^a INFN Sezione di Trieste, Trieste, Italy

^b Università di Trieste, Trieste, Italy

B. Kim, D.H. Kim, G.N. Kim, J. Lee, S.W. Lee, C.S. Moon, Y.D. Oh, S.I. Pak, S. Sekmen, D.C. Son, Y.C. Yang
Kyungpook National University, Daegu, Republic of Korea

H. Kim, D.H. Moon, G. Oh

Chonnam National University, Institute for Universe and Elementary Particles, Kwangju, Republic of Korea

B. Francois, T.J. Kim, J. Park

Hanyang University, Seoul, Republic of Korea

S. Cho, S. Choi, Y. Go, D. Gyun, S. Ha, B. Hong, K. Lee, K.S. Lee, J. Lim, J. Park, S.K. Park, Y. Roh, J. Yoo

Korea University, Seoul, Republic of Korea

J. Goh

Kyung Hee University, Department of Physics, Republic of Korea

H.S. Kim

Sejong University, Seoul, Republic of Korea

J. Almond, J.H. Bhyun, J. Choi, S. Jeon, J. Kim, J.S. Kim, H. Lee, K. Lee, S. Lee, K. Nam, M. Oh, S.B. Oh,
B.C. Radburn-Smith, U.K. Yang, H.D. Yoo, I. Yoon, G.B. Yu

Seoul National University, Seoul, Republic of Korea

D. Jeon, H. Kim, J.H. Kim, J.S.H. Lee, I.C. Park, I. Watson

University of Seoul, Seoul, Republic of Korea

Y. Choi, C. Hwang, Y. Jeong, J. Lee, Y. Lee, I. Yu

Sungkyunkwan University, Suwon, Republic of Korea

V. Veckalns³³

Riga Technical University, Riga, Latvia

V. Dudenias, A. Juodagalvis, G. Tamulaitis, J. Vaitkus

Vilnius University, Vilnius, Lithuania

Z.A. Ibrahim, F. Mohamad Idris³⁴, W.A.T. Wan Abdullah, M.N. Yusli, Z. Zolkapli

National Centre for Particle Physics, Universiti Malaya, Kuala Lumpur, Malaysia

J.F. Benitez, A. Castaneda Hernandez, J.A. Murillo Quijada, L. Valencia Palomo

Universidad de Sonora (UNISON), Hermosillo, Mexico

H. Castilla-Valdez, E. De La Cruz-Burelo, I. Heredia-De La Cruz³⁵, R. Lopez-Fernandez,
A. Sanchez-Hernandez

Centro de Investigacion y de Estudios Avanzados del IPN, Mexico City, Mexico

S. Carrillo Moreno, C. Oropeza Barrera, M. Ramirez-Garcia, F. Vazquez Valencia

Universidad Iberoamericana, Mexico City, Mexico

J. Eysermans, I. Pedraza, H.A. Salazar Ibarquen, C. Uribe Estrada

Benemerita Universidad Autonoma de Puebla, Puebla, Mexico

A. Morelos Pineda

Universidad Autónoma de San Luis Potosí, San Luis Potosí, Mexico

J. Mijuskovic, N. Raicevic

University of Montenegro, Podgorica, Montenegro

D. Krofcheck

University of Auckland, Auckland, New Zealand

S. Bheesette, P.H. Butler

University of Canterbury, Christchurch, New Zealand

A. Ahmad, M. Ahmad, Q. Hassan, H.R. Hoorani, W.A. Khan, M.A. Shah, M. Shoaib, M. Waqas

National Centre for Physics, Quaid-I-Azam University, Islamabad, Pakistan

V. Avati, L. Grzanka, M. Malawski

AGH University of Science and Technology Faculty of Computer Science, Electronics and Telecommunications, Krakow, Poland

H. Bialkowska, M. Bluj, B. Boimska, M. Górski, M. Kazana, M. Szleper, P. Zalewski

National Centre for Nuclear Research, Swierk, Poland

K. Bunkowski, A. Byszuk³⁶, K. Doroba, A. Kalinowski, M. Konecki, J. Krolikowski, M. Misiura, M. Olszewski, M. Walczak

Institute of Experimental Physics, Faculty of Physics, University of Warsaw, Warsaw, Poland

M. Araujo, P. Bargassa, D. Bastos, A. Di Francesco, P. Faccioli, B. Galinhas, M. Gallinaro, J. Hollar, N. Leonardo, J. Seixas, K. Shchelina, G. Strong, O. Toldaiev, J. Varela

Laboratório de Instrumentação e Física Experimental de Partículas, Lisboa, Portugal

P. Bunin, Y. Ershov, M. Gavrilenko, A. Golunov, I. Golutvin, I. Gorbunov, A. Kamenev, V. Karjavine, G. Kozlov, A. Lanev, A. Malakhov, V. Matveev^{37,38}, P. Moisezenz, V. Palichik, V. Perelygin, S. Shmatov, S. Shulha, N. Voytishin, B.S. Yuldashev³⁹, A. Zarubin

Joint Institute for Nuclear Research, Dubna, Russia

L. Chtchipounov, V. Golovtsov, Y. Ivanov, V. Kim⁴⁰, E. Kuznetsova⁴¹, P. Levchenko, V. Murzin, V. Oreshkin, I. Smirnov, D. Sosnov, V. Sulimov, L. Uvarov, A. Vorobyev

Petersburg Nuclear Physics Institute, Gatchina (St. Petersburg), Russia

Yu. Andreev, A. Dermenev, S. Gninenko, N. Golubev, A. Karneyeu, M. Kirsanov, N. Krasnikov, A. Pashenkov, D. Tlisov, A. Toropin

Institute for Nuclear Research, Moscow, Russia

V. Epshteyn, V. Gavrilov, N. Lychkovskaya, A. Nikitenko⁴², V. Popov, I. Pozdnyakov, G. Safronov, A. Spiridonov, A. Stepenov, M. Toms, E. Vlasov, A. Zhokin

Institute for Theoretical and Experimental Physics named by A.I. Alikhanov of NRC 'Kurchatov Institute', Moscow, Russia

T. Aushev

Moscow Institute of Physics and Technology, Moscow, Russia

O. Bychkova, R. Chistov⁴³, M. Danilov⁴³, S. Polikarpov⁴³, E. Tarkovskii

National Research Nuclear University 'Moscow Engineering Physics Institute' (MEPhI), Moscow, Russia

V. Andreev, M. Azarkin, I. Dremin, M. Kirakosyan, A. Terkulov

P.N. Lebedev Physical Institute, Moscow, Russia

A. Belyaev, E. Boos, M. Dubinin⁴⁴, L. Dudko, A. Ershov, A. Gribushin, V. Klyukhin, O. Kodolova, I. Lokhtin, S. Obraztsov, S. Petrushanko, V. Savrin, A. Snigirev

Skobeltsyn Institute of Nuclear Physics, Lomonosov Moscow State University, Moscow, Russia

A. Barnyakov⁴⁵, V. Blinov⁴⁵, T. Dimova⁴⁵, L. Kardapoltsev⁴⁵, Y. Skovpen⁴⁵

Novosibirsk State University (NSU), Novosibirsk, Russia

I. Azhgirey, I. Bayshev, S. Bitiukov, V. Kachanov, D. Konstantinov, P. Mandrik, V. Petrov, R. Ryutin, S. Slabospitskii, A. Sobol, S. Troshin, N. Tyurin, A. Uzunian, A. Volkov

Institute for High Energy Physics of National Research Centre 'Kurchatov Institute', Protvino, Russia

A. Babaev, A. Iuzhakov, V. Okhotnikov

National Research Tomsk Polytechnic University, Tomsk, Russia

V. Borchsh, V. Ivanchenko, E. Tcherniaev

Tomsk State University, Tomsk, Russia

P. Adzic⁴⁶, P. Cirkovic, D. Devetak, M. Dordevic, P. Milenovic, J. Milosevic, M. Stojanovic

University of Belgrade: Faculty of Physics and VINCA Institute of Nuclear Sciences, Serbia

M. Aguilar-Benitez, J. Alcaraz Maestre, A. Álvarez Fernández, I. Bachiller, M. Barrio Luna, J.A. Brochero Cifuentes, C.A. Carrillo Montoya, M. Cepeda, M. Cerrada, N. Colino, B. De La Cruz, A. Delgado Peris, C. Fernandez Bedoya, J.P. Fernández Ramos, J. Flix, M.C. Fouz, O. Gonzalez Lopez, S. Goy Lopez, J.M. Hernandez, M.I. Josa, D. Moran, Á. Navarro Tobar, A. Pérez-Calero Yzquierdo, J. Puerta Pelayo, I. Redondo, L. Romero, S. Sánchez Navas, M.S. Soares, A. Triossi, C. Willmott

Centro de Investigaciones Energéticas Medioambientales y Tecnológicas (CIEMAT), Madrid, Spain

C. Albajar, J.F. de Trocóniz, R. Reyes-Almanza

Universidad Autónoma de Madrid, Madrid, Spain

B. Alvarez Gonzalez, J. Cuevas, C. Erice, J. Fernandez Menendez, S. Folgueras, I. Gonzalez Caballero, J.R. González Fernández, E. Palencia Cortezon, V. Rodríguez Bouza, S. Sanchez Cruz

Universidad de Oviedo, Instituto Universitario de Ciencias y Tecnologías Espaciales de Asturias (ICTEA), Oviedo, Spain

I.J. Cabrillo, A. Calderon, B. Chazin Quero, J. Duarte Campderros, M. Fernandez, P.J. Fernández Manteca, A. García Alonso, G. Gomez, C. Martinez Rivero, P. Martinez Ruiz del Arbol, F. Matorras, J. Piedra Gomez, C. Prieels, T. Rodrigo, A. Ruiz-Jimeno, L. Russo⁴⁷, L. Scodellaro, N. Trevisani, I. Vila, J.M. Vizan Garcia

Instituto de Física de Cantabria (IFCA), CSIC-Universidad de Cantabria, Santander, Spain

K. Malagalage

University of Colombo, Colombo, Sri Lanka

W.G.D. Dharmaratna, N. Wickramage

University of Ruhuna, Department of Physics, Matara, Sri Lanka

D. Abbaneo, B. Akgun, E. Auffray, G. Auzinger, J. Baechler, P. Baillon, A.H. Ball, D. Barney, J. Bendavid, M. Bianco, A. Bocci, P. Bortignon, E. Bossini, C. Botta, E. Brondolin, T. Camporesi, A. Caratelli, G. Cerminara, E. Chapon, G. Cucciati, D. d'Enterria, A. Dabrowski, N. Daci, V. Daponte, A. David, O. Davignon, A. De Roeck, N. Deelen, M. Deile, M. Dobson, M. Dünser, N. Dupont, A. Elliott-Peisert, F. Fallavollita⁴⁸, D. Fasanella, S. Fiorendi, G. Franzoni, J. Fulcher, W. Funk, S. Giani, D. Gigi, A. Gilbert, K. Gill, F. Glege, M. Gruchala, M. Guilbaud, D. Gulhan, J. Hegeman, C. Heidegger, Y. Iiyama, V. Innocente, P. Janot, O. Karacheban²¹, J. Kaspar, J. Kieseler, M. Krammer¹, C. Lange, P. Lecoq, C. Lourenço, L. Malgeri, M. Mannelli, A. Massironi, F. Meijers, J.A. Merlin, S. Mersi, E. Meschi, F. Moortgat, M. Mulders, J. Ngadiuba, J. Niedziela, S. Nourbakhsh, S. Orfanelli, L. Orsini, F. Pantaleo¹⁸, L. Pape, E. Perez, M. Peruzzi, A. Petrilli, G. Petrucciani, A. Pfeiffer, M. Pierini, F.M. Pitters, D. Rabady, A. Racz, M. Rovere, H. Sakulin, C. Schäfer, C. Schwick, M. Selvaggi, A. Sharma, P. Silva, W. Snoeys, P. Sphicas⁴⁹, J. Steggemann, S. Summers, V.R. Tavolaro, D. Treille, A. Tsirou, A. Vartak, M. Verzetti, W.D. Zeuner

CERN, European Organization for Nuclear Research, Geneva, Switzerland

L. Caminada⁵⁰, K. Deiters, W. Erdmann, R. Horisberger, Q. Ingram, H.C. Kaestli, D. Kotlinski, U. Langenegger, T. Rohe, S.A. Wiederkehr

Paul Scherrer Institut, Villigen, Switzerland

M. Backhaus, P. Berger, N. Chernyavskaya, G. Dissertori, M. Dittmar, M. Donegà, C. Dorfer, T.A. Gómez Espinosa, C. Grab, D. Hits, T. Klijnsma, W. Lustermaan, R.A. Manzoni, M. Marionneau, M.T. Meinhard, F. Micheli, P. Musella, F. Nessi-Tedaldi, F. Pauss, G. Perrin, L. Perrozzi, S. Pigazzini, M.G. Ratti, M. Reichmann, C. Reissel, T. Reitenspiess, D. Ruini, D.A. Sanz Becerra, M. Schönenberger, L. Shchutska, M.L. Vesterbacka Olsson, R. Wallny, D.H. Zhu

ETH Zurich - Institute for Particle Physics and Astrophysics (IPA), Zurich, Switzerland

T.K. Aarrestad, C. Amsler⁵¹, D. Brzhechko, M.F. Canelli, A. De Cosa, R. Del Burgo, S. Donato, B. Kilminster, S. Leontsinis, V.M. Mikuni, I. Neutelings, G. Rauco, P. Robmann, D. Salerno, K. Schweiger, C. Seitz, Y. Takahashi, S. Wertz, A. Zucchetta

Universität Zürich, Zurich, Switzerland

T.H. Doan, C.M. Kuo, W. Lin, A. Roy, S.S. Yu

National Central University, Chung-Li, Taiwan

P. Chang, Y. Chao, K.F. Chen, P.H. Chen, W.-S. Hou, Y.y. Li, R.-S. Lu, E. Paganis, A. Psallidas, A. Steen

National Taiwan University (NTU), Taipei, Taiwan

B. Asavapibhop, C. Asawatangtrakuldee, N. Srimanobhas, N. Suwonjandee

Chulalongkorn University, Faculty of Science, Department of Physics, Bangkok, Thailand

A. Bat, F. Boran, A. Celik, S. Cerci⁵², S. Damarseckin⁵³, Z.S. Demiroglu, F. Dolek, C. Dozen, I. Dumanoglu, G. Gokbulut, E. Gurbinar Guler⁵⁴, Y. Guler, I. Hos⁵⁵, C. Isik, E.E. Kangal⁵⁶, O. Kara, A. Kayis Topaksu, U. Kiminsu, M. Oglakci, G. Onengut, K. Ozdemir⁵⁷, S. Ozturk⁵⁸, A. Polatoz, A.E. Simsek, U.G. Tok, S. Turkcapar, I.S. Zorbakir, C. Zorbilmez

Çukurova University, Physics Department, Science and Art Faculty, Adana, Turkey

B. Isildak⁵⁹, G. Karapinar⁶⁰, M. Yalvac

Middle East Technical University, Physics Department, Ankara, Turkey

I.O. Atakisi, E. Gülmez, M. Kaya⁶¹, O. Kaya⁶², B. Kaynak, Ö. Özçelik, S. Tekten, E.A. Yetkin⁶³

Bogazici University, Istanbul, Turkey

A. Cakir, K. Cankocak, Y. Komurcu, S. Sen⁶⁴

Istanbul Technical University, Istanbul, Turkey

S. Ozkorucuklu

Istanbul University, Istanbul, Turkey

B. Grynyov

Institute for Scintillation Materials of National Academy of Science of Ukraine, Kharkov, Ukraine

L. Levchuk

National Scientific Center, Kharkov Institute of Physics and Technology, Kharkov, Ukraine

F. Ball, E. Bhal, S. Bologna, J.J. Brooke, D. Burns⁶⁵, E. Clement, D. Cussans, H. Flacher, J. Goldstein, G.P. Heath, H.F. Heath, L. Kreczko, S. Paramesvaran, B. Penning, T. Sakuma, S. Seif El Nasr-Storey, V.J. Smith, J. Taylor, A. Titterton

University of Bristol, Bristol, United Kingdom

K.W. Bell, A. Belyaev⁶⁶, C. Brew, R.M. Brown, D. Cieri, D.J.A. Cockerill, J.A. Coughlan, K. Harder, S. Harper, J. Linacre, K. Manolopoulos, D.M. Newbold, E. Olaiya, D. Petyt, T. Reis, T. Schuh, C.H. Shepherd-Themistocleous, A. Thea, I.R. Tomalin, T. Williams, W.J. Womersley

Rutherford Appleton Laboratory, Didcot, United Kingdom

R. Bainbridge, P. Bloch, J. Borg, S. Breeze, O. Buchmuller, A. Bundock, C.H.A.H.A.L. GurpreetSingh⁶⁷, D. Colling, P. Dauncey, G. Davies, M. Della Negra, R. Di Maria, P. Everaerts, G. Hall, G. Iles, T. James, M. Komm, C. Laner, L. Lyons, A.-M. Magnan, S. Malik, A. Martelli, V. Milosevic, J. Nash⁶⁸, V. Palladino, M. Pesaresi, D.M. Raymond, A. Richards, A. Rose, E. Scott, C. Seez, A. Shtipliyski, M. Stoye, T. Strebler, A. Tapper, K. Uchida, T. Virdee¹⁸, N. Wardle, D. Winterbottom, J. Wright, A.G. Zecchinelli, S.C. Zenz

Imperial College, London, United Kingdom

J.E. Cole, P.R. Hobson, A. Khan, P. Kyberd, C.K. Mackay, A. Morton, I.D. Reid, L. Teodorescu, S. Zahid

Brunel University, Uxbridge, United Kingdom

K. Call, B. Caraway, J. Dittmann, K. Hatakeyama, C. Madrid, B. McMaster, N. Pastika, C. Smith

Baylor University, Waco, USA

R. Bartek, A. Dominguez, R. Uniyal, A.M. Vargas Hernandez

Catholic University of America, Washington, DC, USA

A. Buccilli, S.I. Cooper, C. Henderson, P. Rumerio, C. West

The University of Alabama, Tuscaloosa, USA

D. Arcaro, Z. Demiragli, D. Gastler, S. Girgis, D. Pinna, C. Richardson, J. Rohlf, D. Sperka, I. Suarez, L. Sulak, D. Zou

Boston University, Boston, USA

G. Benelli, B. Burkle, X. Coubez¹⁹, D. Cutts, Y.t. Duh, M. Hadley, J. Hakala, U. Heintz, J.M. Hogan⁶⁹, K.H.M. Kwok, E. Laird, G. Landsberg, J. Lee, Z. Mao, M. Narain, S. Sagir⁷⁰, R. Syarif, E. Usai, D. Yu, W. Zhang

Brown University, Providence, USA

R. Band, C. Brainerd, R. Breedon, M. Calderon De La Barca Sanchez, M. Chertok, J. Conway, R. Conway, P.T. Cox, R. Erbacher, C. Flores, G. Funk, F. Jensen, W. Ko, O. Kukral, R. Lander, M. Mulhearn, D. Pellett, J. Pilot, M. Shi, D. Taylor, K. Tos, M. Tripathi, Z. Wang, F. Zhang

University of California, Davis, Davis, USA

M. Bachtis, C. Bravo, R. Cousins, A. Dasgupta, A. Florent, J. Hauser, M. Ignatenko, N. Mccoll, W.A. Nash, S. Regnard, D. Saltzberg, C. Schnaible, B. Stone, V. Valuev

University of California, Los Angeles, USA

K. Burt, Y. Chen, R. Clare, J.W. Gary, S.M.A. Ghiasi Shirazi, G. Hanson, G. Karapostoli, E. Kennedy, O.R. Long, M. Olmedo Negrete, M.I. Paneva, W. Si, L. Wang, S. Wimpenny, B.R. Yates, Y. Zhang

University of California, Riverside, Riverside, USA

J.G. Branson, P. Chang, S. Cittolin, M. Derdzinski, R. Gerosa, D. Gilbert, B. Hashemi, D. Klein, V. Krutelyov, J. Letts, M. Masciovecchio, S. May, S. Padhi, M. Pieri, V. Sharma, M. Tadel, F. Würthwein, A. Yagil, G. Zevi Della Porta

University of California, San Diego, La Jolla, USA

N. Amin, R. Bhandari, C. Campagnari, M. Citron, V. Dutta, M. Franco Sevilla, L. Gouskos, J. Incandela, B. Marsh, H. Mei, A. Ovcharova, H. Qu, J. Richman, U. Sarica, D. Stuart, S. Wang

University of California, Santa Barbara - Department of Physics, Santa Barbara, USA

D. Anderson, A. Bornheim, O. Cerri, I. Dutta, J.M. Lawhorn, N. Lu, J. Mao, H.B. Newman, T.Q. Nguyen, J. Pata, M. Spiropulu, J.R. Vlimant, S. Xie, Z. Zhang, R.Y. Zhu

California Institute of Technology, Pasadena, USA

M.B. Andrews, T. Ferguson, T. Mudholkar, M. Paulini, J. Russ, M. Sun, I. Vorobiev, M. Weinberg

Carnegie Mellon University, Pittsburgh, USA

J.P. Cumalat, W.T. Ford, A. Johnson, E. MacDonald, T. Mulholland, R. Patel, A. Perloff, K. Stenson, K.A. Ulmer, S.R. Wagner

University of Colorado Boulder, Boulder, USA

J. Alexander, J. Chaves, Y. Cheng, J. Chu, A. Datta, A. Frankenthal, K. Mcdermott, J.R. Patterson, D. Quach, A. Rinkevicius⁷¹, A. Ryd, S.M. Tan, Z. Tao, J. Thom, P. Wittich, M. Zientek

Cornell University, Ithaca, USA

S. Abdullin, M. Albrow, M. Alyari, G. Apollinari, A. Apresyan, A. Apyan, S. Banerjee, L.A.T. Bauerdick, A. Beretvas, J. Berryhill, P.C. Bhat, K. Burkett, J.N. Butler, A. Canepa, G.B. Cerati, H.W.K. Cheung, F. Chlebana, M. Cremonesi, J. Duarte, V.D. Elvira, J. Freeman, Z. Gecse, E. Gottschalk, L. Gray, D. Green, S. Grünendahl, O. Gutsche, Hall AllisonReinsvold, J. Hanlon, R.M. Harris, S. Hasegawa, R. Heller, J. Hirschauer, B. Jayatilaka, S. Jindariani, M. Johnson, U. Joshi, B. Klima, M.J. Kortelainen, B. Kreis, S. Lammel, J. Lewis, D. Lincoln, R. Lipton, M. Liu, T. Liu, J. Lykken, K. Maeshima, J.M. Marraffino, D. Mason, P. McBride, P. Merkel, S. Mrenna, S. Nahn, V. O'Dell, V. Papadimitriou, K. Pedro, C. Pena, G. Rakness, F. Ravera, L. Ristori, B. Schneider, E. Sexton-Kennedy, N. Smith, A. Soha, W.J. Spalding, L. Spiegel, S. Stoynev, J. Strait, N. Strobbe, L. Taylor, S. Tkaczyk, N.V. Tran, L. Uplegger, E.W. Vaandering, C. Vernieri, R. Vidal, M. Wang, H.A. Weber

Fermi National Accelerator Laboratory, Batavia, USA

D. Acosta, P. Avery, D. Bourilkov, A. Brinkerhoff, L. Cadamuro, A. Carnes, V. Cherepanov, D. Curry, F. Errico, R.D. Field, S.V. Gleyzer, B.M. Joshi, M. Kim, J. Konigsberg, A. Korytov, K.H. Lo, P. Ma, K. Matchev, N. Menendez, G. Mitselmakher, D. Rosenzweig, K. Shi, J. Wang, S. Wang, X. Zuo

University of Florida, Gainesville, USA

Y.R. Joshi

Florida International University, Miami, USA

T. Adams, A. Askew, S. Hagopian, V. Hagopian, K.F. Johnson, R. Khurana, T. Kolberg, G. Martinez, T. Perry, H. Prosper, C. Schiber, R. Yohay, J. Zhang

Florida State University, Tallahassee, USA

M.M. Baarmand, M. Hohlmann, D. Noonan, M. Rahmani, M. Saunders, F. Yumiceva

Florida Institute of Technology, Melbourne, USA

M.R. Adams, L. Apanasevich, D. Berry, R.R. Betts, R. Cavanaugh, X. Chen, S. Dittmer, O. Evdokimov, C.E. Gerber, D.A. Hangal, D.J. Hofman, K. Jung, C. Mills, T. Roy, M.B. Tonjes, N. Varelas, J. Viinikainen, H. Wang, X. Wang, Z. Wu

University of Illinois at Chicago (UIC), Chicago, USA

M. Alhousseini, B. Bilki⁵⁴, W. Clarida, K. Dilsiz⁷², S. Durgut, R.P. Gandrajula, M. Haytmyradov, V. Khristenko, O.K. Köseyan, J.-P. Merlo, A. Mestvirishvili⁷³, A. Moeller, J. Nachtman, H. Ogul⁷⁴, Y. Onel, F. Ozok⁷⁵, A. Penzo, C. Snyder, E. Tiras, J. Wetzel

The University of Iowa, Iowa City, USA

B. Blumenfeld, A. Cocoros, N. Eminizer, D. Fehling, L. Feng, A.V. Gritsan, W.T. Hung, P. Maksimovic, J. Roskes, M. Swartz

Johns Hopkins University, Baltimore, USA

C. Baldenegro Barrera, P. Baringer, A. Bean, S. Boren, J. Bowen, A. Bylinkin, T. Isidori, S. Khalil, J. King, G. Krintiras, A. Kropivnitskaya, C. Lindsey, D. Majumder, W. Mcbrayer, N. Minafra, M. Murray, C. Rogan, C. Royon, S. Sanders, E. Schmitz, J.D. Tapia Takaki, Q. Wang, J. Williams, G. Wilson

The University of Kansas, Lawrence, USA

S. Duric, A. Ivanov, K. Kaadze, D. Kim, Y. Maravin, D.R. Mendis, T. Mitchell, A. Modak, A. Mohammadi

Kansas State University, Manhattan, USA

F. Rebassoo, D. Wright

Lawrence Livermore National Laboratory, Livermore, USA

A. Baden, O. Baron, A. Belloni, S.C. Eno, Y. Feng, N.J. Hadley, S. Jabeen, G.Y. Jeng, R.G. Kellogg, J. Kunkle, A.C. Mignerey, S. Nabili, F. Ricci-Tam, M. Seidel, Y.H. Shin, A. Skuja, S.C. Tonwar, K. Wong

University of Maryland, College Park, USA

D. Abercrombie, B. Allen, A. Baty, R. Bi, S. Brandt, W. Busza, I.A. Cali, M. D'Alfonso, G. Gomez Ceballos, M. Goncharov, P. Harris, D. Hsu, M. Hu, M. Klute, D. Kovalskyi, Y.-J. Lee, P.D. Luckey, B. Maier, A.C. Marini, C. Mcginn, C. Mironov, S. Narayanan, X. Niu, C. Paus, D. Rankin, C. Roland, G. Roland, Z. Shi, G.S.F. Stephens, K. Sumorok, K. Tatar, D. Velicanu, J. Wang, T.W. Wang, B. Wyslouch

Massachusetts Institute of Technology, Cambridge, USA

A.C. Benvenuti[†], R.M. Chatterjee, A. Evans, S. Guts, P. Hansen, J. Hiltbrand, Y. Kubota, Z. Lesko, J. Mans, R. Rusack, M.A. Wadud

University of Minnesota, Minneapolis, USA

J.G. Acosta, S. Oliveros

University of Mississippi, Oxford, USA

K. Bloom, D.R. Claes, C. Fangmeier, L. Finco, F. Golf, R. Gonzalez Suarez, R. Kamalieddin, I. Kravchenko, J.E. Siado, G.R. Snow, B. Stieger, W. Tabb

University of Nebraska-Lincoln, Lincoln, USA

G. Agarwal, C. Harrington, I. Iashvili, A. Kharchilava, C. McLean, D. Nguyen, A. Parker, J. Pekkanen, S. Rappoccio, B. Roozbahani

State University of New York at Buffalo, Buffalo, USA

G. Alverson, E. Barberis, C. Freer, Y. Haddad, A. Hortiangtham, G. Madigan, D.M. Morse, T. Orimoto, L. Skinnari, A. Tishelman-Charny, T. Wamorkar, B. Wang, A. Wisecarver, D. Wood

Northeastern University, Boston, USA

S. Bhattacharya, J. Bueghly, T. Gunter, K.A. Hahn, N. Odell, M.H. Schmitt, K. Sung, M. Trovato, M. Velasco

Northwestern University, Evanston, USA

R. Bucci, N. Dev, R. Goldouzian, M. Hildreth, K. Hurtado Anampa, C. Jessop, D.J. Karmgard, K. Lannon, W. Li, N. Loukas, N. Marinelli, I. Mcalister, F. Meng, C. Mueller, Y. Musienko³⁷, M. Planer, R. Ruchti, P. Siddireddy, G. Smith, S. Taroni, M. Wayne, A. Wightman, M. Wolf, A. Woodard

University of Notre Dame, Notre Dame, USA

J. Alimena, B. Bylsma, L.S. Durkin, S. Flowers, B. Francis, C. Hill, W. Ji, A. Lefeld, T.Y. Ling, B.L. Winer

The Ohio State University, Columbus, USA

S. Cooperstein, G. Dezoort, P. Elmer, J. Hardenbrook, N. Haubrich, S. Higginbotham, A. Kalogeropoulos, S. Kwan, D. Lange, M.T. Lucchini, J. Luo, D. Marlow, K. Mei, I. Ojalvo, J. Olsen, C. Palmer, P. Piroué, J. Salfeld-Nebgen, D. Stickland, C. Tully, Z. Wang

Princeton University, Princeton, USA

S. Malik, S. Norberg

University of Puerto Rico, Mayaguez, USA

A. Barker, V.E. Barnes, S. Das, L. Gutay, M. Jones, A.W. Jung, A. Khatiwada, B. Mahakud, D.H. Miller, G. Negro, N. Neumeister, C.C. Peng, S. Piperov, H. Qiu, J.F. Schulte, J. Sun, F. Wang, R. Xiao, W. Xie

Purdue University, West Lafayette, USA

T. Cheng, J. Dolen, N. Parashar

Purdue University Northwest, Hammond, USA

K.M. Ecklund, S. Freed, F.J.M. Geurts, M. Kilpatrick, Arun Kumar, W. Li, B.P. Padley, R. Redjimi, J. Roberts, J. Rorie, W. Shi, A.G. Stahl Leiton, Z. Tu, A. Zhang

Rice University, Houston, USA

A. Bodek, P. de Barbaro, R. Demina, J.L. Dulemba, C. Fallon, T. Ferbel, M. Galanti, A. Garcia-Bellido, O. Hindrichs, A. Khukhunaishvili, E. Ranken, P. Tan, R. Taus

University of Rochester, Rochester, USA

B. Chiarito, J.P. Chou, A. Gandrakota, Y. Gershtein, E. Halkiadakis, A. Hart, M. Heindl, E. Hughes, S. Kaplan, S. Kyriacou, I. Laflotte, A. Lath, R. Montalvo, K. Nash, M. Osherson, H. Saka, S. Salur, S. Schnetzer, S. Somalwar, R. Stone, S. Thomas

Rutgers, The State University of New Jersey, Piscataway, USA

H. Acharya, A.G. Delannoy, G. Riley, S. Spanier

University of Tennessee, Knoxville, USA

O. Bouhali⁷⁶, M. Dalchenko, M. De Mattia, A. Delgado, S. Dildick, R. Eusebi, J. Gilmore, T. Huang, T. Kamon⁷⁷, S. Luo, D. Marley, R. Mueller, D. Overton, L. Perniè, D. Rathjens, A. Safonov

Texas A&M University, College Station, USA

N. Akchurin, J. Damgov, F. De Guio, S. Kunori, K. Lamichhane, S.W. Lee, T. Mengke, S. Muthumuni, T. Peltola, S. Undleeb, I. Volobouev, Z. Wang, A. Whitbeck

Texas Tech University, Lubbock, USA

S. Greene, A. Gurrola, R. Janjam, W. Johns, C. Maguire, A. Melo, H. Ni, K. Padeken, F. Romeo, P. Sheldon, S. Tuo, J. Velkovska, M. Verweij

Vanderbilt University, Nashville, USA

M.W. Arenton, P. Barria, B. Cox, G. Cummings, R. Hirosky, M. Joyce, A. Ledovskoy, C. Neu, B. Tannenwald, Y. Wang, E. Wolfe, F. Xia

University of Virginia, Charlottesville, USA

R. Harr, P.E. Karchin, N. Poudyal, J. Sturdy, P. Thapa

Wayne State University, Detroit, USA

T. Bose, J. Buchanan, C. Caillol, D. Carlsmith, S. Dasu, I. De Bruyn, L. Dodd, F. Fiori, C. Galloni, B. Gomber⁷⁸, H. He, M. Herndon, A. Hervé, U. Hussain, P. Klabbers, A. Lanaro, A. Loeliger, K. Long, R. Loveless, J. Madhusudanan Sreekala, T. Ruggles, A. Savin, V. Sharma, W.H. Smith, D. Teague, S. Trembath-reichert, N. Woods

University of Wisconsin - Madison, Madison, WI, USA

[†] Deceased.

¹ Also at Vienna University of Technology, Vienna, Austria.

² Also at IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette, France.

³ Also at Universidade Estadual de Campinas, Campinas, Brazil.

⁴ Also at Federal University of Rio Grande do Sul, Porto Alegre, Brazil.

⁵ Also at UFMS, Nova Andradina, Brazil.

⁶ Also at Universidade Federal de Pelotas, Pelotas, Brazil.

⁷ Also at Université Libre de Bruxelles, Bruxelles, Belgium.

⁸ Also at University of Chinese Academy of Sciences, Beijing, China.

⁹ Also at Institute for Theoretical and Experimental Physics named by A.I. Alikhanov of NRC 'Kurchatov Institute', Moscow, Russia.

¹⁰ Also at Joint Institute for Nuclear Research, Dubna, Russia.

¹¹ Also at Suez University, Suez, Egypt.

¹² Now at British University in Egypt, Cairo, Egypt.

¹³ Also at Purdue University, West Lafayette, USA.

¹⁴ Also at Université de Haute Alsace, Mulhouse, France.

¹⁵ Also at Tbilisi State University, Tbilisi, Georgia.

¹⁶ Also at Ilia State University, Tbilisi, Georgia.

¹⁷ Also at Erzincan Binali Yildirim University, Erzincan, Turkey.

¹⁸ Also at CERN, European Organization for Nuclear Research, Geneva, Switzerland.

¹⁹ Also at RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany.

²⁰ Also at University of Hamburg, Hamburg, Germany.

²¹ Also at Brandenburg University of Technology, Cottbus, Germany.

²² Also at Institute of Physics, University of Debrecen, Debrecen, Hungary.

²³ Also at Institute of Nuclear Research ATOMKI, Debrecen, Hungary.

²⁴ Also at IIT Bhubaneswar, Bhubaneswar, India.

²⁵ Also at Institute of Physics, Bhubaneswar, India.

²⁶ Also at Shoolini University, Solan, India.

²⁷ Also at University of Visva-Bharati, Santiniketan, India.

²⁸ Also at Isfahan University of Technology, Isfahan, Iran.

²⁹ Now at INFN Sezione di Bari ^a, Università di Bari ^b, Politecnico di Bari ^c, Bari, Italy.

³⁰ Also at Italian National Agency for New Technologies, Energy and Sustainable Economic Development, Bologna, Italy.

³¹ Also at Centro Siciliano di Fisica Nucleare e di Struttura Della Materia, Catania, Italy.

- ³² Also at Scuola Normale e Sezione dell'INFN, Pisa, Italy.
- ³³ Also at Riga Technical University, Riga, Latvia, Riga, Latvia.
- ³⁴ Also at Malaysian Nuclear Agency, MOSTI, Kajang, Malaysia.
- ³⁵ Also at Consejo Nacional de Ciencia y Tecnología, Mexico City, Mexico.
- ³⁶ Also at Warsaw University of Technology, Institute of Electronic Systems, Warsaw, Poland.
- ³⁷ Also at Institute for Nuclear Research, Moscow, Russia.
- ³⁸ Now at National Research Nuclear University 'Moscow Engineering Physics Institute' (MEPhI), Moscow, Russia.
- ³⁹ Also at Institute of Nuclear Physics of the Uzbekistan Academy of Sciences, Tashkent, Uzbekistan.
- ⁴⁰ Also at St. Petersburg State Polytechnical University, St. Petersburg, Russia.
- ⁴¹ Also at University of Florida, Gainesville, USA.
- ⁴² Also at Imperial College, London, United Kingdom.
- ⁴³ Also at P.N. Lebedev Physical Institute, Moscow, Russia.
- ⁴⁴ Also at California Institute of Technology, Pasadena, USA.
- ⁴⁵ Also at Budker Institute of Nuclear Physics, Novosibirsk, Russia.
- ⁴⁶ Also at Faculty of Physics, University of Belgrade, Belgrade, Serbia.
- ⁴⁷ Also at Università degli Studi di Siena, Siena, Italy.
- ⁴⁸ Also at INFN Sezione di Pavia ^a, Università di Pavia ^b, Pavia, Italy, Pavia, Italy.
- ⁴⁹ Also at National and Kapodistrian University of Athens, Athens, Greece.
- ⁵⁰ Also at Universität Zürich, Zurich, Switzerland.
- ⁵¹ Also at Stefan Meyer Institute for Subatomic Physics, Vienna, Austria, Vienna, Austria.
- ⁵² Also at Adiyaman University, Adiyaman, Turkey.
- ⁵³ Also at Şırnak University, Sirnak, Turkey.
- ⁵⁴ Also at Beykent University, Istanbul, Turkey, Istanbul, Turkey.
- ⁵⁵ Also at Istanbul Aydin University, Application and Research Center for Advanced Studies (App. & Res. Cent. for Advanced Studies), Istanbul, Turkey.
- ⁵⁶ Also at Mersin University, Mersin, Turkey.
- ⁵⁷ Also at Piri Reis University, Istanbul, Turkey.
- ⁵⁸ Also at Gaziosmanpasa University, Tokat, Turkey.
- ⁵⁹ Also at Ozyegin University, Istanbul, Turkey.
- ⁶⁰ Also at Izmir Institute of Technology, Izmir, Turkey.
- ⁶¹ Also at Marmara University, Istanbul, Turkey.
- ⁶² Also at Kafkas University, Kars, Turkey.
- ⁶³ Also at Istanbul Bilgi University, Istanbul, Turkey.
- ⁶⁴ Also at Hacettepe University, Ankara, Turkey.
- ⁶⁵ Also at Vrije Universiteit Brussel, Brussel, Belgium.
- ⁶⁶ Also at School of Physics and Astronomy, University of Southampton, Southampton, United Kingdom.
- ⁶⁷ Also at IPPP Durham University, Durham, United Kingdom.
- ⁶⁸ Also at Monash University, Faculty of Science, Clayton, Australia.
- ⁶⁹ Also at Bethel University, St. Paul, Minneapolis, USA, St. Paul, USA.
- ⁷⁰ Also at Karamanoğlu Mehmetbey University, Karaman, Turkey.
- ⁷¹ Also at Vilnius University, Vilnius, Lithuania.
- ⁷² Also at Bingol University, Bingol, Turkey.
- ⁷³ Also at Georgian Technical University, Tbilisi, Georgia.
- ⁷⁴ Also at Sinop University, Sinop, Turkey.
- ⁷⁵ Also at Mimar Sinan University, Istanbul, Istanbul, Turkey.
- ⁷⁶ Also at Texas A&M University at Qatar, Doha, Qatar.
- ⁷⁷ Also at Kyungpook National University, Daegu, Korea, Daegu, Korea.
- ⁷⁸ Also at University of Hyderabad, Hyderabad, India.