



Letter

Measurement of the transverse-momentum fraction of strange hadrons from jet-like correlation structures in pp collisions at $\sqrt{s} = 13$ TeVThe ALICE Collaboration¹

ARTICLE INFO

Editor: Dr. M. Doser

Keywords:

Strangeness

Hadronization

Jet fragmentation

Transverse-momentum fraction

Two-particle correlations

ABSTRACT

The first measurements of the average transverse-momentum fraction ($\langle z \rangle$) as a function of transverse momentum (p_T) for strange baryons (Λ and $\bar{\Lambda}$) and strange mesons (K_S^0), produced in mini-jets defined through angular correlations in pp collisions at $\sqrt{s} = 13$ TeV, are reported by the ALICE Collaboration at the LHC. The observable is obtained using a novel method, where the angular correlation between the strange hadrons and inclusive charged hadrons is weighted by the p_T of correlated particles at small angular distance. As a function of strange particles' p_T , the results reveal a flat trend for strange mesons and a decreasing trend for strange baryons in the measured p_T region, indicating distinct hadronization mechanisms for K_S^0 and Λ ($\bar{\Lambda}$). The measurements are compared to Monte Carlo models, namely PYTHIA 8 (with both Monash and Color Rope tunes) and the AMPT (A Multi-Phase Transport) model with string melting. None of these models provides a satisfactory description of the $\langle z \rangle$ distributions at low and intermediate p_T .

1. Introduction

Hadronic collisions at the LHC (pp, p-Pb, and Pb-Pb) provide the opportunity to study particle production across a wide range of charged-particle multiplicities and corresponding initial energy densities [1–6]. One of the most intriguing discoveries at the LHC is that phenomena traditionally associated with heavy-ion collisions, and attributed to the formation of quark–gluon plasma (QGP), exhibit a continuous onset as a function of multiplicity across all collision systems. These features include collective fluid-like behavior [7–9], strangeness enhancement, and baryon-to-meson yield enhancement in the intermediate p_T range [10–12].

In the so-called “small systems”, such as pp and p-Pb collisions, the size of the interaction region is approximately one femtometer. At LHC energies, partonic systems produced in multiparton interactions (MPIs) are expected to overlap within this small region. Given that the range of the strong force is of comparable size, interactions among strings produced nearby at early times may induce effects on the final-state kinematics and hadronization beyond those expected from a simple incoherent superposition of MPIs. In Pb-Pb collisions, final-state interactions are well established and give rise to strong collective effects.

The presence of collective effects in small systems raises the intriguing possibility that droplets of QGP may already be forming in such systems. Furthermore, understanding how MPI-driven effects manifest themselves in small systems can shed light on collective phenomena ob-

served in Pb-Pb collisions. Addressing these issues is important for developing a common understanding of particle production mechanisms across small and large collision systems, and for identifying the conditions that give rise to QGP-like behavior.

The enhancement of strange baryon-to-meson hadron yield ratios (Λ/K_S^0 , Ξ/K_S^0 , Ω/K_S^0) in the intermediate p_T region, observed across all collision systems and progressively increasing with multiplicity, has been attributed to the interplay between radial flow and parton recombination [13–15]. To further investigate the origin of this effect, ALICE has measured these ratios within charged-particle jets with p_T above 10 GeV/c and for particles in the underlying event of these jets [16,17]. The results show that, within the precision of the measurements, the enhancement is small or absent within jets, while in the underlying event, it matches the inclusive results. However, a limitation of this approach is that the production of a particle outside a jet with a minimum energy threshold does not necessarily indicate it was not produced in a hard-scattering process. For example, the increase of the ratios between particle yields measured at $\sqrt{s} = 13$ TeV and 7 TeV with rising p_T suggests that hard scatterings begin to dominate particle production already for $p_T > 2$ GeV/c [18]. This raises the need for a complementary observable to study particle production mechanisms. Assuming that particle production is predominantly governed by parton fragmentation, the fraction of the parton's transverse momentum carried by the final-state particle (p_T/p_T^{jet}) provides complementary insight into the fragmentation process and is therefore a more informative observable than the hadron p_T alone.

Contact: alice-publications@cern.ch.

¹ Authors are listed at the end of the paper.

In the low- and intermediate- p_T region ($p_T < 5 \text{ GeV}/c$), the distinction between jets and soft bulk particle production becomes ambiguous, effectively precluding reliable event-by-event jet reconstruction. Therefore, the present analysis does not rely on associating strange particles with reconstructed jet cones. Instead, the average momentum fraction ($\langle z \rangle$) of Λ ($\bar{\Lambda}$) and K_S^0 hadrons in mini-jets is measured. The mini-jets are defined via $\Delta\phi$ – $\Delta\eta$ angular correlations between these strange hadrons and primary charged particles, where $\Delta\phi$ and $\Delta\eta$ denote the differences in azimuthal angle and pseudorapidity, respectively. The angular differential distribution is weighted by the p_T of the associated particles in order to estimate the p_T of the originating partons.

Jet-like structures manifest themselves as a short-range near-side peak centered at $\Delta\phi = \Delta\eta = 0$ in the angular correlation distribution. After normalizing the p_T -weighted angular correlation distribution by the number of trigger particles and subtracting the uncorrelated background, the integral below the short-range near-side peak measures the summed p_T correlated with the strange particles ($\sum^{\text{NS}} p_T$). A proxy for the original parton p_T can then be obtained by adding the p_T of the trigger particle to $\sum^{\text{NS}} p_T$. This approach provides essential complementary information for the understanding of particle production dynamics and the role of parton fragmentation in the intermediate p_T region.

2. Experimental setup and data samples

The ALICE detector [19] has been optimized for the study of heavy-ion collisions at the LHC. The main objective of ALICE is to explore the physics of strongly-interacting matter in proton-proton collisions and the QGP produced in the extreme conditions of temperature and density created in nucleus-nucleus collisions. The ALICE detector and its performance during Run 2 of the LHC are described in Refs. [20,21]. Its setup consists of central-barrel detectors covering the pseudorapidity range $|\eta| < 0.9$, forward detectors, and the magnet systems.

The central-barrel detectors – Inner Tracking System (ITS), Time Projection Chamber (TPC), Transition Radiation Detector (TRD), Time Of Flight (TOF), Photon Spectrometer (PHOS), Electromagnetic Calorimeter (EMCal), and High Momentum Particle Identification Detector (HMPID) – are surrounded by the L3 solenoid, which provides a magnetic field of up to 0.5 T in strength along the beam direction. These detectors are optimized for tracking and particle identification. The Photon Multiplicity Detector (PMD), Forward Multiplicity Detector (FMD), V0, T0, and Zero Degree Calorimeter (ZDC) – the forward detectors – provide triggering and information on global event properties. The measurements presented in this paper use the information from V0 [21], ITS [22], TPC [23], and TOF [24,25], which are described in more detail in the following paragraphs.

The V0 is used to provide event triggering information. It is composed of two scintillator arrays, V0A and V0C, placed around the beam pipe on each side of the interaction point, covering the pseudorapidity intervals $2.8 < \eta < 5.1$ and $-3.7 < \eta < -1.7$, respectively.

The ITS is the detector closest to the interaction region. The ITS used in Run 2 was made of six cylindrical layers of silicon detectors. The two innermost layers comprise the Silicon Pixel Detector (SPD), which mainly provides hit position information for vertexing and tracking. The intermediate two layers form the Silicon Drift Detectors (SDD), and the outermost two layers form the Silicon Strip Detectors (SSD). They mainly provide tracking information and particle identification via the measurement of ionization energy loss.

The TPC is the main tracking detector of ALICE. It is a cylindrical detector with a volume of about 90 m^3 , designed to be filled with a gas mixture of 90% Ne and 10% CO_2 . By measuring ionization energy loss, the TPC is also used for particle identification.

The TOF, placed outside of the TPC, is made of Multigap Resistive Plate Chambers (MRPC). Complementary to the ITS and the TPC, the TOF provides particle identification at the intermediate momentum range (from 0.5 GeV/c to 3–4 GeV/c) [25] via time-of-flight measurements.

The present analysis is performed with pp collision data at $\sqrt{s} = 13 \text{ TeV}$, which were collected by ALICE during the LHC Run 2 from 2016 to 2018. The events used for the analysis are those satisfying the minimum bias (MB) trigger condition requiring at least one hit in both V0A and V0C. Pileup events are rejected using vertex and tracking information. The events are accepted only if the reconstructed primary vertex position in the beam direction is located in the range $\pm 10 \text{ cm}$ from the ALICE nominal interaction point. Approximately 1.7×10^9 events pass these selection criteria.

Monte Carlo (MC) data samples have been generated using PYTHIA 8 with the Monash tune [26,27] as the primary event generator for pp collisions. The detector response is simulated by transporting the primary events through a model of the ALICE detector using GEANT 3 [28]. These MC samples are used to evaluate the tracking efficiency and the secondary contamination contribution.

3. Analysis procedure

The goal of the present analysis is to study an observable that can approximate the transverse-momentum fraction z of strange particles in low-energy jets. For reconstructed jets, z is defined as $z = p_T^s/p_T^{\text{jet}}$, where p_T^s is the transverse momentum of the strange particle and p_T^{jet} the transverse momentum of the jet. Very low-energy jets ($p_T^{\text{jet}} \ll 5 \text{ GeV}/c$) cannot be reconstructed event-by-event over the fluctuating background of the underlying event. This is particularly true in high-multiplicity pp collisions. An alternative method to assess the fragmentation properties of low-energy jets is the study of $\Delta\phi$ – $\Delta\eta$ angular correlations between particles, given that jet-like correlations produce a distinct peak in the correlation function centered at $\Delta\phi = \Delta\eta = 0$, the so-called short-range near-side peak. In the angular correlation study, $\Delta\phi$ and $\Delta\eta$ are defined as the azimuthal angle and pseudorapidity difference between the trigger and the associated particles, respectively. In this measurement, the strange particles are used as trigger particles, and all primary charged particles are used as associated particles. Primary charged particles are defined as charged particles with a mean proper lifetime τ larger than 1 cm/c, which are produced either promptly at the primary interaction vertex or from decays of particles with $\tau < 1 \text{ cm}/c$ restricted to decay chains leading to the interaction [29]. For the present analysis, the long-lived charged-strange baryons ($\Sigma^\pm$, Ξ^- , Ω^- , and their antiparticles) are excluded from the primary charged-particle definition in order to avoid biases, as their reconstruction requires dedicated topological selections beyond those used for primary charged particles.

3.1. Particle reconstruction

The strange particles, K_S^0 , Λ , and $\bar{\Lambda}$, are reconstructed through their decay daughters in the central barrel region via the $K_S^0 \rightarrow \pi^+ + \pi^-$, $\Lambda \rightarrow p + \pi^-$, and $\bar{\Lambda} \rightarrow \bar{p} + \pi^+$ decay channels, with the branching ratios of $69.20 \pm 0.05\%$, $64.1 \pm 0.5\%$, and $64.1 \pm 0.5\%$, respectively [30]. The reconstruction is performed in the kinematic range $0.6 < p_T < 20.0 \text{ GeV}/c$ and $|\eta| < 0.75$. They are selected based on their characteristic V-shaped decay topology. Being electrically neutral, these strange particles are also referred to as V^0 particles. The selection criteria include the V^0 decay radius in the transverse plane, the distance of the closest approach (DCA) between the daughter tracks, the cosine of the pointing angle (CPA), which is the angle between the line connecting the PV to the decay vertex and the direction of the momentum vector. Additional selection criteria are applied to the daughter tracks of the V^0 , including pseudorapidity requirements, the DCA of each daughter to the primary vertex (PV), and PID and track-quality selections based on the TPC information. These criteria follow those presented in an earlier ALICE publication [17] and are summarized in Table 1. To suppress misidentification between K_S^0 and Λ ($\bar{\Lambda}$) candidates, the invariant mass is also computed under the competing mass hypothesis, and a selection criterion is applied accordingly. In addition, in order to reject strange

Table 1
V⁰ candidate selection criteria.

V ⁰ selection	K _s ⁰	Λ ($\bar{\Lambda}$)
$ \eta_{V^0} $	< 0.75	< 0.75
V ⁰ 2D decay radius (cm)	> 0.5	> 0.5
Proper lifetime (cm/c)	< 20	< 30
DCA between V ⁰ daughter tracks (Nσ)	< 1	< 1
CPA of V ⁰	> 0.97	> 0.995
Competing mass (GeV/c ²)	> 0.005	> 0.010
V ⁰ daughters selection		
$ \eta_{\text{trk}} $	< 0.8	
V ⁰ daughter track DCA to PV (cm)	> 0.06	
TPC dE/dx (Nσ)	< 5	
TPC refit flag	true	
Number of TPC crossed rows	> 70	
TPC crossed rows / findable ratio	> 0.8	

particle candidates from out-of-bunch pileup, at least one daughter track is required to satisfy the ITS refit condition, exploiting the superior time resolution of the ITS.

Primary charged particles are reconstructed in the kinematic range $0.2 < p_T < 20$ GeV/c and $|\eta| < 0.8$ using the ITS and the TPC. Contamination from secondary particles is reduced by applying the selection criteria described in Ref. [31]. Selected primary charged tracks are required to have at least 70 crossed pad rows in the TPC (out of a maximum of 159), and a ratio of crossed rows to the number of findable clusters of at least 0.8. Only tracks with a fraction of shared clusters with other tracks smaller than 0.4 are accepted. The DCA to the primary vertex in the transverse plane (xy-plane) and the beam direction (z-direction) are required to be within 2.4 cm and 3.2 cm, respectively. The tracks are also required to have a good tracking fit quality in both the TPC and the ITS, characterized by goodness-of-fit values χ^2 per cluster smaller than 4 and 36 in the TPC and the ITS, respectively. Only tracks with at least one hit in the two innermost layers of the ITS are selected. Tracks originating from kink topologies, where a charged particle decays into another charged particle and a neutral particle, are rejected.

3.2. p_T -weighted two-particle correlation function

A novel p_T -weighted two-particle correlation method is employed to obtain the average $\langle z(p_T^s) \rangle$ of strange particles in any given p_T^s interval. The corresponding correlation function is defined as:

$$C_{p_T\text{-weighted}}(\Delta\eta, \Delta\phi) = \frac{1}{N_{\text{trig}}} \frac{d^2 \sum p_T}{d\Delta\phi d\Delta\eta}, \quad (1)$$

where N_{trig} is the number of trigger strange particles. The term $\frac{d^2 \sum p_T}{d\Delta\phi d\Delta\eta}$ represents the summed p_T of all the associated particles in any given $\Delta\phi - \Delta\eta$ interval. Both N_{trig} and $\frac{d^2 \sum p_T}{d\Delta\phi d\Delta\eta}$ are summed over all selected events.

The correlation function is corrected for acceptance, efficiency, and uncorrelated background, which will be described in detail in the next section. Finally, the summed p_T of the associated particles per trigger particle $\langle \sum^{\text{NS}} p_T \rangle$ is calculated by integrating the correlation function, and the average $\langle z(p_T^s) \rangle$ is computed as

$$\langle z(p_T^s) \rangle = \frac{p_T^s}{(\sum^{\text{NS}} p_T) + p_T^s}. \quad (2)$$

As will be explained in the following paragraphs in more detail, $\langle z \rangle$ is obtained as a function of the strange particle p_T^s by constructing the correlation function and applying the corrections independently in each p_T^s interval. Due to the steeply falling trigger-particle p_T^s spectra and the finite width of the p_T^s intervals, especially at high transverse momenta, the p_T^s in Eq. (2) is calculated as the average trans-

verse momentum in each p_T interval instead of the value of the interval center.

It is important to emphasize the differences between the present approach and a standard jet analysis. In the latter, jets are reconstructed event by event and the jet p_T is required to be either above a threshold value imposed by the jet reconstruction efficiency or within a specific range. The z is thus obtained event by event, and the resulting z distribution might be affected by the jet p_T selection. In contrast, the present analysis does not rely on explicit jet reconstruction and therefore does not impose any jet- p_T selection. Since the jets are not reconstructed individually, an event-by-event determination of z is not possible. Instead, the observable $\langle z \rangle$ is obtained as an event-averaged quantity, calculated as $p_T^s / \langle p_T^{\text{jet}} \rangle$ in intervals of the trigger-particle p_T^s .

Let $D(z)$ denote the fragmentation function, the probability for a particle to carry a fraction z of the originating parton momentum. Assuming that the parton transverse momentum spectrum falls as $(1/p_T^{\text{parton}})^n$, where typical values of n are between 5 and 6 at high p_T [2]. At fixed trigger-particle transverse momentum p_T , and assuming that z is measured on an event-by-event basis, the probability density of z is proportional to $D(z)z^{n-2}$ [32,33]. The factor z^{n-2} biases the distribution towards larger values of z compared to what would be obtained at fixed p_T^{parton} . This effect is known as the leading particle bias or trigger bias [32,33]. In the present analysis, one instead averages over $\sum^{\text{NS}} p_T$ which is proportional to p_T^s/z . As a result, the weighting factor is reduced to z^{n-3} . This reduced trigger bias represents an advantage of the present method.

3.3. Corrections

The reconstructed strange particles consist of both real signal particles and combinatorial background (fake strange particles). To obtain the correlation function for the signal particles, the contribution to the p_T -weighted correlation function from the combinatorial background is removed using the sideband subtraction method [12,31]. The invariant mass distributions of the strange particle candidates in each p_T^s interval are fitted with a Gaussian function for the signal combined with a linear function describing the combinatorial background. Using the mean and variance (σ) of the Gaussian, the signal and the sideband regions are defined. The signal region is defined as the $\pm 3\sigma$ range centered at the mean value, while the sideband regions, used to estimate the combinatorial background, are located between 6σ and 9σ symmetric around the mean value. The p_T -weighted correlation functions are constructed using triggers from the signal and sideband regions, respectively. The sideband correlation distributions, which provide an estimation of the combinatorial background, are subtracted from the signal correlation distributions.

To account for tracking efficiency, a dedicated efficiency correction is applied to the associated particles. The tracking efficiency (ϵ) is obtained from MC simulations and is defined as the ratio of the number of reconstructed particles to the number of generated particles satisfying the primary charged-particle definition (see Section 1). In this work, the tracking efficiency increases from approximately 70% at the lowest p_T , and reaches a plateau of 85% above $p_T \approx 1.5$ GeV/c. This correction is achieved by using ϵ as another part of the weight when filling the p_T -weighted correlation function. Not all charged tracks passing the selection criteria are primary tracks. The secondary contamination fraction (f_{sec}) is defined as the ratio of secondary particles to the total number of particles passing the selection, and is also determined using MC simulations. The estimated secondary fraction decreases from approximately 3.8% at low p_T to below 1% for $p_T > 1$ GeV/c. To correct for both effects, an additional weight factor beyond the associated particle's p_T , namely $(1 - f_{\text{sec}})/\epsilon$, is introduced when calculating the correlation functions.

The corrected p_T -weighted correlation functions are still strongly affected by the limited and non-uniform acceptance of the detector. In particular, the finite pseudorapidity acceptance leads to a characteristic

triangular shape in the $\Delta\eta$ direction, which is a purely geometrical effect rather than a consequence of physical correlations. To correct for this non-physical modification, the event-mixing technique is applied. In this method, with the same selection criteria, the trigger particles and the associated particles are chosen from different events to construct the mixed-event correlation function, $M(\Delta\eta, \Delta\phi) = \frac{d^2 N_{\text{mix}}}{d\Delta\phi d\Delta\eta}$, where N_{mix} denotes the number of trigger-associate pairs in mixed events. The events are mixed only if the difference in the z -vertex position is less than 2 cm. Since trigger and associated particles come from different events, no physical correlation occurs; thus, the mixed-event distribution reflects only the detector effects. For each trigger p_T^s interval, the p_T -weighted correlation function $C_{p_T\text{-weighted}}(\Delta\eta, \Delta\phi)$ and the mixed-event distribution $M(\Delta\eta, \Delta\phi)$ are constructed independently in different associated particle p_T intervals. By dividing $C_{p_T\text{-weighted}}(\Delta\eta, \Delta\phi)$ and $M(\Delta\eta, \Delta\phi)$, the corrected p_T -weighted correlation function is obtained as

$$C_{p_T\text{-weighted}}^{\text{corrected}}(\Delta\eta, \Delta\phi) = \frac{C_{p_T\text{-weighted}}(\Delta\eta, \Delta\phi)}{\alpha M(\Delta\eta, \Delta\phi)}, \quad (3)$$

where α is the normalization factor for the mixed-event correlation function given by the inverse of the average along the $\Delta\phi$ -axis of bin contents for the bins where $\Delta\eta = 0$. This normalization is motivated by the fact that, at the limit of very small angular distance, the acceptance is expected to approach unity.

Finally, the corrected correlation functions from different associated particle p_T intervals are summed to obtain the final p_T -weighted correlation functions in each p_T^s interval. Examples of the fully corrected p_T -weighted correlation functions for K_S^0 and Λ ($\bar{\Lambda}$) are shown in Fig. 1. The broader structure along $\Delta\eta$ for Λ ($\bar{\Lambda}$) suggests a weaker collimation with respect to the jet axis compared to K_S^0 , possibly related to the different production mechanisms of baryons and mesons.

To compute the $\langle \sum^{\text{NS}} p_T \rangle$ from the jet-like correlations, the combined, corrected correlation function is projected onto the $\Delta\phi$ axis within $|\Delta\eta| < 1.2$. This “short-range” projection is assumed to contain the full jet signal and the contribution from the uncorrelated background (underlying event). Similarly, the projection within $1.2 < |\Delta\eta| < 1.4$ is also obtained. This “long-range” projection is dominated by the underlying event and therefore can be used to provide an estimation of the uncorrelated background in the “short-range”. For this purpose, the “long-range” projection needs to be normalized per unit of rapidity and be subtracted from the “short-range” projection. Examples are shown in Fig. 2, where the “short-range” projection is shown in green points and the normalized “long-range” projection is drawn in the red. Then, $\langle \sum^{\text{NS}} p_T \rangle$ can be extracted from the jet-like correlations as the integral of the near-side peak region. In most cases, the integration is performed within the $(-0.5\pi, 0.5\pi)$ region in the $\Delta\phi$ in order to include all jet contributions. However, for $0.6 < p_T^s < 1.0$ GeV/c for K_S^0 and $0.6 < p_T^s < 0.8$ GeV/c for Λ ($\bar{\Lambda}$), a broader near-side peak is observed and the integration range is therefore extended to $(-0.6\pi, 0.6\pi)$.

4. Systematic uncertainties

The systematic uncertainties are evaluated by varying the selection criteria or other analysis parameters. In each p_T^s interval, the individual systematic uncertainties are considered uncorrelated, and the total systematic uncertainty is calculated as the square root of the sum of their squares. For each source, the difference Δ between the varied and nominal results is divided by the corresponding statistical uncertainty, $\sigma_\Delta = \sqrt{|\sigma_{\text{var}}^2 - \sigma_{\text{nom}}^2|}$, which is computed taking into account the correlation between the nominal and varied results. The symbols σ_{var} and σ_{nom} denote the statistical uncertainties associated with the varied and nominal results, respectively [34]. If $N_\sigma = \Delta/\sigma_\Delta < 2$, the varied result is considered statistically compatible with the nominal result, and no systematic uncertainty for this source is assigned. Otherwise, the con-




Fig. 1. Example of the fully corrected p_T -weighted correlation functions for K_S^0 -primary charged-particles correlation (top) and for Λ ($\bar{\Lambda}$)-primary charged-particles correlation (bottom).

tribution is added to the total systematic uncertainty. The summary of all systematic uncertainties is given in Table 2.

The uncertainty due to the primary vertex z -position selection is estimated by tightening the allowed range from $|z_{\text{vtx}}| < 10$ cm to $|z_{\text{vtx}}| < 7$ cm, resulting in a more uniform detector acceptance. For K_S^0 , the systematic uncertainty is negligible compared to the statistical one, and for Λ ($\bar{\Lambda}$), the uncertainty rises to 0.6% in the highest p_T interval.

The uncertainty due to the V^0 signal extraction is determined by varying the signal and sideband regions, which accounts for the effect of the combinatorial background subtraction. The definition of the signal region is varied between 3σ and 7σ centered at the mean of the Gaussian function, and the corresponding lower and upper bounds of the sideband regions are varied between 4σ – 7σ and 8σ – 14σ , respectively. For this source, the uncertainty is less than 0.4% for K_S^0 and Λ ($\bar{\Lambda}$) in most p_T intervals. For Λ ($\bar{\Lambda}$), it goes up to 1.2% in the highest p_T interval.

To assess the systematic uncertainty of the geometrical acceptance correction, the scale factor α for the mixed-event distribution is computed using the inverse of the bin content centered at $\Delta\phi = \Delta\eta = 0$, rather than the default procedure, which uses the average evaluated at $\Delta\eta = 0$. The systematic uncertainty from this source is below 0.5% for K_S^0 and Λ ($\bar{\Lambda}$) in most p_T intervals, rising to 1.2% in the highest p_T interval for K_S^0 .

The systematic uncertainty related to the primary charged track selection is estimated by using a tighter DCA requirement. Tracks need to satisfy a p_T -dependent selection $|DCA_{xy}| < 0.0105 + 0.0350/p_T^{1.1}$ cm in the xy -plane and $|DCA_z| < 2$ cm in the z -direction. For this source, the uncertainty is less than 0.4% for K_S^0 and Λ ($\bar{\Lambda}$).




Fig. 2. Examples of $\Delta\phi$ projections of the fully corrected p_T -weighted correlation functions for K_S^0 –primary charged-particles correlation (left) and for $\Lambda(\bar{\Lambda})$ –primary charged-particles correlation (right). The “short-range” ($|\Delta\eta| < 1.2$) and the normalized “long-range” ($1.2 < |\Delta\eta| < 1.4$) projections are shown in green and red points, respectively.

The systematic uncertainty associated with the choice of the short-range region peak and the control region used for the subtraction of the uncorrelated background is assessed by changing the boundaries of these regions. The following boundaries for the short-range peak region (control region) are used for the estimation, $|\Delta\eta| < 1.1$ ($1.1 < |\Delta\eta| < 1.4$), $|\Delta\eta| < 1.3$ ($1.3 < |\Delta\eta| < 1.4$), $|\Delta\eta| < 1.3$ ($1.3 < |\Delta\eta| < 1.5$), and $|\Delta\eta| < 1.4$ ($1.4 < |\Delta\eta| < 1.5$). For this source, the systematic uncertainty is less than 2.5% in most p_T intervals for K_S^0 and $\Lambda(\bar{\Lambda})$.

By default, the V^0 candidates are required to have at least one decay track with ITS information to reject the out-of-bunch pileup. As a variation of this selection, the candidates are required to have at least one decay track with information in ITS and TOF. For this source, the uncertainty is less than 1% in most intervals for K_S^0 . For $\Lambda(\bar{\Lambda})$, the systematic uncertainty from this source is negligible compared to the statistical one.

To take into account the systematic uncertainty related to the residual contamination from in-bunch (IB) pileup, the full sample is divided into two sets of equal size but different μ , the average number of proton-proton interactions per bunch crossing. The average μ value for the full sample is 0.0178, and for the two sub-samples, the corresponding values are 0.0085 and 0.0271, respectively. For this source, the uncertainty is less than 2% for K_S^0 and for most p_T intervals for $\Lambda(\bar{\Lambda})$.

The imperfect description of the ALICE detector in the MC simulation [35] leads to systematic biases in the estimation of the tracking efficiency and therefore of the $\sum^{\text{NS}} p_T$. The corresponding systematic uncertainty of $\langle z \rangle$ is calculated by varying the tracking efficiency by $\pm 3\%$ [36,37] and is found to be less than 1.4% for K_S^0 and $\Lambda(\bar{\Lambda})$.

Strange baryon Λ and $\bar{\Lambda}$ candidates can also originate from the decay of $\Sigma^\pm$, $\Xi^\pm$, $\Omega^\pm$, and their corresponding antiparticles. For the systematic uncertainty related to this source, the $\langle z \rangle$ of primary- and feed-down $\Lambda(\bar{\Lambda})$ are studied in simulations. The study indicates that $\langle z \rangle$ is not highly sensitive to the origin of the baryon. For this source, a 2% uncertainty uncorrelated with p_T is assigned.

5. Results and discussion

The average transverse-momentum fractions $\langle z \rangle$ as a function of strange particle $p_T^\perp$ in pp collisions at $\sqrt{s} = 13$ TeV for K_S^0 and Λ produced in the $|\eta| < 0.75$ are shown in Fig. 3. The summed transverse momentum of the associated particles ($\sum^{\text{NS}} p_T$) has been measured for particles with $0.2 < p_T < 20$ GeV/c and no extrapolation outside this fiducial region has been performed.

At high $p_T^\perp$ (> 6 GeV/c), $\langle z \rangle$ is approximately 0.6 for both particle species. This relatively large value compared to typical momentum fractions of light hadrons in jets ($\langle z \rangle \approx 0.3$) [38] is expected due to the trigger bias, characteristic of particles originating from the fragmentation of

Table 2

Systematic uncertainties of $\langle z \rangle$ for K_S^0 and $\Lambda(\bar{\Lambda})$. The uncertainties vary within the indicated ranges depending on the $p_T^\perp$ interval.

Source	Systematic uncertainties (%)	
	K_S^0	$\Lambda(\bar{\Lambda})$
z_{vis} selection	negligible	0.0 – 0.6
V^0 signal extraction	0.0 – 0.4	0.1 – 1.2
Mixed-event scale	0.0 – 0.5	0.0 – 1.2
Primary charged track selection	0.0 – 0.1	0.0 – 0.4
Uncorrelated background subtraction	0.3 – 2.5	0.5 – 2.1
Out-of-bunch pileup	0.0 – 1.1	negligible
In-bunch pileup	0.0 – 1.6	0.1 – 2.4
Material budget	1.1 – 1.3	0.7 – 1.4
Feed-down	N/A	2.0
Total	1.5 – 2.9	2.3 – 4.0

partons, whose p_T -differential production rate follows a steeply falling power-law spectrum (see discussion in Section 3.2).

For K_S^0 mesons, there is no indication of a significant change in the production mechanism over the measured $p_T^\perp$ range as $\langle z \rangle$ remains stable at ≈ 0.6 within uncertainties. On the other hand, interestingly, for the $\Lambda(\bar{\Lambda})$ baryons, $\langle z \rangle$ increases as $p_T^\perp$ decreases, showing a rise from intermediate to low p_T and deviating from that of the K_S^0 below approximately 4 GeV/c. For the lowest $p_T^\perp$ interval, $\langle z \rangle = 0.78$, which is around 30% higher than the high $p_T^\perp$ limit. Qualitatively, a rise can be expected from a steeper than power-law parent-particle p_T spectrum in the intermediate $p_T^\perp$ region, which would worsen the aforementioned bias. However, if this were the only effect, it should be similar for K_S^0 and $\Lambda(\bar{\Lambda})$. Hence, the increase of $\langle z \rangle$ for $\Lambda(\bar{\Lambda})$ is either because $\Lambda(\bar{\Lambda})$ take a larger momentum fraction of the originating partons, or because there is a larger contribution from the isolated $\Lambda(\bar{\Lambda})$ production in soft processes, which corresponds to $z = 1$. The observed difference in $\langle z \rangle$ between K_S^0 and $\Lambda(\bar{\Lambda})$ may also be attributed to strangeness and baryon number conservation. For $\Lambda(\bar{\Lambda})$, balancing can mainly occur via strange baryons, none of which contribute to $\sum^{\text{NS}} p_T$, while K_S^0 can be balanced by kaons, including the charged ones contributing to $\sum^{\text{NS}} p_T$.

In order to investigate potential differences in the hadronization mechanisms, the results are compared with several Monte Carlo model calculations employing different hadronization schemes, including PYTHIA 8.3 [39] (Monash and Color Rope tunes) and AMPT (with string melting) [40], as shown in Fig. 4. The same kinematic and acceptance selections are applied in the simulation. All primary charged particles are used as associated particles, excluding the long-lived charged strange baryons discussed in Section 3, and the same analysis




Fig. 3. The average transverse-momentum fraction ($\langle z \rangle$) for $\Lambda(\bar{\Lambda})$ (blue) and K_S^0 (red) as a function of p_T^s in minimum bias pp collisions at $\sqrt{s} = 13$ TeV. The p_T^s ranges from 0.6 to 20 GeV/c. Statistical and systematic uncertainties of $\langle z \rangle$ are represented by vertical error bars and empty boxes, respectively. Data points are drawn at the corresponding average p_T^s values in each interval. The horizontal bars represent the p_T^s interval widths. The statistical and systematic uncertainties of the average p_T^s values are negligible.

procedure is followed. Unlike in data, the particle identity is known at generator level and no detector effects are involved. Therefore, neither the side-band subtraction nor the additional weight $(1 - f_{\text{sec}})/\epsilon$ is required for the correction on reconstruction inefficiency and secondary-particle contamination. The band widths represent the quadratic sum of the statistical uncertainties and the systematic uncertainties associated with the uncorrelated background subtraction, as described in Section 4. The hadronization mechanism in PYTHIA8 is based on the Lund string fragmentation model. Coherence between strings from different parton-parton interactions is implemented via the so-called color reconnection (CR) mechanism. Partons from different strings or parton showers are allowed to reconnect their color flow, potentially forming shorter or more energetically favorable strings before hadronization. The mechanism affects the final-state multiplicity and improves the description of the increase of the average p_T with multiplicity. Since version 8.2, Monash 2013 is the default tune. It is a general-purpose tune that provides a consistent description of many observables measured at LHC and previous collider data [27]. Since CR alone is not able to describe strangeness enhancement, the Color Rope mechanism has been introduced. Overlapping strings merge or interact forming “ropes”—color flux tubes with higher color charge. These ropes have an increased effective string tension compared to normal strings which enhances strangeness production. In the AMPT model, the initial state is generated using HIJING [41], followed by partonic scatterings described by the Zhang’s Parton Cascade model [42]. The hadronization process is then implemented through quark coalescence, where partons close in phase space are recombined to form hadrons.

As expected, in the high p_T^s region, the observed $\langle z \rangle$ is well reproduced by the models. In the intermediate p_T^s region, the models underestimate the data by approximately 10%. For K_S^0 , they predict a minimum close to 2 GeV/c and a steep rise towards lower p_T^s . For $\Lambda(\bar{\Lambda})$, the calculations show a steep rise below 2 GeV/c, without a significant minimum, unlike the case for K_S^0 . In contrast, the measurement exhibits a smooth and continuous behavior over the measured p_T^s interval.

For the PYTHIA tunes, primary V^0 production is compared to production including K_S^0 triggers originating from the decays of the resonances $K^*(892)^0$, $K^*(892)^\pm$, and $\phi(1020)$, and $\Lambda(\bar{\Lambda})$ triggers coming from the decays of $\Sigma^\pm$, $\Xi^\pm$, $\Omega^\pm$ and their corresponding antiparticles, as shown in Fig. 4 with solid-filled bands and hatched bands, respectively. In the




Fig. 4. The average transverse-momentum fraction ($\langle z \rangle$) for K_S^0 (top) and $\Lambda(\bar{\Lambda})$ (bottom) in data compared with model calculations. The data are compared to PYTHIA 8 with the Monash tune (red band), the Color Rope tune (green band) and the AMPT model with the string melting (blue band). For both PYTHIA 8 tunes, the hatched bands represent the results excluding K_S^0 triggers coming from decays of the resonances $K^*(892)^0$, $K^*(892)^\pm$, and $\phi(1020)$, and $\Lambda(\bar{\Lambda})$ triggers coming from decays of $\Sigma^\pm$, $\Xi^\pm$, $\Omega^\pm$ and their corresponding antiparticles.

case of $\Lambda(\bar{\Lambda})$, $\langle z \rangle$ does not change significantly when excluding the feed-down contribution, while for K_S^0 , $\langle z \rangle$ increases by at most about 10%.

6. Conclusion

The first measurement of the average transverse-momentum fraction $\langle z \rangle$ as a function of strange particle p_T for the strange baryon $\Lambda(\bar{\Lambda})$ and the strange meson K_S^0 within mini-jets in pp collisions at $\sqrt{s} = 13$ TeV is reported. The results show different behavior for K_S^0 and $\Lambda(\bar{\Lambda})$ in the low and intermediate p_T regions, suggesting different hadronization properties for mesons and baryons in these ranges. The higher $\langle z \rangle$ of $\Lambda(\bar{\Lambda})$ observed in the intermediate p_T region could be attributed to either a harder fragmentation or a larger production of isolated Λ . This could be related to the enhancement of strange-baryon-to-meson yield ratios (Λ/K_S^0) observed in previous measurements, in this p_T region [18]. For K_S^0 , there is no strong p_T dependence of $\langle z \rangle$, which may indicate that there is no significant change of the production mechanism in the intermediate p_T region with respect to the high p_T region. The PYTHIA 8 Monash and Color Rope tunes, as well as the AMPT model, do not describe the K_S^0 data. For $\Lambda(\bar{\Lambda})$, the models overestimate the values at low p_T but reproduce the overall decreasing trend toward higher p_T , with

AMPT showing the closest behavior in the intermediate region. Overall, all used models fail to satisfactorily reproduce the measured $\langle z \rangle$. They predict a different trend and strongly underestimate the $\langle z \rangle$ of K_S^0 at the intermediate p_T , while for Λ ($\bar{\Lambda}$), they show a steeper decrease than observed in the data at low and intermediate p_T . Future studies will extend the analysis to multi-strange baryons and investigate the multiplicity dependence of the transverse-momentum fraction. This will provide further information on the origin of the baryon enhancement in the intermediate p_T^s region.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

The ALICE Collaboration would like to thank all its engineers and technicians for their invaluable contributions to the construction of the experiment and the CERN accelerator teams for the outstanding performance of the LHC complex. The ALICE Collaboration gratefully acknowledges the resources and support provided by all Grid centres and the Worldwide LHC Computing Grid (WLCG) collaboration. The ALICE Collaboration acknowledges the following funding agencies for their support in building and running the ALICE detector: A. I. Alikhanyan National Science Laboratory (Yerevan Physics Institute) Foundation (ANSI), State Committee of Science and World Federation of Scientists (WFS), Armenia; Austrian Academy of Sciences, Austrian Science Fund (FWF): [M 2467-N36] and Nationalstiftung für Forschung, Technologie und Entwicklung, Austria; Ministry of Communications and High Technologies, National Nuclear Research Center, Azerbaijan; Rede Nacional de Física de Altas Energias (Renafae), Financiadora de Estudos e Projetos (Finep), Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP) and The Sao Paulo Research Foundation (FAPESP), Brazil; Bulgarian Ministry of Education and Science, within the National Roadmap for Research Infrastructures 2020–2027 (object CERN), Bulgaria; Ministry of Education of China (MOEC), Ministry of Science & Technology of China (MSTC) and National Natural Science Foundation of China (NSFC), China; Ministry of Science and Education and Croatian Science Foundation, Croatia; Centro de Aplicaciones Tecnológicas y Desarrollo Nuclear (CEADEN), Cubaenergía, Cuba; Ministry of Education, Youth and Sports of the Czech Republic, Czech Republic; The Danish Council for Independent Research | Natural Sciences, the VIL-LUM FONDEN and Danish National Research Foundation (DNRF), Denmark; Helsinki Institute of Physics (HIP), Finland; Commissariat à l’Energie Atomique (CEA) and Institut National de Physique Nucléaire et de Physique des Particules (IN2P3) and Centre National de la Recherche Scientifique (CNRS), France; Bundesministerium für Forschung, Technologie und Raumfahrt (BMFTR) and GSI Helmholtzzentrum für Schwerionenforschung GmbH, Germany; National Research, Development and Innovation Office, Hungary; Department of Atomic Energy Government of India (DAE), Department of Science and Technology, Government of India (DST), University Grants Commission, Government of India (UGC) and Council of Scientific and Industrial Research (CSIR), India; National Research and Innovation Agency - BRIN, Indonesia; Istituto Nazionale di Fisica Nucleare (INFN), Italy; Japanese Ministry of Education, Culture, Sports, Science and Technology (MEXT) and Japan Society for the Promotion of Science (JSPS) KAKENHI, Japan; Consejo Nacional de Ciencia (CONACYT) y Tecnología, through Fondo de Cooperación Internacional en Ciencia y Tecnología (FONCICYT) and Dirección General de Asuntos del Personal Académico (DGAPA), Mexico; Nederlandse Organisatie voor Wetenschappelijk Onderzoek (NWO), Netherlands; The Research Council of Norway, Norway; Pontificia Universidad Católica del Perú, Peru; Ministry of Science and Higher Education, National Science Centre and WUT ID-UB, Poland; Korea Institute of

Science and Technology Information and National Research Foundation of Korea (NRF), Republic of Korea; Ministry of Education and Scientific Research, Institute of Atomic Physics, Ministry of Research and Innovation and Institute of Atomic Physics and Universitatea Nationala de Stiinta si Tehnologie Politehnica Bucuresti, Romania; Ministerstvo školstva, výskumu, vývoja a mládeže SR, Slovakia; National Research Foundation of South Africa, South Africa; Swedish Research Council (VR) and Knut & Alice Wallenberg Foundation (KAW), Sweden; European Organization for Nuclear Research, Switzerland; Suranaree University of Technology (SUT), National Science and Technology Development Agency (NSTDA) and National Science, Research and Innovation Fund (NSRF via PMU-B B05F650021), Thailand; Turkish Energy, Nuclear and Mineral Research Agency (TENMAK), Turkey; National Academy of Sciences of Ukraine, Ukraine; Science and Technology Facilities Council (STFC), United Kingdom; National Science Foundation of the United States of America (NSF) and United States Department of Energy, Office of Nuclear Physics (DOE NP), United States of America. In addition, individual groups or members have received support from: Czech Science Foundation (grant no. 23-07499S), Czech Republic; FORTE project, reg. no. CZ.02.01.01/00/22_008/0004632, Czech Republic, co-funded by the European Union, Czech Republic; European Research Council (grant nos. 101220549, 950692), European Union; Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) “Neutrinos and Dark Matter in Astro- and Particle Physics” (grant no. SFB 1258), Germany; CONVECS project, CUP C97H23001700002 FESR 2021–2027 program, Italy.

Data availability

This manuscript has associated data in a HEPData repository at: <https://www.hepdata.net/record/ins3132536>.

The ALICE Collaboration

D.A.H. Abdallah¹³⁴, I.J. Abualrob¹¹², S. Acharya⁴⁹, K. Agarwal^{1,23}, G. Aglieri Rinella³², L. Aglietta²⁴, N. Agrawal²⁵, Z. Ahammed¹³², S. Ahmad¹⁵, I. Ahuja³⁶, Z. Akbar⁷⁹, V. Akishina³⁸, M. Al-Turany⁹⁴, B. Alessandro⁵⁵, A.R. Alfarasyi¹⁰¹, R. Alfaro Molina⁶⁶, B. Ali¹⁵, A. Alici²⁵, J. Alme²⁰, G. Alocco²⁴, T. Alt⁶³, I. Altsybeev⁹², C. Andrei⁴⁴, N. Andreou¹¹¹, A. Andronic¹²³, M. Angeletti³², V. Anguelov⁹¹, F. Antinori⁵³, P. Antonioli⁵⁰, N. Apadula⁷¹, H. Appelshäuser⁶³, S. Arcelli²⁵, R. Arnaldi⁵⁵, I.C. Arsene¹⁹, M. Arslanod¹³⁵, A. Augustinus³², R. Averbeck⁹⁴, M.D. Azmi¹⁵, H. Baba¹²¹, A.R.J. Babu¹³⁴, A. Badalà⁵², J. Bae¹⁰⁰, Y. Bae¹⁰⁰, Y.W. Baek¹⁰⁰, X. Bai¹¹⁶, R. Bailhache⁶³, Y. Bailung¹²⁵, R. Bala⁸⁸, A. Baldissari¹²⁷, B. Balis², S. Bangalia¹¹⁴, V. Barbasova³⁶, F. Barile³¹, L. Barioglio⁵⁵, M. Barlou²⁴, B. Barman⁴⁰, G.G. Barnaföldi⁴⁵, L.S. Barnby¹¹¹, E. Barreau⁹⁹, V. Barret¹²⁴, L. Barreto¹⁰⁶, K. Barth³², E. Bartsch⁶³, N. Bastid¹²⁴, G. Batigne⁹⁹, D. Battistini^{34,92}, B. Batyunya¹³⁹, L. Baudino^{11,24}, D. Bauri⁴⁶, J.L. Bazo Alba⁹⁸, I.G. Bearden⁸⁰, P. Becht⁹⁴, D. Behera^{47,77}, S. Behera⁴⁶, M.A.C. Behling⁶³, I. Belikov¹²⁶, V.D. Bella¹²⁶, F. Bellini²⁵, R. Bellwied¹¹², L.G.E. Beltran¹⁰⁵, Y.A.V. Beltran⁴³, G. Bencedi⁴⁵, O. Benchikh⁷³, A. Bensaoula¹¹², S. Beole²⁴, A. Berdnikova⁹¹, L. Bergmann⁷¹, L. Bernardinis²³, L. Betev³², P.P. Bhaduri¹³², T. Bhalla⁸⁷, A. Bhasin⁸⁸, B. Bhattacharjee⁴⁰, B. Bianchi²⁴, J. Bielčik³⁴, J. Bielčíková⁸³, A. Bilandzic⁹², A. Binoy¹¹⁴, G. Biro⁴⁵, S. Biswas⁴, M.B. Blidaru⁹⁴, N. Bluhme³⁸, C. Blume⁶³, F. Bock⁸⁴, T. Bodova²⁰, L. Boldizsár⁴⁵, M. Bombara³⁶, P.M. Bond³², G. Bonomi^{54,131}, H. Borel¹²⁷, A. Borissov¹³⁹, A.G. Borquez Carcamo⁹¹, E. Botta²⁴, N. Bouchhar¹⁷, Y.E.M. Bouziani⁶³, D.C. Brandibur⁶², L. Bratrud⁶³, P. Braun-Munzinger⁹⁴, M. Bregant¹⁰⁶, M. Broz³⁴, G.E. Bruno^{31,93}, V.D. Buchachiev³⁵, M.D. Buckland⁸², G.F. Budiski¹⁰⁶, H. Buesching⁶³, S. Bufalino²⁹,

P. Buhler⁷³, N. Burmasov¹³⁹, Z. Buthelezi^{67,120}, A. Bylinkin²⁰, O.B. Bylund¹²⁸, J.C. Cabanillas Noris¹⁰⁵, M.F.T. Cabrera¹¹², H. Caines¹³⁵, A. Caliva²⁸, E. Calvo Villar⁹⁸, J.M.M. Camacho¹⁰⁵, P. Camerini²³, M.T. Camerlingo⁴⁹, F.D.M. Canedo¹⁰⁶, S. Cannito²³, S.L. Cantway¹³⁵, M. Carabas¹⁰⁹, F. Carnesecchi³², C. Carr⁹⁷, L.A.D. Carvalho¹⁰⁶, J. Castillo Castellanos¹²⁷, M. Castoldi³², F. Catalano¹¹², S. Cattaruzzi²³, R. Cerri²⁴, I. Chakaberia⁷¹, P. Chakraborty¹³³, J.W.O. Chan¹¹², S. Chandra¹³², S. Chapeland³², M. Chartier¹¹⁵, S. Chattopadhyay¹³², M. Chen³⁹, T. Cheng⁶, M.I. Cherciu⁶², C. Cheshkov¹²⁵, D. Chiappara²⁷, V. Chibante Barroso³², D.D. Chinellato⁷³, F. Chinua²⁴, J. Cho⁵⁷, S. Cho⁵⁷, P. Chochula³², Z.A. Chochulska^{113,133}, C. Choi¹⁶, P. Christakoglou⁸¹, P. Christiansen⁷², T. Chujo¹²², B. Chytla¹³³, M. Ciacco²⁴, C. Cicalo⁵¹, G. Cimdor^{24,32}, F. Cindolo⁵⁰, F. Colamaria⁴⁹, D. Colella³¹, A. Colelli³¹, M. Colocci²⁵, M. Concas³², G. Conesa Balbastre⁷⁰, Z. Conesa del Valle¹²⁸, G. Contin²³, J.G. Contreras³⁴, M.L. Coquet⁹⁹, P. Cortese^{55,130}, M.R. Cosentino¹⁰⁸, F. Costa³², S. Costanza²¹, P. Crochet¹²⁴, M.M. Czarnynoga¹³³, A. Dainese⁵³, E. Dall'occo³², G. Dange³⁸, M.C. Danisch¹⁶, A. Danu⁶², A. Daribayeva³⁸, P. Das³², S. Das⁴, A.R. Dash¹²³, S. Dash⁴⁶, A. De Caro²⁸, G. de Cataldo⁴⁹, J. de Cuveland³⁸, A. De Falco²², D. De Gruttola²⁸, N. De Marco⁵⁵, C. De Martin²³, S. De Pasquale²⁸, R. Deb¹³¹, R. Del Grande³⁴, L. Dello Stritto³², G.G.A. de Souza^{11,106}, P. Dhanke¹⁸, D. Di Bari³¹, M. Di Costanzo²⁹, A. Di Mauro³², B. Di Ruzza^{49,129}, B. Diab³², Y. Ding⁶, J. Ditzel⁶³, R. Divià³², U. Dmitrieva⁵⁵, A. Dobrin⁶², B. Dönigus⁶³, L. Döpper⁴¹, L. Drzensla², J.M. Dubinski¹³³, A. Dubla⁹⁴, P. Dupieux¹²⁴, N. Dzalaiova¹³, T.M. Eder¹²³, E.C. Ege⁶³, R.J. Ehlers⁷¹, F. Eisenhut⁶³, R. Ejima⁸⁹, D. Elia⁴⁹, B. Erasmus⁹⁹, F. Ercolessi²⁵, B. Espagnon¹²⁸, G. Eulisse³², D. Evans⁹⁷, L. Fabbietti⁹², G. Fabbri⁵⁰, M. Faggin³², J. Faivre⁷⁰, W. Fan¹¹², Y. Fan⁶, T. Fang⁶, A. Fantoni⁴⁸, A. Feliciello⁵⁵, W. Feng⁶, A. Fernández Téllez⁴³, B. Fernando¹³⁴, L. Ferrandi¹⁰⁶, A. Ferrero¹²⁷, C. Ferrero^{5,55}, A. Ferretti²⁴, F.M. Fionda⁵¹, A.N. Flores¹⁰⁴, S. Foertsch⁶⁷, I. Fokin⁹¹, U. Follo^{5,55}, R. Forynski¹¹¹, E. Fragiaco⁵⁶, H. Friberg⁹², U. Fuchs³², D. Fuligno²³, N. Funicello²⁸, C. Furget⁷⁰, T. Fusayasu⁹⁵, J.J. Gaardhøje⁸⁰, M. Gagliardi²⁴, A.M. Gago⁹⁸, T. Gahlaut⁴⁶, C.D. Galvan¹⁰⁵, S. Gami⁷⁷, C. Garabatos⁹⁴, J.M. Garcia⁴³, E. Garcia-Solis⁹, S. Garetti¹²⁸, C. Gargiulo³², P. Gasik⁹⁴, A. Gautam¹¹⁴, M.B. Gay Ducati⁶⁵, M. Germain⁹⁹, R.A. Gernhaeuser⁹², M. Giacalone³², G. Gioachin²⁹, S.K. Giri¹³², P. Giubellino⁵⁵, P. Giubilato²⁷, P. Gläsel⁹¹, E. Glimos¹¹⁹, M.G.F.S.A. Gomes⁹¹, L. Gonella²³, V. Gonzalez¹³⁴, M. Gorgon², K. Goswami⁴⁷, S. Gotovac³³, V. Grabski⁶⁶, L.K. Graczykowski¹³³, E. Grecka⁸³, A. Grelli⁵⁸, C. Grigoras³², S. Grigoryan^{1,139}, O.S. Groetvik³², M. Gronbeck⁴¹, F. Grosa³², S. Gross-Bölting⁹⁴, J.F. Grosse-Oetringhaus³², R. Grosso⁹⁴, D. Grund³⁴, N.A. Grunwald⁹¹, R. Guernane⁷⁰, M. Guilbaud⁹⁹, K. Gulbrandsen⁸⁰, J.K. Gumprecht⁷³, T. Gündem⁶³, T. Gunji¹²¹, J. Guo¹⁰, W. Guo⁶, A. Gupta⁸⁸, R. Gupta⁸⁸, R. Gupta⁴⁷, K. Gwizdziel¹³³, L. Gyulai⁴⁵, T. Hachiya⁷⁵, C. Hadjidakis¹²⁸, F.U. Haider⁸⁸, S. Haidlova³⁴, M. Haldar⁴, W. Ham¹⁰⁰, H. Hamagaki⁷⁴, Y. Han¹³⁷, R. Hannigan¹⁰⁴, J. Hansen⁷², J.W. Harris¹³⁵, A. Harton⁹, M.V. Hartung⁶³, A. Hasan¹¹⁸, H. Hassan¹¹³, D. Hatzifotiadiou⁵⁰, P. Hauer⁴¹, L.B. Havener¹³⁵, E. Hellbär³², H. Helstrup³⁷, M. Hemmer⁶³, S.G. Hernandez¹¹², G. Herrera Corral⁸, K.F. Hetland³⁷, B. Heybeck⁶³, H. Hillemanns³², B. Hippolyte¹²⁶, I.P.M. Hobus⁸¹, F.W. Hoffmann³⁸, B. Hofman⁵⁸, Y. Hong⁵⁷, A. Horzyk², Y. Hou^{11,94}, P. Hristov³², L.M. Huhta¹¹³, T.J. Humanic⁸⁵, V. Humlova³⁴, M. Husar⁸⁶, A. Hutson¹¹², D. Hutter³⁸, M.C. Hwang¹⁸, M. Inaba¹²², A. Isakov⁸¹, T. Isidori¹¹⁴, M.S. Islam⁴⁶, M. Ivanov⁹⁴, M. Ivanov¹³, K.E. Iversen⁷², M. Jablonski², B. Jacak^{18,71}, N. Jacazio¹³⁰, P.M. Jacobs⁹⁹, A. Jádlovská¹⁰², S. Jádlovská¹⁰², S. Jaelani⁷⁹, J.N. Jager⁶³, C. Jahnke¹⁰⁷, M.J. Jakubowska¹³³, E.P. Jamroz², D.M. Janik³⁴, M.A. Janik¹³³, C.A. Jauch⁹⁴, S. Ji¹⁶, Y. Ji⁹⁴, S. Jia⁸⁰, T. Jiang¹⁰, A.A.P. Jimenez⁶⁴, S. Jin¹⁰, Z. Jolesz⁴⁵, F. Jonas⁷¹, D.M.

Jones¹¹⁵, J.M. Jowett^{32,94}, J. Jung⁶³, M. Jung⁶³, A. Junique³², J. Juračka³⁴, J. Kaewjai^{101,115}, A. Kaiser^{32,94}, P. Kalinak⁵⁹, A. Kalweit³², H. Kang¹², A. Karasu Uysal¹³⁶, N. Karatzenis⁹⁷, T. Karavicheva¹³⁹, M.J. Karwowska¹³³, V. Kashyap⁷⁷, M. Keil³², B. Ketzer⁴¹, J. Keul⁶³, S.S. Khade⁴⁷, A. Khuntia⁵⁰, Z. Khuranova⁶³, B. Kileng³⁷, B. Kim¹⁰⁰, D.J. Kim¹¹³, D. Kim¹⁰⁰, E.J. Kim⁶⁸, G. Kim⁵⁷, H. Kim⁵⁷, J. Kim¹³⁷, J. Kim⁵⁷, J. Kim¹³⁷, J. Kim³², M. Kim¹⁶, M. Kim¹⁸, S. Kim¹⁷, T. Kim¹³⁷, J.T. Kinner¹²³, I. Kisel³⁸, A. Kisel¹³³, J.L. Klay⁵, J. Klein³², S. Klein⁷¹, C. Klein-Bösing¹²³, M. Kleiner⁶³, A. Kluge³², M.B. Knuesel¹³⁵, C. Kobdaj¹⁰¹, R. Kohara¹²¹, A. Kondratyev¹³⁹, J. Konig⁶³, P.J. Konopka³², G. Kornakov¹³³, M. Korwieser⁹², C. Koster⁸¹, A. Kotliarov⁸³, N. Kovacic⁸⁶, M. Kowalski¹⁰³, V. Kozhuharov³⁵, G. Kozlov³⁸, I. Králik⁵⁹, A. Kravčáková³⁶, M.A. Krawczyk³², L. Krcaj³², F. Krizek⁸³, K. Krizkova Gajdosova³⁴, C. Krug⁶⁵, M. Krüger⁶³, E. Kryshen¹³⁹, V. Kučera⁴⁹, C. Kuhn¹²⁶, D. Kumar¹³², L. Kumar⁸⁷, N. Kumar⁸⁷, S. Kumar³², S. Kundu³², M. Kuo¹²², P. Kurashvili⁷⁶, S. Kurita⁸⁹, S. Kushpil⁸³, A. Kuznetsov¹³⁹, M.J. Kweon⁵⁷, Y. Kwon¹³⁷, S.L. La Pointe³⁸, P. La Rocca²⁶, A. Lakrathok¹⁰¹, S. Lambert⁹⁹, A.R. Landou⁷⁰, R. Langoy¹¹⁸, P. Larionov³², E. Laudi³², L. Lautner⁹², R.A.N. Laveaga¹⁰⁵, R. Lavicka⁷³, R. Lea^{54,131}, J.B. Lebert³⁸, H. Lee¹⁰⁰, S. Lee⁵⁷, I. Legrand⁴⁴, G. Legras¹²³, A.M. Lejeune³⁴, T.M. Lelek², I. León Monzón¹⁰⁵, M.M. Lesch⁹², P. Lévai⁴⁵, M. Li⁶, P. Li¹⁰, X. Li¹⁰, B.E. Liang-Gilman¹⁸, J. Lien¹¹⁸, R. Lietava⁹⁷, I. Likmeta¹¹², B. Lim⁵⁵, H. Lim¹⁶, S.H. Lim¹⁶, Y.N. Lima¹⁰⁶, S. Lin¹⁰, V. Lindenstruth³⁸, C. Lippmann⁹⁴, D. Liskova¹⁰², D.H. Liu⁶, J. Liu¹¹⁵, Y. Liu⁶, G.S.S. Liveraro¹⁰⁷, I.M. Lofnes^{20,37}, C. Loizides²⁰, S. Lokos¹⁰³, J. Lömker⁵⁸, X. Lopez¹²⁴, E. López Torres⁷, C. Lotteau¹²⁵, P. Lu¹¹⁶, W. Lu⁶, Z. Lu¹⁰, O. Lubyets⁹⁴, G.A. Lucia²⁹, F.V. Lugo⁶⁶, J. Luo³⁹, G. Luparello⁵⁶, J. M. Friedrich⁹², Y.G. Ma³⁹, V. Machacek⁸⁰, M. Mager³², M. Mahlein⁹², A. Maire¹²⁶, E. Majerz², M.V. Makariev³⁵, G. Malfattore⁵⁰, N.M. Malik⁸⁸, N. Malik¹⁵, S.D. Mallick¹²⁸, N. Mallick¹¹³, G. Mandaglio^{30,52}, S. Mandal⁷⁷, S.K. Mandal⁷⁶, A. Manea⁶², R. Manhart⁹², A.K. Manna⁴⁷, F. Manso¹²⁴, G. Mantzaridis⁹², V. Manzari⁴⁹, Y. Mao⁶, R.W. Marcjan², G.V. Margagliotti²³, A. Margotti⁵⁰, A. Marín⁹⁴, C. Markert¹⁰⁴, P. Martinengo³², M.I. Martínez⁴³, M.P.P. Martins^{32,106}, S. Masciocchi⁹⁴, M. Masera²⁴, A. Masoni⁵¹, L. Massacrier¹²⁸, O. Massen⁵⁸, A. Mastroserio^{49,129}, L. Mattei^{24,124}, S. Mattiazzo²⁷, A. Matyjá¹⁰³, J.L. Mayo¹⁰⁴, F. Mazzaschi³², M. Mazzilli³¹, Y. Melikyan⁴², M. Melo¹⁰⁶, A. Menchaca-Rocha⁶⁶, J.E.M. Mendez⁶⁴, E. Meninno⁷³, M.W. Menzel^{32,91}, P.M. Meredith¹⁰⁴, M. Meres¹³, L. Micheletti⁵⁵, D. Mihai¹⁰⁹, D.L. Mihaylov⁹², A.U. Mikalsen²⁰, K. Mikhaylov¹³⁹, L. Millot⁷⁰, N. Minafra^{11,114}, D. Miśkowiec⁹⁴, A. Modak⁵⁶, B. Mohanty⁷⁷, M. Mohisin Khan^{11,15}, M.A. Molander⁴², M.M. Mondal⁷⁷, S. Monira¹³³, D.A. Moreira De Godoy¹²³, A. Morsch³², C. Moscatelli²³, T. Mrnjavac³², S. Mrozinski⁶³, V. Muccifora⁴⁸, S. Muhuri¹³², A. Mulliri²², M.G. Munhoz¹⁰⁶, R.H. Munzer⁶³, L. Musa³², J. Musinsky⁵⁹, J.W. Myrcha¹³³, B. Naik¹²⁰, A.I. Nambrath¹⁸, B.K. Nandi⁴⁶, R. Nania⁵⁰, E. Nappi⁴⁹, A.F. Nassirpour¹⁷, V. Nastase¹⁰⁹, A. Nath⁹¹, N.F. Nathanson⁸⁰, A. Neagu¹⁹, L. Nellen⁶⁴, R. Nepeivoda⁷², S. Nese¹⁹, N. Nicassio³¹, B.S. Nielsen⁸⁰, E.G. Nielsen⁸⁰, F. Noferini⁵⁰, H. Noh⁵⁷, S. Noh¹², P. Nomokonov¹³⁹, J. Norman¹¹⁵, N. Novitzky⁸⁴, J. Nystrand²⁰, M.R. Ockleton¹¹⁵, M. Ogino⁷⁴, J. Oh¹⁶, S. Oh¹⁷, A. Ohlson⁷², M. Oida⁸⁹, L.A.D. Oliveira¹⁰⁷, C. Oppedisano⁵⁵, A. Ortiz Velasquez⁶⁴, H. Osanai⁷⁴, J. Otwinowski¹⁰³, M. Oya⁸⁹, K. Oyama⁷⁴, S. Padhan¹³¹, D. Pagano^{54,131}, V. Pagliarino⁵⁵, G. Paic⁶⁴, A. Palasciano^{49,93}, I. Panasenکو⁷², P. Panigrahi⁴⁶, C. Pantouvakis²⁷, H. Park¹²², J. Park¹⁶, J. Park¹²², S. Park¹⁰⁰, T.Y. Park¹³⁷, J.E. Parkkila¹³³, P.B. Pati⁸⁰, Y. Patley⁴⁶, R.N. Patra⁴⁹, J. Patter⁴⁷, B. Paul¹³², F. Pazdic⁹⁷, H. Pei⁶, T. Peitzmann⁵⁸, X. Peng^{11,53}, S. Perciballi²⁴, G.M. Perez⁷, M. Petrovici⁴⁴, S. Piano⁵⁶, M. Pika¹³, P. Pillot⁹⁹, O. Pinazza^{32,50}, C. Pinto³², S. Pisano⁴⁸, M. Płoskon⁷¹, A. Plachta¹³³, M. Planinic⁸⁶, D.K. Plociennik²,

S. Politano³², N. Poljak⁸⁶, A. Pop⁴⁴, S. Porteboeuf-Houssais¹²⁴, J.S. Potgieter¹¹⁰, I.Y. Pozos⁴³, K.K. Pradhan⁴⁷, S.K. Prasad⁴, S. Prasad^{45,47}, R. Preghenella⁵⁰, F. Prino⁵⁵, C.A. Pruneau¹³⁴, M. Puccio³², S. Pucillo²⁸, S. Pulawski¹¹⁷, L. Quaglia²⁴, A.M.K. Radhakrishnan⁴⁷, S. Ragoni¹⁴, A. Rai¹³⁵, A. Rakotozafindrabe¹²⁷, N. Ramasubramanian¹²⁵, L. Ramello^{55,130}, C.O. Ramírez-Álvarez⁴³, E. Rao¹⁸, M. Rasa²⁶, S.S. Räsänen⁴², R. Rath⁹⁴, M.P. Rauch²⁰, I. Ravasenga³², M. Razza²⁵, K.F. Read^{84,119}, C. Reckziegel¹⁰⁸, A.R. Redelbach³⁸, K. Redlich^{88,76}, H.D. Regules-Medel⁴³, A. Rehman²⁰, F. Reidt³², H.A. Reme-Ness³⁷, K. Reygers⁹¹, M. Richter²⁰, A.A. Riedel⁹², W. Riegler³², A.G. Riffero²⁴, M. Rignanes²⁷, C. Ripoli²⁸, C. Ristea⁶², M. Rodríguez Cahuantzi⁴³, K. Røed¹⁹, E. Rogochaya¹³⁹, D. Rohr³², D. Röhrich²⁰, S. Rojas Torres³⁴, P.S. Rokita¹³³, G. Romanenko²⁵, F. Ronchetti³², D. Rosales Herrera⁴³, K. Roslon¹³³, A. Rossi⁵³, A. Roy⁴⁷, A. Roy¹¹⁸, S. Roy⁴⁶, N. Rubini⁵⁰, O. Rubza¹⁵, J.A. Rudolph⁸¹, D. Ruggiano¹³³, R. Rui²³, P.G. Russek², A. Rustamov⁷⁸, A. Rybicki¹⁰³, L.C.V. Ryder¹¹⁴, G. Ryu⁶⁹, J. Ryu¹⁶, W. Rzesza⁹², B. Sabiu⁵⁰, R. Sadek⁷¹, S. Sadhu⁴¹, A. Saha³¹, S. Saha^{46,77}, B. Sahoo⁴⁷, R. Sahoo⁴⁷, D. Sahu⁶⁴, P.K. Sahu⁶⁰, J. Saini¹³², S. Sakai¹²², S. Sambyal⁸⁸, D. Samitz⁷³, I. Sanna³², D. Sarkar⁸⁰, V. Sarritzu²², V.M. Sarti⁹², M.H.P. Sas⁸¹, U. Savino²⁴, S. Sawan⁷⁷, E. Scapparone⁵⁰, J. Schambach⁸⁴, H.S. Scheid³², C. Schiaua⁴⁴, R. Schicker⁹¹, F. Schlepper^{32,91}, A. Schmah⁹⁴, C. Schmidt⁹⁴, M. Schmidt⁹⁰, J. Schoengarth⁶³, R. Schotter⁷³, A. Schröter³⁸, J. Schukraft³², K. Schweda⁹⁴, G. Scioli²⁵, E. Scomparin⁵⁵, J.E. Seger¹⁴, D. Sekihata¹²¹, M. Selina⁸¹, I. Selyuzhenkov⁹⁴, S. Senyukov¹²⁶, J.J. Seo⁹¹, L. Serkin^{IX,64}, L. Šerkšnytė³², A. Sevcenco⁶², T.J. Shaba⁶⁷, A. Shabetai⁹⁹, R. Shahoyan³², B. Sharma⁸⁸, D. Sharma⁴⁶, H. Sharma⁵³, M. Sharma⁸⁸, S. Sharma⁸⁸, T. Sharma⁴⁰, U. Sharma⁸⁸, O. Sheibani¹³⁴, K. Shigaki⁸⁹, M. Shimomura⁷⁵, Q. Shou³⁹, S. Siddhanta⁵¹, T. Siemiarz⁷⁶, L.L.D. Silva¹⁰⁶, T.F. Silva¹⁰⁶, W.D. Silva¹⁰⁶, D. Silvermyr⁷², T. Simantathammakul¹⁰¹, R. Simeonov³⁵, B. Singh⁴⁶, B. Singh⁸⁸, K. Singh⁴⁷, R. Singh⁷⁷, R. Singh⁵³, S. Singh¹⁵, T. Sinha⁹⁶, B. Sitar¹³, M. Sitta^{55,130}, T.B. Skaali¹⁹, G. Skorodumovs⁹¹, N. Smirnov¹³⁵, K.L. Smith¹⁶, F.M.A. Smits¹¹³, R.J.M. Snellings⁵⁸, E.H. Solheim¹⁹, S. Solokhin⁸¹, C. Sonnabend^{32,94}, J.M. Sonneveld⁸¹, F. Soramel²⁷, A.B. Soto-Hernandez⁸⁵, L.E. Spencer¹⁰⁴, R. Spijkers⁸¹, C. Sporer¹¹³, I. Sputowska¹⁰³, J. Staa⁷², J. Stachel⁹¹, L.L. Stahl¹⁰⁶, I. Stan⁶², A.G. Stejskal¹¹⁴, T. Stellhorn¹²³, S.F. Stiefelmaier⁹¹, D. Stocco⁹⁹, I. Storehaug¹⁹, M.M. Stortvedt³⁷, N.J. Strangmann⁶³, P. Stratmann¹²³, S. Strazzi²⁵, A. Sturniolo^{30,52,115}, Y. Su⁶, A.A.P. Suaide¹⁰⁶, C. Suire¹²⁸, A. Suiui¹⁰⁹, M. Suljić³², V. Sumberia⁸⁸, S. Sumowidagdo⁷⁹, P. Sun¹⁰, N.B. Sundstrom⁵⁸, L.H. Tabares⁷, A. Tabikh⁷⁰, S.F. Taghavi⁹², J. Takahashi¹⁰⁷, M.A. Talamantes Johnson⁴³, G.J. Tambave⁷⁷, Z. Tang¹¹⁶, J. Tanwar⁸⁷, J.D. Tapia Takaki¹¹⁴, N. Tapus¹⁰⁹, L.A. Tarasovicova³⁶, M.G. Tarzila⁴⁴, A. Tauro³², A. Tavora García^{104,128}, G. Tejeda Muñoz⁴³, L. Terlizzi²⁴, C. Terrevoli⁴⁹, D. Thakur⁵⁵, S. Thakur⁴, M. Thogersen¹⁹, D. Thomas¹⁰⁴, A.M. Tiekoetter¹²³, N. Tiltmann^{32,123}, A.R. Timmins¹¹², A. Toia⁶³, R. Tokumoto⁸⁹, S. Tomassini²⁵, K. Tomohiro⁸⁹, Q. Tong⁶, V.V. Torres⁹⁹, A. Trifiro^{30,52}, T. Triloki⁹³, A.S. Triolo³², S. Tripathy⁷², T. Tripathy¹²⁴, S. Trogolo²⁴, V. Trubnikov³, W.H. Trzaska¹¹³, T.P. Trzcinski¹³³, C. Tsolanta¹⁹, R. Tu³⁹, R. Turrisi⁵³, T.S. Tveter¹⁹, K. Ullaland²⁰, B. Ulukutlu⁹², S. Upadhyaya¹⁰³, A. Uras¹²⁵, M. Urioni²³, G.L. Usai²², M. Vaid⁸⁸, M. Vala³⁶, N. Valle⁵⁴, L.V.R. van Doremalen⁵⁸, M. van Leeuwen⁸¹, R.J.G. van Weelden⁸¹, D. Varga⁴⁵, Z. Varga¹³⁵, P. Vargas Torres⁶⁴, O. Vázquez Doce⁴⁸, O. Vazquez Rueda¹¹², G. Vecil^{II,23}, P. Veen¹²⁷, E. Vercellin²⁴, R. Verma⁴⁶, R. Vértési⁴⁵, M. Verweij⁵⁸, L. Vickovic³³, Z. Vilakazi¹²⁰, A. Villani²³, C.J.D. Villiers⁶⁷, T. Virgili²⁸, M.M.O. Virta^{42,80}, A. Vodopyanov¹³⁹, M.A. Völkl⁹⁷, S.A. Voloshin¹³⁴, G. Volpe³¹, B. von Haller³², I. Vorobyev³², J. Vrláková³⁶, J. Wan³⁹, C. Wang³⁹, D. Wang³⁹, Y. Wang¹¹⁶, Y. Wang³⁹, Y. Wang⁶, Z.

Wang³⁹, F. Weiglhofer³², S.C. Wenzel³², J.P. Wessels¹²³, P.K. Wiacek², J. Wiechula⁶³, J. Wikne¹⁹, G. Wilk⁷⁶, J. Wilkinson⁹⁴, G.A. Willems¹²³, N. Wilson¹¹⁵, B. Windelband⁹¹, J. Witte⁹¹, M. Wojnar², C.I. Worek², J.R. Wright¹⁰⁴, C.-T. Wu^{6,27}, W. Wu⁹², Y. Wu¹¹⁶, K. Xiong³⁹, Z. Xiong¹¹⁶, L. Xu^{6,125}, R. Xu⁶, Z. Xue⁷¹, A. Yadav⁴¹, A.K. Yadav¹³², Y. Yamaguchi⁸⁹, S. Yang⁵⁷, S. Yang²⁰, S. Yano⁸⁹, Z. Ye⁷¹, E.R. Yeats¹⁸, J. Yi⁶, R. Yin³⁹, Z. Yin⁶, I.-K. Yoo¹⁶, J.H. Yoon⁵⁷, H. Yu¹², S. Yuan²⁰, A. Yuncu⁹¹, V. Zaccaro²³, C. Zampolli³², N. Zardoshti³², P. Závada⁶¹, B. Zhang⁹¹, C. Zhang¹²⁷, M. Zhang^{6,124}, M. Zhang^{6,27}, S. Zhang³⁹, X. Zhang⁶, Y. Zhang¹¹⁶, Y. Zhang¹¹⁶, Z. Zhang⁶, M. Zhao¹⁰, D. Zhou⁶, Y. Zhou⁸⁰, Z. Zhou³⁹, J. Zhu³⁹, S. Zhu^{94,116}, Y. Zhu⁶, A. Zingaretti²⁷, S.C. Zugravel⁵⁵, N. Zurlo^{54,131}

Affiliation Notes

^I Also at INFN Trieste, Trieste, Italy

^{II} Also at Fondazione Bruno Kessler (FBK), Trento, Italy

^{III} Also at Czech Technical University in Prague, Prague, Czech Republic

^{IV} Also at Instituto de Física da Universidade de São Paulo

^V Also at Dipartimento DET del Politecnico di Torino, Turin, Italy

^{VI} Also at University College of Dublin, Dublin, Ireland

^{VII} Also at Department of Applied Physics, Aligarh Muslim University, Aligarh, India

^{VIII} Also at Institute of Theoretical Physics, University of Wrocław, Wrocław, Poland

^{IX} Also at Facultad de Ciencias, Universidad Nacional Autónoma de México, Mexico City, Mexico

^X Deceased

Collaboration Institutes

¹ A.I. Alikhanyan National Science Laboratory (Yerevan Physics Institute) Foundation, Yerevan, Armenia

² AGH University of Krakow, Cracow, Poland

³ Bogolyubov Institute for Theoretical Physics, National Academy of Sciences of Ukraine, Kyiv, Ukraine

⁴ Bose Institute, Department of Physics and Centre for Astroparticle Physics and Space Science (CAPSS), Kolkata, India

⁵ California Polytechnic State University, San Luis Obispo, California, United States

⁶ Central China Normal University, Wuhan, China

⁷ Centro de Aplicaciones Tecnológicas y Desarrollo Nuclear (CEADEN), Havana, Cuba

⁸ Centro de Investigación y de Estudios Avanzados (CINVESTAV), Mexico City, Mérida, Mexico

⁹ Chicago State University, Chicago, Illinois, United States

¹⁰ China Nuclear Data Center, China Institute of Atomic Energy, Beijing, China

¹¹ China University of Geosciences, Wuhan, China

¹² Chungbuk National University, Cheongju, Republic of Korea

¹³ Comenius University Bratislava, Faculty of Mathematics, Physics and Informatics, Bratislava, Slovak Republic

¹⁴ Creighton University, Omaha, Nebraska, United States

¹⁵ Department of Physics, Aligarh Muslim University, Aligarh, India

¹⁶ Department of Physics, Pusan National University, Pusan, Republic of Korea

¹⁷ Department of Physics, Sejong University, Seoul, Republic of Korea

¹⁸ Department of Physics, University of California, Berkeley, California, United States

¹⁹ Department of Physics, University of Oslo, Oslo, Norway

²⁰ Department of Physics and Technology, University of Bergen, Bergen, Norway

²¹ Dipartimento di Fisica, Università di Pavia, Pavia, Italy

²² Dipartimento di Fisica dell'Università, and Sezione INFN, Cagliari, Italy

- ²³ Dipartimento di Fisica dell'Università and Sezione INFN, Trieste, Italy
- ²⁴ Dipartimento di Fisica dell'Università and Sezione INFN, Turin, Italy
- ²⁵ Dipartimento di Fisica e Astronomia dell'Università and Sezione INFN, Bologna, Italy
- ²⁶ Dipartimento di Fisica e Astronomia dell'Università and Sezione INFN, Catania, Italy
- ²⁷ Dipartimento di Fisica e Astronomia dell'Università and Sezione INFN, Padova, Italy
- ²⁸ Dipartimento di Fisica È.R. Caianiello' dell'Università and Gruppo Collegato INFN, Salerno, Italy
- ²⁹ Dipartimento DISAT del Politecnico and Sezione INFN, Turin, Italy
- ³⁰ Dipartimento di Scienze MIFT, Università di Messina, Messina, Italy
- ³¹ Dipartimento Interateneo di Fisica 'M. Merlin' and Sezione INFN, Bari, Italy
- ³² European Organization for Nuclear Research (CERN), Geneva, Switzerland
- ³³ Faculty of Electrical Engineering, Mechanical Engineering and Naval Architecture, University of Split, Split, Croatia
- ³⁴ Faculty of Nuclear Sciences and Physical Engineering, Czech Technical University in Prague, Prague, Czech Republic
- ³⁵ Faculty of Physics, Sofia University, Sofia, Bulgaria
- ³⁶ Faculty of Science, P.J. Šafárik University, Košice, Slovak Republic
- ³⁷ Faculty of Technology, Environmental and Social Sciences, Bergen, Norway
- ³⁸ Frankfurt Institute for Advanced Studies, Johann Wolfgang Goethe-Universität Frankfurt, Frankfurt, Germany
- ³⁹ Fudan University, Shanghai, China
- ⁴⁰ Gauhati University, Department of Physics, Guwahati, India
- ⁴¹ Helmholtz-Institut für Strahlen- und Kernphysik, Rheinische Friedrich-Wilhelms-Universität Bonn, Bonn, Germany
- ⁴² Helsinki Institute of Physics (HIP), Helsinki, Finland
- ⁴³ High Energy Physics Group, Universidad Autónoma de Puebla, Puebla, Mexico
- ⁴⁴ Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest, Romania
- ⁴⁵ HUN-REN Wigner Research Centre for Physics, Budapest, Hungary
- ⁴⁶ Indian Institute of Technology Bombay (IIT), Mumbai, India
- ⁴⁷ Indian Institute of Technology Indore, Indore, India
- ⁴⁸ INFN, Laboratori Nazionali di Frascati, Frascati, Italy
- ⁴⁹ INFN, Sezione di Bari, Bari, Italy
- ⁵⁰ INFN, Sezione di Bologna, Bologna, Italy
- ⁵¹ INFN, Sezione di Cagliari, Cagliari, Italy
- ⁵² INFN, Sezione di Catania, Catania, Italy
- ⁵³ INFN, Sezione di Padova, Padova, Italy
- ⁵⁴ INFN, Sezione di Pavia, Pavia, Italy
- ⁵⁵ INFN, Sezione di Torino, Turin, Italy
- ⁵⁶ INFN, Sezione di Trieste, Trieste, Italy
- ⁵⁷ Inha University, Incheon, Republic of Korea
- ⁵⁸ Institute for Gravitational and Subatomic Physics (GRASP), Utrecht University/Nikhef, Utrecht, Netherlands
- ⁵⁹ Institute of Experimental Physics, Slovak Academy of Sciences, Košice, Slovak Republic
- ⁶⁰ Institute of Physics, Homi Bhabha National Institute, Bhubaneswar, India
- ⁶¹ Institute of Physics of the Czech Academy of Sciences, Prague, Czech Republic
- ⁶² Institute of Space Science (ISS), Bucharest, Romania
- ⁶³ Institut für Kernphysik, Johann Wolfgang Goethe-Universität Frankfurt, Frankfurt, Germany
- ⁶⁴ Instituto de Ciencias Nucleares, Universidad Nacional Autónoma de México, Mexico City, Mexico
- ⁶⁵ Instituto de Física, Universidade Federal do Rio Grande do Sul (UFRGS), Porto Alegre, Brazil
- ⁶⁶ Instituto de Física, Universidad Nacional Autónoma de México, Mexico City, Mexico
- ⁶⁷ iThemba LABS, National Research Foundation, Somerset West, South Africa
- ⁶⁸ Jeonbuk National University, Jeonju, Republic of Korea
- ⁶⁹ Korea Institute of Science and Technology Information, Daejeon, Republic of Korea
- ⁷⁰ Laboratoire de Physique Subatomique et de Cosmologie, Université Grenoble-Alpes, CNRS-IN2P3, Grenoble, France
- ⁷¹ Lawrence Berkeley National Laboratory, Berkeley, California, United States
- ⁷² Lund University Department of Physics, Division of Particle Physics, Lund, Sweden
- ⁷³ Marietta Blau Institute, Vienna, Austria
- ⁷⁴ Nagasaki Institute of Applied Science, Nagasaki, Japan
- ⁷⁵ Nara Women's University (NWU), Nara, Japan
- ⁷⁶ National Centre for Nuclear Research, Warsaw, Poland
- ⁷⁷ National Institute of Science Education and Research, Homi Bhabha National Institute, Jatni, India
- ⁷⁸ National Nuclear Research Center, Baku, Azerbaijan
- ⁷⁹ National Research and Innovation Agency - BRIN, Jakarta, Indonesia
- ⁸⁰ Niels Bohr Institute, University of Copenhagen, Copenhagen, Denmark
- ⁸¹ Nikhef, National institute for subatomic physics, Amsterdam, Netherlands
- ⁸² Nuclear Physics Group, STFC Daresbury Laboratory, Daresbury, United Kingdom
- ⁸³ Nuclear Physics Institute of the Czech Academy of Sciences, Husinec-Řež, Czech Republic
- ⁸⁴ Oak Ridge National Laboratory, Oak Ridge, Tennessee, United States
- ⁸⁵ Ohio State University, Columbus, Ohio, United States
- ⁸⁶ Physics department, Faculty of science, University of Zagreb, Zagreb, Croatia
- ⁸⁷ Physics Department, Panjab University, Chandigarh, India
- ⁸⁸ Physics Department, University of Jammu, Jammu, India
- ⁸⁹ Physics Program and International Institute for Sustainability with Knotted Chiral Meta Matter (WPI-SKCM²), Hiroshima University, Hiroshima, Japan
- ⁹⁰ Physikalisches Institut, Eberhard-Karls-Universität Tübingen, Tübingen, Germany
- ⁹¹ Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany
- ⁹² Physik Department, Technische Universität München, Munich, Germany
- ⁹³ Politecnico di Bari, Bari, Italy
- ⁹⁴ Research Division and ExtreMe Matter Institute EMMI, GSI Helmholtzzentrum für Schwerionenforschung GmbH, Darmstadt, Germany
- ⁹⁵ Saga University, Saga, Japan
- ⁹⁶ Saha Institute of Nuclear Physics, Homi Bhabha National Institute, Kolkata, India
- ⁹⁷ School of Physics and Astronomy, University of Birmingham, Birmingham, United Kingdom
- ⁹⁸ Sección Física, Departamento de Ciencias, Pontificia Universidad Católica del Perú, Lima, Peru
- ⁹⁹ SUBATECH, IMT Atlantique, Nantes Université, CNRS-IN2P3, Nantes, France
- ¹⁰⁰ Sungkyunkwan University, Suwon City, Republic of Korea
- ¹⁰¹ Suranaree University of Technology, Nakhon Ratchasima, Thailand
- ¹⁰² Technical University of Košice, Košice, Slovak Republic
- ¹⁰³ The Henryk Niewodniczanski Institute of Nuclear Physics, Polish Academy of Sciences, Cracow, Poland
- ¹⁰⁴ The University of Texas at Austin, Austin, Texas, United States
- ¹⁰⁵ Universidad Autónoma de Sinaloa, Culiacán, Mexico
- ¹⁰⁶ Universidade de São Paulo (USP), São Paulo, Brazil
- ¹⁰⁷ Universidade Estadual de Campinas (UNICAMP), Campinas, Brazil
- ¹⁰⁸ Universidade Federal do ABC, Santo Andre, Brazil

- ¹⁰⁹ Universitatea Nationala de Stiinta si Tehnologie Politehnica Bucuresti, Bucharest, Romania
- ¹¹⁰ University of Cape Town, Cape Town, South Africa
- ¹¹¹ University of Derby, Derby, United Kingdom
- ¹¹² University of Houston, Houston, Texas, United States
- ¹¹³ University of Jyväskylä, Jyväskylä, Finland
- ¹¹⁴ University of Kansas, Lawrence, Kansas, United States
- ¹¹⁵ University of Liverpool, Liverpool, United Kingdom
- ¹¹⁶ University of Science and Technology of China, Hefei, China
- ¹¹⁷ University of Silesia in Katowice, Katowice, Poland
- ¹¹⁸ University of South-Eastern Norway, Kongsberg, Norway
- ¹¹⁹ University of Tennessee, Knoxville, Tennessee, United States
- ¹²⁰ University of the Witwatersrand, Johannesburg, South Africa
- ¹²¹ University of Tokyo, Tokyo, Japan
- ¹²² University of Tsukuba, Tsukuba, Japan
- ¹²³ Universität Münster, Institut für Kernphysik, Münster, Germany
- ¹²⁴ Université Clermont Auvergne, CNRS/IN2P3, LPC, Clermont-Ferrand, France
- ¹²⁵ Université de Lyon, CNRS/IN2P3, Institut de Physique des 2 Infinis de Lyon, Lyon, France
- ¹²⁶ Université de Strasbourg, CNRS, IPHC UMR 7178, F-67000 Strasbourg, France, Strasbourg, France
- ¹²⁷ Université Paris-Saclay, Centre d'Etudes de Saclay (CEA), IRFU, Département de Physique Nucléaire (DPHN), Saclay, France
- ¹²⁸ Université Paris-Saclay, CNRS/IN2P3, IJCLab, Orsay, France
- ¹²⁹ Università degli Studi di Foggia, Foggia, Italy
- ¹³⁰ Università del Piemonte Orientale, Vercelli, Italy
- ¹³¹ Università di Brescia, Brescia, Italy
- ¹³² Variable Energy Cyclotron Centre, Homi Bhabha National Institute, Kolkata, India
- ¹³³ Warsaw University of Technology, Warsaw, Poland
- ¹³⁴ Wayne State University, Detroit, Michigan, United States
- ¹³⁵ Yale University, New Haven, Connecticut, United States
- ¹³⁶ Yildiz Technical University, Istanbul, Turkey
- ¹³⁷ Yonsei University, Seoul, Republic of Korea
- ¹³⁸ Affiliated with an institute formerly covered by a cooperation agreement with CERN
- ¹³⁹ Affiliated with an international laboratory covered by a cooperation agreement with CERN

References

- [1] ALICE Collaboration, S. Acharya, et al., Multiplicity dependence of light-flavor hadron production in pp collisions at $\sqrt{s} = 7$ TeV, Phys. Rev. C 99 (2) (2019) 024906. [arXiv:1807.11321](#) [nucl-ex], <https://doi.org/10.1103/PhysRevC.99.024906>
- [2] ALICE Collaboration, S. Acharya, et al., Charged-particle production as a function of multiplicity and transverse sphericity in pp collisions at $\sqrt{s} = 5.02$ and 13 TeV, Eur. Phys. J. C 79 (10) (2019) 857. [arXiv:1905.07208](#) [nucl-ex], <https://doi.org/10.1140/epjc/s10052-019-7350-y>
- [3] ALICE Collaboration, J. Adam, et al., Centrality dependence of particle production in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, Phys. Rev. C 91 (6) (2015) 064905. [arXiv:1412.6828](#) [nucl-ex], <https://doi.org/10.1103/PhysRevC.91.064905>
- [4] ALICE Collaboration, B. Abelev, et al., Centrality dependence of charged particle production at large transverse momentum in Pb–Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, Phys. Lett. B 720 (2013) 52–62. [arXiv:1208.2711](#) [hep-ex], <https://doi.org/10.1016/j.physletb.2013.01.051>
- [5] B. Abelev, et al., ALICE Collaboration, Pion, kaon, and proton production in central Pb–Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, Phys. Rev. Lett. 109 (2012) 252301. [arXiv:1208.1974](#) [hep-ex], <https://doi.org/10.1103/PhysRevLett.109.252301>
- [6] ALICE Collaboration, S. Acharya, et al., The ALICE experiment: a journey through QCD, Eur. Phys. J. C 84 (8) (2024) 813. [arXiv:2211.04384](#) [nucl-ex], <https://doi.org/10.1140/epjc/s10052-024-12935-y>
- [7] ALICE Collaboration, S. Acharya, et al., Observation of partonic flow in proton-proton and proton-nucleus collisions (2024). [arXiv:2411.09323](#) [nucl-ex].
- [8] CMS Collaboration, V. Khachatryan, et al., Observation of long-range near-side angular correlations in proton-proton collisions at the LHC, JHEP 09 (2010) 091. [arXiv:1009.4122](#) [hep-ex], [https://doi.org/10.1007/JHEP09\(2010\)091](https://doi.org/10.1007/JHEP09(2010)091)
- [9] G. Aad, et al., ATLAS Collaboration, Observation of long-range elliptic azimuthal anisotropies in $\sqrt{s} = 13$ and 2.76 TeV pp collisions with the ATLAS detector, Phys. Rev. Lett. 116 (17) (2016) 172301. [arXiv:1509.04776](#) [hep-ex], <https://doi.org/10.1103/PhysRevLett.116.172301>
- [10] ALICE Collaboration, J. Adam, et al., Enhanced production of multi-strange hadrons in high-multiplicity proton-proton collisions, Nat. Phys. 13 (2017) 535–539. [arXiv:1606.07424](#) [nucl-ex], <https://doi.org/10.1038/nphys4111>
- [11] ALICE Collaboration, B.B. Abelev, et al., Multiplicity dependence of pion, kaon, proton and lambda production in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, Phys. Lett. B 728 (2014) 25–38. [arXiv:1307.6796](#) [nucl-ex], <https://doi.org/10.1016/j.physletb.2013.11.020>
- [12] ALICE Collaboration, S. Acharya, et al., Multiplicity dependence of (multi-)strange hadron production in proton-proton collisions at $\sqrt{s} = 13$ TeV, Eur. Phys. J. C 80 (2) (2020) 167. [arXiv:1908.01861](#) [nucl-ex], <https://doi.org/10.1140/epjc/s10052-020-7673-8>
- [13] B. Muller, J. Schukraft, B. Wyslouch, First results from Pb + Pb collisions at the LHC, Ann. Rev. Nucl. Part. Sci. 62 (2012) 361–386. [arXiv:1202.3233](#) [hep-ex], <https://doi.org/10.1146/annurev-nucl-102711-094910>
- [14] R.J. Fries, B. Muller, C. Nonaka, S.A. Bass, Hadronization in heavy ion collisions: recombination and fragmentation of partons, Phys. Rev. Lett. 90 (2003) 202303. [arXiv:nucl-th/0301087](#), <https://doi.org/10.1103/PhysRevLett.90.202303>
- [15] P. Bozek, Hydrodynamic flow from RHIC to LHC, Acta Phys. Polon. B 43 (2012) 689. [arXiv:1111.4398](#) [nucl-th], <https://doi.org/10.5506/APhysPolB.43.689>
- [16] ALICE Collaboration, S. Acharya, et al., Production of Λ and K_S^0 in jets in p–Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV and pp collisions at $\sqrt{s} = 7$ TeV, Phys. Lett. B 827 (2022) 136984. [arXiv:2105.04890](#) [nucl-ex], <https://doi.org/10.1016/j.physletb.2022.136984>
- [17] S. Acharya, et al., ALICE Collaboration, Production of K_S^0 , Λ ($\bar{\Lambda}$), $\Xi^\pm$ and $\Omega^\pm$ in jets and in the underlying event in pp and p–Pb collisions, JHEP 07 (2023) 136. [arXiv:2211.08936](#) [nucl-ex], [https://doi.org/10.1007/JHEP07\(2023\)136](https://doi.org/10.1007/JHEP07(2023)136)
- [18] ALICE Collaboration, S. Acharya, et al., Production of light-flavor hadrons in pp collisions at $\sqrt{s} = 7$ and $\sqrt{s} = 13$ TeV, Eur. Phys. J. C 81 (3) (2021) 256. [arXiv:2005.11120](#) [nucl-ex], <https://doi.org/10.1140/epjc/s10052-020-08690-5>
- [19] ALICE Collaboration, K. Aamodt, et al., The ALICE experiment at the CERN LHC, JINST 3 (2008) S08002. <https://doi.org/10.1088/1748-0221/3/08/S08002>
- [20] B.B. Abelev, et al., ALICE Collaboration, Performance of the ALICE experiment at the CERN LHC, Int. J. Mod. Phys. A 29 (2014) 1430044. [arXiv:1402.4476](#) [nucl-ex], <https://doi.org/10.1142/S0217751X14300440>
- [21] ALICE Collaboration, E. Abbas, et al., Performance of the ALICE VZERO system, JINST 8 (2013) P10016. [arXiv:1306.3130](#) [nucl-ex], <https://doi.org/10.1088/1748-0221/8/10/P10016>
- [22] ALICE Collaboration, B. Abelev, et al., Technical design report for the upgrade of the ALICE inner tracking system, J. Phys. G 41 (2014) 087002. <https://doi.org/10.1088/0954-3899/41/8/087002>
- [23] ALICE Collaboration, G. Dellacasa, et al., ALICE: technical design report of the time projection chamber (2000). CERN-LHCC-2000-001.
- [24] ALICE Collaboration, G. Dellacasa, et al., ALICE technical design report of the time-of-flight system (TOF) (2000). CERN-LHCC-2000-012.
- [25] ALICE Collaboration, S. Strazzi, 15 Years young: the ALICE TOF detector in the LHC run 3, Nucl. Instrum. Meth. A 1077 (2025) 170521. <https://doi.org/10.1016/j.nima.2025.170521>
- [26] T. Sjöstrand, S. Ask, J.R. Christiansen, R. Corke, N. Desai, P. Ilten, S. Mrenna, P. Prestel, C.O. Rasmussen, P.Z. Skands, An introduction to PYTHIA 8.2, Comput. Phys. Commun. 191 (2015) 159–177. [arXiv:1410.3012](#) [hep-ph], <https://doi.org/10.1016/j.cpc.2015.01.024>
- [27] P. Skands, S. Carrazza, J. Rojo, Tuning PYTHIA 8.1: the Monash 2013 tune, Eur. Phys. J. C 74 (8) (2014) 3024. [arXiv:1404.5630](#) [hep-ph], <https://doi.org/10.1140/epjc/s10052-014-3024-y>
- [28] R. Brun, F. Bruyant, F. Carminati, S. Giani, M. Maire, A. McPherson, G. Patrick, L. Urban, GEANT detector description and simulation tool (1994). CERN-W5013. <https://doi.org/10.17181/CERN.MUHF.DMJ1>
- [29] ALICE Collaboration, S. Acharya, et al., The ALICE definition of primary particles (2017). ALICE-PUBLIC-2017-005.
- [30] S. Navas, et al., Particle Data Group Collaboration, Review of particle physics, Phys. Rev. D 110 (3) (2024) 030001. <https://doi.org/10.1103/PhysRevD.110.030001>
- [31] ALICE Collaboration, S. Acharya, et al., K_S^0 and (anti-) Λ -hadron correlations in pp collisions at $\sqrt{s} = 13$ TeV, Eur. Phys. J. C 81 (10) (2021) 945. [arXiv:2107.11209](#) [nucl-ex], <https://doi.org/10.1140/epjc/s10052-021-09678-5>
- [32] S.D. Ellis, M. Jacob, P.V. Landshoff, Jets and correlations in large p(T) reactions, Nucl. Phys. B 108 (1976) 93. [https://doi.org/10.1016/0550-3213\(76\)90126-7](https://doi.org/10.1016/0550-3213(76)90126-7)
- [33] M.J. Tannenbaum, Review of hard scattering and jet analysis, PoS CFRNC2006 (2006) 001. [arXiv:nucl-ex/0611008](#), <https://doi.org/10.22323/1.030.0001>
- [34] R. Barlow, Systematic errors: facts and fictions, in: Conference on Advanced Statistical Techniques in Particle Physics, 2002, pp. 134–144. [arXiv:hep-ex/0207026](#)
- [35] ALICE Collaboration, S. Acharya, et al., Data-driven precision determination of the material budget in ALICE, JINST 18 (11) (2023) P11032. [arXiv:2303.15317](#) [physics.ins-det], <https://doi.org/10.1088/1748-0221/18/11/P11032>
- [36] ALICE Collaboration, S. Acharya, et al., Measurement of inclusive charged-particle jet production in pp and p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, JHEP 05 (2024) 041. [arXiv:2307.10860](#) [nucl-ex], [https://doi.org/10.1007/JHEP05\(2024\)041](https://doi.org/10.1007/JHEP05(2024)041)
- [37] ALICE Collaboration, S. Acharya, et al., Measurement of charged jet cross section in pp collisions at $\sqrt{s} = 5.02$ TeV, Phys. Rev. D 100 (9) (2019) 092004. [arXiv:1905.02536](#) [nucl-ex], <https://doi.org/10.1103/PhysRevD.100.092004>
- [38] ALICE Collaboration, S. Acharya, et al., Multiplicity dependence of charged-particle intra-jet properties in pp collisions at $\sqrt{s} = 13$ TeV, Eur. Phys. J. C 84 (10) (2024) 1079. [arXiv:2311.13322](#) [hep-ex], <https://doi.org/10.1140/epjc/s10052-024-13228-0>

- [39] C. Bierlich, et al., A comprehensive guide to the physics and usage of PYTHIA 8.3, SciPost Phys. Codeb. 2022 (2022) 8. [arXiv:2203.11601](https://arxiv.org/abs/2203.11601) [hep-ph], <https://doi.org/10.21468/SciPostPhysCodeb.8>
- [40] Z.-W. Lin, C.M. Ko, B.-A. Li, B. Zhang, S. Pal, A multi-phase transport model for relativistic heavy ion collisions, Phys. Rev. C 72 (2005) 064901. [arXiv:nucl-th/0411110](https://arxiv.org/abs/nucl-th/0411110). <https://doi.org/10.1103/PhysRevC.72.064901>
- [41] X.-N. Wang, M. Gyulassy, HIJING: a Monte Carlo model for multiple jet production in pp, pA and AA collisions, Phys. Rev. D 44 (1991) 3501–3516. <https://doi.org/10.1103/PhysRevD.44.3501>
- [42] B. Zhang, ZPC 1.0.1: a parton cascade for ultrarelativistic heavy ion collisions, Comput. Phys. Commun. 109 (1998) 193–206. [arXiv:nucl-th/9709009](https://arxiv.org/abs/nucl-th/9709009), [https://doi.org/10.1016/S0010-4655\(98\)00010-1](https://doi.org/10.1016/S0010-4655(98)00010-1)