



Barnacle taphonomy and ultrastructure: comparing modern and fossil shells from the western Mediterranean

GIULIA BOSIO, ALBERTO COLLARETA, LUCA PELLEGRINO, GIOVANNI BIANUCCI, GIANNI INSACCO AND GIOVANNI COLETTI

LETHAIA



Fossil acorn barnacles are excellent palaeoenvironmental proxies in shallow-marine settings. However, the impact of post-depositional and diagenetic alterations on the barnacle shell remains poorly understood. Opercular plate preservation and observations on abrasion and fragmentation of the wall plates are the most valued taphonomic indicators. Although the shell microstructure of marine invertebrates usually provides useful insights for understanding these processes, conventional microscopic observations are often insufficient for assessing the degree of micro- and ultrastructural alteration in fossil barnacles. This study compares the ultrastructure of modern and fossil (Pliocene and Pleistocene) barnacle shells from the western Mediterranean to explore their ultrastructure and investigate the impact of taphonomic and diagenetic processes. Under Scanning Electron Microscopy (SEM), modern barnacles show a porous 'dragon-scale pattern'. SEM and cathodoluminescence analyses reveal microstructural changes in fossil barnacles, including dissolution features and diagenetic imprints. Evidence of microbial activity is also observed, in both modern and fossil specimens, in the form of microbioerosion traces made by cyanobacteria. Despite their stable low-magnesium calcite composition, barnacle shells from coarse-grained deposits usually show significant diagenetic alteration. In contrast, specimens from fine-grained sediments typically exhibit minimal alteration, preserving ultrastructural features almost comparable to those of their modern counterparts. This study underscores the importance of high-resolution analytical approaches in palaeontological research to advance our understanding of barnacle ultrastructure, taphonomy and diagenesis. □ *Fossil-diagenesis, microstructure, cathodoluminescence, Chelonibia, Concavus, Perforatus*

Giulia Bosio ✉ [giulia.bosio.giulia@gmail.com], Dipartimento di Scienze della Terra, Università di Pisa, Via Santa Maria 53, 56126 Pisa, Italy; Dipartimento di Scienze dell'Ambiente e della Terra, Università degli Studi di Milano-Bicocca, Piazza della Scienza 1, 20126 Milano, Italy; Alberto Collareta [alberto.collareta@unipi.it], Dipartimento di Scienze della Terra, Università di Pisa, Via Santa Maria 53, 56126 Pisa, Italy; Luca Pellegrino [lu.pellegrino@unito.it], Dipartimento di Scienze della Terra, Università degli Studi di Torino, Via Valperga Caluso 35, 10125 Torino, Italy; Giovanni Bianucci [giovanni.bianucci@unipi.it], Dipartimento di Scienze della Terra, Università di Pisa, Via Santa Maria 53, 56126 Pisa, Italy; Gianni Insacco [g.insacco@comune.comiso.rg.it], Museo Civico di Storia Naturale di Comiso, Via degli Studi 9, 97013 Comiso, Italy; Giovanni Coletti [giovanni.p.m.coletti@gmail.com], Dipartimento di Scienze dell'Ambiente e della Terra, Università di Milano-Bicocca, Piazza della Scienza 4, 20126 Milano, Italy; manuscript received on 12/02/2025; manuscript accepted on 02/04/2025; manuscript published on 12/08/2025 in Lethaia 58(3).

Fossil acorn barnacles (Cirripedia, Balanomorpha) are valuable proxies for palaeoecological and palaeoenvironmental studies in shallow-marine settings due to their wide distribution and abundance, high growth rates, and low-magnesium calcite shell.

Today, large acorn barnacle accumulations are typically found at middle and high latitudes (Raymond & Stetson 1932; Hoskin & Nelson 1969; Newman & Ross 1971; Milliman 1972; Müller & Milliman 1973; Farrow *et al.* 1978; Hottinger 1983; Domack 1988; Nelson *et al.* 1988; Scoffin 1988; Wilson 1988; Henrich *et al.* 1995; Frank *et al.* 2014), though they can also occur in low-latitude regions, especially in nutrient-rich water (Glynn & Wellington 1983; Halfar *et al.* 2006;

Westphal *et al.* 2010; Michel *et al.* 2011; Reijmer *et al.* 2012; Klicpera *et al.* 2013; Reymond *et al.* 2016; Bialik *et al.* 2023). In spite of a general paucity of palaeontological studies on barnacles, these sessile crustaceans are common in the Cenozoic fossil record, especially in Neogene and Quaternary deposits (Darwin 1854; De Alessandri 1895; Davadie-Suaudeau 1952; Newman *et al.* 1969; Menesini 1971; Pajaud 1976; Foster & Buckeridge 1987; Menesini & Casella 1988; Donovan 1989; Doyle *et al.* 1996; Collins *et al.* 2014; Buckeridge 2015; Gale & Sørensen 2015; Coletti *et al.* 2018, 2021, 2024; Collareta *et al.* 2019; Buckeridge *et al.* 2020; Hoffmann *et al.* 2020; Gale 2021; Gale *et al.* 2021; Kočí *et al.* 2024a).

Most barnacle shells exhibit relatively high growth rates (Newman *et al.* 1967; Bourget & Crisp 1975; Crisp & Bourget 1985; Anderson 1994; Lopez *et al.* 2012), enabling them to effectively record short-term fluctuations of the environmental conditions such as seasonal variations (Buckeridge 1975; Killingley & Newman 1982). A fast growth rate can also lead to the accumulation of high numbers of shells in short time spans (Hoskin & Reed 1984; Buckeridge 1999, 2015). Consequently, barnacle shells serve as valuable and detailed indicators of environmental dynamics, providing insights into the ecological conditions during their growth (e.g. Burgess *et al.* 2010; Coletti *et al.* 2018, 2024), provided that diagenesis does not alter the original shell structure.

Barnacle shells are composed of low-Mg calcite, which is diagenetically more stable and less soluble compared to aragonite and high-Mg calcite (Clarke & Wheeler 1917; Chave 1954; Hall *et al.* 1967; Bourget 1974; Hover *et al.* 2001; Borromeo *et al.* 2017; Ullmann *et al.* 2018). Its structure consists of thin layers of calcite crystals embedded within an organic matrix – a complex, hierarchically organized structure (Bourget 1987; Khalifa *et al.* 2011; Lewis *et al.* 2014; Gal *et al.* 2015; Checa *et al.* 2019, 2020; Shaw *et al.* 2024). Biomineralization is controlled by this organic matrix that determines the type, size and orientation of the crystals that form the shell (Rodríguez-Navarro *et al.* 2006; Khalifa *et al.* 2011). This layered arrangement is not homogeneous, but rather varies in response to seasonal shifts and ontogenetic development, both of which influence the shell growth rate (Nishizaki & Carrington 2015), and thus its geochemical characteristics. Killingley & Newman (1982) found that the calcite of recent balanomorph barnacle shells exhibits systematically higher $\delta^{18}\text{O}$ values than those expected under equilibrium conditions, whereas Burgess *et al.* (2010), Bojar *et al.* (2018) and Ullmann *et al.* (2018) suggested that the cirripede shells form at isotopic equilibrium with the surrounding seawater. Furthermore, the mineralogy of barnacle shells is extremely variable not only in comparison to other marine invertebrates, but also within the Cirripedia, which also includes species with shells made of chitin and/or hydroxyl-apatite (Carpenter & Lohmann 1992; Buckeridge & Newman 2006; Khalifa *et al.* 2011; Iglíowska *et al.* 2018; Ullmann *et al.* 2018).

Therefore, although acorn barnacles present an attractive opportunity for palaeoenvironmental studies, the aforementioned ‘grey areas’ highlight how the few microstructural and geochemical analyses that have been conducted on extant (Murdock & Currey 1978; Dougherty *et al.* 1990; Fernández *et al.* 2001; Rodríguez-Navarro *et al.* 2006; Khalifa *et al.* 2011; Raman & Kumar 2011; Lewis *et al.* 2014; Bojar *et al.*

2018; Mitchell *et al.* 2019; Checa *et al.* 2019, 2020; Shaw *et al.* 2024) and fossil forms (e.g. Buckeridge 1975; Roskowski *et al.* 2010; Gale & Schweigert 2016; Collareta *et al.* 2018; Taylor *et al.* 2019; Bosio *et al.* 2020; Paces *et al.* 2023; Coletti *et al.* 2024) severely limits the potential of barnacles as (palaeo)environmental indicators, which contrasts with the attention that is given to the micro- and ultrastructure of other groups of fossil invertebrates (e.g. Barbin & Gaspard 1995; Barbin 2000; Machel 2000; Vinn *et al.* 2008a,b; Vinn & Kupriyanova 2011; Angiolini *et al.* 2012; Vinn 2013, 2020; Crippa *et al.* 2016, 2020; Casella *et al.* 2018; Bosio *et al.* 2020; Sanfilippo *et al.* 2021; Guido *et al.* 2024).

Furthermore, only few taphonomic studies are available (Donovan 1988; Donovan 1993; Doyle *et al.* 1996; Nielsen & Funder 2003; Nomura & Maeda 2008; Coletti *et al.* 2024). Based on macroscopic observations, Doyle *et al.* (1996) and then Nomura & Maeda (2008) recognized seven and four different types of macroscopic taphonomic preservation of barnacle shells, respectively. Nielsen & Funder (2003) described three grades of alteration of barnacle shells based on the extent of abrasion and fragmentation. Overall, well-preserved, complete barnacle shells with articulated or associated opercular plates seems to be typically associated with sheltered, low-energy environments, while disarticulated, fragmented and abraded specimens are related to high-energy settings (e.g. Nielsen & Funder 2002; Coletti *et al.* 2024).

To address these gaps in knowledge, this work analyses modern and fossil barnacles specimens from the Western Mediterranean using cathodoluminescence (CL) and scanning electron microscopy (SEM), with the aim of (i) investigating the barnacle ultrastructure and (ii) developing a standardized taphonomic protocol in fossil barnacles. The fossil specimens come from various Pliocene-Pleistocene barnacle-rich deposits of central and northwestern Italy, and represent different (palaeo)environmental settings (ranging from coastal and very shallow to open shelf), burial types (fine-grained versus coarse-grained deposits), geological time intervals (including the Early and Late Pliocene, and Early Pleistocene), barnacle taxa (numbering four species in two families), and ecological adaptations (including epizoic as well as non-epizoic forms). Together with fossil samples, modern barnacles from the central Mediterranean Sea were analysed for comparison.

Geographical and stratigraphical setting

The fossils analysed in this work come from deposits that span the Lower Pliocene to Lower Pleistocene




Fig. 1. Geographic position of the sampled localities. A, location of the studied Pliocene and Pleistocene outcrops (red circles) and sampling sites of all modern barnacles but *Chelonibia testudinaria* (orange circles). B, location of the study area in the Mediterranean region and sampling site of the modern *C. testudinaria* specimens.

(Fig. 1). The oldest such deposits are those of the Lower Pliocene succession cropping out at the western Ligurian locality of Pairoia (Imperia Province) (Fig. 2A, B) (Giammarino *et al.* 2010; Dalla Giovanna 2016; Coletti *et al.* 2021). Three main coarse-grained, mixed siliciclastic-bioclastic facies are identified in the studied succession, which are mainly composed of calcirudites: a barnacle-dominated facies that developed in a shallow (less than 15 m deep), nearshore palaeoenvironment; a foraminifera-dominated facies from deeper waters (ranging between 40–100 m); and an intermediate facies situated in-between (Coletti *et al.* 2021). The barnacle-rich layers date back to the early-middle Zanclean and originated in a semi-enclosed, high-energy coastal environment at water depths of less than 30 m (Coletti *et al.* 2021).

The mid-Pliocene succession of Certaldo (Florence Province) is located in the Valdelsa Basin of central Tuscany (Benvenuti *et al.* 2014). The deposits comprise mainly shelf mudstones belonging to the Argille Azzurre Formation, where several shell beds host barnacles as well as molluscs and fish remains (Fig. 2C, D). These deposits have been referred to the upper Zanclean – lower Piacenzian interval, and interpreted as resulting from offshore deposition at depths of a few tens metres (Benvenuti *et al.* 2014; Dominici *et al.* 2018).

The succession exposed at Case Nuove (also known as Casenuove - Pozzale; Florence Province) (Fig. 2E) includes mid-Piacenzian paralic sequences that deposited along the northern edge of the Valdelsa Basin (Dominici *et al.* 1995; Benvenuti *et al.* 2014) (Collareta *et al.* 2020, 2021; Merella *et al.* 2023). Within this succession, a balaenid skeleton was preserved along with many epizoic barnacles (Bianucci 1996; Collareta *et al.* 2016). The palaeoenvironment is reconstructed as a very shallow setting, likely less than 10 metres water depth, and possibly an interdistributary bay (Dominici *et al.* 1995).

Another Upper Pliocene succession is found at Montefollonico (Siena Province), in the Siena-Radicofani Basin of eastern Tuscany (Martini *et al.* 2021). The fossil-rich calcirudites exposed at this locality (Fig. 2F) are thought to have deposited in a low-energy setting along a sheltered coast (Nalin *et al.* 2016). Here, a barnacle-dominated facies has been recognized (Coletti *et al.* 2024).

Finally, the Lower Pleistocene Fauglia succession is located in the northern sector of the Tora-Fine Basin, one of the westernmost Neogene extensional basins of Tuscany (Bossio *et al.* 1999). At the Montalto Quarry, highly fossiliferous marine siltstones and sandstones are exposed (Fig. 2G). They are attributed to the Calabrian Sabbie di Nugola Vecchia Formation (Bosio




Fig. 2. Field photos of the studied outcrops. A, the Lower Pliocene succession exposed at Pairola. B, close-up of the barnacle-dominated facies of Pairola. C, the mid-Pliocene succession exposed at Certaldo. D, barnacle-rich shell bed cropping out at Certaldo. E, the Case Nuove locality, where an Upper Pliocene succession crops out in an abandoned quarry. F, detail of the Upper Pliocene, barnacle-dominated facies exposed at Montefollonico. G, the Lower Pleistocene succession cropping out at Fauglia. H, detail of a barnacle-encrusted oyster from the barnacle-rich oyster reef exposed at Fauglia.

et al. 2021). These deposits indicate a protected, marginal-marine setting, featuring a *Posidonia oceanica* meadow, a *Cladocora caespitosa* bank and an *Ostrea edulis* reef (Bosio *et al.* 2021; Mariani *et al.* 2022). The barnacle-rich layer coincides with the oyster reef horizon (Fig. 2H), which originated at some 5–10 m water depth (Bosio *et al.* 2021; Collareta *et al.* 2024).

Material

Fossil specimens of *Concavus concavus* (Bronn, 1831) were collected at Pairola (43.94°N; 8.09°E), Certaldo (43.56°N; 11.03°E) and Montefollonico (43.13°N; 11.73°E) (Fig. 1). Specimens of *Perforatus perforatus* (Bruguière, 1789) were collected at Fauglia (43.56°N; 10.53°E) (Fig. 1). Both the extinct *C. concavus* and the extant *P. perforatus* belong in the same subfamily of Balanidae, namely, Concavinae (Menesini 1965; Newman 1982; Zullo 1992). In addition, specimens of the common turtle barnacle, *Chelonibia testudinaria* (Linnaeus, 1758), were collected at Case Nuove (43.69°N; 10.95°E) (Fig. 1, 4) (Table 1).

Concavus concavus specimens were collected from the Lower Pliocene rudstones and packstones of Pairola (Fig. 4A), the oldest studied succession. According to Coletti *et al.* (2024), they are often fragmented, and usually exhibit a poor preservation state, corresponding to Type 7 of Doyle *et al.* (1996) and Type D of Nomura & Maeda (2008). Some shells are found in small clusters of relatively well-preserved specimens that have been dislodged from their substrate (Fig. 4A), matching Type 2 of Doyle *et al.* (1996) and Type C of Nomura & Maeda (2008). Opercular plates are very rare and invariably separated from their corresponding shells (Coletti *et al.* 2024).

Barnacle specimens from the mid-Pliocene deposits of Certaldo were collected from open shelf mudstones (Coletti *et al.* 2024) (Fig. 4B). According to Coletti *et al.* (2024), these barnacles are mostly found as displaced individuals, which conforms to Type 2 (Doyle *et al.* 1996) and Type C (Nomura & Maeda, 2008). In addition, rare individuals occur attached to their original shelly substrate, though the latter are no longer found in life position (see Fig. 4B; Type B of Nomura & Maeda, 2008). Disarticulated

Table 1. List of the analysed barnacle-rich and barnacle-dominated facies and localities. Modified after Coletti *et al.* (2024)

Site	Age	Type of barnacle accumulation	Setting	Water depth	Lithology	Barnacle species
Pairola	lower-middle Zanclean	Barnacle-dominated facies	High-energy, nearshore coastal setting	0–15 m	Coarse-grained, mixed siliciclastic bioclastic	<i>Concavus concavus</i>
Certaldo	upper Zanclean-lower Piacenzian	Barnacle-rich layer	Low-energy, open shelf setting	> 40 m	Shell-rich calcareous mudstone	<i>Perforatus perforatus</i>
Case Nuove	middle Piacenzian	Barnacle-rich layer	Low-energy, nearshore coastal embayment	0–10 m	Shell-rich calcareous sandy siltstone	<i>Chelonibia testudinaria</i>
Montefollonico	Piacenzian	Barnacle-dominated facies	Moderate-energy, nearshore coastal setting	20–40 m	Coarse-grained, mixed siliciclastic bioclastic	<i>Perforatus perforatus</i>
Fauglia	Calabrian	Barnacle-rich oyster reef	Moderate-energy, nearshore coastal setting (possibly a coastal embayment)	5–10 m	Coarse-grained, mixed siliciclastic bioclastic	<i>Perforatus perforatus</i>

and fragmented plates, representing Type 7 of Doyle *et al.* (1996) and Type D of Nomura & Maeda (2008), were also collected. In all these cases, whole barnacle shells as well as shell fragments are typically well preserved, retaining the original pigmentation and showing no signs of abrasion (Fig. 4B), being classified in the Grade 0 of Nielsen & Funder (2003). In addition, the fragile opercular plates (both scuta and terga) are

usually abundant, being still articulated to the corresponding shells in most cases.

In the Upper Pliocene deposits of Case Nuove, a bioclastic sandstone to siltstone layer hosts a partial mysticete (baleen whale) skeleton, in close proximity of which tens of turtle barnacle shells have been found (Fig. 4E). This association suggests that the barnacles lived as epibionts on the baleen whale, from




Fig. 3. Macroscopic images of the barnacle shells. A, generalized structure of a balanid barnacle shell in external view (modified after Buckeridge 1983). B, modern *Amphibalanus amphitrite* specimen from Imperia. C, interpretative structure of the barnacle shown in panel B. D, modern *A. amphitrite* specimen from Livorno. E, modern *Perforatus perforatus* specimens from Imperia. F, modern *P. perforatus* specimen from Marina del Boccale. G, H, modern *Chelonibia testudinaria* specimen from Pozzallo. I, generalized structure of a chelonibiine shell modified from Collareta et al. (2022).

which they detached after death as the cetacean soft tissues decayed (Collareta 2020; Coletti et al. 2024). The specimens are found both as disarticulated valves and as complete or nearly complete shells (Fig. 4E); however, the taphonomic classifications of Doyle et al. (1996) and Nomura & Maeda (2008) cannot be applied due to the epizoic nature and membraneous base of *C. testudinaria*. In any case, the shell morphology is very well preserved and evidence of abrasion was not detected (Fig. 4E), thus conforming to Grade 0 of Nielsen & Funder (2003) (Coletti et al. 2024).

At Montefollonico, *C. concavus* specimens are found in Late Miocene fine-grained calcirudite deposits. The analysed specimens usually exhibit a moderate to good preservation state, with some retaining the original pigmentation of the shell. According to Coletti et al. (2024), they are usually found as

displaced clusters, but some are still attached to the corresponding substrate (e.g. pebbles and rhodoliths), though not in life position, thus corresponding to Type 2 of Doyle et al. (1996) and Type C of Nomura & Maeda (2008), respectively (Fig. 4C). Disarticulated and fragmented plates are also common, conforming to Type 5 of Doyle et al. (1996) and Type D of Nomura & Maeda (2008). Shells also exhibit some evidence of abrasion and fragmentation, allowing for identifying Grades 0 and 1 of Nielsen & Funder (2003). Opercular plates (mainly scuta) are also commonly found sparse within the sediment.

Coletti et al. (2024) recorded *P. perforatus* from the Lower Pleistocene oyster reef of Fauglia, from which some specimens were essentially complete, including the delicate opercula (Fig. 4D). In light of this, they can be classified as Type 1 of Doyle et al. (1996) and Type A of Nomura & Maeda (2008)




Fig. 4. A, cluster of fossil *Concavus concavus* specimens from Pairola. B, fossil *C. concavus* specimen from Certaldo, retaining its pink pigmentation. C, fossil *C. concavus* specimens from Montefollonico. D, cluster of fossil *Perforatus perforatus* specimens from Fauglia. E, fossil *Chelonibia testudinaria* specimen from Case Nuove. For detailed images of the opercular plates see Coletti *et al.* (2024).

(Coletti *et al.* 2024). Moreover, shells do not display any evidence of abrasion or fragmentation (Grade 0 of Nielsen & Funder, 2003). Though the macroscopic preservation state is good overall, shells are distinctly compressed diagenetically (Fig. 4D).

As a reference point, Recent specimens of *Amphibalanus amphitrite* (Darwin, 1854) were collected from Imperia, western Liguria (43.88°N, 8.05°E), and Livorno, southern Tuscany (43.56°N, 10.31°E) (Figs 1, 3A–D). *Perforatus perforatus* was also collected at Imperia, western Liguria (43.88°N, 8.05°E) and at Marina del Boccale, southern Tuscany (43.47°N, 10.32°E) (Figs 1, 3E,F). Finally, *C. testudinaria* was obtained at Pozzallo, southern Sicily (36.72°N, 14.85°E) (Figs 1, 3G–I). All specimens apart from *C. chelonibia* – which were removed from a beached loggerhead turtle, *Caretta caretta* (Linnaeus, 1758) – come from water depths of less than one metre.

Analytical methods

Twenty-three thick and thin sections were prepared following the method outlined by Coletti *et al.* (2019) and Collareta *et al.* (2019), embedding repeatedly the barnacle shell in epoxy resin, and then proceeding with cutting transversally the shell at various heights (near the base of the shell, just above the base of the sheath and close to the apex of the shell) by using a Buehler IsoMet, low-speed, precision-cutting machine at the Università degli Studi di Milano-Bicocca.

Thin sections were observed under a Leica transmitted-light optical microscope at the Università

degli Studi di Milano-Bicocca. Seven such sections, including both modern and fossil specimens from all the studied localities, were examined using a CITL Optical CL microscope stage at the Università degli Studi di Milano-Bicocca. The microscope was operated at a voltage of 6.3 kV, with a current intensity of about 1.1 mA, under a vacuum of about 7.3 Pa.

Following Crippa *et al.* (2016), thick sections were first polished with 400- and 1000 SiC abrasive powder, and then briefly etched with diluted HCl for less than 15 seconds for highlighting structures. These sections were carbon-coated and observed for morphological analyses through a Zeiss FEG Gemini 500 SEM operating at 3.0 kV at the Università degli Studi di Milano-Bicocca. SEM images were examined in order to characterize and compare the micro- and ultrastructure of fossil and modern specimens as well as for identifying possible evidences of erosion, dissolution and recrystallization during biostratinomic and diagenetic processes.

Results

Microscopic structure

All studied barnacle shells feature an inner and an outer lamina (Fig. 5). Between the inner and outer laminae are distinct longitudinal tubes. The longitudinal tubes are separated from each other by septa, which in transverse section can be seen to form interlamine figures (Fig. 5). Except for *Chelonibia*, these reflect the complex interlocking between the wall plates and the calcareous base, which articulate with each other without being cemented to each other to allow for the growth of the shell (Bourget 1977). The parietal tubes are partially or entirely infilled secondarily (during life) by concentric rings of biogenic carbonate. The sheath usually exhibits concentric laminations (Fig. 5F, G.) The internal sutural edges, where the wall plates meet each other, do typically display interlocking tooth-like projections with different degrees of complexity. These structures provide strength and durability to the shell while permitting the growth of the body chamber. Although these first-order structural elements are shared by all the investigated specimens, both recent and fossil, some microtaphonomic considerations are still possible. In the barnacle specimens from Montefollonico, Pairola and Fauglia, the large parietal tubes are diagenetically filled by drusy crystals of carbonate cement (Fig. 5D, E, H), which suggests that they have been rather severely affected by diagenesis. In contrast, specimens from Case Nuove

and Certaldo are typified by empty parietal tubes, suggesting that diagenetic alteration was minor overall (Fig. 5G, I, J). In addition, modern specimens of both *Chelonibia testudinaria* and *Perforatus perforatus* and one fossil specimen from Certaldo have the outer lamina perforated by microborings, which however are best visible under the SEM (see paragraph below).

Ultrastructure

SEM observation of modern shells reveal that different species of epizoic and non-epizoic barnacles exhibit a similar ultrastructure, without variations at different shell heights or among the inner lamina, the outer lamina and the sheath. In particular, the modern species *Amphibalanus amphitrite* from Livorno




Fig. 5. Recent and fossil barnacle shells in transverse thin section. A, microstructure of a modern *Perforatus perforatus* specimen from Imperia. B, internal structure of the wall plate and the sheath as interpreted by the transverse section shown in panel A. C, microstructure of a modern *Amphibalanus amphitrite* specimen from Livorno. D, microstructure of a fossil *P. perforatus* specimen from Fauglia. E, microstructure of a fossil *Concavus concavus* specimen from Pairola. F, microstructure of a modern *Chelonibia testudinaria* specimen from Pozzallo. G, microstructure of a fossil *C. testudinaria* specimen from Case Nuove. H, microstructure of a fossil *C. concavus* specimen from Montefollonico. I, microstructure of a fossil *C. concavus* specimen from Certaldo. J, microstructure of the same *C. concavus* specimen as in panel I, observed along a cut slightly closer to the apex of the barnacle shell.

(Fig. 6A–C) and Imperia (Fig. 6D–F), and *Perforatus perforatus* from Imperia (Fig. 6G–I) and Marina del Boccale (Fig. 6J–L) exhibit the same crystal pattern in transverse section. At the micro-scale, the lamellar structure is revealed to consist of loosely packed, scale-like elements measuring from 1 to 5 μm in length and about 1 μm in thickness (Fig. 6). These scales are arranged in a complex 3D disposition with irregular spaces and crevices ('dragon-scale pattern' *sensu* Coletti *et al.* 2024). This structural arrangement results in the growth lines that are visible under an optical microscope at low magnification. At the nano-scale, porosity can be detected both in-between the scales and within the scales themselves (Fig. 6), thus confirming the porous nature of the barnacle shell at all levels.

Low-magnification (1.4k $\times$) SEM observations reveal no significant differences between the modern and fossil specimens (Figs 6–8). However, at higher magnification (2.8k and 9.3k $\times$), specimens of *Concavus concavus* from Certaldo (Fig. 7A–C) and Montefollonico (Fig. 7D–F) possess bulkier scales with more poorly defined margins than modern specimens. Fossil specimens from these localities show indeed the best-preserved shell structure at different magnifications. On the other side of the preservational spectrum, in the *C. concavus* specimens from Pairola, the dragon-scale pattern is entirely obliterated, with no discernible microstructural and ultrastructural patterns (Fig. 7G–I). The *P. perforatus* specimens from Fauglia exhibit variable degrees of ultrastructural preservation, with some displaying a




Fig. 6. SEM images of recent barnacles. A–C, ultrastructure of the wall plate of a modern specimen of *Amphibalanus amphitrite* from Livorno at different magnifications. D–F, ultrastructure of the wall plate of a modern specimen of *A. amphitrite* from Imperia at different magnifications. G–I, ultrastructure of the wall plate of a modern specimen of *Perforatus perforatus* from Imperia at different magnifications. J–L, ultrastructure of the sheath of a modern specimen of *P. perforatus* from Marina del Boccale at different magnifications.




Fig. 7. SEM images of well-preserved and poorly preserved fossil barnacles. A–C, ultrastructure of the wall plate of a well-preserved fossil specimen of *Concavus concavus* from Certaldo at different magnifications. D–F, ultrastructure of the sheath of a well-preserved fossil specimen of *C. concavus* from Montefollonico at different magnifications. G–I, ultrastructure of the wall plate of a poorly preserved fossil specimen of *P. perforatus* from Fauglia at different magnifications. J–L, ultrastructure of the wall plate of a poorly preserved fossil specimen of *P. perforatus* from Fauglia at different magnifications.




Fig. 8. SEM images of recent and fossil *Chelonibia testudinaria* shells. A–C, ultrastructure of the wall plate of a modern specimen from Pozzallo at different magnifications. D–F, ultrastructure of the wall plate of a moderately well-preserved fossil specimen from Case Nuove at different magnifications.

complete obliteration of the micro- and ultrastructure (Fig. 7J–L), thus resembling the Pairola specimens. The dragon-scale pattern is also clearly present in the modern *Chelonibia testudinaria* specimen from Pozzallo, resembling the non-epizoic barnacle samples in terms of both scale arrangement and scale size (Fig. 8A–C). In the fossil *C. testudinaria* specimen from Case Nuove, the ultrastructure is poorly preserved, but not totally obliterated. Scales are larger, more irregular, and have poorly defined margins compared to its modern counterpart (Fig. 8D–F). Consistent with these microstructural observation, the modern specimens exhibit intact interlaminate figures and empty parietal tubes (Fig. 9G), whereas some fossil samples such as the *P. perforatus* specimen from Fauglia show widespread recrystallization and parietal tubes that are partly filled by very large crystals in a drusy-like fashion (Fig. 9H). SEM images reveal interesting shell modification patterns. The modern specimens of *C. testudinaria* from Pozzallo (Fig. 9A–D) and *P. perforatus* from Imperia (Fig. 9E) feature the pervasive occurrence of several resin-casts of microborings that penetrate into the outer wall. They occur as filamentous structures, 2–5 µm across and up to hundreds microns long, ranging from almost straight to strongly bent in shape, which may (or may not) branch (Fig. 9A–E). Thinner microboring casts, down to less than 1 µm across, can also be observed in one specimen of *C. concavus* from Certaldo (Fig. 9F).

Cathodoluminescence

Along with the SEM observations, the cathodoluminescence analyses allowed for the recognition of different degrees of alteration in the fossil specimens compared to their modern counterparts. Recent barnacle shells, including both epizoic and non-epizoic forms, display a very weak luminescence lining the interlaminate figures, the inner wall of the parietal tubes, the outer surface of the inner lamina and the growth lines of both laminae, ranging in colour from blue to very light reddish (Figs 10A–D, 11E,F). In particular, *Perforatus perforatus* from Imperia shows a predominant low blue and purple luminescence lining the interlaminate figures and the parietal tubes (Fig. 10A, B), whereas the modern *Amphibalanus amphitrite* from Livorno exhibits a general low blue luminescence and a low reddish luminescence that concentrates around the inner wall of the parietal tubes (Fig. 10C, D). As regards the epizoic forms, the modern specimen of *Chelonibia testudinaria* from Pozzallo shows a predominant low blue luminescence, similar to that observed for the non-epizoic barnacles (Fig. 11E, F).

Fossil specimens span across a broad spectrum of luminescence types. The least luminescent sample, namely, the *C. concavus* specimen from Certaldo, displays a low purple luminescence of the interlaminate figures and a moderate red luminescence of the outer wall (Fig. 10E, F). The conspecific fossil specimen from Montefollonico shows a moderate purple to red luminescence along the inner lamina and the interlaminate figures, and luminescent (red-coloured) parietal tubes that have been filled by diagenetic calcite (Fig. 10G, H). On the other hand, the *C. concavus* specimen from Pairola displays a high, fully red-coloured luminescence, and a diagenetic filling of the parietal tubes consisting of sparitic and drusy calcite crystals (Fig. 11A, B). The *P. perforatus* specimen from Fauglia is the most luminescent: it displays a very strong, red to orange and yellow luminescence throughout the shell and parietal tubes, the latter being filled by drusy sparitic calcite (Fig. 11C, D). Similar to the moderately well-preserved barnacle specimens from Certaldo and Montefollonico, the fossil *C. testudinaria* shell from Case Nuove exhibits a moderate purple to red luminescence that concentrates along the growth lines and interlaminate figures (Fig. 11G, H).

Discussion

Macroscopic observations provide useful insights into the preservation of barnacle specimens and their depositional environments. Well-preserved, complete specimens, when not related to rapid burial, tend to be associated with sheltered, low-energy settings, whereas poorly preserved, disarticulated, fragmented remains are the rule in more exposed settings (Nielsen & Funder 2002; Coletti *et al.* 2024). A key indicator is provided by the presence/absence of the opercular plates, because these delicate structures are often displaced and lost in high-energy environments. According to Coletti *et al.* (2024), fossil barnacles from coarse-grained deposits characterized by high values of primary porosity, such as sandstones, packstones and rudstones, display a poor (at Pairola and Fauglia) to intermediate (at Montefollonico) preservation at all scales. The *Concavus concavus* specimens from Pairola and those of *Perforatus perforatus* from Montefollonico are mostly fragmented, with poor to moderate macroscopic preservation and some indications of abrasion, thus indicating moderate energy conditions, with the exception of barnacles found attached to their own substrate (at Fauglia). On the contrary, in fine-grained sediments




Fig. 9. SEM images showing taphonomic details of the analysed barnacle shells. A, microborings along the outer wall of the modern *Chelonibia testudinaria* shell from Pozzallo. B, resin-cast of the microborings occurring along the outer wall of the modern *C. testudinaria* shell from Pozzallo; they are tentatively assigned to the ichnogenera *Fascichnus*, *Scolecia* and *Eurygonum*. C, close-up of the microborings in Figure 8B. D, close-up of the microborings in Figure 8C. E, resin-cast microborings characterizing modern *Perforatus perforatus* from Imperia; they are tentatively assigned to the ichnogenera *Fascichnus*, *Scolecia* and *Eurygonum*. F, microborings occurring along the outer wall of a fossil *Concavus concavus* shell from Certaldo. G, shell ultrastructure of *Amphibalanus amphitrite* from Imperia, showing the interlaminar figures and empty parietal tubes. H, parietal tube of a fossil *P. perforatus* specimen from Fauglia filled with a drusy carbonate cement forming very large crystals.

such as those that entombed the *Chelonibia testudinaria* specimens from Case Nuove and the *P. perforatus* specimens from Certaldo, the shell morphology

and original colour are well preserved, with no signs of abrasion. In summary, the macroscopic expression of barnacle preservation is strongly controlled

by the hydrodynamic conditions of the depositional environment as well as by the subsequent diagenetic processes.

At the microscale, all specimens consist of an inner and an outer lamina, in-between are characteristic tree-like interlaminar figures and parietal tubes (Bourget 1977). The preservation of these internal structures has major consequences on all aspects of (acorn) barnacle palaeontology as their highly diagnostic value has been well known since the pioneering studies of Cornwall (1956, 1958, 1959, 1960, 1962) and Davadie (1963).

No significant differences were found among the analysed fossil and recent specimens. However, the barnacles from Montefollonico, Pairola and Fauglia exhibit drusy carbonate crystals filling the parietal tubes, thus highlighting a diagenetic imprint (Fig. 5D, E, H). On the other hand, the shells from Case Nuove and Certaldo display empty parietal tubes that suggests no precipitation from the diagenetic fluids (Fig. 5G, I, J). Though microscopical observations under conventional optical microscopy are not sufficient to differentiate between pristine and altered samples, analyses at higher magnification such as SEM imaging can help in this issue. Previous studies have shown that crystals are arranged massively in the outer shell wall, whereas in the inner wall they form an oriented disposition (Rodríguez-Navarro *et al.* 2006), and the same is true for the opercular plates (Khalifa *et al.* 2011). On the other hand, the radii and alae are formed by elongated fibrous crystals (Murdock & Currey 1978; Mitchell *et al.* 2019; Checa *et al.* 2019, 2020) permeated by a pore network of likely organic channels and pockets involved in the biomineralization process (Mitchell *et al.* 2019). Some variations among barnacle genera might exist (Mitchell *et al.* 2019), but recent investigations suggest that the same heterogeneous mix of crystallite dimensions and morphologies should invariantly occur across different barnacle species (Shaw *et al.* 2024). Our exploration of the ultrastructure of the wall plates reveals a consistent pattern across the studied barnacle species, both extinct and extant. At the ultrastructural level, the modern specimens of both epizoic and non-epizoic barnacles exhibit loosely packed scale-like crystals arranged in the 'dragon-scale pattern' described by Coletti *et al.* (2024) (Figs 6–8). The studied fossil specimens reflect different degrees of preservation of this pattern. The Pliocene specimens of *P. perforatus* from Montefollonico and Certaldo preserve a dragon-scale ultrastructure where the scales retain their original arrangement (Fig. 7). However, they are bulkier and exhibit poorly defined margins compared to the modern samples, thus suggesting a slight alteration due to

diagenesis. In contrast, specimens from the coarse-grained sediments of Pairola and Fauglia are more strongly altered, their original ultrastructural pattern being entirely obliterated in some cases (Fig. 7). This indicates that these fossil shells underwent more extensive diagenetic changes, likely due to exposure to unfavourable post-depositional conditions such as the percolation of diagenetic fluids through the sediment pores. An intermediate degree of preservation is observed for the Case Nuove chelonibiids (Fig. 8).

Erosional microstructures are found in both modern and fossil barnacles. In the first case, they are preserved in both epizoic and non-epizoic forms. In the latter case, they have only been recognized in a well-preserved specimen from Certaldo. The observed features are interpreted as microbial borings, which are produced by a wide array of endolithic prokaryotes and eukaryotes, including phototrophic (e.g. cyanobacteria, algae) as well as heterotrophic (e.g. fungi, foraminifera) forms (e.g. Chacón *et al.* 2006; Glaub *et al.* 2007; Färber *et al.* 2015; Salamon *et al.* 2019; Golubic *et al.* 2019; Kočí *et al.* 2024b). In particular, the microborings observed in the modern examples (Fig. 9A–E) are reminiscent of the morphological diversity of the ichnogenera *Fascichnus*, *Scolecia* and *Eurygonum* (Radtke 1991; Perry & Macdonald, 2002; Försterra *et al.* 2005; Seuss *et al.* 2015; Wisshak *et al.* 2019; Mono *et al.* 2023), which are produced by cyanobacteria (e.g. Radtke & Golubic 2005, 2011). Comparable microbioerosional features have recently been described on barnacles from the polar regions, highlighting the occurrence of microborings of phototrophic euendoliths in shells from the shallow euphotic zone (Meyer *et al.* 2020, 2021). A putative attribution to phototrophic endoliths is supported by the very shallow-water (< 1 m) setting inhabited by one of the microbored modern specimens (*P. perforatus*). The microborings observed in the fossil specimen of Certaldo are similar to those of the modern barnacles, suggesting a well-preserved outer wall and a microbial activity due to photosynthesising bioeroders on a relatively sunlit offshore seafloor.

The cathodoluminescence (CL) response of carbonate biogenic shells correlates with their Mn²⁺ content (Glover 1977, Barbin *et al.* 1991). In barnacle shells, the latter reflects the Mn²⁺ availability in the seawater inhabited by the calcifying animals (Gordon *et al.* 1970; Blanchard & Chastee 1976). Barbin (2000) considered the orange luminescent bands exhibited by barnacle shells to be related to the internal growth bands. However, the author noted that data are insufficient to positively link the CL emission to environmental factors. Based on our results,




Fig. 10. Transmitted light and cathodoluminescence photomicrographs of barnacle shells. A, B, modern specimen of *Perforatus perforatus* from Imperia, showing a predominant low blue luminescence and a low reddish luminescence around the parietal tubes. C, D, modern specimen of *Amphibalanus amphitrite* from Livorno, showing a predominant low blue luminescence and a low reddish luminescence around the parietal tubes. E, F, fossil specimen of *Concavus concavus* from Certaldo, exhibiting a low purple luminescence of the interlaminar figures and a moderate red luminescence of the outer wall. G, H, fossil specimen of *C. concavus* from Montefollonico, exhibiting a moderate purple to red luminescence.

we can conclude that CL analysis have the potential to provide further insights into the diagenetic alteration in fossil barnacle shells. All the modern barnacle

examples exhibit a very weak luminescence, primarily in blue and light reddish tones. This luminescence concentrates along the interlaminar figures, the




Fig. 11. Transmitted light and cathodoluminescence photomicrographs of barnacle shells. A, B, fossil specimen of *Concavus concavus* from Pairola, showing a high red luminescence and a diagenetic cement fill of the parietal tubes. C, D, fossil specimen of *Perforatus perforatus* from Fauglia, exhibiting a very high red to orange luminescence and a diagenetic cement-fill of the parietal tubes. E, F, modern specimen of *Chelonibia testudinaria* from Pozzallo, showing a predominant low blue luminescence. G, H, fossil specimen of *C. testudinaria* from Case Nuove, exhibiting a moderate purple to red luminescence.

inner walls of parietal tubes, the outer surface of the inner lamina and the growth lines (Figs 10, 11). This is recognizable as the intrinsic luminescence of the CL bands, which in turn correlate with the growth

lines (Barbin 1992; Langlet *et al.* 2006). In contrast, the fossil specimens reveal a broader range of luminescence intensities, reflecting the extent of their diagenetic alteration (Figs 10, 11). The *C. concavus*

specimen from Certaldo, which is among the least altered samples, displays a low purple luminescence of the interlaminate figures and a moderate red luminescence of the outer wall. Specimens from Case Nuove and Montefollonico exhibit moderate luminescence, transitioning from purple to red along the inner lamina and interlaminate figures, with luminescent parietal tubes that are partially filled with diagenetic calcite in the case of the latter locality. On the other hand, the specimens from Pairola and Fauglia reveal significant diagenetic imprints, with high-intensity red to orange to yellow luminescence permeating the entire shell, thus suggesting substantial diagenetic infilling by sparitic and drusy calcite. This luminescent response is particularly evident in the parietal tubes, which are filled with diagenetic cement, thus indicating extensive post-depositional diagenetic processes. In conclusion, the best-preserved fossil shells exhibit a low to moderate luminescence, from blue to purple to light reddish in colour, similar to the modern counterparts; whereas the poorly preserved specimens with a high diagenetic imprint display a high red to orange to yellow luminescence.

From the field to the laboratory, macroscopic and microscopic analyses helped to establish a protocol for the taphonomic evaluation of fossil barnacle shells. Overall, the integration of taphonomic macroscopic classifications, optical microscopy, SEM observations on the shell ultrastructure and CL analyses allows for a detailed evaluation of the degree of diagenetic alteration of fossil barnacles. On a general note, our results confirm that burial in fine-grained sediments is much less detrimental to the shell preservation at all scales – from the macroscopic features to the micro- and ultrastructure. This is also supported by a better preservation of the original C and O stable-isotope signatures of shells embedded in fine-grained sediments characterized by low permeability (Coletti *et al.* 2024). Barnacle shells from coarse-grained deposits such as those from Fauglia and Pairola exhibit a more impactful diagenetic imprint, including widespread recrystallization of the shell carbonates and the infilling of parietal tubes with diagenetic carbonate cement. These changes are likely the result of exposure to diagenetic fluids and/or meteoric waters, which can lead to the dissolution and recrystallization of the shell materials (e.g. Flügel 2004; Pederson *et al.* 2019).

Conclusions

This study demonstrates that a combined approach, including a macroscopic taphonomic assessment, microscopic (optical and SEM) ultrastructural

analyses and cathodoluminescence (CL) investigations, provides a robust framework for evaluating fossil barnacle preservation and diagenetic alteration. Macroscopic observations reveal that well-preserved specimens are typically associated with fine-grained sediments indicative of low-energy sedimentary environments, while poorly preserved, fragmented specimens occur in coarse-grained sediments typical of higher-energy depositional settings. Microscopic and SEM analyses of the transverse section of the wall plates reveal that well-preserved fossil specimens share a distinctive ‘dragon-scale’ ultrastructure with their modern counterparts. Diagenetic alteration is more pronounced in specimens from coarse-grained, highly porous sediments, where recrystallization can lead to the obliteration of the original ultrastructure. CL analyses indicate that modern barnacle shells exhibit a weak, predominantly blue luminescence, while the fossil specimens display a broader luminescence spectrum, from blue to purple to red to yellow, reflecting different degrees of diagenesis. Despite their low-magnesium mineralogical stability and seemingly pristine preservation, many barnacle shells from coarse-grained deposits were found to be altered from a micro- and ultrastructural point of view as a likely consequence of the interaction with the diagenetic fluids and/or meteoric waters. Conversely, shells from fine-grained deposits exhibit little signs of diagenetic alteration compared to their modern counterparts. Overall, these results underscore the importance of the depositional environment and diagenetic processes in barnacle preservation, thus enhancing our understanding of the fossilization pathways of cirripede shells.

Acknowledgements.– This paper was funded by a grant of the International Association of Sedimentologists (IAS Post-doctoral Grant Scheme) assigned to GC. Additional funding that supported the detailed taxonomic analysis of barnacles was granted by the LinnéSys: Systematics Research Fund of the Linnean Society and the Systematics Association to GC and AC. The authors would like to thank J. Buckeridge for many helpful discussions on modern and fossil barnacles. Thanks are also due to S. Zanchetta and M. Rocca for their help and assistance with the cathodoluminescence analyses. Special thanks also go to the GAMPS (Gruppo Avis Mineralogia e Paleontologia Scandicci), and especially to S. Casati and A. Di Cencio, for their help during field activities in the Tuscan countryside. The authors are also grateful to Tomáš Kočí and another anonymous reviewer and to the Editor-in-Chief Peter Doyle for fruitful suggestions, which help in improving the quality of this work.

References

- Anderson, D.T. 1994: *Barnacles: Structure, Function, Development and Evolution*, 357 pp. Chapman & Hall, London.
- Benvenuti, M., Bianchi, G., Bruttini, J., Buoniconti, M., Chiarantini, L., Dallai, L., Di Pasquale, G., Donati, A., Grassi, F. & Pescini, V. 2014: Studying the Colline Metallifere mining area

- in Tuscany: an interdisciplinary approach. In Jacquo Silvertanta (ed.): *Proceedings of the Research and Preservation of Ancient Mining Areas, Yearbook of the Institute Europa Subterranea, 9th International Symposium on Archaeological Mining History*, 5–8. MuSe-Trento, Trento.
- Bialik, O.M., Coletti, G., Mariani, L., Commissario, L., Desbiolles, F. & Meroni, A.N. 2023: Availability and type of energy regulate the global distribution of neritic carbonates. *Scientific Reports* 13, 19687. <https://doi.org/10.1038/s41598-023-47029-4>
- Bianucci, G. 1996: A new record of baleen whale from the Pliocene of Tuscany (Italy). *Atti della Società Toscana di Scienze Naturali, Memorie, Serie A* 102, 101–104.
- Blanchard S.C. & Chasteen, N.D. 1976: Determination of manganese (II) in powdered Barnacle shells by electron paramagnetic resonance. *Analytica Chimica Acta* 82, 113–119. [https://doi.org/10.1016/S0003-2670\(01\)82209-1](https://doi.org/10.1016/S0003-2670(01)82209-1)
- Bojar, A.V., Lécuyer, C., Bojar, H.P., Fourel, F. & Vasile, Ş. 2018: Ecophysiology of the hydrothermal vent snail *Ifremeria nautili* and barnacle *Eochionelasmus ohtai manusensis*, Manus Basin, Papua New Guinea: insights from shell mineralogy and stable isotope geochemistry. *Deep Sea Research Part I: Oceanographic Research Papers* 133, 49–58. <https://doi.org/10.1016/j.dsr.2018.02.002>
- Borromeo, L., Zimmermann, U., And'ò, S., Coletti, G., Bersani, D., Basso, D., Gentile, P., Schulz, B. & Garzanti, E. 2017: Raman spectroscopy as a tool for magnesium estimation in Mg-calcite. *Journal of Raman Spectroscopy* 48, 983–992. <https://doi.org/10.1002/jrs.5156>
- Bosio, G., Malinverno, E., Collareta, A., Di Celma, C., Gioncada, A., Parente, M., Berra, F., Marx, F.G., Vertino, A., Urbina, M. & Bianucci, G. 2020: Strontium isotope stratigraphy and the thermophilic fossil fauna from the middle Miocene of the East Pisco Basin (Peru). *Journal of South American Earth Sciences* 97, 102399. <https://doi.org/10.1016/j.jsames.2019.102399>
- Bosio, G., Di Cencio, A., Coletti, G., Casati, S. & Collareta, A. 2021: Exceptionally preserved coral bank and seagrass meadow from the Lower Pleistocene of Fauglia (Tuscany, Italy). *Alpine and Mediterranean Quaternary* 34, 237–256.
- Bossio, A., Foresi, L.M., Liotta, D., Mazzanti, R., Mazzei, R., Salvatoreini, G. & Squarci, P. 1999: Riordino delle conoscenze sul Bacino neogenico del Tora-Fine (Toscana-Italia). *Atti della Società Toscana di Scienze Naturali, Memorie, Serie A* 106, 1–16.
- Bourget, E. 1977: Shell structure in sessile barnacles. *Le Naturaliste canadien* 104, 281–323.
- Bourget, E. 1987: Barnacle shells: composition, structure and growth. In Southward, A.J. (ed.): *Barnacle Biology*, 267–285. Routledge, London. <https://doi.org/10.1201/9781315138053-14>
- Bourget, E. & Crisp, D.J. 1975: An analysis of the growth bands and ridges of barnacle shell plates. *Journal of the Marine Biological Association of the United Kingdom* 55, 439–461. <https://doi.org/10.1017/S0025315400016052>
- Buckeridge, J.S. 1975: *Studies on the Non-pedunculate Lower Tertiary Cirripedia of New Zealand*. Unpublished MSc thesis. University of Auckland, New Zealand.
- Buckeridge, J.S. 1983: Fossil barnacles (Cirripedia: Thoracica) of New Zealand and Australia. *New Zealand Geological Survey, Paleontological Bulletin* 50, 1–151.
- Buckeridge, J.S. 1999: A new deep sea barnacle, *Tetrachaelasma tasmanicum* sp. nov. (Cirripedia: Balanomorpha) from the South Tasman Rise, South Pacific Ocean. *New Zealand Journal of Marine and Freshwater Research* 33, 521–531. <https://doi.org/10.1080/00288330.1999.9516897>
- Buckeridge, J.S. 2015: Revision of Southern Hemisphere taxa referred to *Fosterella* (Crustacea: Cirripedia), and their extinction in response to Pleistocene cooling. *Integrative Zoology* 10, 555–571. <https://doi.org/10.1111/1749-4877.12161>
- Buckeridge, J.S. & Newman, W.A. 2006: A revision of the Iblidae and the pedunculate barnacles (Crustacea: Cirripedia: Thoracica), including new ordinal, familial and generic taxa, and two new species from New Zealand and Tasmanian waters. *Zootaxa* 1136, 1–38. <https://doi.org/10.11646/zootaxa.1136.1.1>
- Buckeridge, J.S., Kočí, T., Gašparič, R. & Kočová Veselská, M. 2020: *Actinobalanus? sloveniensis* (Thoracica, Balanoidea), a new species of cirripede from the Oligocene and Miocene of Slovenia that grew attached to wood substrates. *Neues Jahrbuch für Geologie und Paläontologie-Abhandlungen* 296, 51–65. <https://doi.org/10.1127/njgpa/2020/0902>
- Burgess, S.N., Henderson, G.M. & Hall, B.L. 2010: Reconstructing Holocene conditions under the McMurdo Ice Shelf using Antarctic barnacle shells. *Earth and Planetary Science Letters* 298, 385–393. <https://doi.org/10.1016/j.epsl.2010.08.015>
- Chacón, E., Berrendero, E. & Pichel, F.G. 2006: Biogeological signatures of microboring cyanobacterial communities in marine carbonates from Cabo Rojo, Puerto Rico. *Sedimentary Geology* 185, 215–228. <https://doi.org/10.1016/j.sedgeo.2005.12.014>
- Chave, K.E. 1954: Aspects of the biogeochemistry of magnesium 1. Calcareous marine organisms. *The Journal of Geology* 62, 266–283. <https://doi.org/10.1086/626162>
- Checa, A.G., Salas, C., Rodríguez-Navarro, A.B., Grenier, C. & Lagos, N.A. 2019: Articulation and growth of skeletal elements in balanid barnacles (Balanidae, Balanomorpha, Cirripedia). *Royal Society Open Science* 6, 190458. <https://doi.org/10.1098/rsos.190458>
- Checa, A. G., González-Segura, A., Rodríguez-Navarro, A.B. & Lagos, N.A. 2020: Microstructure and crystallography of the wall plates of the giant barnacle *Austromegabalanus psittacus*: a material organised by crystal growth. *Journal of the Royal Society Interface* 17, 20190743. <https://doi.org/10.1098/rsif.2019.0743>
- Clarke, F.W. & Wheeler, W.C. 1917: The inorganic constituents of marine invertebrates. *US Geological Survey Professional Paper* 124, 62pp. <https://doi.org/10.3133/pp102>
- Coletti, G., Collareta, A., Bosio, G., Urbina-Schmitt, M. & Buckeridge, J. 2019: *Perumegabalanus calziai* gen. et sp. nov., a new intertidal megabalanine barnacle from the early Miocene of Peru. *Neues Jahrbuch für Geologie und Paläontologie-Abhandlungen* 294, 197–212. <https://doi.org/10.1127/njgpa/2019/0856>
- Coletti, G., Bosio, G. & Collareta, A. 2021: Lower Pliocene barnacle facies of western Liguria (NW Italy): a peek into a warm past and a glimpse of our incoming future. *Rivista Italiana di Paleontologia e Stratigrafia* 127, 103–131.
- Coletti, G., Bosio, G., Collareta, A., Bialik, O.M., Regattieri, E., Cornacchia, I., Insacco, G. & Buckeridge, J. 2024: Barnacle-rich facies as a tool for palaeoenvironmental reconstructions. *Palaeogeography, Palaeoclimatology, Palaeoecology* 634, 111914. <https://doi.org/10.1016/j.palaeo.2023.111914>
- Collareta, A. 2020: Discovery of complementary males in a Pliocene accumulation of *Chelonibia testudinaria* (Linnaeus, 1758), with some notes on the evolution of androdioccy in turtle barnacles. *Neues Jahrbuch für Geologie und Paläontologie-Abhandlungen* 297, 193–203. <https://doi.org/10.1127/njgpa/2020/0920>
- Collareta, A., Bosselaers, M. & Bianucci, G. 2016: Jumping from turtles to whales: a Pliocene fossil record depicts an ancient dispersal of *Chelonibia* on mysticetes. *Rivista Italiana di Paleontologia e Stratigrafia* 122, 35–44.
- Collareta, A., Regattieri, E., Zanchetta, G., Lambert, O., Catanzariti, R., Bosselaers, M., Covelo, P., Varola, A. & Bianucci, G. 2018: New insights on ancient cetacean movement patterns from oxygen-isotope analyses of a Mediterranean Pleistocene whale barnacle. *Neues Jahrbuch für Geologie und Paläontologie-Abhandlungen* 288, 143–159. <https://doi.org/10.1127/njgpa/2018/0729>
- Collareta, A., Coletti, G., Bosio, G., Buckeridge, J., de Muizon, C., DeVries, T. J., Varas-Malca, R.M., Altamirano-Sierra, A., Urbina-Schmitt, M. & Bianucci, G. 2019: A new barnacle (Cirripedia: Neobalaniformes) from the early Miocene of Peru: Palaeoecological and palaeobiogeographical implications. *Neues Jahrbuch für Geologie und Paläontologie-Abhandlungen* 292, 321–338. <https://doi.org/10.1127/njgpa/2019/0825>
- Collareta, A., Casati, S., Terranova, E., Nobile, F., Bosio, G., Di Cencio, A. & Coletti, G. 2024: The youngest European record

- of the chelonian family Trionychidae (Calabrian, Central Italy) offers new clues on the Quaternary extirpation history of the softshell turtles. *Geosciences* 14, 239. <https://doi.org/10.3390/geosciences14090239>
- Collareta, A., Merella, M., Mollen, F. H., Casati, S. & Di Cencio, A. 2020: The extinct catshark *Pachyscyllium distans* (Probst, 1879) (Elasmobranchii: Carcharhiniformes) in the Pliocene of the Mediterranean Sea. *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen* 295, 129–139. <https://doi.org/10.1127/njgpa/2020/0877>
- Collareta, A., Mollen, F.H., Merella, M., Casati, S. & Di Cencio, A. 2021: Remarkable multicuspid teeth in a new elusive skate (Chondrichthyes, Rajiformes) from the Mediterranean Pliocene. *Paläontologische Zeitschrift* 95, 117–128. <https://doi.org/10.1007/s12542-020-00542-7>
- Collareta, A., Newman, W.A., Bosio, G. & Coletti, G. 2022: A new chelonibiid from the Miocene of Zanzibar (Eastern Africa) sheds light on the evolution of shell architecture in turtle and whale barnacles (Cirripedia: Coronuloidea). *Integrative Zoology* 17, 24–43. <https://doi.org/10.1111/1749-4877.12554>
- Collins J.S., Donovan S.K. & Mellish C. 2014: An illustrated guide to the fossil barnacles (Cirripedia) from the Crag (Plio-Pleistocene) of East Anglia. *Proceedings of the Geologists' Association* 125, 215–26. <https://doi.org/10.1016/j.pgeola.2014.01.004>
- Cornwall, I.E. 1956: Identifying recent and fossil barnacles by the figures in the shell. *Journal of Paleontology* 30, 646–651.
- Cornwall, I.E. 1958: Identifying recent and fossil barnacles. *Canadian Journal of Zoology* 36, 79–89. <https://doi.org/10.1139/z58-008>
- Cornwall, I.E., 1959: More shell figures and notes on barnacles. *Canadian Journal of Zoology* 37, 401–406. <https://doi.org/10.1139/z59-047>
- Cornwall, I.E. 1960: Barnacle shell figures and repairs. *Canadian Journal of Zoology* 38, 827–832. <https://doi.org/10.1139/z60-085>
- Cornwall, I.E. 1962: The identification of barnacles with further figures and notes. *Canadian Journal of Zoology* 40, 621–629. <https://doi.org/10.1139/z62-057>
- Crippa, G., Ye, F., Malinverno, C. & Rizzi, A. 2016: Which is the best method to prepare invertebrate shells for SEM analysis? Testing different techniques on recent and fossil brachiopods. *Bollettino della Società Paleontologica Italiana* 55, 111–125.
- Crippa, G., Griesshaber, E., Checa, A.G., Harper, E.M., Roda, M.S. & Schmahl, W.W. 2020: Orientation patterns of aragonitic crossed-lamellar, fibrous prismatic and myostracal microstructures of modern *Glycymeris* shells. *Journal of Structural Biology* 212, 107653. <https://doi.org/10.1016/j.jsb.2020.107653>
- Crisp, D.J. & Bourget, E. 1985: Growth in barnacles. *Advances in Marine Biology* 22, 199–244. [https://doi.org/10.1016/S0065-2881\(08\)60052-8](https://doi.org/10.1016/S0065-2881(08)60052-8)
- Dalla Giovanna, G., Fanucci, F., Pellegrini, L., Seno, S., Bonini, L., Decarlis, A., Maino, M., Morelli, D., Vercesi, P.L., Roccati, A., Cobianchi, M. & Mancin, N. 2016: *Note Illustrative Della Carta Geologica d'Italia Alla scala 1:50000, Foglio 259, Imperia*. Servizio Geologico d'Italia, Roma.
- Darwin, C. 1854: *A Monograph of the Sub-class Cirripedia, with Figures of all the Species*, 770 pp Ray Society, London.
- Davadie, C. 1963: *Etude des balanes d'Europe et d'Afrique*, 146 pp. Editions du Centre National de la Recherche Scientifique, Paris
- Davadie-Suaudeau, C. 1952 : Contribution à l'étude des Balanidés Tertiaires de l'Algérie. *Bulletin de le Service de la Carte géologique de l'Algérie, séries 1, Paléontologie* 14, 1–109.
- De Alessandri, G. 1895: Contribuzione allo studio dei Cirripedi fossili d'Italia. *Bollettino della Società Geologica d'Italia* 13, 234–314.
- Domack, E.W. 1988: Biogenic facies in the Antarctic glacial-marine environment: basis for a polar glacial-marine summary. *Palaeogeography, Palaeoclimatology, Palaeoecology* 63, 357–372. [https://doi.org/10.1016/0031-0182\(88\)90105-8](https://doi.org/10.1016/0031-0182(88)90105-8)
- Dominici, S., Rook, L., Benvenuti, M. & Abbazzi, L. 1995: Tapir remains in paralic deposits of Pliocene age in lower Valdarno (Tuscany, Italy): Facies analysis and taphonomy. *Geobios* 28, 131–135. [https://doi.org/10.1016/S0016-6995\(95\)80161-8](https://doi.org/10.1016/S0016-6995(95)80161-8)
- Dominici, S., Danise, S. & Benvenuti, M. 2018: Pliocene stratigraphic paleobiology in Tuscany and the fossil record of marine megafauna. *Earth-Science Reviews* 176, 277–310. <https://doi.org/10.1016/j.earscirev.2017.09.018>
- Donovan, S.K. 1988: Palaeoecology and taphonomy of barnacles from the Plio-Pleistocene Red Crag of East Anglia. *Proceedings of the Geologists' Association* 99, 279–289. [https://doi.org/10.1016/S0016-7878\(88\)80054-3](https://doi.org/10.1016/S0016-7878(88)80054-3)
- Donovan, S.K. 1989: Palaeoecology and significance of barnacles in the mid-Pliocene *Balanus* Bed of Tobago, West Indies. *Geological Journal* 24, 239–250. <https://doi.org/10.1002/gj.3350240402>
- Donovan, S.K. 1993: The ecology of ancient barnacles. *Rocks & Minerals* 68, 115–119. <https://doi.org/10.1080/00357529.1993.9926538>
- Dougherty, W. J. 1990: SEM observations on the interfacial surface of the cement of the adult barnacle, attached to natural and synthetic adherends. *Tissue and Cell* 22, 463–470. [https://doi.org/10.1016/0040-8166\(90\)90075-K](https://doi.org/10.1016/0040-8166(90)90075-K)
- Doyle P., Mather, A.E., Bennett, M.R. & Bussell, M.A. 1996: Miocene barnacle assemblages from Southern Spain and their palaeoenvironmental significance. *Lethaia* 29, 267–274. <https://doi.org/10.1111/j.1502-3931.1996.tb01659.x>
- Färber, C., Wisshak, M., Pyko, I., Bellou, N. & Freiwald, A. 2015: Effects of water depth, seasonal exposure, and substrate orientation on microbial bioerosion in the Ionian Sea (Eastern Mediterranean). *PloS One* 10, e0126495. <https://doi.org/10.1371/journal.pone.0126495>
- Farrow, G.E., Cucci, M. & Scoffin, T.P. 1978: Calcareous sediments on nearshore continental shelf of western Scotland. *Proceedings of the Royal Society of Edinburgh, Section B: Biological Sciences* 76, 55–76. <https://doi.org/10.1017/S0269727000002797>
- Fernández, M.S., Moya, A., Lopez, V. & Arias, J.L. 2001: Secretion pattern, ultrastructural localization and function of extracellular matrix molecules evolved in eggshell formation. *Matrix Biology* 19, 793–803. [https://doi.org/10.1016/S0945-053X\(00\)00128-1](https://doi.org/10.1016/S0945-053X(00)00128-1)
- Flügel, E. 2004: *Microfacies of Carbonate Rocks: Analysis, Interpretation and Application*, 2004 pp. Springer, Berlin. <https://doi.org/10.1007/978-3-662-08726-8>
- Försterra, G., Beuck, L., Häussermann, V. & Freiwald, A. 2005: Shallow-water *Desmophyllum dianthus* (Scleractinia) from Chile: characteristics of the biocoenoses, the bioeroding community, heterotrophic interactions and (paleo)-bathymetric implications. In Freiwald, A. & Roberts, J.M. (eds): *Cold-water corals and ecosystems*. Erlangen Earth Conference Series 937-977. Springer, Berlin, Heidelberg. https://doi.org/10.1007/3-540-27673-4_48
- Foster, B.A. & Buckeridge, J.S. 1987: Barnacle paleontology. In Southward, A.J. (ed.): *Barnacle Biology*. Crustacean Issues 5, 43–62. Balkema, Rotterdam. <https://doi.org/10.1201/9781315138053-2>
- Frank, T.D., James, N.P., Bone, Y., Malcolm, I. & Bobak, L.E. 2014: Late Quaternary carbonate deposition at the bottom of the world. *Sedimentary Geology* 305, 1–16. <https://doi.org/10.1016/j.sedgeo.2014.02.008>
- Gal, A., Weiner, S. & Addadi, L. 2015: A perspective on underlying crystal growth mechanisms in biomineralisation: solution mediated growth versus nanosphere particle accretion. *CrystEngComm* 17, 2606. <https://doi.org/10.1039/C4CE01474J>
- Gale, A.S. 2021: Acorn barnacles (Crustacea, Thoracica, Balanomorphia) from the Eocene and Oligocene of the Isle of Wight and Hampshire (United Kingdom) and Manche, northern France. *Acta Geologica Polonica* 71, 153–174. <https://doi.org/10.24425/agp.2020.132263>
- Gale, A. & Schweigert, G. 2016: A new phosphatic-shelled cirripede (Crustacea, Thoracica) from the Lower Jurassic (Toarcian)

- of Germany – the oldest epiplanktonic barnacle. *Palaeontology* 59, 59–70. <https://doi.org/10.1111/pala.12207>
- Gale, A.S. & Sørensen, A.M. 2015: Origin of the balanomorph barnacles (Crustacea, Cirripedia, Thoracica) – new evidence from the Late Cretaceous (Campanian) of Sweden. *Journal of Systematic Palaeontology* 13, 791–824. <https://doi.org/10.1080/14772019.2014.954824>
- Gale, A.S., Rosso, A. & Vertino, A. 2021: Cirripedes (Crustacea, Thoracica) from the Plio-Pleistocene of Sicily (Italy) and a review of the history of Mediterranean Cainozoic cirripedes. *Cainozoic Research* 21, 3–74.
- Giammarino, S., Fanucci, F., Orezzi, S., Rosti, D., Morelli, D., Cobiainchi, M., Di Stefano, A., Fravega, P., Vannucci, G., Piazza, M. & Finocchiaro, F. 2010: *Note Illustrative Della Carta Geologica d'Italia Alla scala 1:50000, Foglio 258–271, San Remo*. Servizio Geologico d'Italia, Roma.
- Glaub, I., Golubic, S., Gektidis, M., Radtke, G., & Vogel, K. 2007: Microborings and microbial endoliths: geological implications. In Miller, W. III (ed): *Trace Fossils*, 368–381. Elsevier Science. <https://doi.org/10.1016/B978-044452949-7/50147-9>
- Glover, E.D. 1977: Cathodoluminescence, iron and manganese content, and the early diagenesis of carbonates. Unpublished PhD thesis, University of Wisconsin.
- Glynn, P.W. & Wellington, G.M. 1983: *Coral and Coral Reefs of the Galapagos islands*, 330 pp. University of California Press, Berkeley and Los Angeles.
- Golubic, S., Schneider, J., Le Campion-Alsumard, T., Campbell, S.E., Hook, J.E. & Radtke, G. 2019: Approaching microbial bioerosion. *Facies* 65, 1–17. <https://doi.org/10.1007/s10347-019-0568-1>
- Gordon, C.M., Carr, R.A. & Larson, R.E. 1970: The influence of environmental factors on the sodium and manganese content of Barnacle Shells. *Limnology and Oceanography* 15, 461–466. <https://doi.org/10.4319/lo.1970.15.3.0461>
- Guido, A., D'Amico, F., DeVries, T.J., Kočí, T., Collareta, A., Bosio, G. & Sanfilippo, R. 2024: Double-phased controlled and influenced biomineralization in marine invertebrates: the example of Miocene to recent reef-building polychaete cirratulids from southern Peru. *Palaeogeography, Palaeoclimatology, Palaeoecology* 639, 112060. <https://doi.org/10.1016/j.palaeo.2024.112060>
- Halfar, J., Strasser, M., Riegl, B. & Godinez-Orta, L. 2006: Oceanography, sedimentology and acoustic mapping of a bryomol carbonate factory in the northern Gulf of California, Mexico. In Pedley, H.M. & Carannante, G. (eds): *Cool-Water Carbonates: Depositional Systems and Palaeoenvironmental Controls*. Geological Society of London Special Publications 255, 197–215. <https://doi.org/10.1144/GSL.SP.2006.255.01.13>
- Hall, A., Kennedy, W.J. & Taylor, J. H. 1967: Aragonite in fossils. *Proceedings of the Royal Society of London, Series B: Biological Sciences* 168, 377–412. <https://doi.org/10.1098/rspb.1967.0071>
- Henrich, L., Freiwald, A., Betzler, C., Bader, B., Samtleben, C., Brachert, T.C., Wehrmann, A., Zankl, H. & Kühlmann, D.H.H. 1995: Controls on modern carbonate sedimentation on warm-temperate to Arctic coasts, shelves and seamounts in the Northern Hemisphere: implications for fossil counterparts. *Facies* 32, 71–108. <https://doi.org/10.1007/BF02536865>
- Hoffmann, R., Bitner, M. A., Pisera, A., Jäger, M., Auer, G., Giraldo-Gómez, V., Kočí, T., Buckeridge, J., Mueler, M., Stevens, K. & Schneider, S. 2020: Late Miocene biota from the Abad Member of the Carboneras-Níjar Basin (Spain, Andalusia): A bathyal fossil assemblage pre-dating the Messinian salinity crisis. *Geobios*, 59, 1–28. <https://doi.org/10.1016/j.geobios.2020.03.002>
- Hoskin, C.M. & Nelson, R.V. 1969: Modern marine carbonate sediment, Alexander Archipelago, Alaska. *Journal of Sedimentary Petrology* 39, 581–590. <https://doi.org/10.1306/74D71CD5-2B21-11D7-8648000102C1865D>
- Hoskin, C.M. & Reed, J.K. 1984: Barnacle plate sediment production by sheephead, the Indian River, Florida. *Geo-Marine Letters* 4, 55–57. <https://doi.org/10.1007/BF02237975>
- Hottinger L.K. 1983: Neritic Macroid Genesis: an Ecological Approach. In Peryt, T.M. (ed.): *Coated Grains*, 38–55. Springer-Verlag, Berlin. https://doi.org/10.1007/978-3-642-68869-0_5
- Hover, V.C., Walter, L.M. & Peacor, D.R. 2001: Early marine diagenesis of biogenic aragonite and Mg-calcite: new constraints from high-resolution STEM and AEM analyses of modern platform carbonates. *Chemical Geology* 175, 221–248. [https://doi.org/10.1016/S0009-2541\(00\)00326-0](https://doi.org/10.1016/S0009-2541(00)00326-0)
- Khalifa, G.M., Weiner, S. & Addadi, L. 2011: Mineral and matrix components of the operculum and shell of the barnacle *Balanus amphitrite*: calcite crystal growth in a hydrogel. *Crystal Growth & Design* 11, 5122–5130. <https://doi.org/10.1021/cg2010216>
- Killingley, J.S. & Newman, W.A. 1982: ^{18}O fractionation in barnacle calcite: a barnacle paleotemperature equation. *Marine Environmental Research* 40, 893–902.
- Klicpera A., Taylor P.D. & Westphal H. 2013: Bryoliths constructed by bryozoans in symbiotic associations with hermit crabs in a tropical heterozoan carbonate system, Golfe d'Arguin, Mauritania. *Marine Biodiversity* 43, 429–444. <https://doi.org/10.1007/s12526-013-0173-4>
- Kočí T., Veselská M.K., Newman W.A., Buckeridge J.S. & Sklenář J. 2017: *Archaeochionelasmus nevasilovae* gen. et sp. nov. (Cirripedia, Balanomorphia, Chionelasmatoidea) from the Bohemian Cretaceous Basin (Czech Republic): the first bona fide Cretaceous neobalaniform. *Zootaxa* 4294, 2, 181–196. <https://doi.org/10.11646/zootaxa.4294.2.3>
- Kočí, T., Bosio, G., Collareta, A., Sanfilippo, R., Ekrt, B., Urbina, M. & Malinverno, E. 2021: First report on the cirratulid (Annelida, Polychaeta) reefs from the Miocene Chilcatay and Pisco Formations (East Pisco Basin, Peru). *Journal of South American Earth Sciences* 107, 103042. <https://doi.org/10.1016/j.james.2020.103042>
- Kočí, T., Gašparič, R., Buckeridge, J., Kočová Veselská, M. & Šoster, A. 2024a: The first record of a Konservat-Lagerstätten in which early post-settlement stages of fossil archaeobalanids (Cirripedia: Balanomorphia) are preserved. *Integrative Zoology* 19, 200–223. <https://doi.org/10.1111/1749-4877.12728>
- Kočí, T., Veselská, M. K., Goedert, J. L., Buckeridge, J. S., Reitner, J. & Váchová, L. 2024b: First Eocene record of a bangialean rhodophyte (the endolithic microboring *Conchocelichnus seilacheri*) and coralline red algae from the Pacific Coast of North America. *PalZ* 98, 203–222. <https://doi.org/10.1007/s12542-023-00678-2>
- Langlet D., Alunno-Bruscia M., Rafélis de M., Renard M., Roux M., Schein, E. & Buestel D. 2006. Experimental and natural cathodoluminescence in the shell of *Crassostrea gigas* from Thau lagoon (France): ecological and environmental implications. *Marine Ecology Progress Series* 317, 143–156. <https://doi.org/10.3354/meps317143>
- Lewis, A.C., Burden, D.K., Wahl, K.J., & Everett, R.K. 2014: Electron backscatter diffraction (EBSD) study of the structure and crystallography of the barnacle *Balanus amphitrite*. *The Journal of The Minerals, Metals & Materials Society* 66, 143–148. <https://doi.org/10.1007/s11837-013-0793-y>
- López, D.A., López, B.A., Arriagada, S.E., Mora, O.A., Bedecarratz, P.C., Pineda, M.O., Gonzalez, M.L., Andrade, L.I., Uribe, J.M. & Riquelme, V.A. 2012: Diversification of Chilean aquaculture: the case of the giant barnacle *Austromegabalanus psittacus* (Molina, 1782). *Latin American Journal of Aquatic Research* 40, 596–607. <https://doi.org/10.3856/vol40-issue3-fulltext-9>
- Mariani, L., Coletti, G., Mateu-Vicens, G., Bosio, G., Collareta, A., Khokhlova, A., Di Cencio, A., Casati, S. & Malinverno, E. 2022: Testing an indirect palaeo-seagrass indicator: Benthic foraminifera from the lower Pleistocene *Posidonia* meadow of Fauglia (Tuscany, Italy). *Marine Micropaleontology* 173, 102126. <https://doi.org/10.1016/j.marmicro.2022.102126>
- Martini, I., Ambrosetti, E., Brogi, A., Aldinucci, M., Zwaan, F. & Sandrelli, F. 2021: Polyphase extensional basins: interplay between tectonics and sedimentation in the Neogene Siena-Radicofani Basin (Northern Apennines, Italy). *International*

- Journal of Earth Sciences* 110, 1729–1751. <https://doi.org/10.1007/s00531-021-02038-4>
- Menesini, E. 1965: Caratteri morfologici e struttura microscopica di alcune specie di balani neogenici. *Paleontographia Italica* 59, 85–129.
- Menesini, E. 1971: Nuove specie di *Balanus* (Cirripedia) dell'Eocene e dell'Oligocene de Baccina di Parigi. *Bollettino della Società Paleontologica Italiana* 10, 19–34.
- Menesini E. & Casella C. 1988: Balanidi pliocenici della provincia di Almeria (Andalusia orientale - Spagna). Studio sistematico. *Atti della Società Toscana de Scienze Naturali di Pisa, Memorie, Serie A*, 95, 231–269.
- Merella, M., Collareta, A., Casati, S., Di Cencio, A. & Bianucci, G. 2023: Pliocene geotourism: innovative projects for valorizing the paleontological heritage of three different-staged quarries of Tuscany (Central Italy). *Geoheritage* 15, 82. <https://doi.org/10.1007/s12371-023-00838-5>
- Meyer, N., Wisshak, M. & Freiwald, A. 2020: Ichnodiversity and bathymetric range of microbioerosion traces in polar barnacles of Svalbard. *Polar Research* 39, 3766. <https://doi.org/10.33265/polar.v39.3766>
- Meyer, N., Wisshak, M. & Freiwald, A. 2021: Bioerosion ichnodiversity in barnacles from the Ross Sea, Antarctica. *Polar Biology* 44, 667–682. <https://doi.org/10.1007/s00300-021-02825-4>
- Michel J., Mateu-Vicens G. & Westphal H. 2011: Modern Heterozoan carbonates from a eutrophic tropical shelf (Mauritania). *Journal of Sedimentary Research* 81, 641–655. <https://doi.org/10.2110/jsr.2011.53>
- Milliman J.D. 1972: Atlantic Continental Shelf and Slope of the United States-Petrology of the Sand Fraction of Sediments, Northern New Jersey to Southern Florida. *US Geological Survey Professional Paper* 529-J, 1–48. <https://doi.org/10.3133/pp529J>
- Mitchell, R.L., Coleman, M., Davies, P., North, L., Pope, E.C., Pleydell-Pearce, C., Harris, W. & Johnston, R. 2019: Macro-to-nanoscale investigation of wall-plate joints in the acorn barnacle *Semibalanus balanoides*: correlative imaging, biological form and function, and bioinspiration. *Journal of the Royal Society Interface* 16, 20190218. <https://doi.org/10.1098/rsif.2019.0218>
- Mono, P., Hoffmann, R., Wisshak, M., Lokier, S.W., Pederson, C.L., Hennhofer, D., Diaz, M.R., Swart, P.K., Nehrke, G. & Immenhauser, A. 2024: Microborings reveal alternating agitation, resting and sleeping stages of modern marine ooids. *Sedimentology* 71, 359–382. <https://doi.org/10.1111/sed.13149>
- Murdock, G.R. & Currey, J.D. 1978: Strength and design of shells of the two ecologically distinct barnacles, *Balanus balanoides* and *Semibalanus (Balanus) balanoides* (Cirripedia). *The Biological Bulletin* 155, 169–192. <https://doi.org/10.2307/1540874>
- Müller, J. & Milliman, J.D. 1973: Relict carbonate-rich sediments on Southwestern Grand Bank, Newfoundland. *Canadian Journal of Earth Sciences* 10, 1744–1750. <https://doi.org/10.1139/e73-170>
- Nalin, R., Ghinassi, M., Foresi, L.M. & Dallanave, E. 2016: Carbonate deposition in restricted basins: a Pliocene case study from the Central Mediterranean (Northwestern Apennines), Italy. *Journal of Sedimentary Research* 86, 236–267. <https://doi.org/10.2110/jsr.2016.14>
- Nelson, C.S., Keane, S.L. & Head, P.S. 1988: Non-tropical carbonate deposits on the modern New Zealand shelf. *Sedimentary Geology* 60, 71–94. [https://doi.org/10.1016/0037-0738\(88\)90111-X](https://doi.org/10.1016/0037-0738(88)90111-X)
- Newman, W.A. 1982: A review of extant taxa of the 'group of *Balanus concavus*' (Cirripedia, Thoracica) and a proposal for genus-group ranks. *Crustaceana* 43, 25–36. <https://doi.org/10.1163/156854082X00056>
- Newman, W.A. & Ross, A. 1971: Antarctic Cirripedia. *Antarctic Research Series* 14, 1–257. <https://doi.org/10.1029/AR014>
- Newman, W.A., Zullo, V.A. & Stephan A. Wainwright, S.A. 1967: A Critique on Recent Concepts of Growth in Balanomorphs (Cirripedia, Thoracica). *Crustaceana* 12, 167–178. <https://doi.org/10.1163/156854067X00594>
- Newman W.A., Zullo V.A. & Withers T.H. 1969: Cirripedia. In: Moore R.C. (Ed.): *Treatise on Invertebrate Paleontology, Part R, Arthropoda 4, Vol. 1*. University of Kansas, Lawrence and the Geological Society of America, Boulder, R206–295.
- Nielsen, J.K. & Funder, S. 2003: Taphonomy of Eemian marine molluscs and acorn barnacles from eastern Arkhangelsk region, northern Russia. *Palaeogeography Palaeoclimatology Palaeoecology* 191, 139–168. [https://doi.org/10.1016/S0031-0182\(02\)00707-1](https://doi.org/10.1016/S0031-0182(02)00707-1)
- Nishizaki, M.T. & Carrington, E. 2015: The effect of water temperature and velocity on barnacle growth: Quantifying the impact of multiple environmental stressors. *Journal of Thermal Biology* 54, 37–46. <https://doi.org/10.1016/j.jtherbio.2015.02.002>
- Nomura, S. & Maeda, H. 2008: Significance of autochthonous fossil barnacles from the Miocene Natori Group at the Moniwa-Goishi area, Northeast Japan. *Paleontological Research* 12, 63–79. [https://doi.org/10.2517/1342-8144\(2008\)12\[63:SOAFBF\]2.0.CO;2](https://doi.org/10.2517/1342-8144(2008)12[63:SOAFBF]2.0.CO;2)
- Paces, J.B., Minor, S.A., Schmidt, K.M. & Hoffman, J. 2023: Strontium isotope chronostratigraphic age of a sirenian fossil site on Santa Rosa Island, Channel Islands National Park, California. *Scientific Investigations, Report 2023-5026*, US Geological Survey, Boulder. <https://doi.org/10.3133/sir20235026>
- Pajaud, D. 1976: A propos des fossiles du Pliocène d'Aguilas (Sud d'Almeria, Espagne) découverte de Cirripèdes operculés (Crustacés) et discussion sur l'étho-écologie des formes épizoaires. *Géobios* 9, 481–502. [https://doi.org/10.1016/S0016-6995\(76\)80004-6](https://doi.org/10.1016/S0016-6995(76)80004-6)
- Pederson, C., Mavromatis, V., Dietzel, M., Rollion-Bard, C., Nehrke, G., Jöns, N., Jochum, K.P. & Immenhauser, A. 2019: Diagenesis of mollusc aragonite and the role of fluid reservoirs. *Earth and Planetary Science Letters* 514, 130–142. <https://doi.org/10.1016/j.epsl.2019.02.038>
- Perry, C.T. & Macdonald, I.A. 2002: Impacts of light penetration on the bathymetry of reef microboring communities: implications for the development of microendolithic trace assemblages. *Palaeogeography, Palaeoclimatology, Palaeoecology* 186, 101–113. [https://doi.org/10.1016/S0031-0182\(02\)00446-7](https://doi.org/10.1016/S0031-0182(02)00446-7)
- Radtke, G. 1991: Die mikroendolithischen spurenfossilien im Alt-Tertiär West-Europas und ihre palökologische Bedeutung. *CFS* 138, 150pp.
- Radtke, G. & Golubic, S. 2005: Microborings in mollusk shells, Bay of Safaga, Egypt: Morphometry and ichnology. *Facies* 51, 118–134. <https://doi.org/10.1007/s10347-005-0016-02>
- Radtke, G. & Golubic, S. 2011: Microbial Euendolithic Assemblages and Microborings in Intertidal and Shallow Marine Habitats: Insight in Cyanobacterial Speciation. In: *Advances in Stromatolite Geobiology. Lecture Notes in Earth Sciences*, 131. Springer, Berlin, Heidelberg. https://doi.org/10.1007/978-3-642-10415-2_16
- Raman, S. & Kumar, R. 2011: Construction and nanomechanical properties of the exoskeleton of the barnacle, *Amphibalanus reticulatus*. *Journal of Structural Biology* 176, 360–369. <https://doi.org/10.1016/j.jsb.2011.08.015>
- Raymond P.E. & Stetson H.C. 1932: A calcareous beach on the coast of Maine. *Journal of Sedimentary Petrology* 2, 51–62. <https://doi.org/10.1306/D4268DFB-2B26-11D7-8648000102C1865D>
- Reijmer J.J.G., Bauch T. & Schäfer P. 2012: Carbonate facies patterns in surface sediments of upwelling and non-upwelling shelf environments (Panama, East Pacific). *Sedimentology* 59, 32–56. <https://doi.org/10.1111/j.1365-3091.2010.01214.x>
- Reymond C.E., Zihrl K.S., Halfar J., Riegl B., Humphreys A. & Hildegard W. 2016: Heterozoan carbonates from the equatorial rocky reefs of the Galapagos Archipelago. *Sedimentology* 63, 940–958. <https://doi.org/10.1111/sed.12244>
- Rodríguez-Navarro, A.B., CabraldeMelo, C., Batista, N., Morimoto, N., Alvarez-Lloret, P., Ortega-Huertas, M., Fuenzalida, V.M., Arias, J.I., Wiff, J.P. & Arias, J. L. 2006: Microstructure and

- crystallographic-texture of giant barnacle (*Austromegabalanus psittacus*) shell. *Journal of Structural Biology* 156, 355–362. <https://doi.org/10.1016/j.jsb.2006.04.009>
- Roskowski, J.A., Patchett, P.J., Spencer, J.E., Pearthree, P.A., Dettman, D.L., Faulds, J.E. & Reynolds, A.C. 2010: A late Miocene–early Pliocene chain of lakes fed by the Colorado River: evidence from Sr, C, and O isotopes of the Bouse Formation and related units between Grand Canyon and the Gulf of California. *Bulletin* 122, 1625–1636. <https://doi.org/10.1130/B30186.1>
- Salamon, K., Kołodziej, B. & Stefanskyi, V.L. 2019: Simple methods for detection of microborings produced by coral-associated microendoliths. *Facies* 65, 16. <https://doi.org/10.1007/s10347-019-0560-9>
- Scoffin T.P. 1988: The environments of production and deposition of calcareous sediments on the shelf west of Scotland. *Sedimentary Geology* 60, 107–124. [https://doi.org/10.1016/0037-0738\(88\)90113-3](https://doi.org/10.1016/0037-0738(88)90113-3)
- Seuss, B., Hembree, D.I., Wisshak, M., Mapes, R.H. & Landman, N.H. 2015: Taphonomy of backshore versus deep-marine collected *Nautilus macromphalus* conchs (New Caledonia). *Palaios* 30, 503–513. <https://doi.org/10.2110/palo.2014.057>
- Shaw, J., Kang, Y., Triano, C., Hoppe, C.J., Aldred, N., Metzler, R.A. & Dickinson, G.H. 2024: Comparative Assessment of Shell Structural, Mechanical, and Elemental Properties in Adult Acorn Barnacles. *Diversity* 16, 482. <https://doi.org/10.3390/d16080482>
- Taylor, L.D., O'Dea, A., Bralower, T.J. & Finnegan, S. 2019: Isotopes from fossil coronulid barnacle shells record evidence of migration in multiple Pleistocene whale populations. *Proceedings of the National Academy of Sciences USA* 116, 7377–7381. <https://doi.org/10.1073/pnas.1808759116>
- Ullmann, C.V., Gale, A.S., Huggett, J., Wray, D., Frei, R., Korte, C., Broom-Fendley, S., Littler, K. & Hesselbo, S.P. 2018: The geochemistry of modern calcareous barnacle shells and applications for palaeoenvironmental studies. *Geochimica et Cosmochimica Acta* 243, 149–168. <https://doi.org/10.1016/j.gca.2018.09.010>
- Vinn, O. 2013: SEM study of semi-oriented tube microstructures of Serpulidae (Polychaeta, Annelida): Implications for the evolution of complex oriented microstructures. *Microscopy Research and Technique* 76, 453–456. <https://doi.org/10.1002/jemt.22186>
- Vinn, O. 2020: Biomineralization of polychaete annelids in the fossil record. *Minerals* 10, 858. <https://doi.org/10.3390/min10100858>
- Vinn, O. & Kupriyanova, E.K. 2011: Evolution of a dense outer protective tube layer in serpulids (Polychaeta, Annelida). *Carnets de Géologie-Notebooks on Geology CG2011_L05*, 137–147. <https://doi.org/10.4267/2042/43896>
- Vinn, O., Ten Hove, H.A. & Mutvei, H. 2008a: On the tube ultrastructure and origin of calcification in sabellids (Annelida, Polychaeta). *Palaeontology* 51, 295–301. <https://doi.org/10.1111/j.1475-4983.2008.00763.x>
- Vinn, O., Ten Hove, H.A., Mutvei, H. & Kirsimaee, K. 2008b: Ultrastructure and mineral composition of serpulid tubes (Polychaeta, Annelida). *Zoological Journal of the Linnean Society* 154, 633–650. <https://doi.org/10.1111/j.1096-3642.2008.00421.x>
- Westphal H., Halfar J. & Freiwald A. 2010: Heterozoan carbonates in subtropical to tropical settings in the present and past. *International Journal of Earth Sciences* 99, S1, S153–S169. <https://doi.org/10.1007/s00531-010-0563-9>
- Wilson J.B. 1988: A model for temporal changes in the faunal composition of shell gravels during a transgression on the continental shelf around the British Isles. *Sedimentary Geology* 60, 95–105. [https://doi.org/10.1016/0037-0738\(88\)90112-1](https://doi.org/10.1016/0037-0738(88)90112-1)
- Wisshak, M., Seunß, B. & Nützel, A. 2008: Evolutionary implications of an exceptionally preserved Carboniferous microboring assemblage in the Buckhorn Asphalt Lagerstätte (Oklahoma, USA). In: Wisshak, M. & Tapanila, L. (eds) *Current Developments in Bioerosion*, Erlangen Earth Conference Series, Springer, Berlin, 21–54. https://doi.org/10.1007/978-3-540-77598-0_2
- Zullo, V.A. 1992: Revision of the balanid barnacle genus *Concavus* Newman, 1982, with the description of a new subfamily, two new genera, and eight new species. *Memoir (The Paleontological Society)* 27, 1–46. <https://doi.org/10.1017/S0022336000060947>