

Personal private herbaria: a valuable but neglected source of floristic data. The case of Italian collections today

Francesco Roma-Marzio¹, Lorenzo Peruzzi¹, Gianni Bedini¹

¹ *Dipartimento di Biologia, Unità di Botanica, Università di Pisa, Via Derna 1, 56126, Pisa, Italy*

Corresponding author: *Francesco Roma-Marzio* (francesco.romamarzio@for.unipi.it)

Academic editor: *G. Domina* | Received 3 February 2017 | Accepted 9 March 2017 | Published 17 March 2017

Citation: Roma-Marzio F, Peruzzi L, Bedini G (2017) Personal private herbaria: a valuable but neglected source of floristic data. The case of Italian collections today. *Italian Botanist* 3: 7–15. doi: 10.3897/italianbotanist.3.12097

Abstract

Today private collections still represent an important source of information as is evident from the available literature, where, in many cases, private collections are mentioned both in floristic and taxonomic/systematic researches. We present a summary of the main information about private herbaria in Italy, whose collections are currently increasing. Based on our survey, we retrieved information on 34 personal herbaria where 156,361 specimens are preserved at present. Piedmont and Tuscany resulted the two Italian regions with the highest number of collections, whereas for 9 regions we obtained no answer. The most represented families resulted Asteraceae and Poaceae, whereas the most represented genera resulted *Carex*, *Trifolium*, and *Hieracium*. Taken all together, these collections rank 16th among the 68 institutional public herbaria officially recognised in Italy.

Keywords

Herbarium specimens, Italian botanist, plant collection, floristic data

Introduction

Private herbarium collections have represented the starting point for the establishment of many of the still-extant natural history museums around the world. In the 17th and 18th centuries, the fashion for collecting plants resulted in the amassing of large private collections, which later became the basis of institutional collections (Stearn 1971). Thus, the Oxford University herbarium (OXF, acronyms follow Thiers 2016)

was largely formed from the private herbaria of Jacob Bobart the Younger (1641–1719), William Sherard (1658–1728), Henry Barron Fielding (1805–1851) and George Claridge Druce (1850–1932) (Clokier 1964, Stearn 1971). The Natural History Museum of London (BM) owes its foundation to the tireless activity of Sir Hans Sloane (1660–1753) and Sir Joseph Banks (1743–1820), whose houses quickly become too full to accommodate their collections (Beer 1953, Stearn 1971). Likewise, the herbarium of the Royal Botanic Gardens, Kew (K) started to develop from the private herbaria of Sir William Jackson Hooker (1785–1865), George Bentham (1800–1884), and William Arnold Bromfield (1801–1851) (Stearn 1971). In Italy, there are 68 institutional public herbaria (Taffetani 2012), many of them hosting significant collections donated by private collectors. These collections represent an important part of preserved plant specimens. The herbarium of the University of Florence (FI), currently preserving more than five million samples, hosts the private collection of its founder, Filippo Parlatore (1816–1877), composed by about 300,000 specimens (Cuccuini 2009), as well as other private collections. Among these, one of the most important is the herbarium of Philip Barker Webb (1793–1854) (FI-W), counting about 250,000 specimens, and purchased in the 1855, by the then curator Filippo Parlatore (Moggi 1993, Nepi 2009). The herbarium of the University of Pisa (PI) hosts nine main private collections, donated to or purchased by the university and detached from the general collection (Amadei et al. 2007, 2012). Among these collections, noteworthy are the herbarium of Pietro Pellegrini (1867–1957), composed by 22,000 specimens, mainly from the Apuan Alps (Garbari and Del Carratore 1993); the herbarium of Teodoro Caruel (1830–1898), counting about 14,500 samples representing a valuable document related to his “*Prodromo della Flora Toscana*” (Caruel 1860); and the herbarium of Michele Guadagno (1878–1930) with gatherings mainly from southern Italy.

Private collections are preserved also in other Italian public institutions like civic museums or schools. To name but a few examples, the herbarium of Giovanni Montini (1802–1854) is preserved in the Civic Museum of Bassano del Grappa (Lasen and Busnardo 1993); the herbarium of Pietro Zangheri (1889–1983) is currently preserved in a specific section of the Civic Museum of Natural History of Verona (VER) (Viciani 2011); and the three herbaria of Erminio Ferrarini (1919–2002), overall composed by 8655 samples, are now preserved in the high school “Guglielmo Marconi” of Carrara (Tuscany), the natural history museum of the Lunigiana (Aulla, Tuscany) and in FI (Maccioni et al. 2008).

Today, private collections still represent an important source of information. This is evident from the available literature where, in many cases, private collections are mentioned both in floristic (e.g. Bartolucci and Peruzzi 2007, Anzalone et al. 2010, Selvi 2010, Peruzzi et al. 2011, 2016, Roma-Marzio et al. 2016, Ardenghi and Polani 2016) and taxonomic / systematic researches (e.g. Brilli-Cattarini and Gubellini 1986, Iamónico 2015, Peccenini and Polatschek 2016).

In this paper, we make a first attempt to summarize the main information about private herbaria currently preserved in Italy.

Material and methods

We defined as private herbarium a collection of dried vascular plants, continuously increased with new gatherings, and independently managed by a private collector, without support from any public or private research institution. We excluded the institutional herbaria listed in the *Index Herbariorum* (Thiers 2016), and all the herbaria maintained in the public bodies such as educational institutions, regional and national parks, libraries and public archives.

In order to obtain data from Italian private collections of plants, we prepared an on-line questionnaire based on a Google module. The questionnaire included 17 questions (15 with a free-text answer and 2 with a multiple-choice answer), focused on the herbarium name, its owner, the starting date of collection and the number and the main geographic origin of specimens, the address where the herbarium is preserved, the kind of management (including pest control), and other questions about the dissemination of data in the past, and the availability of the owner to share the data (Table 1).

In order to disseminate our initiative we used the mailing list of the working group for floristics, systematics and evolution of the Italian Botanical Society (*Gruppo per la Floristica, Sistematica ed Evoluzione*). In addition we used the most common social networks (e.g. Facebook) and the Acta Plantarum forum (<http://www.actaplantarum.org/>), an Italian website focused on national floristic information; in some cases we also sent e-mail messages to personal addresses.

Results

Based on our survey, we checked 34 personal herbaria, where 156,361 specimens are currently preserved. A list of personal herbaria, and relative informations, can be freely accessed at <https://goo.gl/eJA0Ga>. This online checklist will be continuously updated in case that other collectors will send us information about their collections.

The richest herbarium, with 20,000 estimated samples, resulted the *Herbarium Antonietti* (Piedmont) followed by *Herbarium Soldano* (Piedmont, 18,100 samples) and *Herbarium Branchetti* (Emilia-Romagna, 13,000 samples). On the other hand, the herbarium with the lower number of samples resulted the *Herbarium Caetani* with 38 specimens (Table 2). The oldest herbarium is the *Herb. Hölzl Norbert* started in 1960 and preserved in Andriano (Bolzano), followed by the *Erbario Soldano* started in 1973, whereas the most recent are *Erbario Pascale* and *Herbarium Caetani* started in 2014.

Concerning the geographic location of the herbaria (Figure 1), Piedmont and Tuscany resulted the two Italian regions with the highest number (8 herbaria each), whereas we obtained no answer from nine regions (Veneto, Friuli Venezia Giulia, Marche, Umbria, Lazio, Abruzzo, Molise, Puglia, and Basilicata). Concerning the regions where most of the specimens were collected, these correspond to the regions where each herbarium is preserved.

Table I. Questions proposed in the online survey.

Question	Type of answer
Name of the herbarium	free
Owner and email	free
City where the herbarium is preserved	free
Number of preserved samples (add * for estimated number)	free
Are the specimens organized in a database? In case of positive answer, what kind of database?	free
Are you willing to share your data in a free and online database?	yes/no
Are photo/scan of the specimens available?	yes/no
Starting date of the collection	free
Number of taxa (add * for estimated number)	free
Most represented family (and %)	free
Most represented genus (and %)	free
Main country where the samples were collected (and %)	free
Main Italian region where the samples were collected (and%)	free
Type of organization of the collection (e.g. alphabetical order of genera)	free
Has the herbarium some pests? How it is serious?	free
Are the samples subjected to a pest control? What kind?	free
Was the herbarium mentioned in some scientific publications? Could you cite some?	free


**Figure 1.** Distribution of the 34 surveyed Italian private herbaria.

Table 2. Names and number of specimens preserved in the 34 Italian private herbaria resulted by present study.

Herbarium name as provided by owners	N° of samples
<i>Herb. Antonietti</i>	20,000
<i>Erbario Soldano</i>	18,100
<i>Erbario Branchetti</i>	13,000
<i>Erbario Enzo Bona</i>	13,000
<i>Herb. Hölzl Norbert</i>	12,300
<i>Herb. Ardingo e Franco Picco</i>	8,000
<i>Herbarium A. Ruggero</i>	5,000
<i>Herb. Domina</i>	5,000
<i>Herb. Dellavedova Roberto</i>	5,000
<i>Herbarium Meridianum-Collezione Maiorca-Caprio</i>	4,740
<i>Erbario Merli</i>	4,000
<i>Herb. Calvia</i>	4,000
<i>Herb. Nicola Ardenghi</i>	4,000
<i>Herb. Selvi</i>	3,600
<i>Herb. Croce</i>	3,500
<i>Erbario di Stefano Atzori</i>	3,500
<i>Herb. Cecchi</i>	3,273
<i>Herb. M. Bovio</i>	3,116
<i>Herb. Franco Giordana</i>	3,027
<i>Herbarium Varalda</i>	3,000
<i>Erbario Morelli</i>	2,500
<i>Herbarium Marco La Rosa</i>	2,085
<i>Erbario Gonnelli</i>	2,000
<i>Herbarium Braydense</i>	1,947
<i>HB. G. Pellegrino</i>	1,397
<i>HbGanz</i>	1,300
<i>Herb. Gianguzzi-Palermo</i>	1,200
<i>Herb. Giuseppe Cataldi</i>	1,123
<i>Herb. Tognon</i>	1,000
<i>Herb. Orsenigo</i>	1,000
<i>Erbario Trapanese</i>	800
<i>Herb. Roma-Marzio</i>	600
<i>Erbario Pascale</i>	315
<i>Herbarium Caetani</i>	38

The most represented families resulted Asteraceae (in 24 herbaria) and Poaceae (8 herbaria), whereas the most represented genera resulted *Carex* and *Trifolium* (in 4 herbaria respectively), followed by *Hieracium* (in 3 herbaria).

Regarding the availability to share their own data in a free and online database, 24 collectors (71%) have agreed to this proposal whereas, concerning citations of herbaria in a scientific paper, 29 herbaria (85%) were cited at least once.

Almost all herbaria (93%) have an associated database (41% in Excel, 37% in MS-Access, 7% in FileMaker and 7% in MS Word, 4% in SQL-compliant DBMS and 4% in other systems); for 13 herbaria (38%) digital photos or scans of the specimens are available. As regards the arrangement of the collection, 16 (47%) collectors are adopting an alphabetical order, 10 (29%) follow Pignatti (1982), 3 other floras (1 Baroni 1907, 1 Fiori 1923–1929, and 1 Tutin et al. 1964–1980), 2 collectors (6%) adopted a geographic criterion, and 3 collectors (9%) did not report any criteria of ordination.

Pest problems afflicted 22 herbaria (65%) of which 13 (59%) show negligible, 9 (41%) intermediate, and none serious problems. About the solution adopted by collectors to prevent/control pests, 50% use freezing methods, 29% use chemical products (mainly para-dichlorobenzene and camphor), 6% a combination of freezing and chemical products and 5% do not adopt any pest control.

A cross-examination of these two questions (entity of the pest and adopted solution) revealed that private herbarium without treatment are always hit by medium (60%) to low (40%) pest problems. On the other hand, chemical treatment resulted slightly more efficient than freezing, whereas herbaria treated both with freezing and chemical treatment have no (65%) or only low (< 35%) pest problems (Figure 2).




Figure 2. Stacked bar chart showing the relationship between the degree of pest problem and the treatments to prevent it in the Italian private herbaria.

Conclusions

Based on our investigation, about 156,000 specimens are currently preserved in Italian personal herbaria, making this “virtual” herbarium, taken all together, rank 16th among the 68 institutional public herbaria. A very preliminary bibliographic survey revealed that some private herbaria were recently cited in scientific papers (e.g. Anzalone et al. 2010, Iamonico 2015), although the owners did not answer to our survey. For this reason we infer that the real number of specimens preserved in the private collections in

Italy could reach 200,000 or more. For the same reason, the putative absence of private herbaria in central Italy resulting from our investigation could be an underestimation.

We hope that our survey can represent the starting point to re-evaluate the role and the importance of private plant collections, where a remarkable amount of valuable floristic data is currently preserved.

Acknowledgements

All the following owners of private Italian herbaria who answered to the on-line survey are gratefully acknowledged: Michele Aleo, Aldo Antonietti, Nicola M.G. Ardenghi, Stefano Atzori, Enzo Bona, Maurizio Bovio, Romans Caetani, Giacomo Calvia, Giuseppe Cataldi, Lorenzo Cecchi, Antonio Croce, Roberto Della Vedova, Gianni-antonio Domina, Franco Eugenio Rota, Claudia Ganz, Lorenzo Gianguzzi, Franco Giordana, Vincenzo Gonnelli, Norbert Hölzl, Marco La Rosa, Giovanni Maiorca, Marco Merli, Viliam Morelli, Simone Orsenigo, Marziano Pascale, Giorgio Pellegrino, Franco Picco, Alessandro Ruggero, Federico Selvi, Adriano Soldano, Gianni Tognon, Piero Giuseppe Varalda.

References

- Amadei L, Bedini G, Garbari F, Pistolesi G (2007) Erbari. Conservare piante attraverso i secoli. Grafiche Cappelli, Sesto Fiorentino, Firenze, 80 pp.
- Amadei L, Bedini G, Cocchi L, Maccioni S, Peruzzi L, Vangelisti R (2013) Herbarium Horti Botanici Pisani: its present role in scientific research at regional, national and international level. *Atti della Società Toscana di Scienze Naturali, Memorie, serie B* 119(2012): 119–122.
- Anzalone B, Iberite M, Lattanzi E (2010) La flora vascolare del Lazio. *Informatore Botanico Italiano* 42(1): 187–317.
- Ardenghi NMG, Polani F (2016) La Flora della provincia di Pavia (Lombardia, Italia settentrionale). 1. L'Oltrepò Pavese. *Natural History Sciences* 3(2): 51–79. <https://doi.org/10.4081/nhs.2016.269>
- Baroni E (1907) *Guida Botanica*. Rocca San Casciano, 574 pp.
- Bartolucci F, Peruzzi L (2007) Distribuzione del genere *Gagea* Salisb. (Liliaceae) nell'Appennino centro-settentrionale. *Biogeographia, nuova serie* 28: 205–238. <https://doi.org/10.21426/b6110012>
- Beer de GH (1953) Sir Hans Sloane. *The British Museum Quarterly* 18(1): 2–4. <https://doi.org/10.2307/4422405>
- Brilli-Cattarini AJB, Gubellini L (1986) Una nuova specie di *Cardamine* (Cruciferae) della Penisola Italiana e Sicilia. *Webbia* 39(2): 397–407. <https://doi.org/10.1080/00837792.1986.10670379>
- Caruel T (1860) *Prodromo della flora Toscana*. Firenze, 767 pp.

- Clokier HN (1964) An account of the herbaria of the Department of Botany in the University of Oxford. London, Oxford University Press, 280 pp.
- Cuccuini P (2009) L'Erbario Centrale Italiano (E.C.I.) o Herbarium Centrale Italicum (H.C.I.). In: Raffaelli M (Ed.) Il Museo di Storia Naturale dell'Università di Firenze-Le collezioni botaniche. Firenze University Press, 165–198.
- Fiori A (1923–1929) Nuova flora analitica d'Italia, 2 Voll., Tipografia M. Ricci, Firenze.
- Garbari F, Del Carratore F (1993) The flora of the Apuan Alps and the Herbarium of Pietro Pellegrini at Pisa. *Webbia* 48: 421–429.
- Iamónico D (2015) Taxonomic revision of the genus *Amaranthus* (Amaranthaceae) in Italy. *Phytotaxa* 199(1): 1–84. <https://doi.org/10.11646/phytotaxa.199.1.1>
- Lasen C, Busnardo G (1993) L'Erbario Montini (Bassano del Grappa) ed il suo contributo alla Flora Triveneta. *Webbia*, 48: 387–396.
- Moggi G (1993) Botanical collections in Florence from their origin to the present day. *Webbia* 48: 35–60.
- Maccioni S, Baldini R, Amadei L, Bedini G (2009) Erminio Ferrarini (1919–2002) e i suoi erbari di Carrara, Aulla e villa di Comano. *Atti della Società Toscana di Scienze Naturali, Memorie, serie B* 115 (2008): 69–76.
- Nepi C (2009) L'Erbario Webb. In: Raffaelli M (Ed.) Il Museo di Storia Naturale dell'Università di Firenze-Le collezioni botaniche. Firenze University Press, 103–116.
- Nepi C, Cuccuini P (1993) Collectors and collections in the «Herbarium Centrale Italicum» (Phanerogamic section). *Webbia* 48: 79–84.
- Peccenini S, Polatschek A (2016) The genus *Erysimum* (Brassicaceae) in Italy, part III: Key to the species and description of the new species *E. apenninum*, *E. etruscum*, *E. pignattii*. *Annalen Naturhistorischen Museums Wien*, B 118: 147–166.
- Pignatti S (1982) *Flora d'Italia*, 3 voll. Edagricole, Bologna.
- Peruzzi L, Barbo M, Bartolucci F, Bovio M, Carta A, Ciccarelli D, Conti F, Costalonga S, Di Pietro R, Galasso G, Gestri G, Lattanzi E, Lavezzo P, Marsili S, Peccenini S, Pierini B, Tardella FM, Terzo V, Turrise RE, Bedini G (2011) Contributo alla conoscenza floristica delle Colline Pisane: resoconto dell'escursione del Gruppo di Floristica (S.B.I.) nel 2009. *Informatore Botanico Italiano* 43: 3–27.
- Peruzzi L, Viciani D, Angiolini C, Astuti G, Avanzi A, Baldanzi C, Benesperi R, Bonari G, Bonini I, D'Antraccoli M, Castellani B, Dell'Olmo L, Di Nuzzo L, Domina G, Ercolini P, Ferretti G, Fontana D, Gestri G, Gottschlich G, Grazzini A, Lastrucci L, Lazzaro L, Malfanti F, Marsiaj G, Piazzini S, Pierini B, Roma-Marzio F, Sani A, Selvi F, Vicenti C, Bedini G (2016) Contributi per una flora vascolare di Toscana. VII (357–439). *Atti della Società Toscana di Scienze Naturali, Memorie, Serie B* 122 (2015): 67–78.
- Roma-Marzio F, Bernardo L, Liguori P, Peruzzi L (2016) Vascular flora of Monte Sparviere (Southern Italy, Pollino Massif). *Atti della Società Toscana di Scienze Naturali, Memorie, serie B* 122 (2015): 73–88.
- Selvi F (2010) A critical checklist of the vascular flora of Tuscan Maremma (Grosseto province). *Flora Mediterranea* 20: 47–139.
- Stearn WT (1971) Sources of information about botanic gardens and herbaria. *Biological Journal of Linnean Society* 3: 225–233. <https://doi.org/10.1111/j.1095-8312.1971.tb00184.x>

- Taffetani F (Ed.) (2012) *Herbaria. Il grande libro degli erbari italiani. Per la ricerca tassonomica, la conoscenza ambientale e la conservazione del patrimonio.* Nardini, Firenze, 814 pp.
- Thiers B (2016) *Index Herbariorum: A global directory of public herbaria and associated staff.* New York Botanical Garden's Virtual Herbarium. Available from: <http://sweetgum.nybg.org/ih/> [accessed 10 September 2016]
- Tutin TG, Burges NA, Chater AO, Edmondson JR, Heywood VH, Moore DM, Valentine DH, Walters SM, Webb DA (Eds) (1964–1980) *Flora Europaea*, Vol. 1–5. Cambridge University Press, Cambridge.
- Viciani D (2011) Notulae sulla flora del Parco Nazionale delle Foreste Casentinesi, Monte Falterona e Campigna (Appennino Tosco-Romagnolo): approfondimenti su alcuni campioni critici dell'Erbario Zangheri. *Quaderno di Studi e Notizie di Storia Naturale della Romagna* 34: 1–5.
- Visconti A (1993) Il contributo di Philip Barker Webb alla fondazione e all'incremento dell'Herbarium Centrale Italicum. *Webbia* 48: 71–78.